

# CEE 260/MIE 273: Probability and Statistics in Civil Engineering

## Lecture M3c: Lognormal and Exponential Distributions

**Jimi Oke**

UMassAmherst  

---

College of Engineering

September 26, 2025

# Outline

- ① Introduction
- ② The lognormal distribution
- ③ Exponential distribution
- ④ Outlook

# Recap of normal distribution

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$



# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol  $\Phi$  (“phi”) is used to represent the CDF of the **standard normal distribution**, whose values can be looked up in a table.

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters  $\mu$  and  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- The parameters of a normal distribution  $N(\mu, \sigma^2)$  correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol  $\Phi$  (“phi”) is used to represent the CDF of the **standard normal distribution**, whose values can be looked up in a table.
- In MATLAB, the **normcdf(x, mu, sigma)** and **norminv(p, mu, sigma)** can be used to compute probabilities and inverse CDFs of the normal distribution, respectively.

# Using the standard normal CDF probability table

# Using the standard normal CDF probability table


- First convert the random variable to its  $Z$ -score

# Using the standard normal CDF probability table

- First convert the random variable to its  $Z$ -score
- Find the corresponding value in the table

# Using the standard normal CDF probability table

- First convert the random variable to its Z-score
- Find the corresponding value in the table



# Objectives of today's lecture



# Objectives of today's lecture

- Lognormal distribution

# Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions

# Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions
- Exponential distribution

# Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions
- Exponential distribution
- Memoryless property of exponential distribution

# Lognormal distribution

PDF

# Lognormal distribution

## PDF

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma^2$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma^2)$ ) has the PDF:

# Lognormal distribution

## PDF

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma^2$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma^2)$ ) has the PDF:


$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{\ln(x)-\mu}{\sigma}\right)^2} \quad x \geq 0 \quad (3)$$

# Lognormal distribution

## PDF

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma^2$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma^2)$ ) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{\ln(x)-\mu}{\sigma}\right)^2} \quad x \geq 0 \quad (3)$$






# Lognormal distribution

## PDF

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma^2$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma^2)$ ) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{\ln(x)-\mu}{\sigma}\right)^2} \quad x \geq 0 \quad (3)$$




# Lognormal distribution

## PDF

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma^2$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma^2)$ ) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{\ln(x)-\mu}{\sigma}\right)^2} \quad x \geq 0 \quad (3)$$



# Mean, median and variance of a lognormal distribution

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

Mean

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{\left(\mu + \frac{1}{2}\sigma^2\right)} \quad (4)$$

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (4)$$

## Median



# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (4)$$

## Median

The median of  $X$  is:

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (4)$$

## Median

The median of  $X$  is:

$$\text{Median}(X) = e^{\mu} \quad (5)$$

## Variance

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (4)$$

## Median

The median of  $X$  is:

$$\text{Median}(X) = e^{\mu} \quad (5)$$

## Variance

The variance of  $X$  is given by:

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (4)$$

## Median

The median of  $X$  is:

$$\text{Median}(X) = e^{\mu} \quad (5)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X) =$$

# Mean, median and variance of a lognormal distribution

Let  $X \sim \mathcal{LN}(\mu, \sigma^2)$

## Mean

The mean of  $X$  is given by

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (4)$$

## Median

The median of  $X$  is:

$$\text{Median}(X) = e^{\mu} \quad (5)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (6)$$

# Example 1: Mean and variance of lognormal distribution (1)

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution



# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) =$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$\therefore$  The mean is given by

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21}$$



# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\begin{aligned} \therefore \text{The mean is given by } \mathbb{E}(X) &= e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} \\ &= 5(e^{0.21}) \end{aligned}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21}$$

$$= 5(e^{0.21}) = \boxed{6.17 \text{ days}}$$

# Example 1: Mean and variance of lognormal distribution (2)

# Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

$$\mathbb{V}(X) = (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2])$$

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

$$\begin{aligned}\mathbb{V}(X) &= (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2]) \\ &= (\exp(0.42) - 1)(\exp(2 \ln(5) + 0.42))\end{aligned}$$

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

$$\begin{aligned}\mathbb{V}(X) &= (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2]) \\ &= (\exp(0.42) - 1)(\exp(2 \ln(5) + 0.42)) \\ &= \boxed{19.86 \text{ days}^2}\end{aligned}$$



# Relationship between normal and lognormal distributions

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

$$X \sim \mathcal{N}(\mu, \sigma^2)$$

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

$$X \sim \mathcal{N}(\mu, \sigma^2) \implies$$

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

$$X \sim \mathcal{N}(\mu, \sigma^2) \implies e^X \sim \mathcal{LN}(\mu, \sigma^2) \quad (8)$$



# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

$$X \sim \mathcal{N}(\mu, \sigma^2) \implies e^X \sim \mathcal{LN}(\mu, \sigma^2) \quad (8)$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$

- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

$$X \sim \mathcal{N}(\mu, \sigma^2) \implies e^X \sim \mathcal{LN}(\mu, \sigma^2) \quad (8)$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
- However,  $X \sim \mathcal{LN}(\mu, \sigma^2)$ , then  $\mu = E(\ln(X))$  and  $\sigma^2 = \mathbb{V}(\ln(X))$

# Relationship between normal and lognormal distributions

- A random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma^2$  if  $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma^2) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma^2) \quad (7)$$


- Conversely, a random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$  then  $e^X$  is **lognormally** distributed with the same parameters.

$$X \sim \mathcal{N}(\mu, \sigma^2) \implies e^X \sim \mathcal{LN}(\mu, \sigma^2) \quad (8)$$

- If  $X \sim \mathcal{N}(\mu, \sigma^2)$ , then  $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
- However,  $X \sim \mathcal{LN}(\mu, \sigma^2)$ , then  $\mu = E(\ln(X))$  and  $\sigma^2 = \mathbb{V}(\ln(X))$

# Relationship between normal and lognormal (cont.)

# Relationship between normal and lognormal (cont.)



# Positive skewness of lognormal distribution

# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed

# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median




# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median

# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median



# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) =$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma_X \sqrt{2\pi}}$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma_X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma}$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:



# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b)$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma}$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} e^{-\frac{1}{2} z^2} dz \quad (10)$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} e^{-\frac{1}{2} z^2} dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} e^{-\frac{1}{2} z^2} dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b)$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} e^{-\frac{1}{2} z^2} dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi \left( \frac{\ln b - \mu}{\sigma} \right)$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} e^{-\frac{1}{2} z^2} dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi \left( \frac{\ln b - \mu}{\sigma} \right) -$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b e^{-\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2} dx \quad (9)$$

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} e^{-\frac{1}{2} z^2} dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi \left( \frac{\ln b - \mu}{\sigma} \right) - \Phi \left( \frac{\ln a - \mu}{\sigma} \right) \quad (11)$$



## Example 2: Equipment breakdown

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

### Solution

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

### Solution

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

### Solution

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) =$$

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

### Solution

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

### Solution

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) =$$

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

### Solution

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$



## Example 2: Equipment breakdown (cont.)

## Example 2: Equipment breakdown (cont.)

### Solution

## Example 2: Equipment breakdown (cont.)

### Solution

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 =$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$= 1.792 - 0.495 =$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\begin{aligned}\frac{\ln x_0 - 1.792}{0.30} &= \Phi^{-1}(0.05) \\ \ln x_0 - 1.792 &= 0.30[-\Phi^{-1}(0.95)] \\ \ln x_0 &= 1.792 + 0.30(-1.65) \\ &= 1.792 - 0.495 = 1.297\end{aligned}$$



## Example 2: Equipment breakdown (cont.)

### Solution

$$\begin{aligned}\frac{\ln x_0 - 1.792}{0.30} &= \Phi^{-1}(0.05) \\ \ln x_0 - 1.792 &= 0.30[-\Phi^{-1}(0.95)] \\ \ln x_0 &= 1.792 + 0.30(-1.65) \\ &= 1.792 - 0.495 = 1.297\end{aligned}$$

Therefore, the required inspection interval is:

$$x_0 =$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\begin{aligned}\frac{\ln x_0 - 1.792}{0.30} &= \Phi^{-1}(0.05) \\ \ln x_0 - 1.792 &= 0.30[-\Phi^{-1}(0.95)] \\ \ln x_0 &= 1.792 + 0.30(-1.65) \\ &= 1.792 - 0.495 = 1.297\end{aligned}$$

Therefore, the required inspection interval is:

$$x_0 = e^{1.297} =$$

## Example 2: Equipment breakdown (cont.)

### Solution

$$\begin{aligned}\frac{\ln x_0 - 1.792}{0.30} &= \Phi^{-1}(0.05) \\ \ln x_0 - 1.792 &= 0.30[-\Phi^{-1}(0.95)] \\ \ln x_0 &= 1.792 + 0.30(-1.65) \\ &= 1.792 - 0.495 = 1.297\end{aligned}$$

Therefore, the required inspection interval is:

$$x_0 = e^{1.297} = 3.66 \text{ months}$$

# Modeling probabilities of elapsed times

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

Now consider the variable  $Y$  representing the **elapsed time** between successive arrivals.



# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

Now consider the variable  $Y$  representing the **elapsed time** between successive arrivals.

- What is the probability the time between the third and fourth arrivals is less than  $y$  minutes, for instance?

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

Now consider the variable  $Y$  representing the **elapsed time** between successive arrivals.

- What is the probability the time between the third and fourth arrivals is less than  $y$  minutes, for instance?
- This is modeled by the **exponential distribution** with parameter  $\lambda$ .

# Exponential distribution

# Exponential distribution

## Definition

# Exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

# Exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) =$$

# Exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) = \lambda e^{-\lambda x}$$

# Exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0 \quad (12)$$




# Exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0 \quad (12)$$



# CDF of the exponential distribution

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) =$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x)$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$


$$F_X(x) = 1 - e^{-\lambda x}$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$






# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$




Note that  $P(X \leq x) = 1 - e^{-\lambda x}$ ,

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$



Note that  $P(X \leq x) = 1 - e^{-\lambda x}$ , while  $P(X > x) = 1 - (1 - e^{-\lambda x}) = e^{-\lambda x}$

# Mean and variance of the exponential distribution

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X)$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

## Variance

The variance of  $X$  is given by:



# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X)$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X) =$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X) = \frac{1}{\lambda^2} \quad (14)$$

## Example 3: Waiting for a flight

## Example 3: Waiting for a flight

The delay time  $T$  of a flight is exponentially distributed with  $\lambda = 2$  (delays per hour).

## Example 3: Waiting for a flight

The delay time  $T$  of a flight is exponentially distributed with  $\lambda = 2$  (delays per hour). Answer the following questions:

- (a) What is the mean delay (waiting) time,  $\mathbb{E}(T)$ ?
- (b) What is the variance of the delay time  $\mathbb{V}(T)$ ?
- (c) Find the probability that a flight will be delayed by no more than 10 minutes.
- (d) Assuming you have been waiting for a flight for an hour, what is the probability that the flight will be delayed for an additional 30 minutes? (i.e. Find  $P(T > 1.5 | T > 1)$ ).

## Example 3: Waiting for a flight (cont.)

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by



## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) =$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda} = \frac{1}{2}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2} = \frac{1}{2^2} =$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$E(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2} = \frac{1}{2^2} = \boxed{0.25\text{hr}^2}$$



## Example 3: Waiting for a flight (cont.)

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - e^{-\lambda \cdot \frac{1}{6}}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - e^{-\lambda \cdot \frac{1}{6}} = 1 - e^{-2(\frac{1}{6})}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$\begin{aligned} P\left(T \leq \frac{1}{6}\right) &= 1 - e^{-\lambda \cdot \frac{1}{6}} = 1 - e^{-2(\frac{1}{6})} \\ &= 1 - e^{-\frac{1}{3}} \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$\begin{aligned}
 P\left(T \leq \frac{1}{6}\right) &= 1 - e^{-\lambda \cdot \frac{1}{6}} = 1 - e^{-2\left(\frac{1}{6}\right)} \\
 &= 1 - e^{-\frac{1}{3}} = \boxed{0.283}
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:



## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$P(T > (0.5 + 1) | T > 1) = P(T > 1.5 | T > 1)$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)}
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule})
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}}
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]}
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\ &= e^{-2(0.5)} \quad (= P(T > 0.5)) \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\
 &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\
 &= e^{-1}
 \end{aligned}$$



## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\ &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\ &= e^{-1} = \boxed{0.37} \end{aligned}$$

# Memorylessness of the exponential distribution

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

## Memoryless property

$$P(T > t + s | T > s) =$$

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

## Memoryless property

$$P(T > t + s | T > s) = P(T > t) \quad (15)$$

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

## Memoryless property

$$P(T > t + s | T > s) = P(T > t) \quad (15)$$

That is, it does not matter from which time the waiting begins (i.e. conditioning); the probability of an elapsed time remains the same.

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$   
CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$   
CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$



# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) =$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

PDF:  $f_X(x) =$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

PDF:  $f_X(x) = \lambda e^{-\lambda x},$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) =$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x},$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$



# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Mean:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (20)$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Mean:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (20)$$

Variance:

$$\mathbb{V}(X) = \frac{1}{\lambda^2} \quad (21)$$

# MATLAB Homework

# MATLAB Homework

- Ensure your submission is strictly a script saved with the `.m` extension

# MATLAB Homework


- Ensure your submission is strictly a script saved with the **.m** extension
- MATLAB can only execute a script if it is in the *current folder*.

# MATLAB Homework

- Ensure your submission is strictly a script saved with the **.m** extension
- MATLAB can only execute a script if it is in the **current folder**. Otherwise you may get a message like the one below:


# MATLAB Homework

- Ensure your submission is strictly a script saved with the **.m** extension
- MATLAB can only execute a script if it is in the **current folder**. Otherwise you may get a message like the one below:



# MATLAB Homework

- Ensure your submission is strictly a script saved with the **.m** extension
- MATLAB can only execute a script if it is in the **current folder**. Otherwise you may get a message like the one below:



If so, simply click on **Change Folder** or move the file to the current folder you are in. Finally, always make sure the path of a file being read by a script is valid from its location, otherwise you will have to deal with “File not found” errors.



# Midterm Exam

# Midterm Exam

- 24-hour open-resource examination

# Midterm Exam

- 24-hour open-resource examination
- Available for download via Canvas on Wednesday, October 16th at **10:00 AM**

# Midterm Exam

- 24-hour open-resource examination
- Available for download via Canvas on Wednesday, October 16th at **10:00 AM**
- Due by **October 21st** at **11:59 PM**

# Midterm Exam

- 24-hour open-resource examination
- Available for download via Canvas on Wednesday, October 16th at **10:00 AM**
- Due by **October 21st** at **11:59 PM**
- Exam length will be similar to previous midterms or the practice exam(s) available on Canvas.

# Midterm Exam

- 24-hour open-resource examination
- Available for download via Canvas on Wednesday, October 16th at **10:00 AM**
- Due by **October 21st** at **11:59 PM**
- Exam length will be similar to previous midterms or the practice exam(s) available on Canvas.
- Exam is designed to be completed in 2-3 hours or less.

# Midterm Exam

- 24-hour open-resource examination
- Available for download via Canvas on Wednesday, October 16th at **10:00 AM**
- Due by **October 21st** at **11:59 PM**
- Exam length will be similar to previous midterms or the practice exam(s) available on Canvas.
- Exam is designed to be completed in 2-3 hours or less. The 24-hr window gives you flexibility and time to plan, organize and check your work before submission.

# Midterm Exam

- 24-hour open-resource examination
- Available for download via Canvas on Wednesday, October 16th at **10:00 AM**
- Due by **October 21st** at **11:59 PM**
- Exam length will be similar to previous midterms or the practice exam(s) available on Canvas.
- Exam is designed to be completed in 2-3 hours or less. The 24-hr window gives you flexibility and time to plan, organize and check your work before submission.
- You can use your calculator/computer (Matlab/Python) to compute probabilities (as long as you indicate how you obtained your answer).