

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

$$\begin{aligned}
 \text{Задача 1} \quad \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

Уастб 1

$$\begin{aligned}
 \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^{\alpha} dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

Уастб 1

$$\begin{aligned}
 \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^{\alpha} dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

Уастб 1

$$\begin{aligned}
 \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^{\alpha} dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

Уастб 1

$$\begin{aligned}
 \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^{\alpha} dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

Уастб 1

$$\begin{aligned}
 \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^{\alpha} dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

Уастб 1

$$\begin{aligned}
 \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^{\alpha} dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

$$\begin{aligned}
 \text{Задача 1} \quad \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

$$\begin{aligned}
 \text{Задача 1} \quad \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

$$\begin{aligned}
 \text{Задача 1} \quad \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} + C = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$

8.1.26

$$\begin{aligned}
 \int \frac{5 + \sin^3 x}{\sin^2 x} dx &= \int \left(\frac{5}{\sin^2 x} + \frac{\sin^3 x}{\sin^2 x} \right) dx = \\
 &= 5 \int \frac{1}{\sin^2 x} dx + \int \sin x dx = \\
 &= -5 \operatorname{ctg} x + (-\cos x) + C = -5 \operatorname{ctg} x - \cos x + C
 \end{aligned}$$

D/3

8.1.29

$$\begin{aligned}
 \text{Задача 1} \quad \int \frac{dx}{x^2 \sqrt{x}} &= \int \frac{dx}{x^2 \cdot x^{\frac{1}{2}}} = \int \frac{dx}{x^{5/2}} = \int x^{-5/2} dx = \\
 &= \left[\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + C \right] = \frac{x^{-5/2+1}}{-5/2+1} = \frac{x^{-3/2}}{-3/2} + C = \\
 &= -\frac{2}{3} x^{-3/2} + C = -\frac{2}{3x\sqrt{x}} + C
 \end{aligned}$$

8.1.30

$$\begin{aligned}
 \int \frac{dx}{x^2 + 3} &= \int \frac{dx}{x^2 + (\sqrt{3})^2} = \left[\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctg} \frac{x}{a} + C \right] = \\
 &= \frac{1}{\sqrt{3}} \operatorname{arctg} \frac{x}{\sqrt{3}} + C = \frac{\sqrt{3}}{3} \operatorname{arctg} \left(\frac{\sqrt{3}x}{3} \right) + C
 \end{aligned}$$

8.1.31

$$\begin{aligned}
 \int \frac{1}{5^x} dx &= \int 5^{-x} dx = \frac{5^{-x}}{\ln 5} + C = \\
 \int \frac{1}{5^x} dx &= \int \left(\frac{1}{5} \right)^x dx = \left[\int a^x dx = \frac{a^x}{\ln a} + C \right] = \\
 &= \frac{\left(\frac{1}{5} \right)^x}{\ln \left(\frac{1}{5} \right)} + C = \frac{1}{\ln \left(\frac{1}{5} \right) 5^x} = \frac{1}{\ln(5^{-1}) 5^x} = -\frac{1}{\ln 5 \cdot 5^x}
 \end{aligned}$$