

Template Screening Report (v5.7.1 - Multi-Source AI Evaluator)

Journal Template: Jurnal Teknologi

Submission No: 26132

Author Name: Nursaadah et al.

Article Title: Development of Transformation Construct and Transgenic Tobacco Plants Carrying EgPAL5 Gene

Source File: DEVELOPMENT OF TRANSFORMATION CONSTRUCT AND TRANSGENIC TOBACCO PLANTS CARRYING EgPAL5 GENE .docx

Screened PDF: DEVELOPMENT OF TRANSFORMATION CONSTRUCT AND TRANSGENIC TOBACCO PLANTS CARRYING EgPAL5 GENE .pdf

Screening Date: 2026-05-21 09:42:14

Detection Source Debug Report

Item	DOC Detection	PDF Detection	AI Detection	Final Source
Article Title	NO	YES	YES	PDF + AI
Authors	NO	YES	YES	PDF + AI
Affiliations	NO	YES	YES	PDF + AI
Corresponding Author	YES	YES	SKIPPED	DOC + PDF
Graphical Abstract	YES	YES	SKIPPED	DOC + PDF
English Abstract	YES	YES	SKIPPED	DOC + PDF
Keywords	YES	YES	SKIPPED	DOC + PDF
Malay Abstract (Abstrak)	YES	YES	SKIPPED	DOC + PDF
Kata Kunci	YES	YES	SKIPPED	DOC + PDF
Introduction	YES	NO	YES	DOC + AI
Methodology	YES	NO	YES	DOC + AI
Results and Discussion	YES	YES	SKIPPED	DOC + PDF
Conclusion	YES	NO	YES	DOC + AI
Acknowledgement	YES	YES	SKIPPED	DOC + PDF
References	YES	YES	SKIPPED	DOC + PDF

Summary: DOC detected 12 items | PDF detected 12 items | AI confirmed 6 items

The system uses DOC + PDF + AI voting logic. AI serves as a third checker when DOC and PDF conflict. AI is skipped when DOC and PDF already agree.

Executive Summary

Overall Decision: **NEEDS MANUAL REVIEW**

Main Reason: The manuscript generally follows the template but contains inconsistencies that require manual review.

Critical Failed Items: None

Items Needing Review:

- Article Title
- Authors
- Affiliations
- Introduction
- Methodology

Suggested Action: Editor should review the highlighted issues before proceeding.

Overall Result

Metric	Score / Value
Final Status	NEEDS MANUAL REVIEW
Final Score	84/100
Structure Score	60%
Layout Score	100%
Formatting Score	100%
Detection Method	DOCX-First + Multi-Source AI (v5.7.1)
Decision	MANUAL REVIEW

How Items Are Evaluated

DOC + PDF + AI SKIPPED	PASS	Both deterministic detectors agree
DOC + PDF + AI CONFIRMED	PASS	All three confirm
DOC + AI	NEEDS REVIEW	DOC+AI confirm, but PDF layout conflicts
PDF + AI	NEEDS REVIEW	PDF+AI confirm, but DOC source extraction fails
DOC only	NEEDS REVIEW	DOC detects, but PDF+AI don't confirm
PDF only	NEEDS REVIEW	PDF detects, but DOC+AI don't confirm
AI only	NEEDS REVIEW	Only AI confirms (deterministic detectors failed)
None	FAILED	All three fail to detect required item

Status Explanation

PASS — Item follows the template structure, position, and formatting.

WARNING/NEEDS REVIEW — Item exists but evidence is inconsistent or layout differs from template.

FAILED — Required item truly missing after DOC/PDF/AI checks.

Item Screening Details

Item	DOC	PDF	AI	Final Source	Status	Reason
Article Title	NO	YES	YES	PDF + AI	WARNING	Not detected by PDF and AI, but DOC source text did not co
Authors	NO	YES	YES	PDF + AI	WARNING	Not detected by PDF and AI, but DOC source text did not
Affiliations	NO	YES	YES	PDF + AI	WARNING	Not detected by PDF and AI, but DOC source text did

Corresponding Author	YES	YES	SKIPPED	DOC + PDF	PASS	Corresponding_Author confirmed by both DOC and PDF detection
Graphical Abstract	YES	YES	SKIPPED	DOC + PDF	PASS	Graphical_Abstract confirmed by both DOC and PDF detection.
English Abstract	YES	YES	SKIPPED	DOC + PDF	PASS	Abstract confirmed by both DOC and PDF detection.
Keywords	YES	YES	SKIPPED	DOC + PDF	PASS	Keywords confirmed by both DOC and PDF detection.
Malay Abstract (Abstrak)	YES	YES	SKIPPED	DOC + PDF	PASS	Abstrak confirmed by both DOC and PDF detection.
Kata Kunci	YES	YES	SKIPPED	DOC + PDF	PASS	Kata_Kunci confirmed by both DOC and PDF detection.
Introduction	YES	NO	YES	DOC + AI	WARNING	Introduction confirmed by DOC and AI, but PDF/layout detection failed.
Methodology	YES	NO	YES	DOC + AI	WARNING	Methodology confirmed by DOC and AI, but PDF/layout detection failed.
Results and Discussion	YES	YES	SKIPPED	DOC + PDF	PASS	Results_Discussion confirmed by both DOC and PDF detection.
Conclusion	YES	NO	YES	DOC + AI	WARNING	Conclusion confirmed by DOC and AI, but PDF/layout detection failed.
Acknowledgement	YES	YES	SKIPPED	DOC + PDF	PASS	Acknowledgement confirmed by both DOC and PDF detection.
References	YES	YES	SKIPPED	DOC + PDF	PASS	References confirmed by both DOC and PDF detection.

Detailed Findings

Article Title [title]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Article Title' section must exist.

Found: Matched heading: "DEVELOPMENT OF TRANSFORMATION CONSTRUCT AND TRANSGENIC TOBACCO PLANTS CARRYING EgPAL5 GENE" | Detection method: score_ranking

Confidence: 1.00

Detection Method: score_ranking

DOC Detection: NO

PDF Detection: YES

AI Detection: YES

AI Confidence: 1.00

AI Evidence: PDF snippet shows a full title section with the title 'DEVELOPMENT OF TRANSFORMATION CONSTRUCT AND TRANSGENIC TOBACCO PLANTS CARRYING EgPAL5 GENE' and

Evidence Sources: PDF + AI

Final Source Used: PDF + AI

Rejected false positives: Jurnal Teknologi, Full Paper, Article history

Status: WARNING

Reason: Component found but evidence is inconsistent.

B. Layout / Coordinate Evaluation

Expected: Page 1, left-center area, top area

Found: Page 1, left column, upper area

Position Check: WARNING

Reason: Position deviates from expected region (tolerance: 0.10).

C. Formatting Evaluation

Expected: Heading should be in uppercase style.

Expected: Heading should use bold font.

Formatting Check: PASS

Reason: Formatting matches template expectations.

Overall Item Status: WARNING

Authors [authors]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Authors' section must exist.

Found: Matched heading: "Mohd Nordin Nursaadaha, Nur Syafiqah Shariful Bahria, Omar Nawawia, Mohamad Shafek Hilmana, " | Detection method: score_rank

Confidence: 0.70

Detection Method: score_ranking

DOC Detection: NO

PDF Detection: YES

AI Detection: YES

AI Confidence: 0.85

AI Evidence: PDF snippet includes full author names and affiliations, confirming presence of authors section. DOC snippet is incomplete and lacks authors section,

Evidence Sources: PDF + AI

Final Source Used: PDF + AI

Rejected false positives: Article history, Received, Accepted

Status: WARNING

Reason: Component found but evidence is inconsistent.

B. Layout / Coordinate Evaluation

Expected: Page 1, left-center area, upper area

Found: Page 1, left column, upper area

Position Check: PASS

Reason: Position matches expected template region.

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: PASS

Reason: Formatting matches template expectations.

Overall Item Status: WARNING

Affiliations [affiliations]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Affiliations' section must exist.

Found: Matched heading: "aLaboratory of Plant Genetics and Cell Biology, Faculty of Plantation and Agrotechnology, Univers" |
Detection method: score_rank

Confidence: 0.80

Detection Method: score_ranking

DOC Detection: NO

PDF Detection: YES

AI Detection: YES

AI Confidence: 0.80

AI Evidence: PDF snippet shows full author list with affiliations and addresses; DOC snippet lacks the Affiliations section despite containing other manuscript par

Evidence Sources: PDF + AI

Final Source Used: PDF + AI

Rejected false positives: Accepted

Status: WARNING

Reason: Component found but evidence is inconsistent.

B. Layout / Coordinate Evaluation

Expected: Page 1, left-center area, upper area

Found: Page 1, left column, upper area

Position Check: PASS

Reason: Position matches expected template region.

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: PASS

Reason: Formatting matches template expectations.

Overall Item Status: WARNING

Corresponding Author [corresponding_author]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Corresponding Author' section must exist.

Found: Matched heading: "Corresponding author" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: PASS

Reason: The required component 'Corresponding Author' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left-center area, upper area

Found: Page 1, right-center area, upper area

Position Check: WARNING

Reason: Position deviates from expected region (tolerance: 0.12).

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: PASS

Reason: Formatting matches template expectations.

Overall Item Status: **PASS**

Graphical Abstract [graphical_abstract]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Graphical Abstract' component must exist.

Found: Matched heading: "Graphical abstract" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: **PASS**

Reason: The required component 'Graphical Abstract' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, upper area

Found: Page 1, left-center area, upper-middle area

Position Check: **WARNING**

Reason: Position deviates from expected region (tolerance: 0.12).

C. Formatting Evaluation

Expected: Image should be present.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **PASS**

English Abstract [abstract]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'English Abstract' section must exist.

Found: Matched heading: "Abstract" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: **PASS**

Reason: The required component 'English Abstract' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left-center area, upper area

Found: Not detected

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **PASS**

Keywords [keywords]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Keywords' section must exist.

Found: Matched heading: "Keywords" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: PASS

Reason: The required component 'Keywords' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left-center area, lower-middle area

Found: Page 1, left-center area, lower-middle area

Position Check: PASS

Reason: Position matches expected template region.

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: PASS

Reason: Formatting matches template expectations.

Overall Item Status: PASS

Malay Abstract (Abstrak) [abstrak]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Malay Abstract (Abstrak)' section must exist.

Found: Matched heading: "Abstrak" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: PASS

Reason: The required component 'Malay Abstract (Abstrak)' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, lower area

Found: Page 1, left-center area, lower area

Position Check: WARNING

Reason: Position deviates from expected region (tolerance: 0.15).

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: PASS

Reason: Formatting matches template expectations.

Overall Item Status: PASS

Kata Kunci [kata_kunci]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Kata Kunci' section must exist.

Found: Matched heading: "Kata kunci" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF
Status: PASS
Reason: The required component 'Kata Kunci' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, lower area
Found: Page 1, left-center area, upper area
Position Check: WARNING
Reason: Position deviates from expected region (tolerance: 0.15).

C. Formatting Evaluation

Expected: Standard paragraph formatting.
Formatting Check: PASS
Reason: Formatting matches template expectations.
Overall Item Status: PASS

Introduction [introduction]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Introduction' section must exist.
Found: Matched heading: "1.0 INTRODUCTION" | Detection method: source_docx_text
Confidence: 0.85
Detection Method: source_docx_text
DOC Detection: YES
PDF Detection: NO
AI Detection: YES
AI Confidence: 0.85
AI Evidence: DOC snippet provides a clear and relevant Introduction section with research context and background. PDF snippet lacks any Introduction content, suggest
Evidence Sources: DOC + AI
Final Source Used: DOC + AI
Rejected false positives: 1.0 INTRODUCTION
Status: WARNING
Reason: Component found but evidence is inconsistent.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, top area
Found: Not detected

C. Formatting Evaluation

Expected: Standard paragraph formatting.
Formatting Check: PASS
Reason: Formatting matches template expectations.
Overall Item Status: WARNING

Methodology [methodology]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Methodology' section must exist.
Found: Matched heading: "2.0 METHODOLOGY" | Detection method: source_docx_text
Confidence: 0.85
Detection Method: source_docx_text
DOC Detection: YES
PDF Detection: NO
AI Detection: YES
AI Confidence: 0.85
AI Evidence: DOC snippet provides a comprehensive Methodology excerpt describing experimental design and procedures. PDF snippet contains only header and author in
Evidence Sources: DOC + AI
Final Source Used: DOC + AI
Rejected false positives: 2.0 METHODOLOGY

Status: **WARNING**

Reason: Component found but evidence is inconsistent.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, top area

Found: Not detected

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **WARNING**

Results and Discussion [results_discussion]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Results and Discussion' section must exist.

Found: Matched heading: "3.0 RESULTS AND DISCUSSION" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: **PASS**

Reason: The required component 'Results and Discussion' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, top area

Found: Page 1, center area, lower area

Position Check: **WARNING**

Reason: Position deviates from expected region (tolerance: 0.20).

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **PASS**

Conclusion [conclusion]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Conclusion' section must exist.

Found: Matched heading: "4.0 CONCLUSION" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: NO

AI Detection: YES

AI Confidence: 0.85

AI Evidence: DOC snippet provides a clear Conclusion section discussing study outcomes, gene transformation, and future applications. PDF snippet contains only hea

Evidence Sources: DOC + AI

Final Source Used: DOC + AI

Rejected false positives: 4.0 CONCLUSION

Status: **WARNING**

Reason: Component found but evidence is inconsistent.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, top area

Found: Not detected

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **WARNING**

Acknowledgement [acknowledgement]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'Acknowledgement' component is recommended.

Found: Matched heading: "Acknowledgement" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: **PASS**

Reason: The required component 'Acknowledgement' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, top area

Found: Page 1, center area, upper area

Position Check: **WARNING**

Reason: Position deviates from expected region (tolerance: 0.20).

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **PASS**

References [references]

A. Structure Evaluation (DOC + PDF + AI)

Expected: The 'References' section must exist.

Found: Matched heading: "References" | Detection method: source_docx_text

Confidence: 0.85

Detection Method: source_docx_text

DOC Detection: YES

PDF Detection: YES

AI Detection: SKIPPED

Evidence Sources: DOC + PDF

Final Source Used: DOC + PDF

Status: **PASS**

Reason: The required component 'References' exists.

B. Layout / Coordinate Evaluation

Expected: Page 1, left column, top area

Found: Page 1, left column, upper-middle area

Position Check: **WARNING**

Reason: Position deviates from expected region (tolerance: 0.20).

C. Formatting Evaluation

Expected: Standard paragraph formatting.

Formatting Check: **PASS**

Reason: Formatting matches template expectations.

Overall Item Status: **PASS**

Editor Conclusion

The manuscript generally follows the template but has items that require manual review. These items had inconsistent evidence between DOC detection, PDF layout analysis, or AI verification. Please review the highlighted items before making a final decision.

Structure Score: 60% (Weight: 40%)

Layout Score: 100% (Weight: 30%)

Formatting Score: 100% (Weight: 30%)

Final Score: 84/100

Items Needing Review:

- Article Title — **WARNING**
- Authors — **WARNING**
- Affiliations — **WARNING**
- Introduction — **WARNING**
- Methodology — **WARNING**

This report was automatically generated by Turnitin Assistant v5.7.1 Template Screening Engine (DOCX-First + Multi-Source AI Evaluator).

DEVELOPMENT OF TRANSFORMATION CONSTRUCT AND TRANSGENIC TOBACCO PLANTS CARRYING *EgPAL5* GENE

Mohd Nordin Nursaadah^a, Nur Syafiqah Shariful Bahri^a, Omar Nawawi^a, Mohamad Shafek Hilman^a, Muhammad Ikhwan Mughni^a, Shamsiah Abdullah^a, Tianqi Bai^b, Cuixian Zhang^b, Mat Yunus Abdul Masani^c, Chai-Ling Ho^d, Chong Yu Lok Yusuf^{a*}

Article history
Received
X Month 202Y
Received in revised form
XA Month 202Y
Accepted
XB Month 202Y

*Corresponding author
yusufchong@uitm.edu.my


^aLaboratory of Plant Genetics and Cell Biology, Faculty of Plantation and Agrotechnology, Universiti Teknologi MARA Jasin Campus, 77300 Merlimau, Melaka, Malaysia

^bInstitute of Tropical and Subtropical Cash Crops, Yunnan Academy of Agricultural Sciences, Baoshan 675800, China

^cMalaysian Palm Oil Board (MPOB) No. 6, Persiaran Institusi, Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia

^dDepartment of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

Graphical abstract



Abstract

Tobacco (*Nicotiana tabacum*) is a globally significant industrial crop, yet its yield and productivity are frequently compromised by a diverse range of biotic and abiotic stresses. Developing crops with enhanced genetic potential is a crucial strategy to ensure survival and yield stability. This study reports the development of a novel genetic resource through the integration of the *EgPAL5* gene from oil palm (*Elaeis guineensis*) into the tobacco genome. The 2,115-bp *EgPAL5* gene was amplified, cloned into the pMDC32 binary vector, and introduced into tobacco plants via *Agrobacterium*-mediated transformation. Regenerated shoots were cultured on selective media containing hygromycin and timentin, maintained until rooting, and subsequently acclimatized to soil. Among 42 putative transformants obtained, 10 vigorous lines were selected for PCR verification, all exhibiting the expected amplicon size confirming stable integration of *EgPAL5* transformation. This work successfully demonstrated the feasibility of incorporating the *EgPAL5* gene into tobacco genome for crop improvement purposes, establishing a validated transgenic resource for future biotechnological applications.

Keywords: *Agrobacterium*-mediated transformation, molecular biology, *Nicotiana tabacum*, PAL, tissue culture

Abstrak

Tembakau (*Nicotiana tabacum*) merupakan tanaman industri yang signifikan di peringkat global, namun hasil dan produktiviti tanaman ini sering terjejas oleh pelbagai tekanan biotik dan abiotik. Pembangunan tanaman dengan potensi genetik yang dipertingkatkan adalah strategi kritikal bagi menjamin kelangsungan hidup dan kestabilan hasil. Kajian ini melaporkan pembangunan sumber genetik baharu melalui integrasi gen *EgPAL5* daripada kelapa sawit (*Elaeis guineensis*) ke dalam genom tembakau. Gen *EgPAL5* sepanjang 2,115-bp telah diamplifikasikan, diklon ke dalam vektor binari pMDC32, dan dimasukkan ke dalam pokok tembakau melalui transformasi berperantara *Agrobacterium*. Pucuk teregenerasi dikultur di atas media pemilihan yang mengandungi higromisin dan timentin, dikekalkan sehingga pembentukan akar, dan seterusnya diaklimatisasi ke tanah. D daripada 42 transforman putatif yang diperolehi, 10

titisan vigor telah dipilih untuk verifikasi PCR, di mana kesemuanya memaparkan saiz ampikon yang dijangkakan, sekali gus mengesahkan integrasi stabil transformasi *EgPAL5*. Kajian ini berjaya membuktikan kebolehlaksanaan pengintegrasian gen *EgPAL5* ke dalam genom tembakau untuk tujuan penambahbaikan tanaman, seterusnya menghasilkan sumber transgenik yang sah bagi aplikasi bioteknologi pada masa hadapan.

Kata kunci: Transformasi berperantara *Agrobacterium*, biologi molekular, *Nicotiana tabacum*, PAL, kultur tisu

© 2020@ Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

Tobacco is a perennial herbaceous plant native to South America. It belongs to the Solanaceae family, which encompasses brinjal, chilli, petunia, potato, and tomato. This crop has traditionally been associated with global cigarette production. Therefore, it is now cultivated worldwide because of its economic importance. According to Zhu *et al.* [1], China is the largest producer of tobacco, followed by India and Brazil. Tobacco can grow up to two meters tall and produces large leaves that accumulate high levels of nicotine [2]. In commercial varieties, particularly *Nicotiana tabacum*, nicotine levels can be increased to approximately 6.47% of dry weight through a genetic manipulation approach [2]. As a hybrid between *N. sylvestris* and *N. tomentosiformis*, *N. tabacum* is an allotetraploid plant that contains 48 chromosomes [3]. It can be easily transformed genetically, which makes it a useful system for plant biotechnology studies. Apart from that, tobacco is also used in environmental applications, including phytoremediation [4]. The plant can absorb high levels of heavy metals, including cadmium, from contaminated soils [5]. Such capability enables remediation of polluted sites while producing biomass for non-food applications. Furthermore, tobacco has been considered as a potential renewable resource for biofuel production owing to its high biomass production [6].

More interestingly, biotechnology advances have expanded the use of tobacco in molecular farming for the production of recombinant proteins, vaccines, and industrial enzymes [7]. Unfortunately, tobacco production is frequently affected by environmental stresses such as extreme temperature [8], drought [9], salinity [10], and flooding [11], which reduce yield and crop quality as a consequence. Apart from abiotic stresses, tobacco is also susceptible to infections caused by fungi, viruses, and bacteria. Several examples of the major diseases threatening the plant include Tobacco Black Shank (TBS) caused by the soil-borne oomycete *Phytophthora nicotianae* [12], Fusarium wilt (FW) caused by *Fusarium oxysporum* [13], Granville wilt (GW) caused by *Ralstonia solanacearum* [14], and Tobacco Mosaic (TM) caused by *Tobamovirus tabaci* or commonly known as Tobacco Mosaic Virus (TMV) [15, 16]. TBS is typically characterised by sudden wilting followed by stem blackening. FW and GW often begin with leaf yellowing and root decay. In GW, hollow stems and lesions may be observed, whereas FW often appears as unilateral chlorosis on one side of the plant. Lastly, TM typically induces chlorotic patches appearing as blotches, streaks, or spots, followed by vein discoloration, leaf curling, and morphological distortion [15, 17]. The above-mentioned abiotic and biotic stresses can significantly affect the productivity and economic value of

tobacco crops. Under stress conditions, plants activate complex signalling pathways that redirect metabolic resources toward survival rather than growth [18]. At the structural level, plant resistance to abiotic and biotic stresses is closely associated with lignification.

Lignin forms a strong physical barrier that restricts pathogen entry and strengthens vascular tissues. In plants, lignin biosynthesis is predominantly regulated by phenylalanine ammonia-lyase (PAL), which catalyses the first step of the phenylpropanoid pathway and directs metabolic flux toward the production of various secondary metabolites. PAL is encoded by a multigene family [19], and its members are often expressed in specific tissues and upregulated under stress conditions [20]. Previous studies have shown that PAL contributes to plant growth and defence responses. For example, overexpression of *GmPAL2.1* in soybean (*Glycine max*) enhances resistance against *Phytophthora sojae* by elevating PAL enzyme activity, increasing glyceollin accumulation, and activating salicylic acid (SA)-mediated signalling pathways that induce the expression of SA-responsive defence genes [21]. In wild legume (*Lotus japonicus*), *LjPAL1* regulates lignin remodelling during *Mesorhizobium loti* symbiosis, and gene knockdown leads to excessive infection thread formation and nodule overproliferation [22]. Furthermore, in grass (*Brachypodium distachyon*), *BdPAL1* knockdown reduced virus-induced SA and lignin accumulation [23]. Previously, Yusuf *et al.* [24] suggested that the *EgPAL5* promoter played a major role in regulating the biosynthesis of stress-related secondary metabolites in oil palm. Collectively, these findings implied that heterologous expression of the *EgPAL5* gene could improve stress tolerance in tobacco plants. This prompted us to introduce the *EgPAL5* gene from oil palm into tobacco plants in the present study, producing stable transgenic lines for further studies.

2.0 METHODOLOGY

2.1 Vector Construction

The 2115-bp full-length coding sequence (CDS) of *EgPAL5* was amplified from the pJET1.2-*EgPAL5* plasmid obtained from Universiti Putra Malaysia. The reaction mixture contained 1× Q5 High-Fidelity Buffer, 0.2 mM dNTPs, 0.5 µM each of the forward (5'-TAA GGT ACC ATG GAG AAC GGC GGT GCT-3') and reverse (5'-GCA CTA GTC TAA CAA ATA GGA AGG GG-3') primers, 15 ng of template DNA, 1.2 U of Q5 DNA Polymerase (NEB, USA), and sterile distilled water to a final volume of 60 µL. The PCR was performed using the following thermal cycling profile: 98 °C for 30 s; 32 cycles of 98 °C for 10 s, 67-69 °C for 30 s, 72

°C for 75 s; and 72 °C for 5 min. The PCR product was purified from agarose gel using the PrimeWay Gel Extraction PCR Purification kit (1st BASE, Singapore). Next, the purified PCR product and pMDC32 plasmid were double digested with *KpnI* and *SpeI* restriction enzymes overnight at 37 °C. The insert DNA was ligated with the linearised pMDC32 vector using T4 DNA Ligase (Thermo Scientific, USA), generating the pMDC32-EgPAL5 construct. The ligation product was transformed into *Escherichia coli*, strain DH5 α competent cells, and positive clones were confirmed by colony PCR, followed by Sanger sequencing. The verified transformation construct was subsequently transformed into *Agrobacterium tumefaciens* strain LBA4404 by electroporation.

2.2 Plant and *Agrobacterium* Preparation

Tobacco (*N. tabacum*) seeds were surface-sterilised using 0.5% (v/v) sodium hypochlorite for 10 min, followed by three rinses with sterile distilled water. Approximately 20–25 seeds were sown per plate on Murashige and Skoog (MS) basal medium solidified with 2.5 g/L Gelrite and germinated under a 16/8-h light/dark photoperiod at 25 °C for four weeks to obtain aseptic donor plants. Single colonies of *A. tumefaciens* carrying the T-DNA were selected from LB agar plates supplemented with 25 mg/L rifampicin, 50 mg/L kanamycin, and 100 mg/L streptomycin. The selected colony was inoculated into 25 mL LB broth, containing the same concentration of antibiotics. The culture was incubated overnight at 28 °C with 250 rpm agitation. Prior to transformation, the bacterial culture was pelleted by centrifugation at 4,000 $\times$ g for 10 min at 4 °C. The supernatant was discarded. The pellet was then resuspended in liquid MS basal medium to an OD₆₀₀ of 1.0 as inoculation medium.

2.3 *Agrobacterium*-mediated transformation

Plant transformation was performed according to Hilman *et al.* [25]. Leaf discs ($\approx$ 0.5–1.0 cm) were excised from a young leaf of four-week-old tobacco. Approximately 30–45 discs were immersed in the *Agrobacterium* suspension and gently agitated for 30 min. Suspended leaf discs were blotted dry on sterile filter paper, and placed abaxial side up on co-cultivation medium (MS salts, 2.0 mg/L 6-Benzylaminopurine (BAP), 0.1 mg/L 1-Naphthaleneacetic acid (NAA). After 48 h of dark incubation at 28 °C, explants were transferred to shooting medium I (SM1: MS salts, 2.0 mg/L BAP, 0.1 mg/L NAA, 50 mg/L hygromycin, 250 mg/L timentin) and maintained under 16/8-h light/dark photoperiod at 25 °C. Emerging shoots were excised after two weeks and transferred to shoot elongation medium II (SM2: MS salts, 0.5 mg/L BAP, antibiotics as in SM1). After an additional two weeks, elongated shoots were subcultured onto rooting medium (RM: MS salts, 0.1 mg/L NAA, antibiotics as in shooting media).

2.4 Acclimatisation and Seed Production

Well-rooted plantlets (1–2 cm root length) were transferred to a soil mixture in pots and acclimatised under transparent plastic covers in a growth room with 16/8-h light/dark photoperiod at 24 °C. Plastic covers were progressively removed over two weeks. Plants were maintained under regular irrigation and supplied with nutrient solution three times weekly. Inflorescences were enclosed in pollination

bags prior to flower opening to prevent cross-pollination. Mature seeds were collected from fully brown desiccated capsules, air-dried for approximately 72 h, and stored at 4 °C.

2.5 Molecular Verification

Genomic DNA (gDNA) was extracted from young leaf tissue of putative transgenic plants using a modified CTAB-based protocol adapted from Edward *et al.* [26]. Briefly, approximately 50 mg of tissue was homogenised in a 1.5-mL microcentrifuge tube with 400 μ L of CTAB buffer using micro pestles. The lysate was centrifuged, and then the supernatant was mixed with an equal volume of isopropanol to precipitate DNA. The pellet was washed twice with 70% (v/v) ethanol, air-dried, and resuspended in 50 μ L of nuclease-free water. The isolated gDNA served as the template for PCR amplification. Tables 1 and 2 present the reaction components and thermal cycling profile. A 2115-bp fragment specific to the *EgPAL5* transgene was amplified using primers listed in Section 2.1. The PCR products were electrophoresed on a 1.2% (w/v) agarose gel stained with FluoroSafe (1stBASE, Singapore). Subsequently, the PCR product was recovered from the agarose gel using a commercial kit as described by Shukor *et al.* [27] and cloned into our in-house developed pJET1.2/blunt vector [28, 29]. The resulting recombinant plasmids were isolated from bacterial clones and subjected to DNA sequencing for verification.

Table 1 PCR setup for *EgPAL5* gene amplification

Component	Final Concentration	Volume per Reaction (μ L)
2X PCR Master Mix	1X	10
10 μ M Forward Primer	0.5 μ M	1
10 μ M Reverse Primer	0.5 μ M	1
gDNA		1
dH ₂ O		7
TOTAL		20

Table 2 Thermal cycling profile for the amplification of the *EgPAL5* gene

Step	Temperature (°C)	Time	No. of Cycles
Initial denaturation	95	2 min	1
Denaturation	95	20 s	38
Annealing	60	20 s	
Extension	72	65 s	
Final extension	72	5 min	1

3.0 RESULTS AND DISCUSSION

3.1 Vector Construction

An approximately 2.1-kb PCR product corresponding to the *EgPAL5* gene was successfully amplified and cloned into the pMDC32 vector, generating the pMDC32-EgPAL5 transformation construct with a total size of 12,111 bp. Sanger sequencing confirmed that the amplicon sequence was identical to the *EgPAL5* gene sequence. The transformation construct was designed so that the expression of *EgPAL5* gene is regulated by a constitutive

2× CaMV 35S promoter with a NOS terminator at the 3'-end of the transgene. It contains a hygromycin resistance gene for the selection of transgenic lines. The plasmid map is shown in Figure 1. The pMDC32 vector has previously been utilised in our laboratory for the overexpression of other oil palm phenylpropanoid genes, including *EgCAD2* [25] and *Eg4CL1* [30]. This vector has proven effective for the heterologous expression of target genes in higher plants.




Figure 1 Plasmid map of the pMDC32-EgPAL5 construct developed for delivering the *EgPAL5* gene into tobacco. RB, right border; 2x CaMV 35S promoter; *EgPAL5*, *Elaeis guineensis* phenylalanine ammonia-lyase 5 gene; NOS, nopaline synthase terminator; CaMV 35S, CaMV 35S promoter; Hyg^R, hygromycin B resistance gene; CaMV poly(A) signal; LB, left border

3.2 Plant and *Agrobacterium* Preparation

In this study, surface-sterilisation of tobacco seeds with 0.5% sodium hypochlorite for 10 min effectively eliminated microbial contaminants while preserving viability. The wild-type tobacco seeds germinated within four days on MS medium (Figure 2a), consistent with reports by Hilman *et al.* [25]. As shown in Figure 2b, the seedlings developed uniformly after being sown on the medium. To avoid overcrowding, the tobacco seedlings were cultured at a density of 20–25 seedlings per plate over four weeks to yield robust plant materials for genetic transformation. Meanwhile, *A. tumefaciens* carrying the pMDC32-EgPAL5 construct was streaked onto selective LB agar medium (Figure 2c) and incubated at 28 °C three days before genetic transformation was performed. A single colony was selected to inoculate liquid LB medium supplemented with appropriate antibiotics. The culture was grown overnight with agitation at approximately 250 rpm to maintain adequate aeration and nutrient distribution, ensuring a sufficient yield of actively growing bacterial cells for plant transformation. The culture medium changed from clear to turbid appearance during growth and multiplication of *A. tumefaciens* (Figure 2d). This indicates a high bacterial density and a homogenous population, appropriate for the subsequent infection process.




Figure 2 Preparation of plant material and bacterial culture for genetic transformation (a) surface-sterilised tobacco seeds cultured on MS medium, (b) four-week-old WT tobacco, (c) *A. tumefaciens* LBA4404 harbouring the pMDC32-EgPAL5 construct cultured on LB agar, (d) *A. tumefaciens* liquid culture used for plant transformation

3.3 Transformation and Co-cultivation

The bacterial cells were collected by centrifugation of bacterial culture at 4 °C and 4,000 × g to ensure optimal cell recovery while minimising stress to preserve *Agrobacterium* viability prior to infection (Figure 3a). To enhance bacterial infection occurrence, minor wounds were purposely created during leaf disc excision. As reported by [31], wounding of explants promotes *Agrobacterium* infection and improves transformation efficiency. This is attributed to the release of plant-derived wound signals, which activate bacterial virulence (*vir*) gene expression, thereby facilitating bacterial entry and subsequent T-DNA transfer into the host genome [31]. Leaf discs were immersed in the infection medium for 30 min with gentle agitation to promote uniform bacterial attachment and T-DNA transfer initiation. Post-infection, discs were blotted dry to remove excess bacteria. This is a critical step to prevent *Agrobacterium* overgrowth that could compromise explant viability (Figure 3b). Explants were spaced evenly across the medium, to further suppress bacterial overgrowth (Figure 3c). The co-cultivation period was strictly maintained to avoid *Agrobacterium* overgrowth.




Figure 3 Genetic transformation of WT tobacco (a) bacterial cell pellet after centrifugation, (b) dry-blotting of infected leaf discs

using sterile forceps, (c) explants uniformly distributed on co-cultivation medium

3.4 Regeneration of Putative Transgenics

When cultured on shooting medium I (SM1), the leaf discs exhibited distinct curling initially, and callus formation was observed at the wounded edges (Figure 4a) within 10–14 days, subsequently developing multiple shoot primordia (Figure 4b). Individual shoots were excised and subcultured onto shooting medium II (SM2) to promote nodal development and elongation (Figure 4c). Since the SM2 contained a lower concentration of BAP, the shoot primordia developed into healthy shoots. After two weeks on SM2, healthy individual shoots were transferred to a rooting medium (RM), where robust root systems developed. During the shoot regeneration and rooting steps, subculturing was performed every two weeks. Regular subculturing prevented nutrient depletion and maintained actively growing cultures throughout the regeneration period. A total of 45 plantlets were regenerated with root lengths of 1.5–2.0 cm (Figure 4d), all of which were transferred to soil for acclimatisation (Figure 4e). Forty-two putative transgenic plants were successfully established under controlled conditions, with minimal transplant shock observed as plastic covers were gradually opened during the acclimatisation period. During the vegetative growth stage, all putative transformants exhibited normal morphology, showing healthy leaf expansion, sturdy stems, and consistent structural vigour, indicating stable regeneration under the optimised protocol.




Figure 4 Regeneration of putative transgenic tobacco plants (a) Leaf explants (0.5–1.0 cm²) placed on SM1, (b) shoot organogenesis with multiple shoot buds developing from induced callus, (c) elongated individual shoots isolated into culture vessel, (d) rooted plantlet ready for acclimatisation, (e) acclimatised regenerant in soil substrate (scale bar = 1 cm)

3.5 Molecular Verification of Transgenics

Due to labour and time constraints, only 10 vigorous plants were selected and subjected to molecular verification. To verify the presence of the *EgPAL5* gene in tobacco genome, gDNA samples were first extracted from the selected plants (Figure 5a). Agarose gel electrophoresis analysis of the gDNA samples revealed distinct, high-molecular-weight bands, confirming that the gDNA was intact and suitable for subsequent PCR analysis. As

anticipated, PCR analysis yielded a distinct amplicon of the expected size (about 2.1-kb) from all samples (Figure 5b). The result demonstrated both the specificity of the primers used and the integration of the *EgPAL5* gene in the tobacco genome. To unequivocally confirm the identity and fidelity of the integrated sequence, the PCR product from a representative transgenic line was cloned and subjected to Sanger sequencing. The sequencing result confirmed that the amplified PCR product corresponding to the coding sequence of *EgPAL5*. The chromatogram (Figure 5c) shows that the obtained sequence data was high quality and interpretable.




Figure 5 Molecular verification of *EgPAL5* transgene integration in transgenic tobacco plants (a) Genomic DNA samples isolated from putative transgenic lines. Lanes L1–L10: Lines 1–10, (b) PCR amplification of the *EgPAL5* transgene in transgenic tobacco lines; Lane M: 1 kb DNA ladder (1st BASE, Singapore); Lanes 1–10: transgenic lines L1–L10; Lane 11: positive control (pMDC32-*EgPAL5* plasmid was used as the PCR template), (c) Sanger sequencing chromatogram of the PCR product cloned in pJET1.2 vector. The arrow indicates the start codon (ATG) of the *EgPAL5* gene

Taken together, the results confirmed that transgenic tobacco plants carrying *EgPAL5* gene from oil palm have been successfully produced in this study. The present work focused on the generation and molecular verification of transgenic tobacco harbouring the *EgPAL5* gene, rather than the functional characterisation of the gene. Therefore, no in-depth molecular and biochemical characterisations such as gene expression analysis and enzymatic assays, was performed on the transgenic plants generated in this study. Future studies should prioritise the selection of independent transgenic lines exhibiting elevated PAL activity to ensure the attainment of the desired outcomes. In addition, field evaluations of the stress tolerance of the transgenic tobacco should be performed to verify whether the generated transgenic plants indeed enhanced resilience under stress conditions. Apart from being an elite variety for tobacco production, the transgenic plants produced in this study can also serve as valuable material for elucidating the functional role of the *EgPAL5* gene in future studies. The development of transgenic tobacco varieties with enhanced stress

tolerance may facilitate the translation from laboratory findings into practical field applications.

4.0 CONCLUSION

The present study successfully generated and verified transgenic tobacco lines carrying the oil palm *EgPAL5* gene, demonstrating the feasibility of introducing this gene into tobacco through the *Agrobacterium*-mediated transformation method. As PAL plays a central role in phenylpropanoid biosynthesis, which contributes to lignin formation and plant defence responses, the transgenic lines generated were anticipated to exhibit enhanced resilience under unfavourable conditions. They also serve as useful resources for future functional and metabolic engineering studies. Ultimately, the development of transgenic tobacco followed by the characterisation of *EgPAL5* has the potential to bridge laboratory findings and practical field applications. It will definitely be beneficial for both basic understanding and translational outcomes in plant stress biology.

References

- [1] Zhu, R., He, S., Ling, H., Liang, Y., Wei, B., Yuan, X., Cheng, W., Peng, B., Xiao, J., Wei, J., He, Y., Xiao, H., Wang, Z. 2024, Optimizing tobacco quality and yield through the scientific application of organic-inorganic fertilizer in China: a meta-analysis. *Frontiers in Plant Science*. Volume 15 - 2024. DOI : <https://doi.org/10.3389/fpls.2024.1500544>
- [2] Datiles, M. J., Acevedo-Rodríguez, P. 2022, *Nicotiana tabacum* (tobacco). CABI Compendium. DOI : <https://doi.org/10.1079/cabicompendium.36326>
- [3] Bai, Y., Pattanaik, S., Patra, B., Werkman, J. R., Xie, C. H., Yuan, L. 2011, Flavonoid-related basic helix-loop-helix regulators, NtAn1a and NtAn1b, of tobacco have originated from two ancestors and are functionally active. *Planta*. 234(2): p. 363-75. DOI : <https://doi.org/10.1007/s00425-011-1407-y>
- [4] Selwal, N., Tabassum, Z., Rahayu, F., Yulia, N. D., Sugiono, S., Endarto, O., Rijaya, P. D., Djajadi, D., Khamidah, A., Wani, A. K. 2023, Therapeutic potential and phytoremediation capabilities of the tobacco plant: Advancements through genetic engineering and cultivation techniques. *Biocatalysis and Agricultural Biotechnology*. 52: p. 102845. DOI : <https://doi.org/10.1016/j.bcab.2023.102845>
- [5] Murtaza, G., Chen, Y., Qian, F., Zheng, G., Usman, M., Azam, M., Deng, G., Ahmed, Z., Chen, S., Iqbal, J., Elshikh, M. S., Rizwana, H., Ahmad, S., Iqbal, R., Gurbanova, L., Lackner, M. 2025, Fullerene nanoparticles enhance potential of phytoremediation and safe cultivation of tobacco (*Nicotiana tabacum*) in cadmium contaminated soil: Tobacco proved as fullerene-philic industrial crop. *Industrial Crops and Products*. 233: p. 121428. DOI : <https://doi.org/10.1016/j.indcrop.2025.121428>
- [6] Wu, S., Gao, C., Pan, H., Wei, K., Li, D., Cai, K., Zhang, H. 2021, Advancements in tobacco (*Nicotiana tabacum* L.) seed oils for biodiesel production. *Front Chem*. 9: p. 834936. DOI : <https://doi.org/10.3389/fchem.2021.834936>
- [7] Naeem, M., Han, R., Ahmad, N., Zhao, W., Zhao, L. 2023, Tobacco as green bioreactor for therapeutic protein production: latest breakthroughs and optimization strategies. *Plant Growth Regulation*. 103(2): p. 227-41. DOI : <https://doi.org/10.1007/s10725-023-01106-w>
- [8] Xu, J., Chen, Z., Wang, F., Jia, W., Xu, Z. 2020, Combined transcriptomic and metabolomic analyses uncover rearranged gene expression and metabolite metabolism in tobacco during cold acclimation. *Scientific Reports*. 10(1): p. 5242. DOI : <https://doi.org/10.1038/s41598-020-62111-x>
- [9] Wei, K., Liu, G., Wei, B., Zhang, Q., Wu, S., Li, Z. 2025, Effects of drought stress on the physiological characteristics of flue-cured tobacco during the vigorous growing period. *Phyton-International Journal of Experimental Botany*. 94(4): p. 1287-98. DOI : <https://doi.org/10.32604/phyton.2025.062385>
- [10] Sun, H., Sun, X., Wang, H., Ma, X. 2020, Advances in salt tolerance molecular mechanism in tobacco plants. *Hereditas*. 157(1): p. 5. DOI : <https://doi.org/10.1186/s41065-020-00118-0>
- [11] Cacal, D., Ricafrente, D., Cacayurin, C., Soriano, R. 2024, Effects of tobacco (*Nicotiana tabacum*) applied with gibberellic acid (GA3) in response to different durations of drought and flooding. *Thai Journal of Agricultural Science*. 57(2): p. 109– 19.
- [12] Zhang, M., Ren, Q., Wu, X., Liu, M., Zhang, L., Yuan, Q., Lu, N., Cao, Y. 2025, Biocontrol potential of four *Pseudomonas* strains against tobacco black shank disease. *Antonie Van Leeuwenhoek*. 118(6): p. 83. DOI : <https://doi.org/10.1007/s10482-025-02092-x>
- [13] Yang, L., Yang, W., Zhang, H., Li, L., Liu, J., Li, M., Xu, R., Xiang, H., Li, Y., Yang, Y. 2026, Fusarium wilt disease induced changes in the composition and function of the rhizosphere metabolome and microbiome in tobacco plants. *Scientific Reports*. DOI : <https://doi.org/10.1038/s41598-026-40653-w>

Ethics Statement

This study was conducted in accordance with institutional biosafety guidelines under the oversight of the Universiti Teknologi MARA Institutional Biosafety Committee. The project was reviewed and classified as an exempted activity under the applicable biosafety regulations. All transgenic plant materials were handled within appropriate biosafety containment facilities. No human participants or animal subjects were involved in this study.

Acknowledgement

The pJET1.2-EgPAL5 plasmid was kindly provided by Assoc. Prof. Dr. Mohd Puad Abdullah of Universiti Putra Malaysia (UPM) Serdang, Malaysia. The authors thank the laboratory staff for facility support. This work was funded by the Ministry of Higher Education (MOHE), Malaysia, under the Fundamental Research Grant Scheme (FRGS/1/2022/STG01/UITM/02/13).

- [14] Katawczik, M., Mila, A. L. 2012, Plant age and strain *Ralstonia solanacearum* affect the expression of resistance of tobacco cultivars to Granville wilt. Tobacco Science. 49(49): p. 8-13.
DOI : <https://doi.org/10.3381/11-013r.1>
- [15] Li, C., Zhang, Q., Wang, X., Liu, Z., Zhang, X., Jiang, C., Cheng, L., Yang, A., Li, Y., Ji, Y., Liu, D. 2025, Transcriptomic and metabolomic analyses reveal molecular mechanisms of tobacco mosaic virus (TMV) resistance in *Nicotiana tabacum* L. BMC Plant Biol. 25(1): p. 1029.
DOI : <https://doi.org/10.1186/s12870-025-07053-0>
- [16] Ahmed, H., Al-Kuwaiti, N. 2025, Detection and biocontrol of *Tobamovirus tabaci* infecting tomato in Iraq. Jurnal Hama dan Penyakit Tumbuhan Tropika. 25: p. 158-68.
DOI : <https://doi.org/10.23960/jhptt.125158-168>
- [17] Rifkind, D., Freeman, G. L. 11 - Tobacco mosaic virus. The Nobel Prize Winning Discoveries in Infectious Diseases. London: Academic Press; 2005. p. 81-4.
DOI : <https://doi.org/10.1016/B978-012369353-2/50018-7>
- [18] Sarala, K., Rao, K. P., Nanda, C., Baghyalakshmi, K., Darvishzadeh, R., Gangadhara, K., Rajappa, J. J. Abiotic stress resistance in tobacco: Advances and strategies. In: Kole C, editor. Genomic Designing for Abiotic Stress Resistant Technical Crops. Cham: Springer International Publishing; 2022. p. 329-427.
DOI : https://doi.org/10.1007/978-3-031-05706-9_10
- [19] Vogt, T. 2010, Phenylpropanoid biosynthesis. Mol Plant. 3(1): p. 2-20.
DOI : <https://doi.org/10.1093/mp/ssp106>
- [20] Yadav, V., Wang, Z., Wei, C., Amo, A., Ahmed, B., Yang, X., Zhang, X. 2020, Phenylpropanoid pathway engineering: An emerging approach towards plant defense. Pathogens. 9(4).
DOI : <https://doi.org/10.3390/pathogens9040312>
- [21] Zhang, C., Wang, X., Zhang, F., Dong, L., Wu, J., Cheng, Q., Qi, D., Yan, X., Jiang, L., Fan, S., Li, N., Li, D., Xu, P., Zhang, S. 2017, Phenylalanine ammonia-lyase2.1 contributes to the soybean response towards *Phytophthora sojae* infection. Sci Rep. 7(1): p. 7242.
DOI : <https://doi.org/10.1038/s41598-017-07832-2>
- [22] Chen, Y., Li, F., Tian, L., Huang, M., Deng, R., Li, X., Chen, W., Wu, P., Li, M., Jiang, H., Wu, G. 2017, The phenylalanine ammonia lyase gene *LjPAL1* is involved in plant defense responses to pathogens and plays diverse roles in *Lotus japonicus*-Rhizobium symbioses. Mol Plant Microbe Interact. 30(9): p. 739-53.
DOI : <https://doi.org/10.1094/mpmi-04-17-0080-r>
- [23] Pant, S. R., Irigoyen, S., Liu, J., Bedre, R., Christensen, S. A., Schmelz, E. A., Sedbrook, J. C., Scholthof, K. B. G., Mandadi, K. K. 2021, *Brachypodium* phenylalanine ammonia lyase (PAL) promotes antiviral defenses against *Panicum mosaic virus* and its satellites. mBio. 12(1).
DOI : <https://doi.org/10.1128/mBio.03518-20>
- [24] Yusuf, C. Y. L., Nawawi, O., Azhari, M. F., Hilman, M. S., Bai, T., Zhang, C., Yi, H., Ni, Z., Xie, D., Abdullah, M. O., Abdul Masani, M. Y., Shaharuddin, N. A., Abdullah, M. P., Ong Abdullah, J. 2026, Functional characterisation of the *EgPAL5* promoter from oil palm (*Elaeis guineensis* Jacq.) reveals its potential roles in plant growth and adaptation to drought. The Journal of Horticultural Science and Biotechnology. 101(1): p. 58-75.
DOI : <https://doi.org/10.1080/14620316.2025.2519330>
- [25] Hilman, M. S., Omar, N., Azhari, M. F., Bai, T., Zhang, C., Abdullah, M. P., Masani, M. Y. A., Yusuf, C. Y. L. 2025, A comprehensive method to generating and identifying transgenic tobacco lines with a single transgene integration locus for functional analysis. Pertanika Journal of Tropical Agricultural Science. 48(1).
DOI : <https://doi.org/10.47836/pjtas.48.1.03>
- [26] Edward, K., Johnstone, C., Thompson, C. 1991, A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. Nucleic Acids Research. 19(6): p. 1349.
- [27] Shukor, N. A. A., Yusuf, C. Y. L., Kumar, S. M., Abiri, R., Abdullah, M. P. 2023, Molecular dissection and an *in-silico* approach of a novel *Gibberellin 20-oxidase* gene of *Hibiscus cannabinus* L. Pakistan Journal of Botany. 55(2): p. 489-500.
DOI : [http://dx.doi.org/10.30848/PJB2023-2\(11\)](http://dx.doi.org/10.30848/PJB2023-2(11))
- [28] Nawawi, O., Abdullah, M. P., Yusuf, C. Y. L. 2022, A strategy for in-house production of a positive selection cloning vector from the commercial pJET1.2/blunt cloning vector at minimal cost. 3 Biotech. 12(9): p. 216.
DOI : <https://doi.org/10.1007/s13205-022-03289-x>
- [29] Nawawi, O., Abdullah, M. P., Yusuf, C. Y. L. 2023, A streamlined strategy for self-production of a commercial positive selection vector, the pJET1.2/blunt cloning vector, using common laboratory *E. coli* strains. 3 Biotech. 13(7): p. 224.
DOI : <https://doi.org/10.1007/s13205-023-03647-3>
- [30] Nawawi, O., Hilman, M. S., Xie, D., Zhang, C., Yi, H., Shaharuddin, N. A., Ghazali, M. N., Masani, M. Y. A., Yusuf, C. Y. L. 2026, Production of transgenic *Arabidopsis* carrying the oil palm *Eg4CL1* gene. Jurnal Teknologi. In Press.
- [31] Yadav, S. K., Katikala, S., Yellisetty, V., Kannepalle, A., Narayana, J. L., Maddi, V., Mandapaka, M., Shanker, A. K., Bandi, V., Bharadwaja, K. P. 2012, Optimization of *Agrobacterium* mediated genetic transformation of cotyledonary node explants of *Vigna radiata*. SpringerPlus. 1(1): p. 59.
DOI : <https://doi.org/10.1186/2193-1801-1-59>