



REGIONAL CONTRIBUTIONS OF OCEAN IRON FERTILIZATION TO ATMOSPHERIC CO_2 CHANGES DURING THE LAST GLACIAL TERMINATION

NATALIA OPAZO, FABRICE LAMBERT
PONTIFICAL CATHOLIC UNIVERSITY OF CHILE



INTRODUCTION

Oceans are the largest source of carbon storage. Their internal processes directly affect the concentrations of atmospheric pCO_2 . In the last 2 million years the climate of the Earth has alternated between glacial and interglacial periods, with low and high atmospheric concentrations of CO_2 . Iron fertilization of the oceans is thought to have contributed a pproximately 20 p.p.m.v. to the 80-100 p.p.m.v. Holocene-LGM difference in atmospheric CO_2 concentrations




Figure 1: A schematic of the ocean's “biological pump”, the sequestration of carbon and alkalinity from the surface ocean (light blue) into the ocean interior (dark blue), and their effects on the ocean's carbon chemistry. Source [1].

In this study, we explore the effect of iron fertilization through mineral dust flux to the ocean surface for Holocene and LGM climatic conditions, but also for idealized intermediate dust values. In this way, we can calculate the theoretical relationship between dust fluxes and atmospheric pCO_2 through the modulation of the biological pump in the ocean (**Figure 1**).

RESULTS




Figure 8: Range percentage of pCO_2 reduction of models: Lambert, Albani, Takemura, MIROC-ESM, MRI-CGCM3.

MATERIALS AND METHODS

We use the cGENIE carbon cycle-focused Earth System Model of Intermediate Complexity to simulate the effect of various dust fields on atmospheric CO_2 through the termination. For five different published Holocene and LGM dust flux estimates [2, 3, 4, 5, 6].




Figure 2: Grid with the specification of oceanic regions run with cGENIE

We produced 8 linearly-spaced intermediate dust flux values at each grid point, thus resulting in 10 incremental global dust flux fields from low Holocene to high LGM values. We simulate the response of atmospheric CO_2 through the termination for global dust fluxes, but also isolate specific High-Nutrient Low Chlorophyll (HNLC) regions of the world's ocean (Figure 2) to quantify their individual contribution to the total CO_2 changes.

PCO_2 REDUCTION OBTAINED BY cGENIE SIMULATION FOR DUST FIELDS VARYING FROM HOLOCENE TO LGM




Figure 3: Global fluxes.




Figure 4: North Pacific regional zone.




Figure 5: Central Pacific regional zone.




Figure 6: South Pacific regional zone.




Figure 7: South Atlantic regional zone.

CONCLUSION

There are only small changes in relative CO_2 capture simulated by cGENIE due to different reconstruction or simulations of dust fluxes, both globally and regionally (Fig. 8 and 9a). Note that this small error is largely due to our removal of those model results that did not change southern hemisphere dust source emissions between Holocene and LGM.

The high-latitude oceans contribute most to CO_2 -drawdown during the LGM (Fig. 9b). This was expected since they are the main HNLC regions of the world's ocean and therefore very sensitive to the contribution of aeolian iron flux that influence on ocean biogeochemistry.

CONTACT INFORMATION

Email neopazo@uc.cl

Email2 lambert@uc.cl

REFERENCES

- [1] Mathis P Hain, DM Sigal, and Gerald H Haug. 8.18—the biological pump in the past. *Reference Module in Earth Systems and Environmental Sciences, Treatise on Geochemistry (Second Edition), The Oceans and Marine Geochemistry*, 8:485–517, 2014.
- [2] Fabrice Lambert, Alessandro Tagliabue, Gary Shaffer, Frank Lamy, Gisela Winckler, Laura Farias, Laura Gallardo, De Pol-Holz, et al. Dust fluxes and iron fertilization in holocene and last glacial maximum climates. *Geophysical Research Letters*, 42(14):6014–6023, 2015.
- [3] S Albani, NM Mahowald, AT Perry, RA Scanza, CS Zender, NG Heavens, V Maggi, JF Kok, and BL Otto-Bliesner. Improved dust representation in the community atmosphere model. *Journal of Advances in Modeling Earth Systems*, 6(3):541–570, 2014.
- [4] T Takemura, M Egashira, K Matsuzawa, H Ichijo, R O'ishi, and Abe-Ouchi. A simulation of the global distribution and radiative forcing of soil dust aerosols at the last glacial maximum. *Atmospheric Chemistry & Physics*, 2009.
- [5] S Watanabe, T Hajima, K Sudo, T Nagashima, T Takemura, H Okajima, T Nozawa, H Kawase, M Abe, T Yokohata, et al. Miroc-esm: model description and basic results of cmip5-20c3m experiments. *Geoscientific Model Development Discussions*, 4(2):1063–1128, 2011.
- [6] Seiji Yukimoto, Yukimasa Adachi, Masahiro Hosaka, Tomonori Sakami, Hiro-masa Yoshimura, Mikitoshi Hirabara, Taichu Y Tanaka, Eiki Shindo, Hiroyuki Tsujino, Makoto Deushi, et al. A new global climate model of the meteorological research institute: Mri-cgcm3—model description and basic performance—. *Journal of the Meteorological Society of Japan. Ser. II*, 90:23–64, 2012.