
Fluid monopoly trajectories with forgetting


$p_0 = 0.1$, $p_1 = 0.35$, $p_2 = 0.8$, $c_1 = 0.05$, $c_2 = 0.65$, $R = 1$, $\gamma = 0.98$; mature boundary and numerical HJB policy.



At attractive thresholds, a holding mixture keeps x constant. Low demand can make movement very slow; arrows show its direction.

Fluid monopoly trajectories with forgetting


$p_0 = 0.3$, $p_1 = 0.35$, $p_2 = 0.8$, $c_1 = 0.05$, $c_2 = 0.65$, $R = 1$, $\gamma = 0.98$; mature boundary and numerical HJB policy.



At attractive thresholds, a holding mixture keeps x constant. Low demand can make movement very slow; arrows show its direction.

Fluid monopoly trajectories with forgetting


$p_0 = 0.5$, $p_1 = 0.35$, $p_2 = 0.8$, $c_1 = 0.05$, $c_2 = 0.65$, $R = 1$, $\gamma = 0.98$; mature boundary and numerical HJB policy.



At attractive thresholds, a holding mixture keeps x constant. Low demand can make movement very slow; arrows show its direction.

Fluid monopoly trajectories with forgetting


$p_0 = 0.7$, $p_1 = 0.35$, $p_2 = 0.8$, $c_1 = 0.05$, $c_2 = 0.65$, $R = 1$, $\gamma = 0.98$; mature boundary and numerical HJB policy.



At attractive thresholds, a holding mixture keeps x constant. Low demand can make movement very slow; arrows show its direction.

Fluid monopoly trajectories with forgetting


$p_0 = 0.8$, $p_1 = 0.35$, $p_2 = 0.8$, $c_1 = 0.05$, $c_2 = 0.65$, $R = 1$, $\gamma = 0.98$; mature boundary and numerical HJB policy.



At attractive thresholds, a holding mixture keeps x constant. Low demand can make movement very slow; arrows show its direction.

Fluid monopoly trajectories with forgetting

$p_0 = 0.9$, $p_1 = 0.35$, $p_2 = 0.8$, $c_1 = 0.05$, $c_2 = 0.65$, $R = 1$, $\gamma = 0.98$; mature boundary and numerical HJB policy.



At attractive thresholds, a holding mixture keeps x constant. Low demand can make movement very slow; arrows show its direction.