

# Homotopy-coherent interchange and equivariant little disk operads

Natalie Stewart

*Department of Mathematics  
Harvard University*

INI Equivariant Homotopy Theory in Context  
*Queen's University Belfast, April 9 2025*



# A question you might happen upon

Interchange and  
little V-disks

Natalie Stewart

In trace methods for Real  $K$ -theory,  $\mathrm{THH}$  has a Real analogue:<sup>1</sup>

$$\begin{array}{ccc} \mathrm{Alg}_{\mathbb{E}_1}(\mathrm{Sp}) & & \mathrm{Alg}_{\mathbb{E}_\sigma}(\mathrm{Sp}_{C_2}) \\ \mathrm{THH} \downarrow & \xrightarrow{\text{“Reality”}} & \downarrow \mathrm{THR} \\ \mathrm{Sp} & & \mathrm{Sp}_{C_2} \end{array}$$

In this,  $\sigma$  is the sign representation and  $\mathbb{E}_\sigma$ -algebras are a genuine-equivariant version of rings with anti-involution.

**Question (c.f. Dotto-Moi-Patchkoria-Reeh<sup>2</sup> '17)**

What algebraic structure does  $\mathrm{THR}$  of highly structured  $C_2$ -ring spectra have?

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>1</sup> Emanuele Dotto. *Stable real K-theory and real topological Hochschild homology*. Thesis (Ph.D.)—University of Copenhagen. 2012. arXiv: 1212.4310 [math.AT].

<sup>2</sup> Emanuele Dotto, Kristian Moi, Irakli Patchkoria, and Sune Precht Reeh. “Real topological Hochschild homology”. In: *J. Eur. Math. Soc. (JEMS)* 23.1 (2021), pp. 63–152.

# How to construct structure on THH

Interchange and  
little  $V$ -disks

Natalie Stewart

## Observation

THH can be given a symmetric monoidal structure, so we may lift

$$\begin{array}{ccccc} \mathrm{Alg}_{\mathcal{O} \otimes \mathbb{E}_1}(\mathrm{Sp}) & \simeq & \mathrm{Alg}_{\mathcal{O}} \mathrm{Alg}_{\mathbb{E}_1}^{\otimes}(\mathrm{Sp}) & \dashrightarrow & \mathrm{Alg}_{\mathcal{O}}(\mathrm{Sp}) \\ & & \downarrow U & & \downarrow U \\ & & \mathrm{Alg}_{\mathbb{E}_1}(\mathrm{Sp}) & \xrightarrow{\mathrm{THH}} & \mathrm{Sp} \end{array}$$

## Theorem (Dunn<sup>3</sup> '88, Lurie<sup>4</sup> '09)

$\mathbb{E}_n \simeq \mathbb{E}_{n-1} \otimes \mathbb{E}_1$ ; hence THH takes  $\mathbb{E}_n$ -rings to  $\mathbb{E}_{n-1}$ -rings.

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>3</sup>Gerald Dunn. "Tensor product of operads and iterated loop spaces". In: *J. Pure Appl. Algebra* 50.3 (1988), pp. 237–258.

<sup>4</sup>Jacob Lurie. *Derived Algebraic Geometry VI: Ek Algebras*. 2009. arXiv: 0911.0018 [math. AT].

# How to construct structure on THR

Interchange and  
little V-disks

Natalie Stewart

## Observation (S.<sup>5</sup>)

THR can be given a  $C_2$ -symmetric monoidal structure, so we may lift

$$\begin{array}{ccccc} \mathrm{Alg}_{\mathcal{O} \otimes \mathbb{E}_\sigma}(\mathrm{Sp}_{C_2}) & \simeq & \mathrm{Alg}_{\mathcal{O}} \mathrm{Alg}_{\mathbb{E}_\sigma}^\otimes(\mathrm{Sp}_{C_2}) & \dashrightarrow & \mathrm{Alg}_{\mathcal{O}}(\mathrm{Sp}_{C_2}) \\ & \searrow & & & \downarrow U \\ & & \mathrm{Alg}_{\mathbb{E}_\sigma}(\mathrm{Sp}_{C_2}) & \xrightarrow{\mathrm{THR}} & \mathrm{Sp}_{C_2} \end{array}$$

## Conjecture

In the 1-categorical world, there is an equivalence  $\mathbb{E}_V \otimes \mathbb{E}_W \simeq \mathbb{E}_{V \oplus W}$ .

Cold open

Historical context

$C$ - $\infty$ -categorical  
operads

$C$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>5</sup>Natalie Stewart. *On tensor products of equivariant commutative operads (draft)*. 2025.

# Statement of the additivity theorem

Interchange and  
little V-disks

Natalie Stewart

For this talk, all terms are defined  $\infty$ -categorically.

## Theorem (S.<sup>6</sup>)

Given  $V, W$  orthogonal  $G$ -representations, we have

$$\mathbb{E}_V \otimes \mathbb{E}_W \simeq \mathbb{E}_{V \oplus W};$$

hence there is an equivalence of  $\infty$ -categories

$$\mathrm{Alg}_{\mathbb{E}_V} \mathrm{Alg}_{\mathbb{E}_W}^{\otimes} (\mathrm{Sp}_G) \simeq \mathrm{Alg}_{\mathbb{E}_{V \oplus W}} (\mathrm{Sp}_G).$$

## Corollary

THR of  $\mathbb{E}_{V \oplus \sigma}$ -rings has a natural  $\mathbb{E}_V$ -ring structure, and no more.

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>6</sup>Natalie Stewart. *On homotopical additivity of equivariant little disks operads (forthcoming)*. 2025.

# Myriad versions of this theorem

- 1 *Dunn* '88:<sup>7</sup>  $C_1^{\otimes n} \simeq C_n$  w.r.t. a **point-set** tensor product.
- 2 *Brinkmeier* '00:<sup>8</sup>  $C_n \otimes C_m \simeq C_{n+m}$  w.r.t. a **point-set** tensor product.
- 3 *Lurie* '09:<sup>9</sup>  $E_n^{\otimes} \otimes^{BV} E_m^{\otimes} \simeq E_{n+m}^{\otimes}$  with respect to a **homotopical** tensor product. Also, tangential structures.
- 4 *Fiedorowicz-Vogt* '15:<sup>10</sup> Dunn-Brinkmeier's result extends to **arbitrary cofibrant  $E_n$ -operads**.
- 5 *Szczesny* '24:<sup>11</sup>  $D_V \otimes D_W \simeq D_{V \oplus W}$  w.r.t. a **point-set** tensor product.
- 6 *S.* '25:<sup>12</sup>  $E_V^{\otimes} \otimes^{BV} E_W^{\otimes} \simeq E_{V \oplus W}^{\otimes}$  with respect to a **homotopical** tensor product. Also, stalkwise-linearizable tangential structures.

<sup>7</sup> Gerald Dunn. "Tensor product of operads and iterated loop spaces". In: *J. Pure Appl. Algebra* 50.3 (1988), pp. 237–258.

<sup>8</sup> Michael Brinkmeier. *On Operads*. Thesis (Ph.D.)—Universität Osnabrück. 2000.

<sup>9</sup> Jacob Lurie. *Derived Algebraic Geometry VI: Ek Algebras*. 2009. arXiv: 0911.0018 [math. AT].

<sup>10</sup> Z. Fiedorowicz and R. M. Vogt. "An additivity theorem for the interchange of  $E_n$  structures". In: *Adv. Math.* 273 (2015), pp. 421–484.

<sup>11</sup> Ben Szczesny. *Equivariant Framed Little Disk Operads are Additive*. 2024. arXiv: 2410.20235 [math. AT].

<sup>12</sup> Natalie Stewart. *On homotopical additivity of equivariant little disks operads (forthcoming)*. 2025.

Interchange and  
little V-disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# A heavily abridged history of $G$ - $\infty$ -categorical operads

Interchange and  
little  $V$ -disks

Natalie Stewart

- 0 Hill-Hopkins '16:<sup>13</sup>  $G$ -commutative monoids are **semi-Mackey functors**.
- 1 Nardin-Shah '22:<sup>14</sup>  $G$ -operads are a type of fibration over a genuine equivariant version  $\underline{\mathbb{F}}_{G,*}$  of Segal's category  $\Gamma^{\text{op}}$ .
- 2 Barkan-Haugsgeng-Steinebrunner '22:<sup>15</sup>  $G$ -operads are also **fibrations over the effective Burnside 2-category  $\text{Span}(\mathbb{F}_G)$** .
- 3 S. '25:<sup>16</sup> homotopy-coherent interchange is corepresented by **BV tensor products** and  $G$ -operads are monadic over  $G$ -symmetric sequences.
- 4 S. '25:<sup>17</sup> Algebras in **(co)cartesian  $G$ -symmetric monoidal structures** have concrete descriptions and  $\mathcal{N}_{I\infty} \otimes \mathcal{N}_{J\infty} \simeq \mathcal{N}_{IV\infty}$

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>13</sup>Michael A. Hill and Michael J. Hopkins. *Equivariant symmetric monoidal structures*. 2016. arXiv: 1610.03114 [math.AT].

<sup>14</sup>Denis Nardin and Jay Shah. *Parametrized and equivariant higher algebra*. 2022. arXiv: 2203.00072 [math.AT].

<sup>15</sup>Shaul Barkan, Rune Haugseng, and Jan Steinebrunner. *Envelopes for Algebraic Patterns*. 2022. arXiv: 2208.07183 [math.CT].

<sup>16</sup>Natalie Stewart. *Equivariant operads, symmetric sequences, and Boardman-Vogt tensor products*. 2025. arXiv: 2501.02129 [math.CT].

<sup>17</sup>Natalie Stewart. *On tensor products of equivariant commutative operads (draft)*. 2025.

# A quasi-definition of $G$ -operads

Interchange and  
little  $V$ -disks

Natalie Stewart

## Definition

- A  **$G$ -preoperad** is a functor  $\mathcal{O}^\otimes \rightarrow \text{Span}(\mathbb{F}_G)$  with cocartesian lifts over backwards maps.
- A  **$G$ -operad** is required to satisfy “Segal conditions.”
- An  **$\mathcal{O}$ -algebra in  $\mathcal{C}^\otimes$**  is a functor preserving cocartesian arrows :

$$\begin{array}{ccc} \mathcal{O}^\otimes & \xrightarrow{\quad \varphi \quad} & \mathcal{C}^\otimes \\ & \searrow \quad \swarrow & \\ & \text{Span}(\mathbb{F}_G) & \end{array}$$

$$\begin{array}{ccc} & \text{Lop}_G & \\ \text{Op}_G & \xleftarrow{\quad} & \text{POp}_G \\ & \text{⊥} & \\ & \xrightarrow{\quad} & \end{array} \quad \text{Cat}_{\infty, / \text{Span}(\mathbb{F}_G)}^{\text{Backwards-cocart}}$$

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References



# The underlying $G$ -symmetric sequence

Interchange and  
little  $V$ -disks

Natalie Stewart

## Construction

Given  $\mathcal{O} \in \text{Op}_G$ ,  $H \subset G$ , and  $S \in \mathbb{F}_G$ , define the **structure space**

$$\begin{array}{ccc} \mathcal{O}(S) & \xrightarrow{\quad} & \text{Mor}(\mathcal{O}^{\otimes}) \\ \downarrow & \lrcorner & \downarrow \\ \{\text{Ind}_H^G S \rightarrow [G/H]\} & \longrightarrow & \text{Mor}(\text{Span}(\mathbb{F}_G)) \end{array}$$

## Proposition (S.<sup>18</sup>)

If  $\mathcal{O}^{\otimes}$  has “one color” then is conservatively identified by  $(\mathcal{O}(S))_{\substack{H \subset G \\ S \in \mathbb{F}_H}}.$

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>18</sup>Natalie Stewart. *Equivariant operads, symmetric sequences, and Boardman-Vogt tensor products*. 2025. arXiv: 2501.02129 [math.CT].

## Example

The **little V-disks G-operad** has S-ary structure space given by **H-equivariant configurations of S in V**:

$$\mathbb{E}_V(S) := \text{Emb}^{H, \text{Affine}}(S \cdot D(V), D(V)) \simeq \text{Conf}_S^H(V).$$

## Example

Given a G-symmetric monoidal category  $\mathcal{C}^\otimes : \text{Span}(\mathbb{F}_G) \rightarrow \text{Cat}_\infty$ , its unstraightening is a G-operad. Given  $X \in \mathcal{C}(G/G)$ , there is an **endomorphism G-operad** with

$$\text{End}_X(S) \simeq \text{Map}_{\mathcal{C}_H} \left( (\text{Res}_H^G X)^{\otimes S}, \text{Res}_H^G X \right)$$

Cold open

Historical context

G-∞-categorical  
operads

G-(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

In particular, an  $\mathbb{E}_V$ -algebra in  $\mathcal{C}$  consists of an object  $X \in \mathcal{C}_G$  and homotopy-coherently compatible maps

$$\mathrm{Conf}_S^H(V) \rightarrow \mathrm{Map}_{\mathcal{C}_H} \left( (\mathrm{Res}_H^G X)^{\otimes S}, \mathrm{Res}_H^G X \right).$$

## Example (Horev-Klang-Zou<sup>19</sup> '20)

Let  $\underline{\mathcal{S}}_G^{G-\times}$  be the *cartesian structure* on  $G$ -spaces. Then, for all  $X \in \mathcal{S}_G$ , we have  $\Omega^V X \in \mathrm{Alg}_{\mathbb{E}_V}(\mathcal{S}_G)$ .

## Example (Horev-Klang-Zou '20, loc. cit.)

Let  $\underline{\mathrm{Sp}}_G^{\otimes}$  be the HHR  $G$ -symmetric monoidal structure. If  $f: \Omega^V X \rightarrow \underline{\mathrm{Pic}}(\underline{\mathrm{Sp}}_G)$  is a **V-loop map**, then  $\mathrm{Th}(f) \in \mathrm{Alg}_{\mathbb{E}_V}(\mathrm{Sp}_G)$ .

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>19</sup>Jeremy Hahn, Asaf Horev, Inbar Klang, Dylan Wilson, and Fofing Zou. *Equivariant nonabelian Poincaré duality and equivariant factorization homology of Thom spectra*. 2024. arXiv: 2006.13348 [math.AT].

# Modelling G-operadic constructions, two ways

*Model categorist:* **(co)fibrantly replace**, then apply a construction to monoids in G-symmetric sequences.

*$\infty$ -categorist:* apply a G-preoperadic construction, then  **$L_{\text{Op}_G}$ -localize**.

*Shared goal:* model corepresenting object for **pairings** (aka interchanging algebras, bifunctors, etc.) akin to May.<sup>20</sup>

$$\begin{array}{ccc} \mathcal{O}^{\otimes} \times \mathcal{P}^{\otimes} & \xrightarrow{\text{"pairing"}} & \mathcal{Q}^{\otimes} \\ \downarrow \pi & & \downarrow \pi \\ \text{Span}(\mathbb{F}_G) \times \text{Span}(\mathbb{F}_G) & \xrightarrow{\wedge} & \text{Span}(\mathbb{F}_G) \end{array}$$

Today, we are  $\infty$ -categorists.

<sup>20</sup>J. P. May. "Pairings of categories and spectra". In: *J. Pure Appl. Algebra* 19 (1980), pp. 299–346.

Interchange and  
little V-disks

Natalie Stewart

Cold open

Historical context

G- $\infty$ -categorical  
operads

G-(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# The Boardman-Vogt tensor product

Interchange and  
little V-disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

## Definition

$$\mathcal{O}^{\otimes} \overset{\text{BV}}{\otimes} \mathcal{P}^{\otimes} := L_{\text{Op}_G} \left( \mathcal{O}^{\otimes} \times \mathcal{P}^{\otimes} \longrightarrow \text{Span}(\mathbb{F}_G)^2 \xrightarrow{\wedge} \text{Span}(\mathbb{F}_G) \right)$$

$\text{Alg}_{\mathcal{O}}(\mathcal{C})$  lifts to a “pointwise”  $G$ -symmetric monoidal category  $\underline{\text{Alg}}_{\mathcal{O}}^{\otimes}(\mathcal{C})$ .

## Proposition (S.<sup>21</sup>)

$(-) \overset{\text{BV}}{\otimes} \mathcal{O}^{\otimes}$  is left adjoint to  $\underline{\text{Alg}}_{\mathcal{O}}^{\otimes}(-)$ , so

$$\text{Alg}_{\mathcal{O}} \underline{\text{Alg}}_{\mathcal{P}}^{\otimes}(\mathcal{C}) \simeq \text{Alg}_{\mathcal{O} \otimes \mathcal{P}}(\mathcal{C}).$$

<sup>21</sup>Natalie Stewart. *Equivariant operads, symmetric sequences, and Boardman-Vogt tensor products*. 2025. arXiv: 2501.02129 [math.CT].

# The Dunn map

Interchange and  
little V-disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations


Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References



# Preoperad models for 1-colored $G$ -operads

Interchange and  
little  $V$ -disks

Natalie Stewart

Given  $\mathcal{P}^\otimes \in \mathbf{POp}_G$ , We let  $\mathcal{P}_{/P}^{\text{act}}$  be the  $\infty$ -category of arrows  $P' \rightarrow P$  projecting to a forward map  $T = T \rightarrow S$ .

## Definition

A map of  $G$ -preoperads  $\alpha: \mathcal{P}^\otimes \rightarrow \mathcal{O}^\otimes$  with  $\mathcal{O}^\otimes$  a “one color”  $G$ -operad is a **weak approximation** if

- 1 The  $G$ - $\infty$ -category of colors  $U\mathcal{P}$  **has a terminal  $G$ -object**, and
- 2 For all  $P \in \mathcal{P}^\otimes$  and  $T \rightarrow \pi(P)$ , **the map of spaces**

$$B\left(\mathcal{P}_{/P}^{\text{act}} \times_{\mathbb{F}_{G,/\pi(P)}} \{T \rightarrow \pi(P)\}\right) \rightarrow \prod_{[H/K] \subset \pi(P)} \mathcal{O}(T \times_{\pi(P)} [H/K])$$

**is a weak equivalence.**

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# Preoperad models for 1-colored $G$ -operads

Interchange and  
little  $V$ -disks

Natalie Stewart

## Proposition (Harpaz<sup>22</sup> '19 + reinterpretation)

If  $\alpha: \mathcal{P}^{\otimes} \rightarrow \mathcal{O}^{\otimes}$  is a weak approximation, pullback is fully faithful

$$\mathrm{Alg}_{\mathcal{O}}(\mathcal{S}_G) \subset \mathrm{Alg}_{\mathcal{P}}(\mathcal{S}_G)$$

with image the  $\mathcal{P}$ -monoids whose “color”  $G$ -functors  $U\mathcal{P} \rightarrow \underline{\mathcal{S}}_G$  are constant.

Weak approximations can be made to have many colors; a weak approximation  $\alpha$  is a **strong approximation** if  $U\mathcal{P} \rightarrow U\mathcal{O}$  is an equivalence.

## Proposition (S.<sup>23</sup>)

$\mathrm{Alg}_{(-)}(\mathcal{S}_G)$  detects  $L_{\mathrm{Op}_G}$ -equivalences when  $U\mathcal{P} \rightarrow U\mathcal{O}$  is an equivalence; in particular, **strong approximations are  $L_{\mathrm{Op}_G}$ -equivalences**.

<sup>22</sup>Yonatan Harpaz. *Little cubes algebras and factorization homology (course notes)*.

<sup>23</sup>Natalie Stewart. *On tensor products of equivariant commutative operads (draft)*. 2025.

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References



# The $L_{\text{Op}_G}$ -conservativity argument, in short

Interchange and  
little  $V$ -disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

## Proof sketch.

Examine the free  $\mathcal{O}$ - $G$ -space monad:<sup>24</sup>

$$(T_{\mathcal{O}}X)^H \simeq \coprod_{S \in \mathbb{F}_H} \left( \mathcal{O}(S) \times (X^S)^H \right)_{h \text{Aut}_H S}.$$

The inclusion  $\text{Aut}_H(S) \subset \text{End}_H(S) = (S^S)^H$  yields natural splitting

$$(T_{\mathcal{O}}S)^H \simeq \mathcal{O}(S) \oplus \text{Junk}.$$

Use monadicity of  $\text{Alg}_{\mathcal{O}}(\mathcal{S}_G) \rightarrow \mathcal{S}_G$  and conservativity of  $(\mathcal{O}(S))$ . □

<sup>24</sup>Natalie Stewart. *Equivariant operads, symmetric sequences, and Boardman-Vogt tensor products*. 2025. arXiv: 2501.02129 [math.CT].

## Proposition (Dugger-Isaksen<sup>25</sup> '01)

If  $X$  is a topological space and  $\mathfrak{D} \subset \mathcal{P}(X)$  a basis of contractible open subsets, then we get a weak equivalence

$$B\mathfrak{D} \xrightarrow{\sim} X$$

## Corollary

Let  $\mathfrak{D}_S^H(V) \subset \text{Conf}_S^H(V)$  be the basis of configurations in affinely  $\coprod_S D(V)$ -shaped invariant subspaces of  $D(V)$ . We get a weak equivalence

$$B\mathfrak{D}_S^H(V) \xrightarrow{\sim} \text{Conf}_S^H(V) \simeq \mathbb{E}_V(S).$$

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

<sup>25</sup>Daniel Dugger and Daniel C. Isaksen. "Topological hypercovers and  $A^1$ -realizations". In: *Math. Z.* 246.4 (2004), pp. 667–689.

$$\begin{array}{ccccccc}
 \mathbb{P}_V^\otimes \overset{BV}{\otimes} \mathbb{P}_W^\otimes & \xleftarrow{L_P} & \mathbb{P}_V^\otimes \times \mathbb{P}_W^\otimes & \xrightarrow{M_P} & \mathbb{P}_V^\otimes \wr \mathbb{P}_W^\otimes & \xrightarrow{\tilde{\varphi}_P} & \mathbb{P}_{V|W}^\otimes \longleftrightarrow \mathbb{P}_{V\oplus W}^\otimes \\
 \alpha_V \otimes \alpha_W \downarrow & & \alpha_V \times \alpha_W \downarrow & & \alpha_V \wr \alpha_W \downarrow & & \alpha_{V|W} \downarrow \qquad \qquad \downarrow \alpha_{V\oplus W} \\
 \mathbb{E}_V^\otimes \overset{BV}{\otimes} \mathbb{E}_W^\otimes & \xleftarrow{L_E} & \mathbb{E}_V^\otimes \times \mathbb{E}_W^\otimes & \xrightarrow{M_E} & \mathbb{E}_V^\otimes \wr \mathbb{E}_W^\otimes & \xrightarrow{\varphi_E} & \mathbb{E}_{V\oplus W}^\otimes = \mathbb{E}_{V\oplus W}^\otimes
 \end{array}$$

We define a  $G$ -preoperad  $\mathbb{P}_V^\otimes$  such that  $\mathbb{P}_{V,/P}^{\text{act}} \simeq \mathfrak{D}_S^H(V)$ , yielding a weak approximation  $\alpha_V: \mathbb{P}_V^\otimes \rightarrow \mathbb{E}_V^\otimes$ . Then, we define a  $\mathbb{P}$ -Dunn map fitting into the above diagram.

Here,  $\mathbb{P}_{V|W}^\otimes$  is the “ $G$ -preoperadic image, i.e. “decomposed little disks.”

We’re tasked with verifying that  $\varphi_E \circ M_E$  induces an equivalence

$$\text{Alg}_{\mathbb{E}_{V\oplus W}}(\mathcal{S}_G) \xrightarrow{\sim} \text{Alg}_{\mathbb{E}_V \times \text{Alg}_{\mathbb{E}_W}}(\mathcal{S}_G) \xleftarrow{\sim} \text{Alg}_{\mathbb{E}_V \otimes \mathbb{E}_W}(\mathcal{S}_G)$$

$$\begin{array}{ccccccc}
 P_V^\otimes \overset{BV}{\otimes} P_W^\otimes & \xleftarrow{L_P} & P_V^\otimes \times P_W^\otimes & \xrightarrow{M_P} & P_V^\otimes \wr P_W^\otimes & \xrightarrow{\tilde{\varphi}_P} & P_{V|W}^\otimes \longleftrightarrow P_{V\oplus W}^\otimes \\
 \alpha_V \otimes \alpha_W \downarrow & & \alpha_V \times \alpha_W \downarrow & & \alpha_V \wr \alpha_W \downarrow & & \alpha_{V|W} \downarrow \quad \alpha_{V\oplus W} \downarrow \\
 E_V^\otimes \overset{BV}{\otimes} E_W^\otimes & \xleftarrow{L_E} & E_V^\otimes \times E_W^\otimes & \xrightarrow{M_E} & E_V^\otimes \wr E_W^\otimes & \xrightarrow{\varphi_E} & E_{V\oplus W}^\otimes = E_{V\oplus W}^\otimes
 \end{array}$$

- 1  $L_P$  and  $L_E$  are  $L_{Op_C}$ -equivalences by fiat.
- 2 A variant of Harpaz's strategy<sup>26</sup> shows that  $M_P$  and  $M_E$  are  $L_{Op_C}$ -equivalences.
- 3 Simple  $\infty$ -category theory shows that  $\alpha_V \times \alpha_W$  is a weak approximation.
- 4 A variant of Dunn's strategy<sup>27</sup> shows that  $\alpha_{V|W}$  is a weak approximation.
- 5 Explicit 1-category theory shows that  $\tilde{\varphi}_P$  is a strong approximation.
- 6 Routine bookkeeping then shows that  $\text{Alg}_{E_{V\oplus W}}(\mathcal{S}_C) \xrightarrow{\sim} \text{Alg}_{E_V \times E_W}(\mathcal{S}_C)$ .

<sup>26</sup>Yonatan Harpaz. *Little cubes algebras and factorization homology (course notes)*.

<sup>27</sup>Gerald Dunn. "Tensor product of operads and iterated loop spaces". In: *J. Pure Appl. Algebra* 50.3 (1988), pp. 237–258.

Cold open

Historical context

 $G$ - $\infty$ -categorical  
operads $G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# Stalkwise-linearizable tangential structures

Define  $\mathbb{E}_{B_G^{\text{linTop}(n)}}^{\otimes}$  to have colors the linearizable  $G$ -actions on  $\mathbb{R}^n$  and operations the *topological* embeddings.

Given a  $G$ -space  $X$  with stalkwise-linearizable equivariant  $\mathbb{R}^n$ -bundle  $T_{\bullet}: X \rightarrow B_G^{\text{linTop}(n)}$ , we define the assembly

$$\mathbb{E}_X^{\otimes} := \text{Lop}_G \left( \mathbb{E}_{B_G^{\text{linTop}(n)}}^{\otimes} \times_{B_G^{\text{linTop}(n)}^{G-\sqcup}} X^{G-\sqcup} \right).$$

## Theorem (S.<sup>28</sup>)

There is a natural equivariant colimit expression of operads

$$\underline{\text{colim}}_{x \in X} \mathbb{E}_{T_x}^{\otimes} \xrightarrow{\sim} \mathbb{E}_X^{\otimes}$$

<sup>28</sup> Natalie Stewart. *On homotopical additivity of equivariant little disks operads (forthcoming)*. 2025.

Interchange and  
little  $V$ -disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# Stalkwise-linearizable tangential structures

Interchange and  
little V-disks

Natalie Stewart

## Corollary (S.)

Given stalkwise-linearizable equivariant  $\mathbb{R}^k$ -bundles  $X \rightarrow B_G^{\text{lin}}\text{Top}(n)$  and  $Y \rightarrow B_G^{\text{lin}}\text{Top}(m)$ , fixing the direct sum structure

$$X \times Y \longrightarrow B_G^{\text{lin}}\text{Top}(n) \times B_G^{\text{lin}}\text{Top}(m) \xrightarrow{\oplus} B_G^{\text{lin}}\text{Top}(n+m),$$

there is an equivalence.

$$\mathbb{E}_X^{\otimes} \otimes^{\text{BV}} \mathbb{E}_Y^{\otimes} \xrightarrow{\sim} \mathbb{E}_{X \times Y}^{\otimes}.$$

Other cases that are covered by the original argument:

- (Equivariant) swiss cheese
- Stratified little disks

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# A curious corollary about geometric topology

Let  $\underline{\text{Disk}}_G^{X\text{-fr}, \sqcup} \subset \underline{\text{Mfld}}_G^{X\text{-fr}, \sqcup}$  be the  $G$ -symmetric monoidal full category whose objects are disjoint unions of  $X$ -framed  $G$ -disks and whose mapping spaces are  $X$ -framed disk embeddings.

## Corollary

There is a  $G$ -symmetric monoidal equivalence

$$\underline{\text{Disk}}_G^{X\text{-fr}, \sqcup} \square \underline{\text{Disk}}_G^{Y\text{-fr}, \sqcup} \simeq \underline{\text{Disk}}_G^{X \times Y\text{-fr}, \sqcup}.$$

Here,  $\square$  is the box product of semi-Mackey functors.

Interchange and  
little  $V$ -disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little  $V$ -disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References

# References

- [BHS22] Shaul Barkan, Rune Haugseng, and Jan Steinebrunner. *Envelopes for Algebraic Patterns*. 2022. arXiv: 2208 . 07183 [math . CT] .
- [Bri00] Michael Brinkmeier. *On Operads*. Thesis (Ph.D.)—Universität Osnabrück. 2000.
- [Dot12] Emanuele Dotto. *Stable real K-theory and real topological Hochschild homology*. Thesis (Ph.D.)—University of Copenhagen. 2012. arXiv: 1212 . 4310 [math . AT] .
- [DMPR21] Emanuele Dotto, Kristian Moi, Irakli Patchkoria, and Sune Precth Reeh. "Real topological Hochschild homology". In: *J. Eur. Math. Soc. (JEMS)* 23.1 (2021), pp. 63–152.
- [Dug05] Daniel Dugger. "An Atiyah-Hirzebruch spectral sequence for KR-theory". In: *K-Theory* 35.3–4 (2005), 213–256 (2006).
- [DI04] Daniel Dugger and Daniel C. Isaksen. "Topological hypercovers and  $A^1$ -realizations". In: *Math. Z.* 246.4 (2004), pp. 667–689.
- [Dun88] Gerald Dunn. "Tensor product of operads and iterated loop spaces". In: *J. Pure Appl. Algebra* 50.3 (1988), pp. 237–258.
- [FV15] Z. Fiedorowicz and R. M. Vogt. "An additivity theorem for the interchange of  $E_n$  structures". In: *Adv. Math.* 273 (2015), pp. 421–484.
- [HHKWZ24] Jeremy Hahn, Asaf Horev, Inbar Klang, Dylan Wilson, and Foling Zou. *Equivariant nonabelian Poincaré duality and equivariant factorization homology of Thom spectra*. 2024. arXiv: 2006 . 13348 [math . AT] .
- [Har] Yonatan Harpaz. *Little cubes algebras and factorization homology (course notes)*.
- [HH16] Michael A. Hill and Michael J. Hopkins. *Equivariant symmetric monoidal structures*. 2016. arXiv: 1610 . 03114 [math . AT] .
- [DAG-VI] Jacob Lurie. *Derived Algebraic Geometry VI: Ek Algebras*. 2009. arXiv: 0911 . 0018 [math . AT] .
- [May80] J. P. May. "Pairings of categories and spectra". In: *J. Pure Appl. Algebra* 19 (1980), pp. 299–346.
- [NS22] Denis Nardin and Jay Shah. *Parametrized and equivariant higher algebra*. 2022. arXiv: 2203 . 00072 [math . AT] .
- [Ste25a] Natalie Stewart. *Equivariant operads, symmetric sequences, and Boardman-Vogt tensor products*. 2025. arXiv: 2501 . 02129 [math . CT] .
- [Ste25b] Natalie Stewart. *On homotopical additivity of equivariant little disks operads (forthcoming)*. 2025.
- [Ste25c] Natalie Stewart. *On tensor products of equivariant commutative operads (draft)*. 2025.
- [Szc24] Ben Szczesny. *Equivariant Framed Little Disk Operads are Additive*. 2024. arXiv: 2410 . 20235 [math . AT] .

This presentation was made in Beamer, with figures via tikz-cd and Inkscape, presented via Impressive. The tex is on my website. The title slide is  $H_{C_2}^{\star}(*_{C_2}; \mathbb{Z})$  [Dug05].

Interchange and  
little V-disks

Natalie Stewart

Cold open

Historical context

$G$ - $\infty$ -categorical  
operads

$G$ -(pre)operads, algebras

BV-tensor products

The Dunn map

Weak approximations

Little V-disks

Interlude: the homotopy type of  
a smooth manifold

The strategy

With tangential structure

References