

Business Analytics & Machine Learning

Homework sheet 1: Statistics

Prof. Dr. Martin Bichler

Julius Durmann, Markus Ewert, Yutong Chao, Dr. Mete Ahunbay

Exercise H1.1 *Population mean*

Determine (with $\alpha = 0.05$) if the following sample was obtained from a population with zero mean:

2, 3, 2, 4, 2, 4, 5, 2, 1, 4, 3, 0, 3, 2, 4, 5, 3, 3, 0, 1.

Exercise H1.2 *Caloric intake*

32 individuals take part in a study about nutritional behavior. One aspect of the study is comparing carnivore diets to non-carnivore diets in terms of daily caloric intake. The research hypothesis states, that the daily average caloric intake of individuals following a non-carnivore diet is lower, compared to individuals following a carnivore diet. Out of 32 participants, 12 adhere to a non-carnivore diet, yielding an average caloric intake of $\bar{x}_1 = 1780$ kcal. In contrast, the remaining 20 participants following a carnivore diet average to $\bar{x}_2 = 1900$ kcal per day. The respective estimated standard deviations result in $s_1 = 230$, and $s_2 = 250$. The daily caloric intake of an individual is assumed to be a normally distributed variable.

- Give a 95% confidence interval of the average daily caloric intake for each of the groups.
- Which conclusions can be drawn from the computed confidence intervals?
- Identify and apply a suitable hypothesis test using a significance level of $\alpha = 0.05$.