

Problem Set 2

Natasha Watkins

Exercise 4.4

By 4.3, the characteristic polynomial of $A^H = A$ is

$$p(\lambda) = \lambda^2 - \text{tr}(A)\lambda + \det(A)$$

where A is a 2×2 matrix of the form

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

The diagonal elements of a Hermitian matrix are real. So the diagonal elements of A are also real. We can check that the roots of the characteristic polynomial are real by finding the discriminant.

$$\begin{aligned} [\text{tr}(A)]^2 - 4\det(A) &= (a + d)^2 - 4(ad - bc) \\ &= a^2 + 2ad + d^2 - 4ad + 4bc \\ &= a^2 - 4ad + d^2 + 4bc \\ &= (a - d)^2 + 4bc \end{aligned}$$

Part i)

As A^H is Hermitian, $b = \bar{c}$, so $4bc \geq 0$. Therefore the discriminant is positive and the roots are real.

Part ii)

If $A^H = -A$, then $b = -\bar{c}$, so $4bc \leq 0$. Therefore, the discriminant is negative and the roots are imaginary.

Exercise 4.6

Consider a matrix A that is upper triangular. We know that the determinant of an upper triangular matrix is the product of its diagonal elements, ie. $\det(A) = \prod_{i=1}^n a_{ii}$.

The eigenvalues of a matrix are such that $\det(\lambda I - A) = 0$.

Therefore,

$$\begin{aligned}\det(\lambda I - A) &= \prod_{i=1}^n (\lambda_i - a_{ii}) = 0 \\ &= (\lambda_1 - a_{11}) \cdot (\lambda_2 - a_{22}) \cdots (\lambda_n - a_{nn}) = 0\end{aligned}$$

The roots of the characteristic polynomial (ie. the eigenvalues) are given by the diagonal elements of A .

Exercise 4.13

Solving for the spectrum of A gives $\sigma(A) = \{\frac{2}{5}, 1\}$, with corresponding eigenvectors of $[1, -1]^T$ and $[2, 1]^T$. Let P be the transition matrix, where

$$P = \begin{bmatrix} 1 & 1 \\ -1 & 2 \end{bmatrix}, \quad D = \begin{bmatrix} \frac{2}{5} & 0 \\ 0 & 1 \end{bmatrix}, \quad P^{-1} = \frac{1}{3} \begin{bmatrix} 1 & 1 \\ -2 & 1 \end{bmatrix}$$

Solving $P^{-1}AP$ gives

$$\begin{bmatrix} 1 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 0.8 & 0.4 \\ 0.2 & 0.6 \end{bmatrix} \frac{1}{3} \begin{bmatrix} 1 & 1 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} \frac{2}{5} & 0 \\ 0 & 1 \end{bmatrix} = D$$

Exercise 4.18

We know that λ satisfies $A\mathbf{x} = \lambda\mathbf{x}$ for some \mathbf{x} . Taking the transpose, we see $\mathbf{x}^T A^T = \lambda\mathbf{x}^T$. This is equivalent to $\det(A^T - \lambda I) = 0$.

The determinant of a matrix A is equal to the determinant of its transpose, so

$$\det(A^T - \lambda I) = \det(A - \lambda I) = 0$$

As we know $\det(A - \lambda I) = 0$ holds for some \mathbf{x} , there exists a vector \mathbf{x}^T such that $\det(A^T - \lambda I) = 0$ holds.

Exercise 4.20

If A^H and B are orthornormally similar, then there is U such that $B = U^H A^H U$.

$$B^H = (U^H A^H U)^H = U^H A U = U^H A^H U = B$$

Therefore, B is also Hermitian.

Exercise 4.27

A is a positive definite matrix, so it is Hermitian. The diagonal elements of a Hermitian matrix are real, as they are their own complex conjugate.

It also is true that $\langle \mathbf{x}, A\mathbf{x} \rangle = \mathbf{x}^H A \mathbf{x} > 0$ for any vector \mathbf{x} . Consider the vector $e_1 = [1 \ 0 \ 0 \ \cdots \ 0]^T$. $e_1^H A e_1$ selects the first diagonal element of A . Likewise, the vector e_2 selects the second diagonal element of A . As $\langle A\mathbf{x}, \mathbf{x} \rangle = \mathbf{x}^H A \mathbf{x} > 0$ holds for all vectors, the diagonal elements are all positive.