

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/267032487>

Neuropathogenesis of Delirium: Review of Current Etiologic Theories and Common Pathways

Article in *The American journal of geriatric psychiatry: official journal of the American Association for Geriatric Psychiatry* · December 2013

DOI: 10.1016/j.jagp.2013.09.005

CITATIONS

379

READS

5,899

1 author:



Jose Maldonado

Stanford Medicine

116 PUBLICATIONS 3,215 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Prediction and Treatment of AWS in the medically ill [View project](#)



Biomarkers of Chronic Fatigue Syndrome [View project](#)

Neuropathogenesis of Delirium: Review of Current Etiologic Theories and Common Pathways

José R. Maldonado, M.D., F.A.P.M., F.A.C.F.E.

Delirium is a neurobehavioral syndrome caused by dysregulation of neuronal activity secondary to systemic disturbances. Over time, a number of theories have been proposed in an attempt to explain the processes leading to the development of delirium. Each proposed theory has focused on a specific mechanism or pathologic process (e.g., dopamine excess or acetylcholine deficiency theories), observational and experiential evidence (e.g., sleep deprivation, aging), or empirical data (e.g., specific pharmacologic agents' association with postoperative delirium, intraoperative hypoxia). This article represents a review of published literature and summarizes the top seven proposed theories and their interrelation. This review includes the "neuroinflammatory," "neuronal aging," "oxidative stress," "neurotransmitter deficiency," "neuroendocrine," "diurnal dysregulation," and "network disconnectivity" hypotheses. Most of these theories are complementary, rather than competing, with many areas of intersection and reciprocal influence. The literature suggests that many factors or mechanisms included in these theories lead to a final common outcome associated with an alteration in neurotransmitter synthesis, function, and/or availability that mediates the complex behavioral and cognitive changes observed in delirium. In general, the most commonly described neurotransmitter changes associated with delirium include deficiencies in acetylcholine and/or melatonin availability; excess in dopamine, norepinephrine, and/or glutamate release; and variable alterations (e.g., either a decreased or increased activity, depending on delirium presentation and cause) in serotonin, histamine, and/or γ -aminobutyric acid. In the end, it is unlikely that any one of these theories is fully capable of explaining the etiology or phenomenologic manifestations of delirium but rather that two or more of these, if not all, act together to lead to the biochemical derangement and, ultimately, to the complex cognitive and behavioral changes characteristic of delirium. (Am J Geriatr Psychiatry 2013; 21:1190–1222)

Key Words: Aging, delirium, encephalopathy, physiologic stress, neuroinflammation, large neutral amino acids, neuroendocrine, oxidative stress

Received May 23, 2012; revised September 10, 2013; accepted September 13, 2013. From the Departments of Psychiatry, Internal Medicine & Surgery and the Psychosomatic Medicine Service, Stanford University School of Medicine, and Board of Directors, American Delirium Society, Stanford, CA. Send correspondence and reprint requests to José R. Maldonado, M.D., F.A.P.M., F.A.C.F.E., Departments of Psychiatry, Internal Medicine & Surgery, and the Psychosomatic Medicine Service, Stanford University School of Medicine, 401 Quarry Rd., Office #2317, Stanford, CA 94305. e-mail: jrm@stanford.edu

© 2013 American Association for Geriatric Psychiatry

<http://dx.doi.org/10.1016/j.jagp.2013.09.005>

DELIRIUM: SCOPE OF THE PROBLEM

Delirium is a neurobehavioral syndrome caused by dysregulation of baseline neuronal activity secondary to systemic disturbances,¹ characterized by an alteration in the level of attention and awareness, which develops over a relative short period of time, and represents a change from the subject's baseline.² Clinically, delirium is an acute or subacute organic mental syndrome characterized by a disturbance in attention (i.e., reduced ability to direct, focus, sustain, and shift attention) and awareness (reduced orientation to the environment), cognition (e.g., memory deficit, disorientation), language, visuospatial ability, or perception that is not better explained by a preexisting, established, or other evolving neurocognitive disorder.^{2–5}

A recent study at a large general hospital found that the point prevalence of delirium is one in every five inpatients, with some variation in certain medicosurgical units and populations.⁶ Studies have found that between 14% and 24% of elderly patients are admitted to the hospital with delirium^{7,8} and that delirium occurs in up to 50% of elderly inpatients.⁹ In the emergency department, the prevalence in elderly patients is 9.6%.¹⁰ The reported rate of postoperative delirium ranges between 10% and 74% depending on the type of surgery and population under study.^{11–15} Studies have demonstrated that up to 87% of critically ill patients develop delirium during their intensive care unit (ICU) stay.¹⁶ This is important considering that the proportion of patients in the ICU aged at least 65 years is 56%.¹⁷ Furthermore, studies predict that by 2015 the rate of elderly aged 80 years and older admitted to the ICU will increase by 72%, representing roughly one in four admissions to the ICU.¹⁸

DELIRIUM CONSEQUENCES

Delirium has been reported to be one of the six leading causes of preventable conditions in hospitalized elderly patients.¹⁹ After controlling for demographics, apparent illness severity, age, and medical comorbidities, patients who develop delirium fare much worse than their nondelirious counterparts. The mortality rate for elderly patients in acute care hospitals is much higher among those with delirium than those without delirium: 8% versus

1% in a study of 229 medically ill, hospitalized elderly patients.²⁰ Even years after it occurred, delirium continues to have a long-lasting impact, with a 3-year mortality of 75% for patients who had developed in-hospital delirium versus 51% for control patients (risk ratio: 2.24), even after accounting for prehospital measures of global cognition, physical functioning, and medical comorbidity.²¹ The number of days of delirium older patients experience during an ICU admission is significantly associated with mortality up to 1 year after admission after controlling for severity of illness ($p = 0.001$).²²


Delirious patients, when compared with patients suffering from the same medical problem who do not develop delirium, experience prolonged hospital stays, on average 5–10 days longer.^{20,23–26} Among elderly medically ill inpatients incident delirium was associated with an excess stay after diagnosis of 7.78 days, even after controlling for covariates.²⁷ Studies have demonstrated that delirium in the emergency department was an independent predictor of prolonged hospital length of stay (i.e., twice as long) compared with nondelirious patients, even after adjusting for confounding factors.²⁸ Significantly more patients who develop delirium in the hospital ultimately require institutional postacute care, such as placement in a skilled nursing facility (e.g., 16% versus 3%), even after controlling for illness severity, activities of daily living status, prior cognitive impairment, and fever.^{20,26}

A review of available empirical evidence suggested that a substantial proportion of patients who survive delirium were much more likely to experience long-term cognitive impairment^{4,29–40} and to require institutional postacute care, such as placement in a skilled nursing facility (e.g., 16% versus 3%), even after controlling for illness severity, activities of daily living status, prior cognitive impairment, and fever.^{26,41} The increased morbidity and extended hospital care associated with delirium have been associated with greater care costs.^{24,42,43} The total annual direct healthcare costs attributable to delirium in the United States has been estimated to be as high as \$152 billion.⁴⁴

NEUROPATHOGENESIS OF DELIRIUM

Despite its high prevalence and high morbidity, much is not understood about delirium. Known risk

FIGURE 1. Theories on the development of delirium. Schematics of the interrelationship of current theories on the pathophysiology of delirium and how they may relate to each other. Each proposed theory has focused on a specific mechanism or pathologic process (e.g., DA excess or Ach deficiency theories), observational and experiential evidence (e.g., sleep deprivation, aging), or empirical data (e.g., specific pharmacologic agents' association with postoperative delirium, intraoperative hypoxia). Most of these theories are complementary rather than competing, with many areas of intersection and reciprocal influence. In the end, it is unlikely that any one of these theories is fully capable of explaining the etiology or phenomenologic manifestations of delirium but rather that their interaction lead to the biochemical derangement and, ultimately, to the complex cognitive and behavioral changes characteristic of delirium.



factors for delirium include advanced age, preexisting cognitive impairment (including dementia), medications (especially those with high anticholinergic potential), sleep deprivation, hypoxia and anoxia, metabolic abnormalities, and a history of alcohol or drug abuse.⁴ For years it has been predicted that metabolic processes must be involved in its etiology, as described by Engel and Romano over 50 years ago: "We thus arrive at the proposition that a derangement in functional metabolism underlies all instances of delirium and that this is reflected at the clinical level by the characteristic disturbance in cognitive functions."^{45 (p. 532)} Therefore, we begin with the premise that delirium is a neurobehavioral syndrome caused by the dysregulation of neuronal activity secondary to systemic disturbances.^{1,45–47}

Over the years, a number of theories have been proposed in an attempt to explain the processes

leading to the development of delirium.¹ Each proposed theory has focused on a specific mechanism or pathologic process (e.g., dopamine [DA] excess or acetylcholine [Ach] deficiency theories), observational and experiential evidence (e.g., sleep deprivation, aging), or empirical data (e.g., specific pharmacologic agents' association with postoperative delirium, intraoperative hypoxia). To date, however, no single unitary pathophysiologic mechanism has been identified.

This article summarizes the seven most prominent theories hypothesized to explain the phenomenon of delirium and highlights areas of commonality with the intent of increasing an appreciation of the underlying pathophysiologic mechanisms, help focus future research, and assist in developing prophylactic and treatment strategies. Most of these theories are complementary rather than competing (Fig. 1). Thus,

TABLE 1. Mechanisms Mediating Delirium and Cognitive Impairment

1. A number of factors and mechanisms leading to delirium may also directly cause CNS damage and neuronal dysfunction and thus mediate both the manifestations of delirium and long-term cognitive impairment (e.g., cytokine release and other neuroinflammatory mediators; decrease perfusion and oxygenation leading to decreased cerebral oxidative metabolism; changes in BBB permeability; hypercatabolic states; water and electrolyte imbalances; excessive GC levels and other HPA axis dysfunctions; melatonin and sleep–wake cycle abnormalities).
2. Pharmacologic agents used either to treat the underlying causes of the delirium (e.g., steroids, calcineurin inhibitors, other immunosuppressant agents, DA) or those agents used to treat delirium (e.g., DA blocking agents, benzodiazepines) may themselves lead to neuronal damage in a fragile brain.
3. Any mechanism listed above may themselves lead to alterations in neurotransmitter concentration or receptor sensitivity, which itself may underlie the different symptoms and clinical presentations of delirium and/or long-term cognitive dysfunction. Thus, the same mechanisms that cause the substrate for delirium may mediate the cognitive impairments observed after the acute presentation of delirium has resolved.
4. It is possible that instead of causing cognitive deficits or dementia, delirium (and its underlying causes) only serve as a catabolic agent leading to an acceleration of normal physiologic cerebral aging mechanisms leading to dementia.
5. It is also possible that an episode of delirium simply unmasks subtle cognitive deficits already present but not yet identified.


Notes: Source: Maldonado²²²

it is likely that none of these theories by themselves is fully capable of explaining the etiology or phenomenologic manifestations of delirium but rather that two or more of these, if not all, act together to lead to the biochemical derangement we know as delirium (Table 1).

NEUROINFLAMMATORY HYPOTHESIS

The Neuroinflammatory Hypothesis (NIH) proposes that acute peripheral inflammatory stimulation (from infectious, surgical, or traumatic etiologies) induces activation of brain parenchymal cells and expression of proinflammatory cytokines and inflammatory mediators in the central nervous system (CNS) that induce neuronal and synaptic dysfunction and subsequent neurobehavioral and cognitive symptoms characteristic of delirium (Fig. 2).^{48–53} Others have previously demonstrated how the brain monitors the presence of peripheral inflammation and how upon exposure to infection or inflammation sick individuals develop nonspecific physiologic (e.g., fever, pain, malaise, fatigue, and anorexia) and behavioral


FIGURE 2. Schematic illustration of neural immune connections. Immune signaling of the CNS via systemic routes and the vagus nerve (Vagus n.) and CNS regulation of immunity via the HPA, hypothalamic–pituitary–thyroid, and hypothalamic–pituitary–gonadal axes and the sympathetic nervous system (SNS) and parasympathetic nervous system (PNS). Cytokine expression within the brain is represented by asterisks within the brain. Dotted lines represent negative regulatory pathways, and solid lines represent positive regulatory pathways. CRH: corticotrophin-releasing hormone; AVP: arginine vasopressin; TRH: thyrotropin-releasing hormone; GnRH: gonadotropin-releasing hormone; ACTH: adrenocorticotrophin hormone; TSH: thyroid-stimulating hormone; T₄: thyroxine; T₃: triiodothyronine; LH: luteinizing hormone; FSH: follicle-stimulating hormone; LC: locus coeruleus; A1, C1, A2, C2: brainstem adrenergic nuclei. (From Marques-Deak et al.³⁷⁷).



(e.g., lethargy, depressed mood, social withdrawal, cognitive loss, and anhedonia) changes known as “sickness behavior.”^{54–59}

Thus, ultimately the NIH suggests that delirium may represent the CNS manifestation of a systemic disease state that has indeed crossed the blood–brain

FIGURE 3. Functional anatomy of the inflammatory reflex. Afferent (sensory) neural signals to the brain stem are relayed by the vagus nerve to the nucleus of the solitary tract (nucleus tractus solitaries [NTS]). Polysynaptic relays then connect to the outflow centers of the autonomic nervous system, the rostral ventrolateral medullary (RVLM) sympathoexcitatory neurons and the vagal motor neurons in the nucleus ambiguus (NA), and the dorsal vagal motor nucleus. Outflow arrives at the coeliac ganglion from either the vagus nerve or the preganglionic efferent nerves, which originate in the sympathetic trunk. Stimulating the vagus nerve suppresses innate immune responses and down-regulates proinflammatory cytokine release in the spleen through a mechanism that depends on nicotinic ACh receptor subunit $\alpha 7$ ($\alpha 7nAChR$). Note that after the activation of the inflammatory reflex by sensory input to the brainstem, the signals are also relayed to the nuclei controlling the function of the HPA axis, which increases GC hormone release by the adrenal gland. This provides an important connection between the neural networks that can acutely provide compensatory signals to adjust immune responses and the humoral anti-inflammatory mechanisms that can more chronically modulate innate and adaptive immune responses. (From Tracey³⁷⁸).



Nature Reviews | Immunology

barrier (BBB).¹ Many of the circumstances associated with a high incidence of delirium (e.g., infections, postoperative states) may be also associated with BBB integrity compromise. It is theorized that several illness processes (i.e., trauma, infections) and surgical procedures may introduce triggering factors leading to the activation of the inflammatory cascade: use of anesthetic agents, extensive tissue trauma, presence of foreign organisms or substances, elevated hormone levels, blood loss and anemia, blood transfusions, use of extracorporeal circulation, hypoxia, ischemia and reperfusion, formation of heparin–protamine complexes, and microemboli formation and migration (reviewed by Maldonado¹).

Others have demonstrated two pathways through which immune signals from the periphery are transduced to the brain (Fig. 3): the neural pathway and the humoral pathway.⁶⁰ In the neural pathway,

peripherally produced pathogen-associated molecular patterns and cytokines activate primary afferent nerves, such as the vagus nerve. The humoral pathway involves circulating pathogen-associated molecular patterns that reach the brain at the level of the choroid plexus and the circumventricular organs where pathogen-associated molecular patterns induce the production and release of proinflammatory cytokines by macrophage-like cells expressing Toll-like receptors. Several studies have demonstrated that patients develop delirium during acute medical hospitalizations experienced elevation of C-reactive protein, interleukin (IL)-6, tumor necrosis factor- α , IL-1RA, IL-10, and IL-8 as compared with patients who did not have delirium, even after adjusting for infection, age, and cognitive impairment, suggesting an association between proinflammatory cytokines and the pathogenesis of delirium.^{61–63}

A study of adult patients admitted to medicine wards showed that those who developed delirium had significantly elevated levels of IL-6 (53% versus 31%) and IL-8 (45% versus 22%) when compared with patients who did not develop delirium, even after adjusting for infection, age, and cognitive impairment.⁶¹ This is the first study to show a relationship between peripherally measured cytokine levels and delirium as a symptom of sickness behavior in acutely admitted elderly. It also showed that cognitive function can be impaired by a systemic infection in patients with a neurodegenerative disorder such as Alzheimer disease and that this cognitive decline is preceded by raised serum levels of IL-1h. Similarly, a recent study found that high preoperative neopterin levels predicted delirium after cardiac surgery in older adults, suggesting that plasma neopterin levels may be a candidate biomarker for delirium among this patient population.⁶⁴ Neopterin is produced by human monocytes/macrophages upon stimulation with the cytokine interferon- γ and thus can serve as a marker for immune system activation.⁶⁵


CNS resident cells react to the presence of peripheral immune signals, leading to production of cytokines and other mediators in the brain, cell proliferation, and activation of the hypothalamics–pituitary–adrenal (HPA) axis (see Neuroendocrine Hypothesis, below) through a complex system of interactions. These neuroinflammatory changes cause BBB permeability disruption (as suspected by elevations of S100 beta) and changes in synaptic transmission, neural excitability, and cerebral blood flow, leading to the neurobehavioral and cognitive symptoms characteristic of delirium (e.g., disruption in behavior and cognitive functions).⁴⁸ S100 beta is a calcium-binding protein with cytokine-like properties, secreted primarily by astrocytes under metabolic stress, and is considered a putative biomarker of CNS damage: Increased cerebrospinal fluid (CSF) and serum S100 beta are linked with adverse CNS outcomes. Several studies have reported findings of elevated S100 beta levels among patients suffering from either dementia^{66–68} or delirium.^{69–72} Furthermore, aging and neurodegenerative disorders exaggerate brain microglial responses after stimulation by systemic immune stimuli such as peripheral inflammation and/or infection (see Neuronal Aging Hypothesis, below).

During or after various disease processes or trauma/surgery, leukocytes adhere to endothelial cells, which make up the bulk of the blood–brain barrier and become activated, leading to degranulation and the release of free oxygen radicals and enzymes, which in turn leads to endothelial cell membrane destruction, disruption of cell–cell adhesions, and increased endothelial permeability.^{73,74} This in turn causes extravascular fluid shifts and the development of perivascular edema in cerebral tissue, which leads to decreased perfusion and longer diffusion distance for oxygen.^{48,75,76} These processes may lead to such extensive perfusion impairment that the blood flow in individual capillaries becomes disrupted; thus, systemic inflammation as a response to trauma or illness leads to microcirculatory impairment and subsequent ischemic injury (see Oxidative Stress Hypothesis, below). Among the pertinent neurotransmitters, Ach synthesis and release may be the most sensitive to this type of hypoxic injury and other homeostatic changes in the brain (see Neurotransmitter Hypothesis, below).⁷⁷ Similarly, neuroinflammatory injuries have also been associated with imbalances in other neurotransmitters including DA, serotonin (5HT), and norepinephrine (NE).⁷⁸

In response to traumatic and systemic events, the systemic inflammatory response is activated, causing monocytes and macrophages to produce neopterin, cytokines, and reactive oxygen species, which can be found in the plasma, urine, and CSF of delirious patients.^{1,48,63,64,79–82} In addition, disruptions of the endothelial cells (as described above) may also lead to enhanced cytokine transport across the disrupted BBB and infiltration of leukocytes and cytokines into the CNS, producing ischemia and neuronal apoptosis (Fig. 4).^{48,83,84}

Systemic inflammation is common in liver failure, and its acquisition is a predictor of hepatic encephalopathy (HE) severity. Studies provide convincing evidence for a role of neuroinflammation in liver failure; this evidence includes activation of microglia, together with increased synthesis in situ of proinflammatory cytokines (i.e., TNF, IL-1 β , and IL-6). The proposed “liver–brain signaling mechanisms” in liver failure include direct effects of systemic proinflammatory molecules, recruitment of monocytes after microglial activation, brain accumulation of ammonia, lactate and manganese, and altered

FIGURE 4. Effect of Inflammation on the development of delirium. Recognition and propagation of peripheral immune stimuli in the CNS. The initial interaction of circulating inflammatory mediators (e.g., cytokines and lipopolysaccharide) with the neurovascular unit occurs through a vast number of receptors and is associated with an increased paracellular permeability of the BBB. In addition to systemic inflammation, other factors affect the integrity of BBB including hypoxia, ischemia, and pain. Recognition of peripheral inflammatory stimuli in the BBB is followed by a cascade of events leading to microglia activation and subsequent modulation of adjacent cells including astrocytes and neurons (represented with dashed reciprocal arrows). (From Cerejeira et al.⁴⁸).



permeability of the BBB.^{85,86} This provides another intersection of the NIH with the Neurotransmitter Hypothesis (NTH) as the above changes may contribute to the alteration in neurotransmitter functioning in cases of HE (e.g., increased DA, 5HT, and γ -aminobutyric acid [GABA]).


There is also mounting evidence that some proinflammatory cytokines that induce sickness behavior also enhance activity of the ubiquitous indoleamine-2,3-dioxygenase (Fig. 5).^{87,88} Activation of indoleamine-2,3-dioxygenase results in decreased tryptophan (TRP) levels, thus a reduction in 5HT and melatonin production, whereas there is a shift to the production of kynurenine and other TRP-derived metabolites that have neurotoxic effects.^{89–91}

Systemic inflammation, such as that caused by injury (including surgery) or infection, has long been recognized as trigger for episodes of delirium, particularly in elderly or demented patients, even though their deliriogenic effect seems to be lessened in younger and nondemented patients.^{51,62,74,92–99}

Yet, even in younger and nondemented adult patients, the severity of the patient's injury or underlying medical problem is significantly directly correlated with the development of delirium.^{100,101}

These facts suggest that either a low dose of precipitant in a vulnerable patient or a high dose in the nonvulnerable may overwhelm the system and lead to delirium. Thus, it seems likely that more severe systemic inflammatory responses are more likely to induce delirium, but pre-existing pathology in cognitive circuitry is a stronger predictor; thus, the interaction between these two factors is key.¹⁰² The NIH intersects with the Neuronal Aging Hypothesis (NAH) because it has been shown that microglia, the major macrophage population of the brain, are primed by prior neurodegenerative pathology to respond more robustly to systemic inflammatory signals.^{53,103} There is also an interaction with the HPA axis (see Neuroendocrine Hypothesis, below) as hormones help coordinate an animal's physiology and behavior to match its environment and maximize

FIGURE 5. Metabolic Pattern of TRP, 5HT, melatonin, and kynurenic and quinolinic acid. Evidence suggests that some proinflammatory cytokines cannot only induce sickness behavior but also enhance activity of the ubiquitous indoleamine-2,3-dioxygenase, leading to deficient TRP levels, thus a reduction in 5HT and melatonin production, and a shift to the production of kynurenine and other neurotoxic TRP-derived metabolites. (Adapted from Stone et al.⁸⁷ & Darlington et al.⁸⁸).




survival; among the most important processes influenced by endocrine hormones are the immunologic and behavioral responses to infection.^{104,105}

NEURONAL AGING HYPOTHESIS

The NAH suggests that the aging process and accompanying physiologic changes constitute an independent risk factor for delirium. The concept of homeostenosis implies that a functional elderly person may maintain health into old age but may become increasingly vulnerable to stress and illness

because of a lack of physiologic reserve.¹⁰⁶ Accordingly, aging is associated with age-related cerebral changes in stress-regulating neurotransmitters, brain–blood flow decline, decreased vascular density, neuron loss (particularly in locus coeruleus and substantia nigra), and intracellular signal transduction systems.^{107–122} This likely explains why the aging process itself is associated with some degree of cognitive deficits and an increased risk of dementia (Table 1). The NAH may also explain why the elderly seem to experience a greater chance of developing delirium when challenged by physiologic distress that is better tolerated by younger individuals.

FIGURE 6. Age and transition to delirium. An estimation of the probability of transitioning to delirium by age indicates that the incremental risk is large for patients aged 65 years and older. The probability of transitioning to delirium increased dramatically (by 2%) for each year of life after 65 years. Adjusted odds ratio: 1.01 [(1.00, 1.02)], $p = 0.03$. (Pandharipande P et al.)¹⁰¹



Although the estimated annual incidence of Alzheimer disease in the population is 0.6% for persons aged 65 and older,¹²³ among elderly patients in the ICU, the probability of developing delirium increases by 2% per year of age for each year after age 65 (Fig. 6).¹⁰¹ Other studies confirm that older age is an independent risk factor among medically ill and surgical patients, with the increase in rate per year dependent on the specific context in which incidence is measured.^{124–128} The data also suggest a reciprocal relationship between delirium and cognitive decline: Dementia is the strongest risk factor for delirium among older patients,^{27,92,129,130} and the development of delirium appears to increase the risk of cognitive decline, including dementia.³²

It is well established that individuals with compromised cognitive ability preoperatively (e.g., dementia) are at greater risk of delirium.^{131–134} A study of elderly subjects undergoing orthopedic surgery demonstrated that the presence of dementia increased the incidence of postoperative delirium from 32% to 100%.³³ However, evidence suggests that decrements in higher-order cognitive functions, such as executive function (e.g., problem solving,

processing speed, planning, complex sequencing, and reasoning), may also predict postoperative delirium in the absence of frank cognitive impairment.^{135,136} A study of nondemented elderly patients undergoing elective orthopedic surgery demonstrated that subtle preoperative attention deficits, as tested by digit vigilance and reaction time testing, were closely associated with postoperative delirium.¹³⁷ In fact, these subtle changes predicted a four- to fivefold increased risk of postoperative delirium for subjects more than one standard deviation above the sample means on these variables.

Conversely, studies have demonstrated that among elderly surgical patients, delirium is a strong independent predictor of cognitive impairment and the occurrence of severe dependency in activities of daily living. In fact, 38 months after discharge from hospital, 53.8% of surviving patients with postoperative delirium continued to experience cognitive impairment, as compared with only 4.4% of nondelirious subjects.³⁵ Similarly, a prospective, matched, controlled cohort study of elderly hip surgery patients demonstrated that the risk of dementia or minimal mild cognitive impairment over a 30-month follow-up almost doubled in inpatients with postoperative delirium compared with those without delirium.³⁶

The data suggest that many older hospital patients do not recover from delirium and that persistence of delirium is associated with adverse outcomes. In some studies, the long-term outcomes (e.g., mortality, nursing home placement, cognition, function) of patients with persistent delirium were consistently worse than the outcomes of patients who had recovered from delirium.¹³⁸ These findings suggest that delirium does not simply persist for a certain time but also predicts a future cognitive decline with an increased risk of dementia. Table 2 provides a list of potential mechanisms associated with the increased risk of delirium in the elderly.

The observed increased levels of circulating inflammatory mediators (e.g., cytokines, acute phase proteins) suggest that chronic neurodegeneration is accompanied by an inflammatory response characterized by chronic, but selective, activation of CNS microglial cells that are “primed” to produce exaggerated inflammatory responses to immunologic challenges.^{48,53} Age-related changes in the immune system, known as immunosenescence, and increased secretion of cytokines by adipose tissue represent the

TABLE 2. Potential Mechanisms Associated with the Increased Risk of Delirium in the Elderly

- Neuronal loss particularly in locus coeruleus and substantia nigra^{110,111}
- Changes in various neurotransmitter systems¹¹¹
- Age-related decline in white matter integrity, observed as increases in water diffusion and volume of hyperintense white matter lesions, intergyral spans, and reduction in fractional anisotropy of water diffusion, correlated with a decline in the global and regional cerebral glucose uptake¹⁰⁹
- Age-related decline in regional cerebral blood flow, particularly in the anterior cingulate gyrus, bilateral basal ganglia, left prefrontal, left lateral frontal and left superior temporal and insular cortex, as measured by single-photon emission tomography¹¹⁹
- Age-related changes in cerebral blood flow, likely associated with brain microvascular pathologies:^{112–116,118}
 - Rarefaction of the microvasculature in some regions of the brain
 - Decreased vascular density
 - Damaged microvessels, with associated microinfarcts and microhemorrhages, likely due to peripheral arterial aging leading to stiffening and dilation of the proximal aorta with transmission of flow pulsations downstream into the brain
 - Decline in cerebrovascular angiogenesis
 - Impaired cerebral blood flow due to tortuous arterioles and deposition of excessive collagen in veins and venules
- Age-related decline in cerebral metabolic rate of oxygen more markedly in bilateral putamen, left supratemporal, left infrafrontal, and left parietal cortices¹²¹
- Decreased oxygen supply (e.g., hypoxia) leading to a decrease in redox activity, resulting in decreased ACh production^{158,194}
- Decreased cerebral oxidative metabolism¹⁵⁸
- Age-related changes in brain neurochemical activity:¹²²
 - There is a significant increase in soluble hexokinase activity with age, due to an increased release of mitochondrially bound hexokinase.
 - There is a negative correlation of the activity of fructose-6-phosphate kinase with age, particularly in brain cortex and putamen.
 - There is a significant decline of carbonic anhydrase (important in the regulation of the P_{O_2}/P_{CO_2} ratio in the brain tissue) with increasing age. Thus, P_{CO_2} -dependent regulation of tissue pH, ionic transport processes, and cerebral blood flow regulation have the tendency to become more and more unstable.
 - There is a progressive, age-dependent decline in cAMP-dependent activity, most significantly in brain cortex and thalamus, followed by hippocampus, amygdala, and globus pallidus.
- Age-related cerebral changes in stress-regulating neurotransmitter and intracellular signal transduction systems⁵³
- Decreased ACh levels in plasma and CSF^{181,185,187,192,195,196}
 - This is likely due to
 - Decreased volume of ACh-producing cells associated with normal aging (see Neurotransmitter Hypothesis)¹⁵⁸
 - Decreased ACh synthesis associated with aging¹⁵⁸
 - Increase in baseline levels of circulating inflammatory mediators including cytokines and acute phase proteins (see Neuroinflammatory Hypothesis)^{48,52,60,61,108,191}

major causes of chronic inflammation, a phenomenon known as “inflammaging.”¹³⁹ This inflammation may contribute to disease progression through the production of inflammatory mediators. The aging process is associated with a two- to fourfold increase

in baseline levels of circulating inflammatory mediators, including cytokines and acute phase proteins.^{52,60,61} Other factors may influence the medically ill elderly patient, such as lower cognitive reserves, lower metabolic capacity, increased sensitivity to medications, and lower threshold to a medication’s anticholinergic effects. In fact, even though high levels of postoperative pain and high opioid use was found to increase the risk for postoperative delirium in all elderly patients, the highest incidence of delirium occurred among patients who had high preoperative risk for delirium (e.g., lower baseline cognitive status, comorbid psychiatric symptoms).¹⁴⁰

Finally, there has been some controversy regarding the contribution of apolipoprotein E (APOE) to the development of postoperative cognitive dysfunction. Of the nine articles published to date on the topic, four found that *APOE-E4* genotype was not associated with postoperative cognitive dysfunction,^{141–144} whereas five suggested a positive relationship^{145–149} (Table 3). At least one study found a positive relationship between delirium with APOE genotype, interferon- γ , and insulin-like growth factor-I and that recovery was significantly associated with lack of APOE4 allele and higher initial interferon- γ .¹⁴⁵ The only published meta-analysis on the matter found a positive association between postoperative cognitive dysfunction and the APOE σ 4 allele.¹⁵⁰

OXIDATIVE STRESS HYPOTHESIS

Hypoperfusion appears to induce chronic oxidative damage in tissues and cells, largely due to the generation of reactive oxygen and reactive nitrogen species. Any condition that outpaces the capacity of endogenous redox systems to neutralize such toxic intermediates could lead to a system imbalance or to major compensatory adjustments that rebalance the system. This new redox state is generally referred to as “oxidative stress” (Fig. 7).¹⁵¹ The Oxidative Stress Hypothesis (OSH), initially proposed by Engel and Romano,⁴⁵ proposes that a number of physiologic processes, such as tissue damage, hypoxia, severe illness, and infections, may give rise to increased oxygen consumption and/or decreased oxygen availability, with associated increased energy expenditure and reduced cerebral oxidative metabolism, leading to cerebral

TABLE 3. Association Between APOE genotype and POCD

Study	Population	Findings	p
Abildstrom et al., 2004 ¹⁴¹ Prospective, N = 967	Patients aged 40 y, older undergoing noncardiac surgery	<ul style="list-style-type: none"> One week after surgery, the incidence of POCD was 11.7% in patients with the $\epsilon 4$ allele and 9.9% in patients without the $\epsilon 4$ allele. Conclusion: unable to show a significant association between APOE genotype and POCD. 	0.41
Adamis et al., 2007 ¹⁴⁵ Prospective, N = 164	≥ 70 y/o hospitalized medically ill patients	<ul style="list-style-type: none"> Recovery was significantly ($p < 0.05$) associated with lack of APOE$\epsilon 4$ allele and higher initial IFN-γ. A model incorporating gender, APOE $\epsilon 4$ status, and insulin-like growth factor-I levels predicted recovery or not from delirium in 76.5% of cases, with a sensitivity of 0.77 and specificity of 0.75. It further found a positive relationship between delirium with APOE genotype, IFN-γ, and insulin-like growth factor-I but not with IL-6, IL-1, TNF-α, and leukemia inhibitory factor. 	<0.05
Leung et al., 2007 ¹⁴⁷ Prospective, N = 190	Patients aged >65 y scheduled to undergo major non-cardiac surgery requiring anesthesia	<ul style="list-style-type: none"> The presence of one copy of the $\epsilon 4$ allele was associated with an increased risk of early postoperative delirium (28.3% vs. 11.1%; $p = 0.005$). Even after adjusting for covariates, patients with one copy of the $\epsilon 4$ allele were still more likely to have an increased risk of early postoperative delirium (OR: 3.64; 95% CI: 1.51–8.77) compared with those without the $\epsilon 4$ allele. 	0.005
Tagarakis et al., 2007 ¹⁴³	Elderly adults undergoing cardiac bypass surgery; excluded dementia	<ul style="list-style-type: none"> Study confirmed the high incidence of cognitive decline and delirium after coronary surgery but did not support the role of the APOE $\epsilon 4$ allele in the occurrence of delirium. 	NS
Van Munster et al., 2007 ¹⁴⁴ N = 415	Acutely admitted patients aged ≥ 65 y to the Department of Medicine	<ul style="list-style-type: none"> The OR for carriers of an APOE $\epsilon 4$ allele compared with patients without an APOE $\epsilon 4$ allele for developing delirium was 1.17 (95% CI: 0.49–2.78) in the cognitively intact patients and 0.42 (95% CI: 0.14–1.30) in the cognitively impaired patients. No relation existed between the total number of APOE $\epsilon 4$ alleles and the different delirium subtypes. 	0.12
Ely et al., 2007 ¹⁴⁶ N = 53	Mechanically ventilated intensive care unit patients	<ul style="list-style-type: none"> Using multivariable regression analysis to adjust for age, admission diagnosis of sepsis or acute respiratory distress syndrome or pneumonia, severity of illness, and duration of coma, the presence of APOE$\epsilon 4$ allele was the strongest predictor of delirium duration (OR: 7.32; 95% CI: 1.82–29.51). 	0.005
Zhang et al., 2008 ¹⁴⁹ N = 196	Elderly patients (>60 y/o) scheduled for major abdominal surgery requiring general anesthesia	<p>The presence of the $\epsilon 4$ allele and low level of education were both associated with an increased risk of EA (36.9% vs. 15.8%, $p = 0.005$; 30% vs. 14.3%, $p = 0.01$). After adjustment for covariates, the patients with the copy of $\epsilon 4$ allele were shown to have a greater likeliness of an</p>	0.01

(Continued)

TABLE 3. (Continued)

Study	Population	Findings	p
Van Munster et al., 2009 ¹⁴⁸ Meta-analysis, N = 656+	Medical Department and Orthopedic/ Traumatology Department of University Hospital from 2003 to 2007	increased risk of EA (OR: 4.32; 95% CI: 1.75–10.05). The OR for delirium adjusted for age, cognitive, and functional impairment of $\sigma 4$ carriers compared with non- $\sigma 4$ carriers was 1.7 (95% CI: 1.1–2.6). Four studies were added to the meta-analysis, which included 1,099 patients in total. The OR for delirium in the meta-analysis was 1.6 (95% CI: 0.9–2.7) of $\sigma 4$ carriers compared with non- $\sigma 4$ carriers. This study and meta-analysis suggest an association between delirium and the APOE $\sigma 4$ allele.	0.04
Bryson et al., 2011 ¹⁴² N = 88	Patients ≥ 60 y/o undergoing open aortic repair	Delirium predicted POCD at discharge (OR: 2.86; 95% CI: 0.99–8.27) but not at 3 months. APOE $\epsilon 4$ genotype was not associated with either delirium or POCD after adjustment for covariates.	0.625

Source: Maldonado²²²

Notes: POCD: postoperative cognitive dysfunction; IFN- γ : interferon- γ ; NS: nor significant; OR: odds ratio; CI: confidence interval.

dysfunction and associated cognitive and behavioral symptoms of delirium. In other words “[delirium is] the clinical expression of a cerebral metabolic defect.”^{45 (p. 531)}


Some have found that oxidative stress and/or antioxidant deficiencies increase damage to cerebral tissue and lead to cognitive decline with irreversible degeneration as sequelae of delirium.^{30,152–154} Among patients undergoing cardiopulmonary bypass surgery, researchers found alterations in the levels of markers of oxidative stress. In this sample, patients who developed delirium demonstrated significantly lower preoperative levels of the antioxidant enzyme catalase (normally produced by the body as the primary endogenous defense against free radical-induced injury) compared with non-postoperative delirium patients.

Also, clinical data exist that correlate poor oxygenation and cerebral dysfunction. A study of medically ill ICU patients demonstrated that oxidative metabolic stress was present within 48 hours before the onset of delirium, as evidenced by abnormalities in three measures of oxygenation (i.e., hemoglobin, hematocrit, and pulse oximetry) among patients who developed delirium.¹⁵⁵ In the same sample, clinical factors associated with greater oxidative stress (e.g., sepsis, pneumonia) occurred more frequently among delirious subjects. Similarly, studies have demonstrated a strong correlation

between intraoperative O₂ saturation and postoperative mental function.^{156,157} Even healthy control subjects may experience delirium after dropping their PaO₂ to 35 mm Hg.¹⁵⁸ In fact, intraoperative cerebral oxygen desaturation was found to be a significant risk factor for postoperative delirium among cardiac surgery patients.¹⁵⁹ In addition, operative cerebral oxygen saturation levels among patients undergoing abdominal surgery was an independent risk factor for postoperative delirium.¹⁶⁰ Similarly, among a group of patients undergoing cardiac surgery, those who developed postoperative delirium had lower pre- and intraoperative cerebral oxygen saturation levels, were older, and had lower preoperative hemoglobin levels compared with nonpostoperative delirium patients.¹⁶¹

Animal studies suggest that neuronal susceptibility to ischemic injury is not uniform. The basal ganglia, thalamus, Purkinje, layer 3 of the cortex, and the pyramidal neurons of the hippocampus are particularly vulnerable to hypoxia, but the degree of damage may vary depending on the cause.¹⁶² A superimposed global mild ischemic injury is often present in critically ill patients, galvanizing the oxidative failure. Indeed, patients in the critical care setting are particularly at risk to suffer the effects of hypoperfusion resulting from a number of potentially controllable extrinsic factors (e.g., intraoperative hypotension).

FIGURE 7. Mechanisms of brain injury after global cerebral ischemia. During cerebral ischemia, excess GLU exits into the extracellular compartment due to cellular depolarization, coupled with its impaired uptake, which results in increases in intracellular Ca^{2+} . The cascade of events responsible for GLU excitotoxicity includes three distinct processes: (1) induction, whereby extracellular GLU efflux is transduced by receptors on the neuronal membrane to cause intracellular Ca^{2+} overload, which leads to lethal intracellular derangements; (2) amplification of the derangement, with an increase in intensity and involvement of other neurons; and (3) expression of cell death triggered by cytotoxic cascades. Excess release of Ca^{2+} and its intracellular influx is thought to be the primary trigger for a variety of complex, deleterious intracellular processes that result from activation of catabolic enzymes such as phospholipases (which lead to cell membrane breakdown, arachidonic acid, and free radical formation) and endonucleases (which lead to fragmentation of genomic DNA and energy failure due to mitochondrial dysfunction). (From Harukuni & Bhardwaj³⁷⁹).



Inadequate oxidative metabolism may be one of the underlying causes of the basic metabolic problems initiating the cascade that leads to the development of delirium, namely the inability to maintain ionic gradients causing cortical spreading depression (i.e., spreading of a self-propagating wave of cellular depolarization in the cerebral cortex);^{163–168} abnormal neurotransmitter synthesis, metabolism, and release;^{169–177} and a failure to effectively eliminate neurotoxic byproducts.^{170,171,175}

The OSH intersects with the NTH because decreased oxygenation causes a failure in oxidative metabolism, which may be one cause of the problems observed in delirium, namely decreased oxygenation, which causes a failure in oxidative metabolism and leads to a failure of the ATPase pump system.¹⁷⁹ When the pump fails, the ionic gradients cannot be maintained, leading to significant influxes of sodium

(Na^+) followed by calcium (Ca^{2+}), whereas potassium (K^+) moves out of the cell.^{178,179} Some have theorized that it is the excess inward flux of Ca^{2+} that precipitates the most significant neurobehavioral disturbances observed in delirious patients.^{232,233} The influx of Ca^{2+} during hypoxic conditions is associated with the dramatic release of several neurotransmitters, particularly glutamate (GLU) and DA.^{178,179} GLU further potentiates its own release as GLU stimulates the influx of Ca^{2+} .^{233,234} and accumulates in the extracellular space as its reuptake and metabolism in glial cells is impeded by the ATPase pump failure.¹⁷⁹ In addition, at least two factors facilitate dramatic increases in DA: First, the conversion of DA to NE, which is oxygen dependent, is significantly decreased, and second, the catechol-o-methyl transferase enzymes, required for degradation of DA, get inhibited by toxic metabolites under

hypoxic conditions, leading to even greater accumulation of DA.²¹⁶ At the same time, 5HT levels fall moderately in the cortex, increase in the striatum, and remain stable in the brainstem.²³⁵ Hypoxia also leads to a reduced synthesis and release of ACh, especially in the basal forebrain cholinergic centers.²³⁶ Indeed, cholinergic neurotransmission is particularly sensitive to metabolic insults, such as diminished availability of glucose and oxygen.²³⁷ ACh synthesis requires acetyl coenzyme A, which is a key intermediate linking the glycolytic pathway and the citric acid cycle. Thus, reduction in cerebral oxygen and glucose supply and deficiencies in enzyme cofactors such as thiamine may induce delirium by impairing ACh production.^{77,238} These changes have been extensively reviewed and discussed elsewhere (Fig. 8).¹

NEUROTRANSMITTER HYPOTHESIS

The NTH was proposed after clinical observations that delirium occurred after the use of substances (e.g., medications, toxins) that alter neurotransmitter function or availability; it was originally used to explain delirium secondary to impaired cholinergic function.^{78,180–185} Some have postulated that delirium may be seen as a temporary psychiatric disorder resulting from a reduced central cholinergic transmission, combined with an increased dopaminergic transmission.^{108,186,187} The NTH stresses the fact that the cholinergic and dopaminergic systems interact not only with each other but with glutamatergic and GABA pathways.¹⁸⁸ In fact, some pharmacologic agents (e.g., opioids) may cause delirium by increasing DA and GLU activity while decreasing ACh availability.¹⁸⁹ In general, the most commonly described neurotransmitter changes associated with delirium are reduced availability of ACh (↓ACh); excess release of DA (↑DA), NE (↑NE), and/or GLU (↑GLU); and alterations (e.g., both a decreased and increased activity depending on circumstances and etiological factors) in 5HT (↓↑5HT), histamine (↓↑H1 and H2), and/or GABA (↓↑GABA) (Table 1), as previously reviewed by others (Table 4).^{1,4,108}


The cholinergic system is one of the most important modulatory neurotransmitter systems in the brain, controlling activities that depend on selective attention, which are an essential component of conscious

awareness¹⁹⁰ (the two key components in *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition*, criteria for diagnosing delirium). Adequate ACh levels are also essential for the regulation of rapid-eye-movement sleep, memory, and synchronization of the electroencephalogram, among others. One of the leading hypotheses is that delirium results from an impairment of central cholinergic transmission^{1,181,191–193} and is considered by some to be “a common denominator” in delirium (or toxic–metabolic encephalopathies).¹⁹⁴ Studies have demonstrated low levels of ACh in plasma and CSF among delirious patients.^{181,185,187,192,195,196} Many have demonstrated a relationship between a drug’s anticholinergic potential and its deliriogenic effects,^{181,184,197–202} thus raising awareness of the potentially significant additive effect of medications commonly thought to have low ACh activity.^{185,203,204}

Similarly, high levels for serum anticholinergic activity¹⁹⁸ have been associated with an increased likelihood of delirium among surgical²⁰⁵ and medical²⁰⁶ inpatients. In fact, serum anticholinergic activity levels may predict delirium,¹⁸¹ and resolving delirium has been correlated to decreasing serum anticholinergic activity levels.²⁰⁷ Of interest, some have found detectable serum anticholinergic activity levels in delirious patients not exposed to known anticholinergic agents, suggesting that endogenous anticholinergic substances may be produced during acute illness and could be implicated in the etiology of delirium.^{195,208} In a group of hospitalized elderly patients, the Anticholinergic Risk Scale was able to predict the all-cause mortality when factored in along with baseline cognitive impairment, in-hospital delirium, place of residence (e.g., home versus nursing home), and length of hospital stay.²⁰⁹

Animal studies have revealed impairment in cholinergic neurotransmission in several models of encephalopathy and delirium, including hypoxia, nitrite poisoning, thiamine deficiency, hepatic failure, carbon monoxide poisoning, anesthesia, selective intracranial atropine injection, physical immobilization, and hypoglycemia.^{186,210,211} Finally, potential animal models mimicking the decreased mobility of critically ill patients have demonstrated that immobilization may cause widespread reduction in ACh levels.^{210,211} Here the NTH intersects with the NAH because aging itself is associated with age-related cerebral changes in stress-regulating hormones and

FIGURE 8. Oxidative stress and neurotransmitter theories of delirium. A basic pathoetiologic model of delirium illustrates the theorized intersections between the OSH and the NTH of the delirium, demonstrating potential common biochemical outcomes, which may explain the complex cognitive and behavioral changes characteristic of delirium. The OSH proposes that a number of physiologic processes (e.g., hypoxia, severe illness, infectious processes) may give rise to increased oxygen consumption and/or decreased oxygen availability, with associated increased energy expenditure and reduced cerebral oxidative metabolism, leading to cerebral dysfunction and associated cognitive and behavioral symptoms of delirium. The NTH was proposed after clinical observations that delirium occurred after the use of substances (e.g., medications, toxins) that alter neurotransmitter function or availability. The OSH intersects with the NTH because decreased oxygenation causes a failure in oxidative metabolism, leading to a failure of the ATPase pump system, which leads to an inability to maintain adequate ionic gradients, which in turn leads to significant electrolyte alterations (e.g., influx of Na^+ and Ca^{2+} ; efflux of K^+) and subsequent alterations (e.g., excess release or decreased availability) of several neurotransmitters (e.g., GLU, DA, Ach). (From Maldonado¹).



intracellular signal transduction systems. Aging is also associated with a decreased volume of ACh producing cells and decreased cerebral oxidative metabolism (see Oxidative Stress Hypothesis, above). Similarly, hypoxia is known to reduce the synthesis and release of ACh.²¹² In addition, delirium is associated not only with an unbalanced inflammatory response but also with a dysfunctional interaction between the cholinergic and immune systems (see Neuroinflammatory Hypothesis, above).²¹³ In fact, acute systemic inflammation is a major trigger for

cholinergic hypoactivity and is thought to be important in cognitive dysfunction during delirium.²¹⁴

Elevations of DA have long been suspected in the development of delirium.^{187,188,191} For example, studies have confirmed elevation of DA's metabolites (i.e., homovanillic acid) in the CSF of patients with fulminant HE.²¹⁵ At least two factors facilitate dramatic increases in DA. First, the conversion of DA to NE, which is oxygen dependent, is significantly decreased (allowing DA to accumulate). Second, the catechol-o-methyl transferase enzymes, required for

TABLE 4. Theorized Neurochemical Mechanisms Associated with Conditions Leading to Delirium

Delirium Source	ACH	DA	GLU	GABA	5HT	NE	Trp	Phe	His	Cytok	HPA axis	NMDA activity	Changes in RBF	EEG	Mel	Inflam	Cort
Anoxia/hypoxia	↓	↑	↑	↑	↓	↓	↔	↑	↑,↓	↑↑	↑	↑	↑	↓	↓	↑	↑
Aging	↓	↓	↓	↓	↓	↓	↓	↓	↓	↑↑	↑	↓	↑	↓	↓	↑	↑
TBI	↑	↑	↑	↑	↑	↑	↑	↑	↓	↑↑	↑	↑	↑	↓	↓	↑↑	↑
CVA	↓	↑	↑	↑	↑	↑	↑	↑	↓	↑↑	↑	↑	↑	↓	↓	↑↑	↑
Hepatic Failure (encephalopathy)	↔	↓	↑	↑	↑	↓	↑	↑	↑	↑↑	↑	↑	↑	↓	↓	↑	↑
Sleep deprivation	↓	↓	↑	↑	↑	↑	↓	↑	↑	↑	↑	↑	↑	↓	↓↑	↑↑	↑
Trauma, Sx, & Post-op	↓	↑	↑	↑	↓	↑	↓	↑	↑	↑	↑	↑	↑	↓	↓	↑	↑
ETOH & CNS-Dep Withdrawal	↑	↑	↑	↓	↑	↑	↓	↑	↑	↑	↑↑	↑	↓	↑	↓	↑	↑
Infection/Sepsis	↓	↓	↑	↑	↓	↓	↓	↓	↓	↑	↑↑	↑↑	↑	↓	↓	↑	↑
Dehydration & Electrolyte Imbalance	↔	↑	↑	↑	↓	↑	?	?	↑	↑	↑	↑	↓	↑	↓	↑↑	↑
Medical Illness	↓	↑	↑	↑	↓	↑	↓	↑	↑	↑	↓	↑	↑	↑	↓	↑	↑

Notes: ↑ = likely to be increased or activated; ↓ = likely to be decreased or slowed; ↔ = no significant changes; (↑↑) = likely a contributor, exact mechanism is unclear; (–) = likely not to be a contributing factor; CVA = cerebro-vascular accident; Sx = surgery; ETOH = alcohol; CNS-Dep = central nervous system depressant agent; ACH = acetylcholine; DA = dopamine; GLU = glutamate; GABA = gamma-aminobutyric acid; 5HT = 5-hydroxytryptamine or serotonin; NE = norepinephrine; Trp = tryptophan; Phe = phenylalanine; His = histamine; Cytok = cytokines; HPA axis = hypothalamic-pituitary-adrenocortical axis; NMDA = N-methyl-D-aspartic acid; RBF = regional blood flow; EEG = electroencephalograph; Mel = melatonin; Inflam = inflammation; Cort = Cortisol. Source: Adapted from Maldonado, J. R. (2008). "Pathoetiologi cal model of delirium: a comprehensive understanding of the neurobiology of delirium and an evidence-based approach to prevention and treatment." *Crit Care Clin* 24(4): 789-856.

degradation of DA, get inhibited by toxic metabolites under hypoxic conditions, leading to even more amassment of DA.²¹⁶

In delirious patients, several metabolic pathways lead to significant increases in DA under impaired oxidative conditions (reviewed elsewhere¹). One such pathway is when significant amounts of DA are released and there is a failure of adequate DA reuptake. A second pathway is when the influx of Ca^{2+} stimulates the activity of tyrosine hydroxylase,²¹⁷ which converts tyrosine to 3,4-dihydroxyphenylalanine and leads to increased DA production and further uncouples oxidative phosphorylation in brain mitochondria.¹⁷⁸ The outcome is a disruption of adenosine triphosphate production and the increased production of toxic metabolites of DA (formed under hypoxic conditions) that inhibit the activity of the oxygen-dependent catechol-O-methyl transferase,^{212,216} the major extracellular deactivator of DA, further leading to high levels of DA. Finally, an increase in the firing rates of catecholamine neurons may further induce tyrosine hydroxylase synthesis, which leads to even more DA production.²¹⁸

Elevation in DA availability may lead to some of the neurobehavioral alterations observed in delirious patients, via DA's direct effects and by potentiating GLU's excitotoxic effects.^{1,187,216} DA may exert its delirigenic effect by one of three mechanisms: (1) the direct excitatory activity of DA (e.g., toxicity with substances known to increase DA release or availability, such as amphetamines, cocaine, and DA); (2) DA enhances GLU-mediated injury;²¹⁶ and (3) excess DA may itself induce apoptosis by mechanisms independent of oxidative stress,²¹⁹ which may explain why DA depletion by α -methyl-paratyrosine may have a neuroprotective effect against hypoxic stress and injury²²⁰ and why DA blockade can be used to reduce hypoxic damage in the hippocampus.²²¹

It is also important to note the growing body of evidence demonstrating that antipsychotic agents are effective not only as treatment of delirium,^{222–227} because a recent meta-analysis demonstrated that antipsychotic agents are one of the few pharmacologic agents demonstrated to prevent delirium.²²⁸ A new meta-analysis revealed no significant differences in efficacy among agents.²²⁹ Furthermore, the authors concluded that the available evidence does not indicate major differences in response rates between clinical subtypes of delirium, suggesting the potential

efficacy of antipsychotic agents in the management and prevention of hypoactive delirium as well as the agitated and mixed types. Thus, it is possible that antipsychotic agents are not only effective in the symptomatic management of the symptoms of delirium, but they also address the underlying massive DA surge associated with some forms of delirium, even the hypoactive type.^{4,222}

GLU is the brain's principal excitatory neurotransmitter, yet excessive activation of *N*-methyl-D-aspartate (NMDA) receptors may lead to neuronal degeneration and cell death.^{230,231} In at least one study of high-risk adults undergoing cardiac surgery, serum concentrations of NMDA, as measured by serum concentrations of NMDA receptor antibodies (NR2Ab), were predictive of severe neurologic adverse events (e.g., delirium, transient ischemic attack, or stroke) postoperatively.

As in the case of DA, increased GLU availability may be due to the massive Ca^{2+} influx described above, associated with a number of physiologic conditions (e.g., hypoxia, hepatic failure), leading to GLU release. Excess GLU further stimulates Ca^{2+} influx, which releases even more GLU.^{232–234} Normally, GLU is released into the synapse and then removed by astrocytes and converted into glutamine, ending its action. Under oxidative conditions, however, GLU accumulates in the extracellular space as its reuptake and metabolism in glial cells is impeded by the ATPase pump failure.¹⁷⁹ It appears that GLU requires the presence of DA to exert some of its toxic effects, namely Ca^{2+} -induced neuronal injury.¹⁷¹ At high levels, DA may cause enough depolarization of neurons to activate the voltage-dependent NMDA receptor, therefore facilitating GLU's neurotoxic effects.²³⁹

It has been known for a while that high ammonia levels are a factor in the pathogenesis of delirium and HE in cirrhotic patients, but it may not be that well known that acute ammonia toxicity is mediated by the activation of NMDA receptors.²⁴⁰ In fact, some have suggested that drug-induced delirium would result from such transient thalamic dysfunction caused by exposure to medications that interfere with central glutamatergic, GABAergic, dopaminergic, and cholinergic pathways at critical sites of action.¹⁸⁸ Furthermore, GLU is metabolized by GLU-decarboxylase into GABA, which itself has been implicated in the development of the delirium.²⁴¹

GABA is the chief inhibitory neurotransmitter in the human CNS and plays a role in regulating neuronal excitability throughout and the regulation of muscle tone. Evidence suggests that GABA activity is increased in some types of delirium but decreased in others. For example, the “increased GABAergic tone” theory of HE, a neuropsychiatric disorder associated with liver failure, proposes the role of increased GABAergic neurotransmission in HE.^{242–245} Some of the supporting evidence comes from clinical experience demonstrating that flumazenil, a highly selective benzodiazepine antagonist at the GABA receptor complex, improved electroencephalographic activity, reversed coma, and improved symptoms of hypoaffective delirium in cirrhotic patients and in some HE patients.^{242,246} In addition, neurosteroids, synthesized in the brain mainly by astrocytes independent of peripheral steroidal sources (i.e., adrenals and gonads), are potent positive allosteric modulators of the GABA_A receptor. As such, neurosteroids stimulate inhibitory neurotransmission in the CNS by increasing GABAergic tone, a suggested pathophysiologic mechanisms in HE.²⁴⁷

Conversely, decreased GABAergic activity has been described in deliria caused by ethanol or CNS-depressant withdrawal²⁴⁸ and antibiotic-induced delirium.²⁴⁹ There is also mounting evidence that some GABAergic substances (e.g., benzodiazepines) may themselves induce delirium due to a variety of mechanisms, including^{101,188,250–252} (1) by interfering with physiologic sleep patterns,²⁵³ (2) by interrupting central cholinergic muscarinic transmission at the level of the basal forebrain and hippocampus,^{254–258} (3) by increasing compensatory up-regulation of NMDA and kainite receptors and Ca²⁺ channels,²⁵⁹ (4) by disrupting thalamic gating function,¹⁸⁸ (5) by causing withdrawal states upon their cessation, and (6) by disrupting the circadian rhythm of melatonin release.²⁶⁰

Acute NE release secondary to hypoxia or ischemia leads to further neuronal injury and the development or worsening of delirium.²⁶¹ Specifically, in cases of alcohol withdrawal, excess noradrenergic activity drives most of the symptoms (e.g., diaphoresis, tachycardia, increased blood pressure, restlessness, anxiety, agitation, tremors). Under normal circumstances the α_2 -receptor inhibits the firing of presynaptic NE neurons. Yet, evidence suggests that during alcohol withdrawal, signaling at the α_2 -receptor may be less sensitive, resulting in an inability of the

noradrenergic system to regulate its firing.^{262,263} In addition, alcohol withdrawal causes an up-regulation of GLU transmission in the locus coeruleus (the major noradrenergic nucleus of the brain), increasing the activity of the noradrenergic system.²⁶⁴ This may contribute to the autonomic instability, behavioral agitation, and psychosis seen during alcohol withdrawal delirium.²⁶⁵ Furthermore, catecholamines can enhance the activity of the bed nucleus of the stria terminalis neurons that may in turn increase the excitability of glutamatergic bed nucleus of the stria terminalis neurons that project to the ventral tegmental area.²⁶⁶

Randomized clinical trials have demonstrated that selective α_2 agonist agents (e.g., dexmedetomidine) substantially decreased the incidence of postoperative delirium compared with GABAergic agents.^{4,14,267,268} Similarly, α_2 agonist agents have shown neuroprotective qualities by suppressing circulating catecholamine levels during cerebral ischemia.²⁶⁹

Alterations in 5-HT activity, both elevated and decreased, have been linked to delirium development in various clinical populations. Normal 5HT synthesis and release in the brain is dependent on the availability of its precursor TRP (Fig. 5). Reduced 5HT levels have been identified in patients suffering from hypoxia, infections and sepsis, alcohol withdrawal delirium, delirium associated with levodopamine use for the treatment of Parkinson disease, immobility, catabolic states, and postoperative delirium, among others.^{191,270} Others have found an association between low 5HT levels associated with hyperactive delirium.²⁷¹ In fact, the sudden discontinuation of 5HT reuptake inhibitors has been associated with various neuropsychiatric syndromes, including delirium.²⁷²

Decreased TRP availability may lead to a reduction in 5HT.^{270,273,274} All large neutral amino acids (i.e., phenylalanine [PHE], TRP, eucine, isoleucine, methionine, tyrosine, valine) compete to enter the brain through the same saturable carrier. Therefore, as the concentration of one increases, CNS entry of other LNAAs conversely decreases.²⁷⁵ PHE has the additional interesting property of conversion to neurotoxic metabolites and competes with TRP for entry into the brain.²⁷⁶ Once it enters the brain, PHE competes with TRP and tyrosine for metabolism, via hydroxylation.²⁷⁰ Studies have demonstrated that elevations of the PHE/LNAA ratio are independently

associated with postoperative delirium.^{274,277,278} Studies of elderly medically ill patients suggest that an elevated plasma PHE/LNAA ratio during acute febrile illness is associated with delirium.^{191,277} Similarly, patients with hepatic²⁷⁹ and septic²⁸⁰ encephalopathy have also been found to experience increased levels of PHE and PHE metabolites in the plasma and CSF. Studies have found that an increased ratio of free-to-bound TRP enhances its availability to brain tissue, which in turn increases 5HT synthesis, thus precipitating HE.²⁸¹

Conversely, elevated 5HT levels have been described among patients suffering from 5HT syndrome, HE, and clozapine-induced delirium.^{191,215,282,283} At least two reports suggest that selective 5HT₃-type receptor antagonists may be effective in the treatment of agitated postoperative delirium.^{284,285} Similarly, hepatic dysfunction may lead to decreased metabolism of precursor amino acids (i.e., PHE, tyrosine), which may lead to increases in TRP availability, which leads to increases in 5HT. In fact, elevation in 5-hydroxyindoleacetic acid levels has been associated with HE and in patients suffering from hypoactive delirium.⁴⁶ Figure 5 shows the relationship between TRP, 5HT, and kynurenic acid metabolism.

Histamine receptors A₁ and A₂ are known to affect the polarity of cortical and hippocampal neurons²⁸⁶ and that both increased and decreased histamine levels may lead to delirium. It is well known that pharmacologic antagonism of either receptor is sufficient to cause delirium.²⁸⁷ Others have suggested that during surgical stress and hypoxia, there may be an excessive release of histamine, which may lead to delirium.²⁸⁸

A detailed review of the studies supporting the various neurotransmitter findings have been published elsewhere and are not repeated here.¹ The NTH intersects with the OSH because the abnormalities in neurotransmitter concentration or receptor sensitivity may underlie the different symptoms and clinical presentations of delirium that are caused by a decreased in cerebral oxidative metabolism.

NEUROENDOCRINE HYPOTHESIS

The Neuroendocrine Hypothesis suggests that delirium represents a reaction to acute stress, mediated by abnormally high glucocorticoid (GC) levels,


which induce a general vulnerability in brain neurons by impairing the ability of neurons to survive after various metabolic insults.^{289,290} GCs are steroid hormones that modulate metabolism, salt balance, development, reproductive processes, and immune function.¹⁰⁴ Although acute elevations of GCs enhance some immune functions, such as leukocyte infiltration at the sites of injury, chronic elevations induce leukocyte apoptosis, reduce proinflammatory cytokine release, and generally suppress immune activity.

Stress activates the HPA axis: Stressors (through inputs from the brainstem nuclei and the amygdala) activate the paraventricular nucleus of the hypothalamus resulting in the release of corticotrophin-releasing hormone, which (through the hypophyseal portal system) acts on the pituitary gland, inducing the release of adrenocorticotrophic hormone, which promotes GC (including cortisol) release from the adrenal cortex. Under normal circumstances, GCs act to aid the body in coping with the demands imposed by stress exposure, mobilizing energy stores, and suppressing nonvital body functions (e.g., inflammatory responses and reproduction) (Fig. 9).^{81,291–293}

However, plenty of scientific evidence demonstrates that GCs, the adrenal steroids secreted during stress, can have a broad range of deleterious effects in the brain.²⁹² Recent data demonstrated that repeated or prolonged exposure to GCs has a negative impact on brain function and provide evidence suggesting such exposure may contribute to age-related cognitive decline.^{81,294} Growing evidence suggests that GCs may have proinflammatory effects in the brain and can enhance neuroinflammation at multiple levels in the pathway that link lipopolysaccharide exposure to inflammation.²⁹⁵ In fact, GCs have been shown to enhance ischemic and seizure-induced neuronal injury.²⁹⁶ A number of mechanisms have been postulated to explain how excess GC release can compromise the neuron's ability to survive various neurologic insults (e.g., seizures, ischemia), which may lead to or exacerbate cell death (Table 5).

The above may explain how GCs contribute to the pathogenesis of delirium, especially in the elderly.^{292,294,297} The hippocampus, derived from medial regions of the telencephalon, is part of the limbic system and plays important roles in information encoding, related to short-term and long-term memory, and spatial navigation.²⁹⁸ The hippocampus also

FIGURE 9. Neuroimmune circuits in delirium. The HPA axis and inflammation. Various stressors can activate the HPA axis. The hypothalamus is stimulated to secrete corticotrophin-releasing hormone (CRH), which leads to adrenocorticotrophic hormone (ACTH) secretion into the peripheral circulation. ACTH in turn triggers adrenal GC release and production. The CRH system is inhibited by GCs in a negative feedback loop. Tumor necrosis factor (TNF)- α and IL-1 are produced from inflammatory sites and are potent activators of the HPA axis. IL-6 acts synergistically with GCs to stimulate the hepatic secretion of acute phase proteins. Although GCs are widely known for their anti-inflammatory actions, “(–),” more recently proinflammatory effects have also repeatedly been reported, “(+)” (+): enhancing; (–): suppressing. (From Dinkel et al.²⁹¹).



happens to contain the highest concentration of GC receptors of any brain region and thus may be a major target for the negative effects of excessive GC levels. Current evidence suggests that hippocampal malfunction occurs relatively early during the metabolic stress environment, leading to delirium,^{102,290,291,293,299–308} and that the highly catabolic GC's induce a general metabolic vulnerability in these neurons and thus compromise their ability to survive various toxic insults.³⁰⁹ Hippocampal injury may explain some of the attentional deficits and memory

TABLE 5. Potential Mechanisms to Explain How Excess GC Release Can Compromise Neuron's Ability to Survive Neurologic Insults that May Lead to or Exacerbate Cell Death

- Inhibiting glucose transport into neuron, thus inducing metabolic vulnerability^{292,301}
- Increase proinflammatory cell migration, cytokine production, and even transcription factor activity in the brain⁸¹
- Amplifying the damaging cascade of GLU excess, calcium (Ca^{2+}) mobilization, and oxygen radical generation^{304,306,309,380,381}
- Inducing spine loss and dendritic atrophy, thus decreasing neuroplasticity³⁸²
- Enhancing oxygen radical-mediated neurotoxicity³⁸³
- Exacerbating the toxicity of other neurotoxins (e.g., adriamycin) whose mechanisms of action overlap GC pathways³⁸⁴
- Impairing long-term potentiation³⁰³
- Reducing hippocampal glial cell activation and proliferation³⁸⁵
- Altering the expression and signaling of neurotrophins, particularly brain-derived neurotrophic factor^{302,386}
- Exacerbating the breakdown of cytoskeletal proteins (i.e., tau)³⁸⁷
- Impairing neurogenesis³⁸⁸

dysfunction and errors in information processing (leading to confabulation) commonly seen in delirious patients.

The hippocampal formation is of prime importance for normal HPA axis shut-off, and a key abnormality related to GCs excess in delirium seems to be an abnormal “shut-off” of the HPA axis as tested by the dexamethasone suppression test.³¹⁰ The loss of normal inhibition of adrenal steroidogenesis results in continuous secretion of peak amounts of corticosteroids, which causes resistance to cortisol feedback inhibition mediated by receptor loss in the hippocampus.^{292,304,309} The increased GC availability associated with illness and trauma (e.g., burns, surgery) or exogenous steroid administration further disrupt hippocampal function, which in turn may further disinhibit the release of GCs, thus sustaining high levels of circulating cortisol.⁴⁶ Studies have found that patients experiencing postoperative delirium had an impaired stress regulating system with significantly elevated mean plasma cortisol levels compared with the preoperative baseline and nondelirious patients.³¹¹ Among patients with delirium triggered by lower respiratory tract infections, 78% were found by the dexamethasone suppression test to be nonsuppressors compared with 14% of those with normal suppression responses.²⁹⁰ Demented patients with delirium exhibited significant differences in basal cortisol levels compared with demented, nondelirious patients.³¹² Furthermore, there was a strong linear

relationship between delirium and dexamethasone suppression test pathology irrespective of age and severity of dementia: The greater the intensity of delirium, the greater the level of nonsuppression.³¹² Similarly, patients with poststroke delirium exhibited significantly greater activation of the HPA system compared with those without.^{310,313}

DIURNAL DYSREGULATION OR MELATONIN DYSREGULATION HYPOTHESIS

This hypothesis suggests that disruptions to the 24-hour circadian cycle and the usual stages of sleep may lead to disturbances in the integrity of sleep and the physiologic sleep architecture.^{1,222} Sleep deprivation has long been linked to the development of delirium³¹⁴ and psychosis.³¹⁵ Hospitalized patients experience severe alterations of the sleep–wake cycle with sleep loss, sleep fragmentation, and sleep–wake cycle disorganization. The 24-hour internal clock (circadian pattern) is maintained by environmental factors, primarily light exposure, which affects melatonin secretion, and its disruption may lead to the development of delirium.³¹⁶

Others have demonstrated that sleep deprivation may lead to significant memory deficits³¹⁷ and to symptoms of emotional imbalance, likely due to disconnection between the amygdala and the prefrontal cortex.³¹⁸ Similarly, “chronic partial sleep deprivation” (i.e., sleeping limited to 4 hours per night for 5 consecutive nights) may lead to cumulative impairment in attention, critical thinking, reaction time, and recall.³¹⁹ Thus, cumulative sleep debt can cause delirium in itself and can aggravate or perpetuate delirium and its associated cognitive deficits.^{320–322} Studies have found sleep deprivation to consistently precede onset of delirium in post-surgical cardiac patients^{323,324} and that ICU patients with sleep deprivation were significantly more likely to develop delirium than patients without sleep deprivation.^{260,325}

Melatonin plays important roles in multiple bodily functions that may have potential implications regarding the development of delirium in the medically ill: Melatonin has significant chronobiotic effect (i.e., affecting aspects of biologic time structure), has

sleep–wake cycle regulatory effects, helps reset circadian rhythm disturbances, is an effective free radical scavenger with extensive antioxidant activity (particularly nuclear and mitochondrial DNA) with strong antiapoptotic signaling function, has extensive anti-inflammatory activity, and possesses some antinociceptive and analgesic effects.^{326–328} In addition, melatonin reduces the affinity of GCs receptors, prevents GCs inhibition of cell proliferation, and reduces the GC-induced neurotoxicity and apoptosis.³²⁹ These qualities may protect natural mechanisms of learning and memory.^{317,326–328,330–336} Melatonin also inhibits the aggregation of the amyloid beta protein into neurotoxic microaggregates responsible for the neurofibrillary tangles characteristic of Alzheimer disease and prevents the hyperphosphorylation of the tau protein.^{333,337–343} Data suggest that melatonin may have potential implications regarding the development of delirium in the medically ill and postoperative patient. Conversely, it has been theorized that the natural, age-related decline in brain melatonin may contribute to a pro-amyloidogenic microenvironment in the aging brain.³³⁷

Current evidence suggests that acute and chronic sleep deprivation is associated with decreased proportions of natural killer cells,³⁴⁴ reduced lymphokine-activated killer activity,³⁴⁵ and reduced IL-2 production.³⁴⁵ Conversely, cytokines may play a role in normal sleep regulation by increasing non-rapid-eye-movement sleep and decreasing rapid-eye-movement sleep, and during inflammatory events, an increase in cytokine levels may intensify their impact on sleep regulation.³⁴⁶ Moreover, sleep deprivation may alter endocrine and metabolic functions, altering the normal pattern of cortisol release and contributing to alterations of GC feedback regulation, glucose tolerance, and insulin resistance.³⁴⁷ We previously described the impact of disruption of GC regulation and delirium and cognitive function (see Neuroendocrine Hypothesis, above).

Studies have demonstrated a relationship between an irregular melatonin circadian rhythm (i.e., abnormally low serum levels of melatonin) and postoperative delirium.^{260,348} The administration of melatonin in ICU patients has been shown to improve quality of sleep and prolongation of sleep time,³⁴⁹ whereas others demonstrated that the

prophylactic administration of low-dose exogenous melatonin may decrease the incidence of delirium.^{350,351} Finally, data suggest that the use of prophylactic melatonin decreases the incidence and severity of “sundowning” and agitated behavior in elderly, demented individuals.³³²

Some have observed a relationship between the motoric delirium subtype and melatonin levels. A study of hospitalized elderly medical patients were evaluated daily using the Dementia Rating Scale and urinary measures of 6-sulphatoxymelatonin, the chief metabolite of melatonin.³⁵² A study found that during periods of hyperactive delirium, subjects had decreased urinary 6-sulphatoxymelatonin levels, whereas patients with hypoactive delirium had raised 6-sulphatoxymelatonin levels.³⁵³

NETWORK DISCONNECTIVITY HYPOTHESIS

The Network Disconnectivity Hypothesis (NDH) suggests that the heterogeneity of delirium presentations is better explained by the action of various factors (e.g., drugs or toxins) acting on specific brain neurochemical systems.³⁵⁴ Thus, this hypothesis postulates that factors affecting different neurotransmitter-specific projections, whether due to aging (e.g., degeneration), disease (e.g., inflammation), or pharmacologic agents (e.g., anticholinergic substances), will lead to different types of delirium presentations (e.g., hyperactive or hypoactive) and that, depending on the degree of reversibility of those changes, the delirium episode would have short or long-term sequelae (or effects). Initially, it immediately recognized two systems as potential culprits: the cholinergic and the GABAergic systems.

The NDH highlights the role of the cortical cholinergic system and associated projections in mediating specific attentional processing (i.e., sustained, selective, and divided attention performance), arousal, and rapid-eye-movement sleep—associated dreaming^{78,355} and that the selectivity of the behavioral effects of cortical ACh is based on close temporal interactions with converging sensory or associational cortical inputs.³⁵⁶ In fact, increased cholinergic activity in prefrontal regions is hypothesized to contribute to the activation of the anterior attention system and associated executive functions.³⁵⁵ Finally, the NDH

suggests that the failure (e.g., hypo- or hyperactivity) of one system will undoubtedly affect others (i.e., the connectivity principle).

The NDH recognizes that the brain is a highly organized and interconnected structure functioning to allow complex integration of sensory information and motor responses and suggests that delirium represents a variable failure in the integration and appropriate processing of sensory information and motor responses. Thus, the NDH proposes that delirium results from an acute breakdown in network connectivity within the brain.³⁵⁷

Furthermore, the NDH suggests that two important factors determine a subject's vulnerability to delirium: (1) the baseline network connectivity (defined as the connectivity of neural networks within the brain before the precipitating insult provoking delirium), which is influenced by most recognized nonmodifiable delirium risk factors (e.g., age, baseline level of cognitive functioning), and (2) the level of inhibitory tone, which will determine the degree of change in network connectivity and is influenced by modifiable risk factors (e.g., metabolic abnormalities, sleep deprivation, infection and inflammation, medication such as benzodiazepines).³⁵⁷ The model suggests that when these two factors affect separate neuronal networks, to different degrees, they produce the various motoric phenotypes described (e.g., hyperactive, hypoactive, mixed). Thus, the form of delirium that ensues will depend on how and which networks breaks down (dependent on both the individual's baseline network connectivity and the degree of change in inhibitory tone produced).

Electroencephalographic and evoked potential data may provide further support for this theory by suggesting that the pathophysiology of at least some forms of delirium may have a subcortical component.³⁵⁸ More recently, a study using functional magnetic resonance imaging scans examined the correlations of blood oxygen levels between various brain regions in resting-state functional magnetic resonance imaging scans during and after the resolution of delirium. Findings demonstrated a long-lasting disruption in reciprocity of the dorsolateral prefrontal cortex with the posterior cingulate cortex and a reversible reduction of functional connectivity of subcortical regions (e.g., thalamus) with the reticular activating system and with nuclei responsible for

forebrain Ach (i.e., the midbrain nucleus basalis) and DA (i.e., the midbrain ventral tegmental area) innervation.³⁵⁹ The persistence of these physiologic disruptions, beyond the resolution of acute delirious symptoms, may account for reported cognitive problems that often outlast the acute episode of delirium.

The NDH intersects with the NTH because GABAergic neurotransmission is implicated in increasing the inhibitory tone, which may contribute to the development of delirium.³⁵⁷ Furthermore, GABAergic agents may further destabilize the sleep–wake cycle, as they suppress orexinergic neural firing in the perifornical nucleus.³⁶⁰ Orexin (hypocretin) is a peptide produced in the lateral hypothalamus that strengthens the ascending reticular activating system, thus maintaining wakefulness and preventing inappropriate transition into sleep.^{17,18,265,361} It also relates to the NIH because systemic inflammation drives an up-regulation in expression of GABA_A receptors and an increased GABA synthesis^{362,363} and suppresses orexinergic neuronal activity during the wakeful period.³⁶⁴

This model also intersects with the NAH because it suggests that aging is associated with gray matter volume, neurotransmission, and white matter integrity (which consists mostly of glial cells and myelinated axons that transmit signals from one region of the cerebrum to another and between the cerebrum and lower brain centers) and thus may represent an anatomic correlate of functional connectivity.^{357,365,366} A study assessing white matter integrity using diffusion tensor imaging in patients who developed delirium after cardiac surgery suggested that abnormalities in the deep white matter and thalamus could have accounted for the delirious patients' vulnerability to postoperative delirium, compared with nondelirious subjects.³⁶⁷ A more recent study, also conducted among elderly cardiac surgery patients, demonstrated that the prevalence of severe cerebral white matter hyperintensities on magnetic resonance imaging was significantly higher in delirious patients and similarly concluded these lesions were likely one of the most important risk factors for the development of delirium after cardiac surgery.³⁶⁸

Aging is also associated with a reduction in GABAergic tone, thus allowing for increased neuronal activity in certain brain regions, primarily

the prefrontal cortex.³⁶⁹ With advanced age there is a down-regulation of several subunits of the GABA_A receptors (i.e., $\alpha 1$, $\alpha 5$, $\beta 3$, $\gamma 2$); therefore, a stimulus that increases the level of inhibitory tone may have a relatively greater effect and further breakdown network connectivity.³⁷⁰ The aged brain also experiences reductions in orexin signaling,^{371–374} which may contribute to the fluctuating arousal level seen in delirious states when exposed to various noxious stimuli (e.g., infection, GABAergic agents, sleep deprivation). Similarly, it intersects with the NTH because it specifically implicates the GABA system (e.g., benzodiazepine or HE-induced delirium) and the cholinergic system (e.g., anticholinergic delirium) as likely principal culprits.

The NDH also intersects with the NTH because data from anesthetic agent studies demonstrate that some core symptoms of delirium likely involve changes in dynamic aspects of neuronal activity affecting brain's ability to integrate information through functional disconnection between different anatomic structures.^{375,376} Finally, it intersects with the NIH because acute neuroinflammatory reactions affect physiologic processes implicated in neuronal and synaptic function with consequent neurochemical disturbances and functional disconnection between different anatomic structures.⁴⁸

CONCLUSION

Delirium is a neurobehavioral syndrome caused by the disruption of neuronal activity secondary to systemic disturbances. To date, no single unitary pathophysiologic mechanism has been identified. Most existing theories on the etiology of delirium are complementary rather than competing. Thus, it is likely that none of these theories by themselves explains the full phenomenon of delirium but rather that two or more of these, if not all, act together to lead to the biochemical derangement we know as delirium.

A review of the available literature suggests a number of factors that lead to a final common pathway associated with alterations in neurotransmitter synthesis and availability that mediates the behavioral and cognitive changes observed in delirium. These alterations in neurotransmitter synthesis appear to provide a relatively satisfactory

explanation of the complex behavioral and cognitive changes observed in delirium. These factors include a wide range of endocrinologic, immunologic, neuro-inflammatory, neurologic (i.e., brain connectivity circuits), and metabolic effects (i.e., oxidative stress) described throughout the article. In addition, certain specific circumstances or phases of life may lower the threshold for such events or exaggerate the body's response, such as the presence of inflammatory or infectious processes, sleep deprivation, aging, and neurodegenerative disorders. The observed changes in neurotransmitters are not uniform. In general, the most commonly described neurotransmitter changes associated with delirium are deficiencies in Ach availability (\downarrow Ach) and melatonin (\downarrow MEL); excess in DA (\uparrow DA), NE (\uparrow NE), and/or GLU (\uparrow GLU) release;

and variable alterations (e.g., both a decreased and increased activity) in 5HT (\downarrow \uparrow 5HT), histamine (\downarrow \uparrow H1 and H2), and/or GABA (\downarrow \uparrow GABA). Ultimately, it is likely that relative neurotransmitter alterations determine the phenotypic presentations of delirium (e.g., hyperactive, hypoactive, or mixed).

Thus, in the end, it is unlikely that a single cascade will explain the phenomena of delirium. Rather, it is more likely that a number of pathways will be shared, based on common systemic vulnerabilities and pathologic processes, yielding varying neurotransmitter deficiencies. The absence of a single common pathway or uniform neurotransmitter abnormality may explain why effective prophylaxis and treatment of delirium has proven so difficult and remained so elusive.

References

- Maldonado JR: Pathoetiological model of delirium: a comprehensive understanding of the neurobiology of delirium and an evidence-based approach to prevention and treatment. *Crit Care Clin* 2008; 24:789–856
- American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Washington, DC, American Psychiatric Association, 2013
- American Psychiatric Association: Guideline watch: practice guideline for the treatment of patients with delirium, in American Psychiatric Association Practice Guidelines. Edited by Cook IA. Washington, DC, American Psychiatric Association, 2004
- Maldonado JR: Delirium in the acute care setting: characteristics, diagnosis and treatment. *Crit Care Clin* 2008; 24:657–722
- Trzepacz PT, Meagher DJ: Neuropsychiatric aspects of delirium, in *Textbook of Neuropsychiatry* 5th ed. Edited by Yudofsky SC, Hales RE. Washington, DC, American Psychiatric Press, 2008, pp 445–518
- Ryan DJ, O'Regan NA, Caoimh RÓ, et al: Delirium in an adult acute hospital population: predictors, prevalence and detection. *BMJ Open* 2013; 3(1). <http://dx.doi.org/10.1136/bmjopen-2012-001772>
- Inouye SK: Delirium in hospitalized older patients. *Clin Geriatr Med* 1998; 14:745–764
- Naughton BJ, Moran MB, Kadam H, et al: Delirium and other cognitive impairment in older adults in an emergency department. *Ann Emerg Med* 1995; 25:751–755
- Cole MG: Delirium in elderly patients. *Am J Geriatr Psychiatry* 2004; 12:7–21
- Elie M, Rousseau F, Cole M, et al: Prevalence and detection of delirium in elderly emergency department patients. *CMAJ* 2000; 163:977–981
- Wiesel O, Klausner J, Soffer D, et al: [Post-operative delirium of the elderly patient—an iceberg?]. *Harefuah* 2011; 150:260–263, 303
- Vaurio LE, Sands LP, Wang Y, et al: Postoperative delirium: the importance of pain and pain management. *Anesth Analg* 2006; 102:1267–1273
- Dyer CB, Ashton CM, Teasdale TA: Postoperative delirium. A review of 80 primary data-collection studies. *Arch Intern Med* 1995; 155:461–465
- Maldonado JR, Wysong A, van der Starre PJ, et al: Dexmedetomidine and the reduction of postoperative delirium after cardiac surgery. *Psychosomatics* 2009; 50:206–217
- Bruce AJ, Ritchie CW, Blizard R, et al: The incidence of delirium associated with orthopedic surgery: a meta-analytic review. *Int Psychogeriatr* 2007; 19:197–214
- Ely EW, Margolin R, Francis J, et al: Evaluation of delirium in critically ill patients: validation of the Confusion Assessment Method for the Intensive Care Unit (CAM-ICU). *Crit Care Med* 2001; 29:1370–1379
- Rosenthal GE, Kaboli PJ, Barnett MJ, et al: Age and the risk of in-hospital death: insights from a multihospital study of intensive care patients. *J Am Geriatr Soc* 2002; 50:1205–1212
- Bagshaw SM, Webb SA, Delaney A, et al: Very old patients admitted to intensive care in Australia and New Zealand: a multi-centre cohort analysis. *Crit Care* 2009; 13:R45
- Rothschild JL, Leape LL: The Nature and Extent of Medical Injury in Older Patients: Executive Summary. Washington, DC, Public Policy Institute, AARP, 2000
- Francis J, Martin D, Kapoor WN: A prospective study of delirium in hospitalized elderly. *JAMA* 1990; 263:1097–1101
- Curyto KJ, Johnson R, TenHave T, et al: Survival of hospitalized elderly patients with delirium: a prospective study. *Am J Geriatr Psychiatry* 2001; 9:141–147
- Pisani MA, Kong SY, Kasl SV, et al: Days of delirium are associated with 1-year mortality in an older intensive care unit population. *Am J Respir Crit Care Med* 2009; 180:1092–1097
- Ely EW, Gautam S, Margolin R, et al: The impact of delirium in the intensive care unit on hospital length of stay. *Intensive Care Med* 2001; 27:1892–1900
- Ely EW, Shintani A, Truman B, et al: Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA* 2004; 291:1753–1762
- Maldonado JR, Dhimi N, Wise L: Clinical implications of the recognition and management of delirium in general medical and surgical wards. *Psychosomatics* 2003; 44:157–158
- O'Keeffe S, Lavan J: The prognostic significance of delirium in older hospital patients. *J Am Geriatr Soc* 1997; 45:174–178

27. McCusker J, Cole MG, Dendukuri N, et al: Does delirium increase hospital stay? *J Am Geriatr Soc* 2003; 51:1539–1546
28. Han JH, Eden S, Shintani A, et al: Delirium in older emergency department patients is an independent predictor of hospital length of stay. *Acad Emerg Med* 2011; 18:451–457
29. Gunther ML, Jackson JC, Ely EW: The cognitive consequences of critical illness: practical recommendations for screening and assessment. *Crit Care Clin* 2007; 23:491–506
30. Jackson JC, Gordon SM, Hart RP, et al: The association between delirium and cognitive decline: a review of the empirical literature. *Neuropsychol Rev* 2004; 14:87–98
31. Macdonald AJ: Can delirium be separated from dementia? *Dement Geriatr Cogn Disord* 1999; 10:386–388
32. Rockwood K, Cosway S, Carver D, et al: The risk of dementia and death after delirium. *Age Ageing* 1999; 28:551–556
33. Wacker P, Nunes PV, Cabrita H, et al: Post-operative delirium is associated with poor cognitive outcome and dementia. *Dement Geriatr Cogn Disord* 2006; 21:221–227
34. Griffiths RD, Jones C: Delirium, cognitive dysfunction and post-traumatic stress disorder. *Curr Opin Anaesthesiol* 2007; 20:124–129
35. Bickel H, Grading R, Kochs E, et al: High risk of cognitive and functional decline after postoperative delirium. A three-year prospective study. *Dement Geriatr Cogn Disord* 2008; 26:26–31
36. Kat MG, Vreeswijk R, de Jonghe JF, et al: Long-term cognitive outcome of delirium in elderly hip surgery patients. A prospective matched controlled study over two and a half years. *Dement Geriatr Cogn Disord* 2008; 26:1–8
37. Fong TG, Tulebaev SR, Inouye SK: Delirium in elderly adults: diagnosis, prevention and treatment. *Nat Rev Neurol* 2009; 5:210–220
38. MacLulich AM, Beaglehole A, Hall RJ, et al: Delirium and long-term cognitive impairment. *Int Rev Psychiatry* 2009; 21:30–42
39. Girard TD, Jackson JC, Pandharipande PP, et al: Delirium as a predictor of long-term cognitive impairment in survivors of critical illness. *Crit Care Med* 2010; 38:1513–1520
40. Mathews SB, Arnold SE, Epperson CN: Hospitalization and cognitive decline: can the nature of the relationship be deciphered? *Am J Geriatr Psychiatry* 2013; doi: 10.1016/j.jagp.2012.08.012. [Epub ahead of print]
41. Francis J, Kapoor WN: Delirium in hospitalized elderly. *J Gen Intern Med* 1990; 5:65–79
42. Inouye SK: Prevention of delirium in hospitalized older patients: risk factors and targeted intervention strategies. *Ann Med* 2000; 32:257–263
43. Millbrandt EB, Deppen S, Harrison PL, et al: Costs associated with delirium in mechanically ventilated patients. *Crit Care Med* 2004; 32:955–962
44. Leslie DL, Marcantonio ER, Zhang Y, et al: One-year health care costs associated with delirium in the elderly population. *Arch Intern Med* 2008; 168:27–32
45. Engel GL, Romano J: Delirium, a syndrome of cerebral insufficiency. *J Chronic Dis* 1959; 9:260–277, 531, 532
46. Brown T: Basic mechanisms in the pathogenesis of delirium, in *The Psychiatric Care of the Medical Patient*. Edited by Fogel BS, Stoudemire A, Greenberg DB. New York, Oxford Press, 2000, pp 571–580
47. Lipowski ZJ: Update on delirium. *Psychiatr Clin North Am* 1992; 15:335–346
48. Cerejeira J, Firmino H, Vaz-Serra A, et al: The neuroinflammatory hypothesis of delirium. *Acta Neuropathol* 2010; 119:737–754
49. Godbout JP, Chen J, Abraham J, et al: Exaggerated neuroinflammation and sickness behavior in aged mice following activation of the peripheral innate immune system. *FASEB J* 2005; 19:1329–1331
50. Godbout JP, Johnson RW: Age and neuroinflammation: a lifetime of psychoneuroimmune consequences. *Immunol Allergy Clin North Am* 2009; 29:321–337
51. Cunningham C: Systemic inflammation and delirium: important co-factors in the progression of dementia. *Biochem Soc Trans* 2011; 39:945–953
52. Cunningham C, Campion S, Lunnon K, et al: Systemic inflammation induces acute behavioral and cognitive changes and accelerates neurodegenerative disease. *Biol Psychiatry* 2009; 65:304–312
53. Cunningham C, Wilcockson DC, Campion S, et al: Central and systemic endotoxin challenges exacerbate the local inflammatory response and increase neuronal death during chronic neurodegeneration. *J Neurosci* 2005; 25:9275–9284
54. Hart BL: Biological basis of the behavior of sick animals. *Neurosci Biobehav Rev* 1988; 12:123–137
55. Dantzer R: [Neurotropic effects of cytokines: at the limits of immunology and neurobiology]. *Pathol Biol (Paris)* 1994; 42:826–829
56. Dantzer R: Cytokine-induced sickness behavior: mechanisms and implications. *Ann N Y Acad Sci* 2001; 933:222–234
57. Dantzer R: Cytokine-induced sickness behaviour: a neuro-immune response to activation of innate immunity. *Eur J Pharmacol* 2004; 500:399–411
58. Dantzer R, et al: Cytokines and sickness behavior. *Ann N Y Acad Sci* 1998; 840:586–590
59. Konsman JP, Parnet P, Dantzer R: Cytokine-induced sickness behaviour: mechanisms and implications. *Trends Neurosci* 2002; 25:154–159
60. Dantzer R, O'Connor JC, Freund GG, et al: From inflammation to sickness and depression: when the immune system subjugates the brain. *Nature reviews. Neuroscience* 2008; 9:46–56
61. de Rooij SE, van Munster BC, Korevaar JC, et al: Cytokines and acute phase response in delirium. *J Psychosom Res* 2007; 62:521–525
62. Beloosesky Y, Hendel D, Weiss A, et al: Cytokines and C-reactive protein production in hip-fracture-operated elderly patients. *J Gerontol A Biol Sci Med Sci* 2007; 62:420–426
63. McGrane S, Girard TD, Thompson JL, et al: Procalcitonin and C-reactive protein levels at admission as predictors of duration of acute brain dysfunction in critically ill patients. *Crit Care* 2011; 15:R78
64. Osse RJ, Fekkes D, Tulen JH, et al: High preoperative plasma neopterin predicts delirium after cardiac surgery in older adults. *J Am Geriatr Soc* 2012; 60:661–668
65. Murr C, Widner B, Wirleitner B, et al: Neopterin as a marker for immune system activation. *Curr Drug Metab* 2002; 3:175–187
66. Mrak RE, Griffin WS: The role of activated astrocytes and of the neurotrophic cytokine S100B in the pathogenesis of Alzheimer's disease. *Neurobiol Aging* 2001; 22:915–922
67. Peskind ER, Griffin WS, Akama KT, et al: Cerebrospinal fluid S100B is elevated in the earlier stages of Alzheimer's disease. *Neurochem Int* 2001; 39:409–413
68. Petzold A, Jenkins R, Watt HC, et al: Cerebrospinal fluid S100B correlates with brain atrophy in Alzheimer's disease. *Neurosci Lett* 2003; 336:167–170
69. Bokesch PM, Izykenova GA, Justice JB, et al: NMDA receptor antibodies predict adverse neurological outcome after cardiac surgery in high-risk patients. *Stroke* 2006; 37:1432–1436
70. Hall RJ, Ferguson KJ, Andrews M, et al: Delirium and cerebrospinal fluid S100B in hip fracture patients: a preliminary study. *Am J Geriatr Psychiatry* 2013; 21:1239–1243

71. van Munster BC, Korevaar JC, Korse CM, et al: Serum S100B in elderly patients with and without delirium. *Int J Geriatr Psychiatry* 2010; 25:234–239
72. van Munster BC, Korse CM, de Rooij SE, et al: Markers of cerebral damage during delirium in elderly patients with hip fracture. *BMC Neurol* 2009; 9:21
73. He F, Yin F, Peng J, et al: Molecular mechanism for change in permeability in brain microvascular endothelial cells induced by LPS. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2010; 35:1129–1137
74. Hala M: Pathophysiology of postoperative delirium: systemic inflammation as a response to surgical trauma causes diffuse microcirculatory impairment. *Med Hypoth* 2007; 68:194–196
75. Bogatcheva NV, Sergeeva MG, Dudek SM, et al: Arachidonic acid cascade in endothelial pathobiology. *Microvasc Res* 2005; 69: 107–127
76. Uchikado H, Akiyama H, Kondo H, et al: Activation of vascular endothelial cells and perivascular cells by systemic inflammation—an immunohistochemical study of postmortem human brain tissues. *Acta Neuropathol* 2004; 107:341–351
77. Hirsch JA, Gibson GE: Selective alteration of neurotransmitter release by low oxygen in vitro. *Neurochem Res* 1984; 9: 1039–1049
78. Hshieh TT, Fong TG, Marcantonio ER, et al: Cholinergic deficiency hypothesis in delirium: a synthesis of current evidence. *J Gerontol A Biol Sci Med Sci* 2008; 63:764–772
79. Belosessky Y, Hendel D, Weiss A, et al: Cytokines and C-reactive protein production in hip-fracture-operated elderly patients. *J Gerontol A Biol Sci Med Sci* 2007; 62:420–426
80. Dantzer R, Capuron L, Irwin MR, et al: Identification and treatment of symptoms associated with inflammation in medically ill patients. *Psychoneuroendocrinology* 2008; 33:18–29
81. Sorrells SF, Caso JR, Munhoz CD, et al: The stressed CNS: when glucocorticoids aggravate inflammation. *Neuron* 2009; 64: 33–39
82. van Munster BC, Bisschop PH, Zwinderman AH, et al: Cortisol, interleukins and S100B in delirium in the elderly. *Brain Cogn* 2010; 74:18–23
83. Huber JD, Hau VS, Borg L, et al: Blood-brain barrier tight junctions are altered during a 72-h exposure to lambda-carrageenan-induced inflammatory pain. *Am J Physiol Heart Circ Physiol* 2002; 283:H1531–H1537
84. Brooks TA, Hawkins BT, Huber JD, et al: Chronic inflammatory pain leads to increased blood-brain barrier permeability and tight junction protein alterations. *Am J Physiol Heart Circ Physiol* 2005; 289:H738–H743
85. Butterworth RF: The liver-brain axis in liver failure: neuroinflammation and encephalopathy. *Nat Rev Gastroenterol Hepatol* 2013; 10:522–528
86. Bemeur C, Butterworth RF: Liver-brain proinflammatory signaling in acute liver failure: role in the pathogenesis of hepatic encephalopathy and brain edema. *Metab Brain Dis* 2013; 28: 145–150
87. Stone TW, Forrest CM, Darlington LG: Kynurenine pathway inhibition as a therapeutic strategy for neuroprotection. *FEBS J* 2012; 279:1386–1397
88. Darlington LG, Forrest CM, Mackay GM, et al: On the biological importance of the 3-hydroxyanthranilic Acid: Anthranilic Acid Ratio. *Int J Tryptophan Res* 2010; 3:51–59
89. Stone TW, Behan WM, Jones PA, et al: The role of kynurenines in the production of neuronal death, and the neuroprotective effect of purines. *J Alzheim Dis* 2001; 3:355–366
90. Behan WM, Stone TW: Role of kynurenines in the neurotoxic actions of kainic acid. *Br J Pharmacol* 2000; 129:1764–1770
91. Forrest CM, Mackay GM, Oxford L, et al: Kynurenine metabolism predicts cognitive function in patients following cardiac bypass and thoracic surgery. *J Neurochem* 2011; 119:136–152
92. Elie M, Cole MG, Primeau FJ, et al: Delirium risk factors in elderly hospitalized patients. *J Gen Intern Med* 1998; 13:204–212
93. Zampieri FG, Park M, Machado FS, et al: Sepsis-associated encephalopathy: not just delirium. *Clinics (Sao Paulo)* 2011; 66: 1825–1831
94. van Gool WA, van de Beek D, Eikelenboom P: Systemic infection and delirium: when cytokines and acetylcholine collide. *Lancet* 2010; 375:773–775
95. Simone MJ, Tan ZS: The role of inflammation in the pathogenesis of delirium and dementia in older adults: a review. *CNS Neurosci Ther* 2011; 17:506–513
96. Lemstra AW, Kalisvaart KJ, Vreeswijk R, et al: Pre-operative inflammatory markers and the risk of postoperative delirium in elderly patients. *Int J Geriatr Psychiatry* 2008; 23:943–948
97. Frederiks JA: Inflammation of the mind. On the 300th anniversary of Gerard van Swieten. *J Hist Neurosci* 2000; 9:307–310
98. van den Boogaard M, Kox M, Quinn KL, et al: Biomarkers associated with delirium in critically ill patients and their relation with long-term subjective cognitive dysfunction; indications for different pathways governing delirium in inflamed and noninflamed patients. *Crit Care* 2011; 15:R297
99. Munster BC, Aronica E, Zwinderman AH, et al: Neuroinflammation in delirium: a postmortem case-control study. *Rejuvenation Res* 2011; 14:615–622
100. Ouimet S, Kavanagh BP, Gottfried SB, et al: Incidence, risk factors and consequences of ICU delirium. *Intensive Care Med* 2007; 33:66–73
101. Pandharipande P, Shintani A, Peterson J, et al: Lorazepam is an independent risk factor for transitioning to delirium in intensive care unit patients. *Anesthesiology* 2006; 104:21–26
102. MacLulich AM, Ferguson KJ, Miller T, et al: Unravelling the pathophysiology of delirium: a focus on the role of aberrant stress responses. *J Psychosom Res* 2008; 65:229–238
103. Murray C, Sanderson DJ, Barkus C, et al: Systemic inflammation induces acute working memory deficits in the primed brain: relevance for delirium. *Neurobiol Aging* 2012; 33:603–616
104. Adelman JS, Martin LB: Immune systems and sickness behavior, in *Encyclopedia of Animal Behavior*. Edited by Breed MD, Moore J. New York, NY, Elsevier, 2010, pp 133–137
105. Berczi I, Szentivanyi A: The Immune-Neuroendocrine Circuitry, in *NeuroImmune Biology*. New York, Elsevier, 2003, pp 561–592
106. Troncale JA: The aging process. Physiologic changes and pharmacologic implications. *Postgrad Med* 1996; 99:111–114; 120–122
107. Coleman PD, Flood DG: Neuron numbers and dendritic extent in normal aging and Alzheimer's disease. *Neurobiol Aging* 1987; 8:521–545
108. van der Mast RC: Pathophysiology of delirium. *J Geriatr Psychiatry Neurol* 1998; 11:138–145; discussion 157–158
109. Kochunov P, Ramage AE, Lancaster JL, et al: Loss of cerebral white matter structural integrity tracks the gray matter metabolic decline in normal aging. *Neuroimage* 2009; 45:17–28
110. Kelly KM, Nadon NL, Morrison JH, et al: The neurobiology of aging. *Epilepsy Res* 2006; 68(suppl 1):S5–S20
111. Juraska JM, Lowry NC: Neuroanatomical changes associated with cognitive aging. *Curr Top Behav Neurosci* 2012; 10: 137–162
112. Chen JJ, Rosas HD, Salat HD: Age-associated reductions in cerebral blood flow are independent from regional atrophy. *Neuroimage* 2011; 55:468–478

113. Brown WR, Thore CR: Review: cerebral microvascular pathology in ageing and neurodegeneration. *Neuropathol Appl Neurobiol* 2011; 37:56–74
114. Bertsch K, Hagemann D, Hermes M, et al: Resting cerebral blood flow, attention, and aging. *Brain Res* 2009; 1267:77–88
115. O'Rourke MF: Arterial aging: pathophysiological principles. *Vasc Med* 2007; 12:329–341
116. Beason-Held LL, Kraut MA, Resnick SM II: Temporal patterns of longitudinal change in aging brain function. *Neurobiol Aging* 2008; 29:497–513
117. Schroeter ML, Schmiedel O, von Cramon DY: Spontaneous low-frequency oscillations decline in the aging brain. *J Cereb Blood Flow Metab* 2004; 24:1183–1191
118. Riddle DR, Sonntag WE, Lichtenwalner RJ: Microvascular plasticity in aging. *Ageing Res Rev* 2003; 2:149–168
119. Van Laere K, Versijpt J, Audenaert K, et al: 99mTc-ECD brain perfusion SPET: variability, asymmetry and effects of age and gender in healthy adults. *Eur J Nucl Med* 2001; 28:873–887
120. Celsis P, Agniel A, Cardebat D, et al: Age related cognitive decline: a clinical entity? A longitudinal study of cerebral blood flow and memory performance. *J Neurol Neurosurg Psychiatry* 1997; 62:601–608
121. Takada H, Nagata K, Hirata Y, et al: Age-related decline of cerebral oxygen metabolism in normal population detected with positron emission tomography. *Neurol Res* 1992; 14(2 suppl):128–131
122. Meier-Ruge W, Iwagoff P, Reichleimer K, et al: Neurochemical findings in the aging brain. *Adv Biochem Psychopharmacol* 1980; 23:323–338
123. Hebert LE, Scherr PA, Beckett LA, et al: Age-specific incidence of Alzheimer's disease in a community population. *JAMA* 1995; 273:1354–1359
124. Milstein A, Pollack A, Kleinman G, et al: Confusion/delirium following cataract surgery: an incidence study of 1-year duration. *Int Psychogeriatr* 2002; 14:301–306
125. Khurana V, Gambhir IS, Kishore D: Evaluation of delirium in elderly: a hospital-based study. *Geriatr Gerontol Int* 2011; 11: 467–473
126. Lee HB, Mears SC, Rosenberg PB, et al: Predisposing factors for postoperative delirium after hip fracture repair in individuals with and without dementia. *J Am Geriatr Soc* 2011; 59: 2306–2313
127. Rudolph JL, Harrington MB, Lucatorto MA, et al: Validation of a medical record-based delirium risk assessment. *J Am Geriatr Soc* 2011; 59(suppl 2):S289–S294
128. Srinonprasert V, Pakdeewongse S, Assanasen J, et al: Risk factors for developing delirium in older patients admitted to general medical wards. *J Med Assoc Thai* 2011; 94(suppl 1):S99–S104
129. McAvay GJ, Van Ness PH, Bogardus ST Jr, et al: Older adults discharged from the hospital with delirium: 1-year outcomes. *J Am Geriatr Soc* 2006; 54:1245–1250
130. Inouye S, Zhang Y, Jones RN, et al: Risk factors for delirium at discharge: development and validation of a predictive model. *Arch Intern Med* 2007; 167:1406–1413
131. Franco JG, Valencia C, Bernal C, et al: Relationship between cognitive status at admission and incident delirium in older medical inpatients. *J Neuropsychiatry Clin Neurosci* 2010; 22: 329–337
132. Litaker D, Locala J, Franco K, et al: Preoperative risk factors for postoperative delirium. *Gen Hosp Psychiatry* 2001; 23:84–89
133. McNicoll L, Pisani MA, Zhang Y, et al: Delirium in the intensive care unit: occurrence and clinical course in older patients. *J Am Geriatr Soc* 2003; 51:591–598
134. Benoit AG, Campbell BI, Tanner JR, et al: Risk factors and prevalence of perioperative cognitive dysfunction in abdominal aneurysm patients. *J Vasc Surg* 2005; 42:884–890
135. Rudolph JL, Jones RN, Grande LJ, et al: Impaired executive function is associated with delirium after coronary artery bypass graft surgery. *J Am Geriatr Soc* 2006; 54:937–941
136. Smith PJ, Attix DK, Weldon BC, et al: Executive function and depression as independent risk factors for postoperative delirium. *Anesthesiology* 2009; 110:781–787
137. Lowery DP, Wesnes K, Ballard CG: Subtle attentional deficits in the absence of dementia are associated with an increased risk of post-operative delirium. *Dement Geriatr Cogn Disord* 2007; 23: 390–394
138. Cole MG, Ciampi A, Belzile E, et al: Persistent delirium in older hospital patients: a systematic review of frequency and prognosis. *Age Ageing* 2009; 38:19–26
139. Michaud M, Balardy L, Moulis G, et al: Proinflammatory cytokines, aging, and age-related diseases. *J Am Med Dir Assoc* 2013; doi: 10.1016/j.jamda.2013.05.009. [Epub ahead of print]
140. Leung JM, Sands LP, Lim E, et al: Does preoperative risk for delirium moderate the effects of postoperative pain and opiate use on postoperative delirium? *Am J Geriatr Psychiatry* 2013; 21: 946–956
141. Abildstrom H, Christiansen M, Siersma, et al: Apolipoprotein E genotype and cognitive dysfunction after noncardiac surgery. *Anesthesiology* 2004; 101:855–861
142. Bryson GL, Wyand A, Wozny D, et al: A prospective cohort study evaluating associations among delirium, postoperative cognitive dysfunction, and apolipoprotein E genotype following open aortic repair. *Can J Anaesth* 2011; 58:246–255
143. Tagarakis GI, Tsolaki-Tagaraki F, Tsolaki M, et al: The role of apolipoprotein E in cognitive decline and delirium after bypass heart operations. *Am J Alzheim Dis Other Dement* 2007; 22: 223–228
144. van Munster BC, Korevaar JC, de Rooij SE, et al: The association between delirium and the apolipoprotein E epsilon4 allele in the elderly. *Psychiatr Genet* 2007; 17:261–266
145. Adamis D, Treloar A, Martin FC, et al: APOE and cytokines as biological markers for recovery of prevalent delirium in elderly medical inpatients. *Int J Geriatr Psychiatry* 2007; 22: 688–694
146. Ely EW, Girard TD, Shintani AK, et al: Apolipoprotein E4 polymorphism as a genetic predisposition to delirium in critically ill patients. *Crit Care Med* 2007; 35:112–117
147. Leung JM, Sands LP, Wang Y, et al: Apolipoprotein E epsilon4 allele increases the risk of early postoperative delirium in older patients undergoing noncardiac surgery. *Anesthesiology* 2007; 107:406–411
148. van Munster BC, Korevaar JC, Zwinderman AH, et al: The association between delirium and the apolipoprotein E epsilon4 allele: new study results and a meta-analysis. *Am J Geriatr Psychiatry* 2009; 17:856–862
149. Zhang F, Lewis M, Yang G, et al: Apolipoprotein E polymorphism, life stress and self-reported health among older adults. *J Epidemiol Commun Health* 2008; 62:e3
150. Adamis D, Van Munster BC, Macdonald AJ: The genetics of deliria. *Int Rev Psychiatry* 2009; 21:20–29
151. Aliev G, Obrenovich ME, Smith MA, et al: Hypoperfusion, mitochondria failure, oxidative stress, and Alzheimer disease. *J Biomed Biotechnol* 2003; 2003:162–163
152. Berr C: Cognitive impairment and oxidative stress in the elderly: results of epidemiological studies. *Biofactors* 2000; 13: 205–209

153. Berr C, Balansard B, Arnaud J, et al: Cognitive decline is associated with systemic oxidative stress: the EVA study. *Etude du Vieillissement Arteriel. J Am Geriatr Soc* 2000; 48:1285–1291
154. Berr C, Richard MJ, Roussel AM, et al: Systemic oxidative stress and cognitive performance in the population-based EVA study. *Etude du Vieillissement Arteriel. Free Radic Biol Med* 1998; 24: 1202–1208
155. Seaman JS, Schillerstrom J, Carroll D, et al: Impaired oxidative metabolism precipitates delirium: a study of 101 ICU patients. *Psychosomatics* 2006; 47:56–61
156. Karlidag R, Unal S, Sezer OH, et al: The role of oxidative stress in postoperative delirium. *Gen Hosp Psychiatry* 2006; 28:418–423
157. Rosenberg J, Kehlet H: Postoperative mental confusion—association with postoperative hypoxemia. *Surgery* 1993; 114: 76–81
158. Gibson GE, Peterson C: Aging decreases oxidative metabolism and the release and synthesis of acetylcholine. *J Neurochem* 1981; 37:978–984
159. Slater JP, Guarino T, Stack J, et al: Cerebral oxygen desaturation predicts cognitive decline and longer hospital stay after cardiac surgery. *Ann Thorac Surg* 2009; 87:36–44; discussion 44–45
160. Morimoto Y, Yoshimura M, Utada K, et al: Prediction of postoperative delirium after abdominal surgery in the elderly. *J Anesth* 2009; 23:51–56
161. Schoen J, Meyerrose J, Paarmann H, et al: Preoperative regional cerebral oxygen saturation is a predictor of postoperative delirium in on-pump cardiac surgery patients: a prospective observational trial. *Crit Care* 2011; 15:R218
162. Ng T, Graham DI, Adams JH, et al: Changes in the hippocampus and the cerebellum resulting from hypoxic insults: frequency and distribution. *Acta Neuropathol* 1989; 78:438–443
163. Basarsky TA, Feighan D, MacVicar BA: Glutamate release through volume-activated channels during spreading depression. *J Neurosci* 1999; 19:6439–6445
164. Iijima T, Shimase C, Iwao Y, et al: Relationships between glutamate release, blood flow and spreading depression: real-time monitoring using an electroenzymatic dialysis electrode. *Neurosci Res* 1998; 32:201–207
165. Moghaddam B, Schenk JO, Stewart, et al: Temporal relationship between neurotransmitter release and ion flux during spreading depression and anoxia. *Can J Physiol Pharmacol* 1987; 65: 1105–1110
166. Shimizu-Sasamata M, Bosque-Hamilton P, Huang PL, et al: Attenuated neurotransmitter release and spreading depression-like depolarizations after focal ischemia in mutant mice with disrupted type I nitric oxide synthase gene. *J Neurosci* 1998; 18: 9564–9571
167. Somjen GG, Aitken PG, Balestrino M, et al: Extracellular ions, hypoxic irreversible loss of function and delayed postischemic neuron degeneration studied in vitro. *Acta Physiol Scand Suppl* 1989; 582:58
168. Somjen GG, Aitken PG, Balestrino M, et al: Spreading depression-like depolarization and selective vulnerability of neurons. A brief review. *Stroke* 1990; 21(11 suppl):III179–III183
169. Busto R, Dietrich WD, Globus MY, et al: Extracellular release of serotonin following fluid-percussion brain injury in rats. *J Neurotrauma* 1997; 14:35–42
170. Busto R, Globus MY, Dietrich WD, et al: Effect of mild hypothermia on ischemia-induced release of neurotransmitters and free fatty acids in rat brain. *Stroke* 1989; 20:904–910
171. Globus MY, Alonso O, Dietrich WD, et al: Glutamate release and free radical production following brain injury: effects of post-traumatic hypothermia. *J Neurochem* 1995; 65:1704–1711
172. Globus MY, Busto R, Dietrich WD, et al: Effect of ischemia on the in vivo release of striatal dopamine, glutamate, and gamma-aminobutyric acid studied by intracerebral microdialysis. *J Neurochem* 1988; 51:1455–1464
173. Globus MY, Busto R, Dietrich WD, et al: Intra-ischemic extracellular release of dopamine and glutamate is associated with striatal vulnerability to ischemia. *Neurosci Lett* 1988; 91:36–40
174. Globus MY, Busto R, Dietrich WD, et al: Direct evidence for acute and massive norepinephrine release in the hippocampus during transient ischemia. *J Cereb Blood Flow Metab* 1989; 9: 892–896
175. Globus MY, Busto R, Martinez E, et al: Ischemia induces release of glutamate in regions spared from histopathologic damage in the rat. *Stroke* 1990; 21(11 suppl):III43–III46
176. Globus MY, Wester P, Busto R, et al: Ischemia-induced extracellular release of serotonin plays a role in CA1 neuronal cell death in rats. *Stroke* 1992; 23:1595–1601
177. Takagi K, Ginsberg MD, Globus MY, et al: Effect of hyperthermia on glutamate release in ischemic penumbra after middle cerebral artery occlusion in rats. *Am J Physiol* 1994; 267(5 Pt 2): H1770–H1776
178. Kirsch JR, Diringer MN, Borel CO, et al: Brain resuscitation. Medical management and innovations. *Crit Care Nurs Clin North Am* 1989; 1:143–154
179. Siesjo BK: Cerebral circulation and metabolism. *J Neurosurg* 1984; 60:883–908
180. Tune LE, Egeli S: Acetylcholine and delirium. *Dement Geriatr Cogn Disord* 1999; 10:342–344
181. Flacker JM, Cummings V, Mach JR Jr, et al: The association of serum anticholinergic activity with delirium in elderly medical patients. *Am J Geriatr Psychiatry* 1998; 6:31–41
182. Itil T, Fink M: Anticholinergic drug-induced delirium: experimental modification, quantitative EEG and behavioral correlations. *J Nerv Ment Dis* 1966; 143:492–507
183. Holinger PC, Klawans HL: Reversal of tricyclic-overdose-induced central anticholinergic syndrome by physostigmine. *Am J Psychiatry* 1976; 133:1018–1023
184. Tune LE, Damlouji NF, Holland A, et al: Association of postoperative delirium with raised serum levels of anticholinergic drugs. *Lancet* 1981; 2:651–653
185. Golinger RC, Peet T, Tune LE: Association of elevated plasma anticholinergic activity with delirium in surgical patients. *Am J Psychiatry* 1987; 144:1218–1220
186. Trzepacz PT, Leavitt M, Ciongoli K: An animal model for delirium. *Psychosomatics* 1992; 33:404–415
187. Trzepacz PT: Is there a final common neural pathway in delirium? Focus on acetylcholine and dopamine. *Semin Clin Neuropsychiatry* 2000; 5:132–148
188. Gaudreau JD, Gagnon P: Psychotogenic drugs and delirium pathogenesis: the central role of the thalamus. *Med Hypoth* 2005; 64:471–475
189. Trzepacz PT: The neuropathogenesis of delirium. A need to focus our research. *Psychosomatics* 1994; 35:374–391
190. Perry E, Walker M, Grace J, et al: Acetylcholine in mind: a neurotransmitter correlate of consciousness? *Trends Neurosci* 1999; 22:273–280
191. Flacker JM, Lipsitz LA: Neural mechanisms of delirium: current hypotheses and evolving concepts. *J Gerontol A Biol Sci Med Sci* 1999; 54:B239–B246
192. Trzepacz PT: Anticholinergic model for delirium. *Semin Clin Neuropsychiatry* 1996; 1:294–303
193. Lipowski ZJ: Delirium: how its concept has developed. *Int Psychogeriatr* 1991; 3:115–120

194. Blass J, Gibson G, Duffy T, et al: Cholinergic dysfunction: a common denominator in metabolic encephalopathies, in *Cholinergic Mechanisms*. Edited by Pepeu G, Ladinsky H. New York, Plenum, 1981, pp 921–928
195. Flacker JM, Wei JY: Endogenous anticholinergic substances may exist during acute illness in elderly medical patients. *J Gerontol A Biol Sci Med Sci* 2001; 56:M353–M355
196. Plaschke K, Thomas C, Engelhardt R, et al: Significant correlation between plasma and CSF anticholinergic activity in pre-surgical patients. *Neurosci Lett* 2007; 417:16–20
197. Tune L, Carr S, Cooper T, et al: Association of anticholinergic activity of prescribed medications with postoperative delirium. *J Neuropsychiatry Clin Neurosci* 1993; 5:208–210
198. Tune LE: Serum anticholinergic activity levels and delirium in the elderly. *Semin Clin Neuropsychiatry* 2000; 5:149–153
199. Flaherty JH: Psychotherapeutic agents in older adults. Commonly prescribed and over-the-counter remedies: causes of confusion. *Clin Geriatr Med* 1998; 14:101–127
200. Clegg A, Young JB: Which medications to avoid in people at risk of delirium: a systematic review. *Age Ageing* 2011; 40:23–29
201. Han L, McCusker J, Cole M, et al: Use of medications with anticholinergic effect predicts clinical severity of delirium symptoms in older medical inpatients. *Arch Intern Med* 2001; 161:1099–1105
202. Praticò C, Quattrone D, Lucanto T, et al: Drugs of anesthesia acting on central cholinergic system may cause post-operative cognitive dysfunction and delirium. *Med Hypoth* 2005; 65:972–982
203. Inouye SK: Predisposing and precipitating factors for delirium in hospitalized older patients. *Dement Geriatr Cogn Disord* 1999; 10:393–400
204. Campbell N, Perkins A, Hui S, et al: Association between prescribing of anticholinergic medications and incident delirium: a cohort study. *J Am Geriatr Soc* 2011; 59(suppl 2):S277–S281
205. Thomas RI, Cameron DJ, Fahs MC: A prospective study of delirium and prolonged hospital stay. Exploratory study. *Arch Gen Psychiatry* 1988; 45:937–940
206. Mussi C, Ferrari R, Ascari S, et al: Importance of serum anticholinergic activity in the assessment of elderly patients with delirium. *J Geriatr Psychiatry Neurol* 1999; 12:82–86
207. Mach JR Jr, Dysken MW, Kuskowski M, et al: Serum anticholinergic activity in hospitalized older persons with delirium: a preliminary study. *J Am Geriatr Soc* 1995; 43:491–495
208. Mulsant BH, Pollock BG, Kirshner M, et al: Serum anticholinergic activity in a community-based sample of older adults: relationship with cognitive performance. *Arch Gen Psychiatry* 2003; 60:198–203
209. Mangoni AA, van Munster BC, Woodman RJ, et al: Measures of anticholinergic drug exposure, serum anticholinergic activity, and all-cause postdischarge mortality in older hospitalized patients with hip fractures. *Am J Geriatr Psychiatry* 2013; 21:785–793
210. Fatranska M, Budai D, Oprsalova Z, et al: Acetylcholine and its enzymes in some brain areas of the rat under stress. *Brain Res* 1987; 424:109–114
211. Takayama H, Mizukawa K, Ota Z, et al: Regional responses of rat brain muscarinic cholinergic receptors to immobilization stress. *Brain Res* 1987; 436:291–295
212. Gibson GE, Peterson C, Sansone J: Decreases in amino acids and acetylcholine metabolism during hypoxia. *J Neurochem* 1981; 37:192–201
213. Cerejeira J, Nogueira V, Luis P, et al: The cholinergic system and inflammation: common pathways in delirium pathophysiology. *J Am Geriatr Soc* 2012; 60:669–675
214. Field RH, Gossen A, Cunningham C: Prior pathology in the basal forebrain cholinergic system predisposes to inflammation-induced working memory deficits: reconciling inflammatory and cholinergic hypotheses of delirium. *J Neurosci* 2012; 32:6288–6294
215. Knell AJ, Davidson AR, Williams R, et al: Dopamine and serotonin metabolism in hepatic encephalopathy. *Br Med J* 1974; 1:549–551
216. Graham DG: Catecholamine toxicity: a proposal for the molecular pathogenesis of manganese neurotoxicity and Parkinson's disease. *Neurotoxicology* 1984; 5:83–95
217. Carmen JS, Wyatt RJ: Calcium and malignant catatonia. *Lancet* 1977; 2:1124–1125
218. Nagatsu T: Genes for human catecholamine-synthesizing enzymes. *Neurosci Res* 1991; 12:315–345
219. Pedrosa R, Soares-da-Silva P: Oxidative and non-oxidative mechanisms of neuronal cell death and apoptosis by L-3,4-dihydroxyphenylalanine (L-DOPA) and dopamine. *Br J Pharmacol* 2002; 137:1305–1313
220. Globus MY, Ginsberg MD, Dietrich WD, et al: Substantia nigra lesion protects against ischemic damage in the striatum. *Neurosci Lett* 1987; 80:251–256
221. Globus MY: Role of dopamine in ischemic neuronal damage. *Stroke* 1989; 20:827–828
222. Maldonado J: *Delirium: Neurobiology, Characteristics and Management*. Edited by Fogel B, Greenberg D. Psychiatric Care of the Medical Patient. Oxford, Oxford University Press, 2013
223. Loneragan E, Britton AM, Luxenberg J, et al: Antipsychotics for delirium. *Cochrane Database Syst Rev* 2007;CD005594
224. Al-Qadheeb NS, O'Connor HH, White AC, et al: Antipsychotic prescribing patterns, and the factors and outcomes associated with their use, among patients requiring prolonged mechanical ventilation in the long-term acute care hospital setting. *Ann Pharmacother* 2013; 47:181–188
225. Larsen KA, Kelly SE, Stern TA, et al: Administration of olanzapine to prevent postoperative delirium in elderly joint-replacement patients: a randomized, controlled trial. *Psychosomatics* 2010; 51:409–418
226. Wang W, Li HL, Wang DX, et al: Haloperidol prophylaxis decreases delirium incidence in elderly patients after noncardiac surgery: a randomized controlled trial. *Crit Care Med* 2012; 40:731–739
227. van den Boogaard M, Schoonhoven L, van Achterberg T, et al: Haloperidol prophylaxis in critically ill patients with a high risk for delirium. *Crit Care* 2013; 17:R9
228. Teslyar P, Stock VM, Wilk CM, et al: Prophylaxis with antipsychotic medication reduces the risk of post-operative delirium in elderly patients: a meta-analysis. *Psychosomatics* 2013; 54:124–131
229. Meagher DJ, McLoughlin L, Leonard M, et al: What do we really know about the treatment of delirium with antipsychotics? Ten key issues for delirium pharmacotherapy. *Am J Geriatr Psychiatry* 2013; 21:1223–1238
230. Rothman S: Synaptic release of excitatory amino acid neurotransmitter mediates anoxic neuronal death. *J Neurosci* 1984; 4:1884–1891
231. Yamada KA, Rothman SM: Diazoxide blocks glutamate desensitization and prolongs excitatory postsynaptic currents in rat hippocampal neurons. *J Physiol* 1992; 458:409–423
232. Choi DW: Calcium-mediated neurotoxicity: relationship to specific channel types and role in ischemic damage. *Trends Neurosci* 1988; 11:465–469
233. Choi DW, Weiss JH, Koh JY, et al: Glutamate neurotoxicity, calcium, and zinc. *Ann N Y Acad Sci* 1989; 568:219–224

234. Katayama Y, Kawamata T, Tamura T, et al: Calcium-dependent glutamate release concomitant with massive potassium flux during cerebral ischemia in vivo. *Brain Res* 1991; 558:136–140
235. Broderick PA, Gibson GE: Dopamine and serotonin in rat striatum during in vivo hypoxic-hypoxia. *Metab Brain Dis* 1989; 4: 143–153
236. Gibson GE, Blass JP: Impaired synthesis of acetylcholine in brain accompanying mild hypoxia and hypoglycemia. *J Neurochem* 1976; 37–42
237. Beresin EV: Delirium in the elderly. *J Geriatr Psychiatry Neurol* 1988; 1:127–143
238. Bertrand N, Bralet J, Beley A: Post-ischemic regional changes in acetylcholine synthesis following transient forebrain ischemia in gerbils. *Neurochem Res* 1992; 17:321–325
239. Lopes da Silva FH, Witter MP, Boeijinga PH, et al: Anatomic organization and physiology of the limbic cortex. *Physiol Rev* 1990; 70:453–511
240. Felipe V, Hermenegildo C, Montoliu C, et al: Neurotoxicity of ammonia and glutamate: molecular mechanisms and prevention. *Neurotoxicology* 1998; 19:675–681
241. Huda A, Guze BH, Thomas A, et al: Clinical correlation of neuropsychological tests with 1H magnetic resonance spectroscopy in hepatic encephalopathy. *Psychosom Med* 1998; 60:550–556
242. Ahboucha S, Butterworth RF: Pathophysiology of hepatic encephalopathy: a new look at GABA from the molecular standpoint. *Metab Brain Dis* 2004; 19:331–343
243. Basile AS, Harrison PM, Hughes RD, et al: Relationship between plasma benzodiazepine receptor ligand concentrations and severity of hepatic encephalopathy. *Hepatology* 1994; 19: 112–121
244. Ferenci P, Schafer DF, Kleinberger G, et al: Serum levels of gamma-aminobutyric-acid-like activity in acute and chronic hepatocellular disease. *Lancet* 1983; 2:811–814
245. Ahboucha S, Talani G, Fanutza T, et al: Reduced brain levels of DHEAS in hepatic coma patients: significance for increased GABAergic tone in hepatic encephalopathy. *Neurochem Int* 2012; 61:48–53
246. Howard CD, Seifert CF: Flumazenil in the treatment of hepatic encephalopathy. *Ann Pharmacother* 1993; 27:46–48
247. Ahboucha S, Butterworth RF: The neurosteroid system: implication in the pathophysiology of hepatic encephalopathy. *Neurochem Int* 2008; 52:575–587
248. Maldonado J: An approach to the patient with substance use and abuse. *Med Clin North Am* 2010; 94:1169–1205; x–i
249. Akaike N, Shirasaki T, Yakushiji T: Quinolones and fenbufen interact with GABAA receptor in dissociated hippocampal cells of rat. *J Neurophysiol* 1991; 66:497–504
250. Marcantonio ER, Juarez G, Goldman L, et al: The relationship of postoperative delirium with psychoactive medications. *JAMA* 1994; 272:1518–1522
251. Kudoh A, Takase H, Takahira Y, et al: Postoperative confusion increases in elderly long-term benzodiazepine users. *Anesth Analg* 2004; 99:1674–1678
252. Tune LE, Bylsma FW: Benzodiazepine-induced and anticholinergic-induced delirium in the elderly. *Int Psychogeriatr* 1991; 3:397–408
253. Sanders RD, Maze M: Contribution of sedative-hypnotic agents to delirium via modulation of the sleep pathway. *Can J Anaesth* 2011; 58:149–156
254. Meuret P, Backman SB, Bonhomme V, et al: Physostigmine reverses propofol-induced unconsciousness and attenuation of the auditory steady state response and bispectral index in human volunteers. *Anesthesiology* 2000; 93:708–717
255. Pain L, Jeltsch H, Lehmann O, et al: Central cholinergic depletion induced by 192 IgG-saporin alleviates the sedative effects of propofol in rats. *Br J Anaesth* 2000; 85:869–873
256. Wang Y, Kikuchi T, Sakai M, et al: Age-related modifications of effects of ketamine and propofol on rat hippocampal acetylcholine release studied by in vivo brain microdialysis. *Acta Anaesthesiol Scand* 2000; 44:112–117
257. Moor E, DeBoer P, Westerink BH: GABA receptors and benzodiazepine binding sites modulate hippocampal acetylcholine release in vivo. *Eur J Pharmacol* 1998; 359:119–126
258. Ancelin ML, Artero S, Portet F, et al: Non-degenerative mild cognitive impairment in elderly people and use of anticholinergic drugs: longitudinal cohort study. *BMJ* 2006; 332:455–459
259. Whittington MA, Lambert JD, Little HJ: Increased NMDA receptor and calcium channel activity underlying ethanol withdrawal hyperexcitability. *Alcohol Alcohol* 1995; 30:105–114
260. Olofsson K, Alling C, Lundberg D, et al: Abolished circadian rhythm of melatonin secretion in sedated and artificially ventilated intensive care patients. *Acta Anaesthesiol Scand* 2004; 48:679–684
261. Bhardwaj A, Brannan T, Martinez-Tica J, et al: Ischemia in the dorsal hippocampus is associated with acute extracellular release of dopamine and norepinephrine. *J Neural Transm Gen Sect* 1990; 80:195–201
262. Borg S, Kvande H, Sedvall G: Central norepinephrine metabolism during alcohol intoxication in addicts and healthy volunteers. *Science* 1981; 213:1135–1137
263. Linnoila M, Mefford I, Nutt D, et al: NIH conference. Alcohol withdrawal and noradrenergic function. *Ann Intern Med* 1987; 107:875–889
264. Engberg G, Hajos M: Ethanol attenuates the response of locus coeruleus neurons to excitatory amino acid agonists in vivo. *Naunyn Schmied Arch Pharmacol* 1992; 345:222–226
265. Dodd PR, Beckmann AM, Davidson MS, et al: Glutamate-mediated transmission, alcohol, and alcoholism. *Neurochem Int* 2000; 37:509–533
266. Silberman Y, Matthews RT, Winder DG: A corticotropin releasing factor pathway for ethanol regulation of the ventral tegmental area in the bed nucleus of the stria terminalis. *J Neurosci* 2013; 33:950–960
267. Pandharipande PP, Pun BT, Herr DL, et al: Effect of sedation with dexmedetomidine vs lorazepam on acute brain dysfunction in mechanically ventilated patients: the MENDS randomized controlled trial. *JAMA* 2007; 298:2644–2653
268. Riker RR, Fraser GL: Altering intensive care sedation paradigms to improve patient outcomes. *Crit Care Clin* 2009; 25:527–538; viii–ix
269. Engelhardt K, Werner C, Eberspacher E, et al: The effect of the alpha 2-agonist dexmedetomidine and the N-methyl-D-aspartate antagonist S(+)-ketamine on the expression of apoptosis-regulating proteins after incomplete cerebral ischemia and reperfusion in rats. *Anesth Analg* 2003; 96:524–531
270. van der Mast RC, Fekkes D, Moleman P, et al: Is postoperative delirium related to reduced plasma tryptophan? *Lancet* 1991; 338:851–852
271. Zivin JA, Stashak J: The effect of ischemia on biogenic amine concentrations in the central nervous system. *Stroke* 1983; 14: 556–562
272. Blum D, Maldonado J, Meyer E, et al: Delirium following abrupt discontinuation of fluoxetine. *Clin Neurol Neurosurg* 2008; 110: 69–70
273. Sahakian BJ, Wurtman RJ, Barr JK, et al: Low tryptophan diet decreases brain serotonin and alters response to apomorphine. *Nature* 1979; 279:731–732

274. Robinson TN, Raeburn CD, Angles EM, et al: Low tryptophan levels are associated with postoperative delirium in the elderly. *Am J Surg* 2008; 196:670–674
275. Fernstrom JD, Wurtman RJ: Brain serotonin content: physiological regulation by plasma neutral amino acids. *Science* 1972; 178:414–416
276. Kaufman S: An evaluation of the possible neurotoxicity of metabolites of phenylalanine. *J Pediatr* 1989; 114:895–900
277. Flacker JM, Lipsitz LA: Large neutral amino acid changes and delirium in febrile elderly medical patients. *J Gerontol A Biol Sci Med Sci* 2000; 55:B249–B252; discussion B253–B254
278. van der Mast RC, van den Broek WW, Fekkes D, et al: Is delirium after cardiac surgery related to plasma amino acids and physical condition? *J Neuropsychiatry Clin Neurosci* 2000; 12:57–63
279. Mizock BA, Sabelli HC, Dubin A, et al: Septic encephalopathy. Evidence for altered phenylalanine metabolism and comparison with hepatic encephalopathy. *Arch Intern Med* 1990; 150: 443–449
280. Takezawa J, Taenaka N, Nishijima MK, et al: Amino acids and thiobarbituric acid reactive substances in cerebrospinal fluid and plasma of patients with septic encephalopathy. *Crit Care Med* 1983; 11:876–879
281. Saleem DM, Haider S, Khan MM, et al: Role of tryptophan in the pathogenesis of hepatic encephalopathy. *J Pakistan Med Assoc* 2008; 58:68–70
282. Sternbach H: The serotonin syndrome. *Am J Psychiatry* 1991; 148:705–713
283. Koponen HJ, Lepola U, Leinonen E: A long-term follow-up study of cerebrospinal fluid 5-hydroxyindoleacetic acid in delirium. *Eur Arch Psychiatry Clin Neurosci* 1994; 244:131–134
284. Bayindir O, Guden M, Akpinar B, et al: Ondansetron hydrochloride for the treatment of delirium after coronary artery surgery. *J Thorac Cardiovasc Surg* 2001; 121:176–177
285. Tagarakis GI, Voucharas C, Tzolaki F, et al: Ondansetron versus haloperidol for the treatment of postcardiotomy delirium: a prospective, randomized, double-blinded study. *J Cardiothorac Surg* 2012; 7:25
286. Schwartz JC, Arrang JM, Garbarg M, et al: Properties and roles of the three subclasses of histamine receptors in brain. *J Exp Biol* 1986; 124:203–224
287. White JM, Rumbold GR: Behavioural effects of histamine and its antagonists: a review. *Psychopharmacology (Berl)* 1988; 95: 1–14
288. Sitter H, Lorenz W, Klotter HJ, et al: Elevated plasma histamine concentration as a sensitive real-time parameter for distinct phases of surgical trauma: a tool for technology assessment. *Agents Actions* 1991; 33:203–207
289. Kral VA: Stress and mental disorders of the senium. *Med Serv J Can* 1962; 18:363–370
290. O'Keeffe ST, Devlin JG: Delirium and the dexamethasone suppression test in the elderly. *Neuropsychobiology* 1994; 30: 153–156
291. Dinkel K, Ogle WO, Sapolsky RM: Glucocorticoids and central nervous system inflammation. *J Neurovirol* 2002; 8:513–528
292. Sapolsky RM: Stress, glucocorticoids, and damage to the nervous system: the current state of confusion. *Stress* 1996; 1:1–19
293. Sapolsky RM: Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Arch Gen Psychiatry* 2000; 57: 925–935
294. Goosens KA, Sapolsky RM: Stress and glucocorticoid contributions to normal and pathological aging. In: *Brain Aging: Models, Methods, and Mechanisms*. Edited by Riddle DR. Boca Raton, FL, CRC Press, 2007
295. Munhoz CD, Sorrells SF, Caso JR, et al: Glucocorticoids exacerbate lipopolysaccharide-induced signaling in the frontal cortex and hippocampus in a dose-dependent manner. *J Neurosci* 2010; 30:13690–13698
296. Tombaugh GC, Yang SH, Swanson RA, et al: Glucocorticoids exacerbate hypoxic and hypoglycemic hippocampal injury in vitro: biochemical correlates and a role for astrocytes. *J Neurochem* 1992; 59:137–146
297. Basavaraju N, Phillips SL: Cortisol deficient state. A cause of reversible cognitive impairment and delirium in the elderly. *J Am Geriatr Soc* 1989; 37:49–51
298. Khalaf-Nazzal R, Francis F: Hippocampal development—old and new findings. *Neuroscience* 2013; 248C:225–242
299. Teeling JL, Felton LM, Deacon RM, et al: Sub-pyrogenic systemic inflammation impacts on brain and behavior, independent of cytokines. *Brain Behav Immun* 2007; 21:836–850
300. MacPherson A, Dinkel K, Sapolsky R: Glucocorticoids worsen excitotoxin-induced expression of pro-inflammatory cytokines in hippocampal cultures. *Exp Neurol* 2005; 194:376–383
301. Horner HC, Packan DR, Sapolsky RM: Glucocorticoids inhibit glucose transport in cultured hippocampal neurons and glia. *Neuroendocrinology* 1990; 52:57–64
302. McLay RN, Freeman SM, Harlan RE, et al: Aging in the hippocampus: interrelated actions of neurotrophins and glucocorticoids. *Neurosci Biobehav Rev* 1997; 21:615–629
303. Pavlides C, Watanabe Y, McEwen BS: Effects of glucocorticoids on hippocampal long-term potentiation. *Hippocampus* 1993; 3: 183–192
304. Sapolsky RM: Glucocorticoids, hippocampal damage and the glutamatergic synapse. *Prog Brain Res* 1990; 86:13–23
305. Seckl JR, Olsson T: Glucocorticoid hypersecretion and the age-impaired hippocampus: cause or effect? *J Endocrinol* 1995; 145:201–211
306. Stein-Behrens BA, Lin WJ, Sapolsky RM: Physiological elevations of glucocorticoids potentiate glutamate accumulation in the hippocampus. *J Neurochem* 1994; 63:596–602
307. Tombaugh GC, Sapolsky RM: Hippocampal glutamine synthetase: insensitivity to glucocorticoids and stress. *Am J Physiol* 1990; 258(5 Pt 1):E894–E897
308. Yusim A, Ajilore O, Bliss T, et al: Glucocorticoids exacerbate insult-induced declines in metabolism in selectively vulnerable hippocampal cell fields. *Brain Res* 2000; 870:109–117
309. Sapolsky RM, Pulsinelli WA: Glucocorticoids potentiate ischemic injury to neurons: therapeutic implications. *Science* 1985; 229:1397–1400
310. Olsson T: Activity in the hypothalamic-pituitary-adrenal axis and delirium. *Dement Geriatr Cogn Disord* 1999; 10:345–349
311. McIntosh TK, Bush HL, Yeston NS, et al: Beta-endorphin, cortisol and postoperative delirium: a preliminary report. *Psychoneuroendocrinology* 1985; 10:303–313
312. Robertsson B, Blennow K, Brane G, et al: Hyperactivity in the hypothalamic-pituitary-adrenal axis in demented patients with delirium. *Int Clin Psychopharmacol* 2001; 16:39–47
313. Fassbender K, Schmidt R, Mössner R, et al: Pattern of activation of the hypothalamic-pituitary-adrenal axis in acute stroke. Relation to acute confusional state, extent of brain damage, and clinical outcome. *Stroke* 1994; 25:1105–1108
314. Lipowski ZJ: Delirium (acute confusional states). *JAMA* 1987; 258:1789–1792
315. Berger M, Vollmann J, Hohagen F, et al: Sleep deprivation combined with consecutive sleep phase advance as a fast-acting therapy in depression: an open pilot trial in medicated and unmedicated patients. *Am J Psychiatry* 1997; 154:870–872

316. BaHammam A: Sleep in acute care units. *Sleep Breath* 2006; 10: 6–15
317. Walker MP, Stickgold R: Sleep, memory, and plasticity. *Annu Rev Psychol* 2006; 57:139–166
318. Yoo SS, Gujar N, Hu P, et al: The human emotional brain without sleep—a prefrontal amygdala disconnect. *Curr Biol* 2007; 17:R877–R878
319. Dinges DF: The state of sleep deprivation: from functional biology to functional consequences. *Sleep Med Rev* 2006; 10:303–305
320. Mistraletti G, Carloni E, Cigada M, et al: Sleep and delirium in the intensive care unit. *Minerva Anestesiol* 2008; 74:329–333
321. Ito H, Harada D, Hayashida K, et al: [Psychiatry and sleep disorders—delirium]. *Seishin Shinkeigaku Zasshi* 2006; 108: 1217–1221
322. Pandharipande P, Ely EW: Sedative and analgesic medications: risk factors for delirium and sleep disturbances in the critically ill. *Crit Care Clin* 2006; 22:313–327; vii
323. Johns MW, Large AA, Masterton JP, et al: Sleep and delirium after open heart surgery. *Br J Surg* 1974; 61:377–381
324. Mundigler G, Delle-Karh G, Koreny M, et al: Impaired circadian rhythm of melatonin secretion in sedated critically ill patients with severe sepsis. *Critical Care Med* 2002; 30:536–540
325. Helton MC, Gordon SH, Nunnery SL: The correlation between sleep deprivation and the intensive care unit syndrome. *Heart Lung* 1980; 9:464–468
326. Reiter RJ: Oxidative damage in the central nervous system: protection by melatonin. *Prog Neurobiol* 1998; 56:359–384
327. Brzezinski A: Melatonin in humans. *N Engl J Med* 1997; 336: 186–195
328. Verster GC: Melatonin and its agonists, circadian rhythms and psychiatry. *Afr J Psychiatry* 2009; 12:42–46
329. Quiros I, Mayo JC, Garcia-Suarez O, et al: Melatonin prevents glucocorticoid inhibition of cell proliferation and toxicity in hippocampal cells by reducing glucocorticoid receptor nuclear translocation. *J Steroid Biochem Mol Biol* 2008; 110:116–124
330. Reiter RJ: Pineal melatonin: cell biology of its synthesis and of its physiological interactions. *Endocr Rev* 1991; 12:151–180
331. Cervantes M, Morali G, Letechipia-Vallejo G: Melatonin and ischemia-reperfusion injury of the brain. *J Pineal Res* 2008; 45: 1–7
332. de Jonghe A, Korevaar JC, van Munster BC, et al: Effectiveness of melatonin treatment on circadian rhythm disturbances in dementia. Are there implications for delirium? A systematic review. *Int J Geriatr Psychiatry* 2010; 25:1201–1208
333. Gunasingh MJ, Philip JE, Ashok BS, et al: Melatonin prevents amyloid protofibrillar induced oxidative imbalance and biogenic amine catabolism. *Life Sci* 2008; 83:96–102
334. Ionov M, Burchell V, Klajnert B, et al: Mechanism of neuroprotection of melatonin against beta-amyloid neurotoxicity. *Neuroscience* 2011; 180:229–237
335. Reiter RJ: Melatonin synthesis: multiplicity of regulation. *Adv Exp Med Biol* 1991; 294:149–158
336. Sun FY, Lin X, Mao LZ, et al: Neuroprotection by melatonin against ischemic neuronal injury associated with modulation of DNA damage and repair in the rat following a transient cerebral ischemia. *J Pineal Res* 2002; 33:48–56
337. Poeggeler B, Miravalle L, Zagorski MG, et al: Melatonin reverses the profibrillogenic activity of apolipoprotein E4 on the Alzheimer amyloid Abeta peptide. *Biochemistry* 2001; 40: 14995–15001
338. Paula-Lima AC, Louzada PR, De Mello FG, et al: Neuroprotection against Abeta and glutamate toxicity by melatonin: are GABA receptors involved? *Neurotox Res* 2003; 5:323–327
339. Pappolla MA, Sos M, Omar RA, et al: Melatonin prevents death of neuroblastoma cells exposed to the Alzheimer amyloid peptide. *J Neurosci* 1997; 17:1683–1690
340. Pappolla MA, Simovich MJ, Bryant-Thomas T, et al: The neuroprotective activities of melatonin against the Alzheimer beta-protein are not mediated by melatonin membrane receptors. *J Pineal Res* 2002; 32:135–142
341. Pappolla M, Bozner P, Soto C, et al: Inhibition of Alzheimer beta-fibrillogenesis by melatonin. *J Biol Chem* 1998; 273:7185–7188
342. Matsubara E, Bryant-Thomas T, Pacheco Quinto J, et al: Melatonin increases survival and inhibits oxidative and amyloid pathology in a transgenic model of Alzheimer's disease. *J Neurochem* 2003; 85:1101–1108
343. Clapp-Lilly KL, Smith MA, Perry G, et al: Melatonin acts as antioxidant and pro-oxidant in an organotypic slice culture model of Alzheimer's disease. *Neuroreport* 2001; 12:1277–1280
344. Ozturk L, Pelin Z, Karadeniz D, et al: Effects of 48 hours sleep deprivation on human immune profile. *Sleep Res Online* 1999; 2:107–111
345. Irwin M, McClintick J, Costlow C, et al: Partial night sleep deprivation reduces natural killer and cellular immune responses in humans. *FASEB J* 1996; 10:643–653
346. Krueger JM, Majde JA: Humoral links between sleep and the immune system: research issues. *Ann N Y Acad Sci* 2003; 992: 9–20
347. Spiegel K, Leproult R, Van Cauter E: Impact of sleep debt on metabolic and endocrine function. *Lancet* 1999; 354: 1435–1439
348. Miyazaki T, Kuwano H, Kato H, et al: Correlation between serum melatonin circadian rhythm and intensive care unit psychosis after thoracic esophagectomy. *Surgery* 2003; 133: 662–668
349. Bourne RS, Mills GH, Minelli C: Melatonin therapy to improve nocturnal sleep in critically ill patients: encouraging results from a small randomised controlled trial. *Crit Care* 2008; 12:R52
350. Sultan SS: Assessment of role of perioperative melatonin in prevention and treatment of postoperative delirium after hip arthroplasty under spinal anesthesia in the elderly. *Saudi J Anaesth* 2010; 4:169–173
351. de Jonghe A, van Munster BC, van Oosten HE, et al: The effects of melatonin versus placebo on delirium in hip fracture patients: study protocol of a randomised, placebo-controlled, double blind trial. *BMC Geriatr* 2011; 11:34
352. Matthews CD, Guerin MV, Wang X: Human plasma melatonin and urinary 6-sulphatoxy melatonin: studies in natural annual photoperiod and in extended darkness. *Clin Endocrinol* 1991; 35:21–27
353. Balan S, Leibovitz A, Zila SO, et al: The relation between the clinical subtypes of delirium and the urinary level of 6-SMT. *J Neuropsychiatry Clin Neurosci* 2003; 15:363–366
354. Ross CA: CNS arousal systems: possible role in delirium. *Int Psychogeriatr* 1991; 3:353–371
355. Sarter M, Bruno JP: Cortical cholinergic inputs mediating arousal, attentional processing and dreaming: differential afferent regulation of the basal forebrain by telencephalic and brainstem afferents. *Neuroscience* 2000; 95:933–952
356. Tremblay N, Warren RA, Dykes RW: Electrophysiological studies of acetylcholine and the role of the basal forebrain in the somatosensory cortex of the cat. II. Cortical neurons excited by somatic stimuli. *J Neurophysiol* 1990; 64:1212–1222
357. Sanders RD: Hypothesis for the pathophysiology of delirium: role of baseline brain network connectivity and changes in inhibitory tone. *Med Hypoth* 2011; 77:140–143

358. Trzepacz PT, Scibassi RJ, Van Thiel DH: Delirium: a subcortical phenomenon? *J Neuropsychiatry Clin Neurosci* 1989; 1: 283–290
359. Choi SH, Lee H, Chung TS, et al: Neural network functional connectivity during and after an episode of delirium. *Am J Psychiatry* 2012; 169:498–507
360. Zecharia AY, et al: The involvement of hypothalamic sleep pathways in general anesthesia: testing the hypothesis using the GABAA receptor beta3N265M knock-in mouse. *J Neurosci* 2009; 29:2177–2187
361. Metnitz PG, Moreno RP, Almeida E, et al: SAPS 3—from evaluation of the patient to evaluation of the intensive care unit. Part 1: Objectives, methods and cohort description. *Intensive Care Med* 2005; 31:1336–1344
362. Xiang YY, Wang S, Liu M, et al: A GABAergic system in airway epithelium is essential for mucus overproduction in asthma. *Nat Med* 2007; 13:862–867
363. Serantes R, Arnalich F, Figueroa M, et al: Interleukin-1beta enhances GABAA receptor cell-surface expression by a phosphatidylinositol 3-kinase/Akt pathway: relevance to sepsis-associated encephalopathy. *J Biol Chem* 2006; 281:14632–14643
364. Gaykema RP, Goehler LE: Lipopolysaccharide challenge-induced suppression of Fos in hypothalamic orexin neurons: their potential role in sickness behavior. *Brain Behav Immun* 2009; 23: 926–930
365. Andrews-Hanna JR, Snyder AZ, Vincent JL, et al: Disruption of large-scale brain systems in advanced aging. *Neuron* 2007; 56: 924–935
366. Yankner BA, Lu T, Loerch P: The aging brain. *Annu Rev Pathol* 2008; 3:41–66
367. Shioiri A, Kurumaji A, Takeuchi T, et al: White matter abnormalities as a risk factor for postoperative delirium revealed by diffusion tensor imaging. *Am J Geriatr Psychiatry* 2010; 18: 743–753
368. Hatano Y, Narumoto J, Shibata K, et al: White-matter hyperintensities predict delirium after cardiac surgery. *Am J Geriatr Psychiatry* 2013; 21:938–945
369. Bishop NA, Lu T, Yankner BA: Neural mechanisms of ageing and cognitive decline. *Nature* 2010; 464:529–535
370. Loerch PM, Lu T, Dakin KA, et al: Evolution of the aging brain transcriptome and synaptic regulation. *PLoS One* 2008; 3:e3329
371. Stanley EM, Fadel J: Aging-related deficits in orexin/hypocretin modulation of the septohippocampal cholinergic system. *Synapse* 2012; 66:445–452
372. Matsumura T, Nakayama M, Nomura A, et al: Age-related changes in plasma orexin-A concentrations. *Exp Gerontol* 2002; 37:1127–1130
373. Kessler BA, Stanley EM, Frederick-Duus D, et al: Age-related loss of orexin/hypocretin neurons. *Neuroscience* 2011; 178:82–88
374. Brownell SE, Conti B: Age- and gender-specific changes of hypocretin immunopositive neurons in C57BL/6 mice. *Neurosci Lett* 2010; 472:29–32
375. Alkire MT: Loss of effective connectivity during general anesthesia. *Int Anesthesiol Clin* 2008; 46:55–73
376. Alkire MT, Hudetz AG, Tononi G: Consciousness and anesthesia. *Science* 2008; 322:876–880
377. Marques-Deak A, Cizza G, Sternberg E: Brain-immune interactions and disease susceptibility. *Mol Psychiatry* 2005; 10:239–250
378. Tracey KJ: Reflex control of immunity. *Nature reviews. Immunology* 2009; 9:418–428
379. Harukuni I, Bhardwaj A: Mechanisms of brain injury after global cerebral ischemia. *Neurol Clin* 2006; 24:1–21
380. Brooke SM, Sapolsky RM: Effects of glucocorticoids in the gp120-induced inhibition of glutamate uptake in hippocampal cultures. *Brain Res* 2003; 972:137–141
381. Moghaddam B, Bolinao ML, Stein-Behrens B, et al: Glucocorticoids mediate the stress-induced extracellular accumulation of glutamate. *Brain Res* 1994; 655:251–254
382. Liston C, Gan WB: Glucocorticoids are critical regulators of dendritic spine development and plasticity in vivo. *Proc Natl Acad Sci USA* 2011; 108:16074–16079
383. McIntosh LJ, Sapolsky RM: Glucocorticoids may enhance oxygen radical-mediated neurotoxicity. *Neurotoxicology* 1996; 17: 873–882
384. McIntosh LJ, Sapolsky RM: Glucocorticoids increase the accumulation of reactive oxygen species and enhance adriamycin-induced toxicity in neuronal culture. *Exp Neurol* 1996; 141: 201–206
385. Crossin KL, Tai MH, Krushel LA, et al: Glucocorticoid receptor pathways are involved in the inhibition of astrocyte proliferation. *Proc Natl Acad Sci USA* 1997; 94:2687–2692
386. Suri D, Vaidya VA: Glucocorticoid regulation of brain-derived neurotrophic factor: relevance to hippocampal structural and functional plasticity. *Neuroscience* 2013; 239:196–213
387. Sapolsky RM: Glucocorticoids, stress, and their adverse neurological effects: relevance to aging. *Exp Gerontol* 1999; 34:721–732
388. Schoenfeld TJ, Gould E: Stress, stress hormones, and adult neurogenesis. *Exp Neurol* 2012; 233:12–21