

Time Series Analysis

Basics of Time Series Analysis: Data Example

Nicoleta Serban, Ph.D.

Professor

Stewart School of Industrial and Systems Engineering

Trend and Seasonality Estimation of
ED Volume Time Series

About This Lesson



Trend Estimation

Equally spaced time points

```
time.pts = c(1:length(Volume))
```

```
time.pts = c(time.pts - min(time.pts))/max(time.pts)
```

Local Polynomial Trend Estimation

```
loc.fit = loess(Volume.tr~time.pts)
```

```
vol.fit.loc = fitted(loc.fit)
```

Splines Trend Estimation

```
library(mgcv)
```

```
gam.fit = gam(Volume.tr~s(time.pts))
```

```
vol.fit.gam = fitted(gam.fit)
```


Is there a trend?

```
plot(dates, Volume.tr, ylab="ED Volume")
```

```
lines(dates, vol.fit.loc, lwd=2, col="brown")
```

```
lines(dates, vol.fit.gam, lwd=2, col="red")
```

Trend Estimation



Trend Estimation (cont'd)

Parametric coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	12.85769	0.02441	526.7	<2e-16


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05

Approximate significance of smooth terms:

	edf	Ref.df	F	p-value
s(time.pts)	8.628	8.96	93.39	<2e-16***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05

R-sq.(adj) = 0.296 Deviance explained = 29.9%



Smooth trend is statistically significant




29.6% of variability explained

Trend and Seasonality Estimation

Model Trend + Monthly Seasonality

```
month = as.factor(format(dates,"%b"))  
gam.fit.seastr.1 = gam(Volume.tr~s(time.pts)+month-1)  
summary(gam.fit.seastr)  
vol.fit.gam.seastr.1 = fitted(gam.fit.seastr.1)  
ggplot(edvoldata, aes(dates, sqrt(Volume+3/8))) + geom_line() + xlab("Time") +  
ylab("Transformed Daily ED Volume")  
lines(dates,vol.fit.gam.seastr.1,lwd=2,col="red")
```

Trend and Seasonality Estimation



Trend and Seasonality Estimation (cont'd)


Add day-of-the-week seasonality

```
week = as.factor(weekdays(dates))  
gam.fit.seastr.2 = gam(Volume.tr~s(time.pts)+month+week)  
summary(gam.fit.seastr.2)  
vol.fit.gam.seastr.2 = fitted(gam.fit.seastr.2)
```

Compare the two fits

```
ggplot(edvoldata, aes(dates, vol.fit.gam.seastr.2)) + geom_line() + xlab("Time") +  
ylab("Seasonality and Trend: Daily ED Volume")  
lines(dates, vol.fit.gam.seastr.1, lwd=2, col="red")
```


Trend and Seasonality Estimation (cont'd)



Trend and Seasonality Estimation (cont'd)

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	12.77772	0.07383	173.063	< 2e-16
monthAug	-0.50061	0.08852	-5.655	1.79e-08
monthDec	0.93273	0.08791	10.611	< 2e-16
monthFeb	0.33589	0.08509	3.948	8.17e-05
monthJan	-0.59767	0.08365	-7.145	1.26e-12
monthJul	-1.53530	0.08831	-17.385	< 2e-16
monthJun	-1.00553	0.08880	-11.324	< 2e-16
monthMar	0.61815	0.08272	7.472	1.18e-13
monthMay	0.05271	0.08272	0.637	0.52403
monthNov	0.94163	0.08900	10.580	< 2e-16
monthOct	0.41557	0.08849	4.696	2.84e-06
monthSep	0.22391	0.08935	2.506	0.01229
weekMonday	0.57169	0.06647	8.601	< 2e-16
weekSaturday	0.04589	0.06641	0.691	0.48967
weekSunday	0.17538	0.06641	2.641	0.00834
weekThursday	-0.16031	0.06647	-2.412	0.01597
weekTuesday	0.08099	0.06647	1.218	0.22323
weekWednesday	-0.11050	0.06647	-1.662	0.09662

Most regression coefficients are statistically significant

Some regression coefficients are statistically significant

Approximate significance of smooth terms:
edf Ref.df F p-value
s(time.pts) 8.792 8.987 151.8 <2e-16 ***

Smooth trend is statistically significant

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05

R-sq.(adj) = 0.627 Deviance explained = 63.2%
GCV = 0.63263 Scale est. = 0.62405 n = 1977

62.7% of variability explained

Trend and Seasonality Estimation (cont'd)

Does the addition of seasonality of day of the week adds predictive power?

```
lm.fit.seastr.1 = lm(Volume.tr~month)
```

```
lm.fit.seastr.2 = lm(Volume.tr~month+week)
```

```
anova(lm.fit.seastr.1,lm.fit.seastr.2)
```

Analysis of Variance Table

Model 1: Volume.tr ~ month

Model 2: Volume.tr ~ month + week

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	1965	2169.9				
2	1959	2071.1	6	98.826	15.58	< 2.2e-16




The seasonality due to day of the week improves the prediction power of the model

Seasonality vs Trend

Compare with & without trend

```
ggplot(edvoldata, aes(dates, vol.fit.gam.seastr.2)) + geom_line() + xlab("Time") +  
ylab("Seasonality and Trend: Daily ED Volume")  
lines(dates, vol.fit.lm.seastr.2, lwd=2, col="blue")  
lines(dates, vol.fit.gam, lwd=2, col="red")
```

Seasonality vs Trend



Summary

