

# Copper [Cu]

Nathaniel Yomogida, SPT

Chloë Kerstein, SPT

## Table of contents

|  |          |
|--|----------|
| <b>1 Overview</b>                      | <b>1</b> |
| <b>2 Macro clinical perspective</b>    | <b>2</b> |
| <b>3 Copper metabolism</b>             | <b>2</b> |
| 3.1 Copper absorption . . . . .        | 2        |
| 3.2 Copper Bioavailability . . . . .   | 2        |
| <b>4 Function</b>                      | <b>2</b> |
| 4.1 Metabolic Functions . . . . .      | 3        |
| 4.2 Iron utilization . . . . .         | 3        |
| 4.3 Vascular function . . . . .        | 3        |
| 4.4 Cardiac function . . . . .         | 3        |
| 4.5 Skeleton . . . . .                 | 3        |
| 4.6 Oxidative damage defense . . . . . | 3        |
| 4.7 Myelination . . . . .              | 3        |
| 4.8 Reproduction . . . . .             | 3        |
| 4.9 Hormone synthesis . . . . .        | 4        |
| <b>5 Multi-Copper Oxidases (MCOs)</b>  | <b>4</b> |
| 5.1 Ceruloplasmin . . . . .            | 4        |
| 5.2 Hephaestin . . . . .               | 4        |
| 5.3 Zyklopen . . . . .                 | 4        |
| <b>References</b>                      | <b>5</b> |

## 1 Overview

- Copper(Cu) is the 26th element in abundance in the crust of the earth<sup>1</sup>.

- 29th periodic element
- 2 stable and 9 radioactive isotopes<sup>1</sup>

## 2 Macro clinical perspective

- Cu is only needed in only trace amounts in humans. Total Cu in the body is >100 mg<sup>1</sup>. The skeleton and muscles account for 1/2 of the Cu in the body<sup>1</sup>.
- Most Cu in the body exists copper almost always exists in biological systems bound to proteins<sup>1</sup>. Free copper in cells and in the body is extremely low<sup>1</sup>
- High [Cu] is related to metabolic activity of organs<sup>1</sup>. Kidney and liver have the highest [Cu], followed by the brain (~5 µg/g), then the heart<sup>1</sup>.

## 3 Copper metabolism

### 3.1 Copper absorption

- Copper absorption is considerably higher than for that of other trace elements (~55-75%)<sup>1</sup>
- Relative amount of copper in the diet seems to be inversely correlated with percent intestinal absorption<sup>1</sup>. Percent absorption increases during states of deficiency<sup>1</sup>.
- Copper absorption occurs mainly in the upper small intestine<sup>1</sup>

### 3.2 Copper Bioavailability

- Dietary factors (i.e. iron, vitamin C, and Zinc) have been reported to exert adverse effects on the bioavailability of copper<sup>1</sup>
- Other factors impact Copper's bioavailability, such as lead poisoning, hemochromatosis, and excessive ingestion of soft drinks<sup>1</sup>

## 4 Function

Through copper's enzymes (Multi-copper oxidases) copper has the unique ability to convert O<sub>2</sub> into H<sub>2</sub>O without producing oxidative "exhaust"<sup>2</sup>. This allows our bodies to manage O<sub>2</sub> without being negatively affected by its toxic and highly reactive nature<sup>2</sup>

## 4.1 Metabolic Functions

Copper plays a crucial role in energy transformation in the body<sup>1</sup>. Copper impacts this process by acting as a cofactor for cytochrome c oxidase (Terminal enzyme in the electron transport chain)<sup>1</sup>

## 4.2 Iron utilization

Copper is important in the normal utilization of iron in the body<sup>1</sup>. From Intestinal iron absorption, iron release from stores (e.g. in macrophages of liver and spleen), iron incorporation into hemoglobin, and even preventing anemia<sup>1</sup>.

## 4.3 Vascular function

- Blood coagulation is assisted by copper<sup>1</sup>.
- Blood pressure control<sup>1</sup>
- Cross-linking of connective tissues in arteries<sup>1</sup>

## 4.4 Cardiac function

- Cross-linking of CT in heart<sup>1</sup>

## 4.5 Skeleton

- Cross-linking of CT in bones<sup>1</sup>

## 4.6 Oxidative damage defense

- Defense against oxidative damage<sup>1</sup>

## 4.7 Myelination

- Myelination of brain and spinal cord<sup>1</sup>

## 4.8 Reproduction

- Copper has a function in reproduction<sup>1</sup>

## 4.9 Hormone synthesis

- Copper plays a role in hormone synthesis<sup>1</sup>

## 5 Multi-Copper Oxidases (MCOs)

3 Multi-Copper oxidases have been detected in humans:<sup>2</sup>

1. Ceruloplasmin<sup>2</sup>
2. Hephaestin<sup>2</sup>
3. Zyklopen<sup>2</sup>

All 3 of these enzymes have a high specificity towards iron with the resulting ferroxidase activity being associated with ferroportin (the only known iron exporter protein in humans.)<sup>2</sup>

Ferroportin exports iron as  $\text{Fe}^{2+}$ , but transferrin, the major iron transporter protein of blood, can bind only  $\text{Fe}^{3+}$  effectively. Iron oxidation in enterocytes is mediated mainly by hephaestin thus allowing dietary iron to enter the bloodstream<sup>2</sup>

### 5.1 Ceruloplasmin

Function: Release of iron from the liver relies on ferroportin and the ferroxidase activity of ceruloplasmin which is found in blood in a soluble form<sup>2</sup>

### 5.2 Hephaestin

Iron oxidation in enterocytes is mediated mainly by hephaestin thus allowing dietary iron to enter the bloodstream<sup>2</sup>

### 5.3 Zyklopen

Zyklopen is involved in iron efflux from placental trophoblasts during iron transfer from mother to fetus<sup>2</sup>

## References

1. Collins JF, Klevay LM. Copper. *Advances in Nutrition (Bethesda, Md)*. 2011;2(6):520-522. doi:[10.3945/an.111.001222](https://doi.org/10.3945/an.111.001222)
2. Vashchenko G, MacGillivray RTA. Multi-copper oxidases and human iron metabolism. *Nutrients*. 2013;5(7):2289-2313. doi:[10.3390/nu5072289](https://doi.org/10.3390/nu5072289)