

Formula for difference of nth powers

We will prove the following identity for all $x, c \in \mathbb{R}, n \in \mathbb{N}$:

$$x^n - c^n = (x - c) \sum_{i=1}^n x^{n-i} c^{i-1}$$

Proof. By induction on n .

$$\text{If } n = 1, \text{ we have } x - c = (x - c) \sum_{i=1}^1 x^{1-i} c^{i-1} = (x - c)(1)$$

Now for some fixed natural number, k , assume:

$$x^{k-1} - c^{k-1} = (x - c) \sum_{i=1}^{k-1} x^{k-1-i} c^{i-1}$$

Multiply both sides of the equation by $x + c$:

$$(x + c)(x^{k-1} - c^{k-1}) = (x + c)(x - c) \sum_{i=1}^{k-1} x^{k-1-i} c^{i-1}$$

Distribute on the left side:

$$x^k - c^k - xc^{k-1} + x^{k-1}c = (x + c)(x - c) \sum_{i=1}^{k-1} x^{k-1-i} c^{i-1}$$

Now distribute the $x - c$ term on the right side:

$$\dots = (x + c) \left(\sum_{i=1}^{k-1} x^{k-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-1-i} c^i \right)$$

And distribute the $x + c$ term on the right side:

$$\dots = \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-i} c^i + \sum_{i=1}^{k-1} x^{k-i} c^i - \sum_{i=1}^{k-1} x^{k-1-i} c^{i+1}$$

Now extract an xc from the third sum and a c^2 from the fourth sum:

$$\dots = \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-i} c^i + (xc) \left(\sum_{i=1}^{k-1} x^{k-1-i} c^{i-1} \right) - (c^2) \left(\sum_{i=1}^{k-1} x^{k-1-i} c^{i-1} \right)$$

The third and fourth sums are now identical, so we can combine them:

$$\dots = \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-i} c^i + c(x - c) \left(\sum_{i=1}^{k-1} x^{k-1-i} c^{i-1} \right)$$

Which, by the induction hypothesis, is equivalent to:

$$\dots = \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-i} c^i + c(x^{k-1} - c^{k-1})$$

Now we bring the left side of the equation back:

$$x^k - c^k - xc^{k-1} + x^{k-1}c = \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-i} c^i + c(x^{k-1} - c^{k-1})$$

And move some terms over to the right, to isolate the $x^k - c^k$:

$$x^k - c^k = xc^{k-1} - x^{k-1}c + \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1} - \sum_{i=1}^{k-1} x^{k-i} c^i + c(x^{k-1} - c^{k-1})$$

Now we turn our attention to the two remaining sums.

For concision and clarity, we make the following designations:

$$\text{Let } \alpha = \sum_{i=1}^{k-1} x^{k+1-i} c^{i-1}$$

$$\text{Let } \beta = \sum_{i=1}^{k-1} x^{k-i} c^i$$

So that we have:

$$x^k - c^k = xc^{k-1} - x^{k-1}c + \alpha - \beta + c(x^{k-1} - c^{k-1})$$

We also introduce two new terms:

$$\text{Let } \gamma = \sum_{i=1}^k x^{k+1-i} c^{i-1}$$

$$\text{Let } \delta = \sum_{i=1}^k x^{k-i} c^i$$

Observe that $\alpha = \gamma - xc^{k-1}$ and $\beta = \delta - c^k$.

We substitute these results into our equation:

$$x^k - c^k = xc^{k-1} - x^{k-1}c + \gamma - xc^{k-1} - \delta - c^k + c(x^{k-1} - c^{k-1})$$

Simplifying and rearranging terms:

$$x^k - c^k = \gamma - \delta - x^{k-1}c - c^k + c(x^{k-1} - c^{k-1})$$

$$x^k - c^k = \gamma - \delta - c(x^{k-1} - c^{k-1}) + c(x^{k-1} - c^{k-1})$$

$$x^k - c^k = \gamma - \delta$$

We replace γ and δ with their definitions:

$$x^k - c^k = \sum_{i=1}^k x^{k+1-i} c^{i-1} - \sum_{i=1}^k x^{k-i} c^i$$

And extract an x and c from the sums to make them identical:

$$x^k - c^k = (x) \sum_{i=1}^k x^{k-i} c^{i-1} - (c) \sum_{i=1}^k x^{k-i} c^{i-1}$$

A final combining of like terms concludes the proof by induction:

$$x^k - c^k = (x - c) \sum_{i=1}^k x^{k-i} c^{i-1}$$

□