

Note on Matrix functions

Nathan Rousselot

August 19, 2023

1 Introduction

In this document we introduce the notion of matrix functions. Say we have a function $f : \mathbb{C} \rightarrow \mathbb{C}$, then we can define the function f on a matrix \mathbf{A} as follows: $f : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$. In the following, we will first provide some theoretical background on matrix functions. Then, we will describe the numerical issues coming with manipulating matrix functions. Finally, we will provide algorithm for efficient computation of matrix functions.

2 Theoretical background

2.1 Natural Definition

Polynomial functions

Let $p : \mathbb{C} \rightarrow \mathbb{C}$ be a polynomial function of degree d :

$$p(t) = \sum_{k=0}^d c_k t^k \quad (2.1)$$

Then, considering a matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$, and posing $\mathbf{A}^0 = I_n$, we can define the polynomial function $p : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$ on a matrix \mathbf{A} as follows:

$$p(\mathbf{A}) = \sum_{k=0}^d c_k \mathbf{A}^k \quad (2.2)$$

Rational functions

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a rational function of the form:

$$f(t) := \frac{p(t)}{q(t)} \quad (2.3)$$

It is not immediate how one would approach this function with a matrix. We want to define

$$f(\mathbf{A}) := q(\mathbf{A})^{-1} p(\mathbf{A}) \quad (2.4)$$

However, this is not well defined if $q(\mathbf{A})$ is singular. In other words, we need to make sure that $q(\mathbf{A})$ is invertible. This is the case if and only if $q(\lambda) \neq 0$ for all eigenvalues λ of \mathbf{A} . This is a very strong condition, and it is not always possible to find a rational function f such that $q(\lambda) \neq 0$ for all eigenvalues λ of \mathbf{A} . We note that a choice in notation has been made in equation 2.4, the other notation is still valid.

Lemma 1. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$, and let $p : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$ and $q : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$, two matrix polynomials. Then,

$$q(\mathbf{A})p(\mathbf{A}) = p(\mathbf{A})q(\mathbf{A}) \quad (2.5)$$

Proof. Let $f(\mathbf{A}) := q(\mathbf{A})p(\mathbf{A})$, and $g(\mathbf{A}) = p(\mathbf{A})q(\mathbf{A})$. Then

$$\begin{aligned} f(\mathbf{A}) &= \left(\sum_{k=0}^d c_k \mathbf{A}^k \right) \left(\sum_{j=0}^m b_j \mathbf{A}^j \right) \\ &= \sum_{k=0}^d \sum_{j=0}^m c_k b_j \mathbf{A}^{j+k} \\ &= \sum_{j=0}^m \sum_{k=0}^d b_j c_k \mathbf{A}^{k+j} \\ &= \left(\sum_{j=0}^m b_j \mathbf{A}^j \right) \left(\sum_{k=0}^d c_k \mathbf{A}^k \right) = g(\mathbf{A}) \end{aligned}$$

□

Theorem 1. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$, and let $p : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$ and $q : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$, two matrix polynomials such that $q(\mathbf{A})$ is non-singular. Then,

$$q(\mathbf{A})^{-1}p(\mathbf{A}) = p(\mathbf{A})q(\mathbf{A})^{-1} \quad (2.6)$$

Proof. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a matrix such that $q(\mathbf{A})$ is non-singular

$$q(\mathbf{A})^{-1}p(\mathbf{A}) = q(\mathbf{A})^{-1}p(\mathbf{A})q(\mathbf{A})q(\mathbf{A})^{-1}$$

Using Lemma 1

$$\begin{aligned} q(\mathbf{A})^{-1}p(\mathbf{A})q(\mathbf{A})q(\mathbf{A})^{-1} &= q(\mathbf{A})^{-1}q(\mathbf{A})p(\mathbf{A})q(\mathbf{A})^{-1} \\ &= p(\mathbf{A})q(\mathbf{A})^{-1} \end{aligned}$$

□

Power Series

Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a function that can be expressed as a power series:

$$f(t) = \sum_{k=0}^{\infty} c_k t^k \quad (2.7)$$

Then, considering a matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$, and posing $\mathbf{A}^0 = I_n$, we can define the power series function $f : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$ on a matrix \mathbf{A} as follows:

$$f(\mathbf{A}) = \sum_{k=0}^{\infty} c_k \mathbf{A}^k \quad (2.8)$$

In the scalar case, we know that the power series converges if $|t| < r$, where r is the radius of convergence. Obviously this translates in the matrix power series

Theorem 2. Let $f : \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}$ be a matrix power series. Then, the series converges if and only if $\rho(\mathbf{A}) < r$, where $\rho(\mathbf{A})$ is the spectral radius of \mathbf{A} , and r is the radius of convergence of the scalar power series.

Proof is provided in Frommer and Simoncini 2008. In the case of a finite-order Laurent Series, *i.e.*:

$$f(t) = \sum_{k=-d}^d c_k t^k \quad (2.9)$$

For the matrix case, we need to ensure convergence (similarly to power series), but also ensure existence of the inverse, as Laurent series do have negative powers. If both of those conditions are satisfied, we can write the Laurent series as a matrix function:

$$f(\mathbf{A}) = \sum_{k=-d}^d c_k \mathbf{A}^k \quad (2.10)$$

2.2 Spectrum-Based Definition

Diagonalizable Matrices

Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a diagonalizable matrix. That means that there exists a matrix $\mathbf{V} \in \mathbb{C}^{n \times n}$ such that \mathbf{V} is invertible, and $\mathbf{A} = \mathbf{V}\mathbf{\Lambda}\mathbf{V}^{-1}$, where $\mathbf{\Lambda}$ is a diagonal matrix containing the eigenvalues of \mathbf{A} :

$$\mathbf{\Lambda} = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{bmatrix} \quad (2.11)$$

Theorem 3. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a diagonalizable matrix. Then we can define the function $f(\mathbf{A})$ as:

$$f(\mathbf{A}) := \mathbf{V}f(\mathbf{\Lambda})\mathbf{V}^{-1} \quad (2.12)$$

with

$$f(\mathbf{\Lambda}) = \begin{bmatrix} f(\lambda_1) & 0 & \cdots & 0 \\ 0 & f(\lambda_2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & f(\lambda_n) \end{bmatrix} \quad (2.13)$$

This property is very handy, as it allows us to compute matrix functions by simply applying the function to the eigenvalues of the matrix. Computationally, this avoids inverting matrices, and lots of matrix products. However, this property is only valid for diagonalizable matrices. This property puts constraints on $f(\mathbf{A})$, as its eigenvectors must form a basis \mathbf{F}^n . Another more practical constraint, that is sufficient but not necessary, is if \mathbf{A} is a full rank matrix, then it is diagonalizable.

Defective Matrices

In some cases, the matrix \mathbf{A} is not diagonalizable, that means the sum of the dimensions of the eigenspaces is less than n , we call that a *Defective Matrix*. In that case, we can generalize the principle of diagonalization using the Jordan canonical form of \mathbf{A} :

$$\mathbf{A} = \mathbf{V}\mathbf{J}\mathbf{V}^{-1} \quad (2.14)$$

where \mathbf{J} is a Jordan matrix, and \mathbf{V} is a matrix containing the generalized eigenvectors of \mathbf{A} . The Jordan matrix is a block diagonal matrix, where each block is a Jordan block. A Jordan block is a matrix of the form:

$$\mathbf{J}_k(\lambda) = \begin{bmatrix} \lambda & 1 & 0 & \cdots & 0 \\ 0 & \lambda & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda & 1 \\ 0 & 0 & \cdots & 0 & \lambda \end{bmatrix} \quad (2.15)$$

Theorem 4. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a defective matrix. Then we can define the function $f(\mathbf{A})$ as:

$$f(\mathbf{A}) := \mathbf{V}f(\mathbf{J})\mathbf{V}^{-1} \quad (2.16)$$

with

$$f(\mathbf{J}) = \begin{bmatrix} f(\mathbf{J}_1(\lambda_1)) & 0 & \cdots & 0 \\ 0 & f(\mathbf{J}_2(\lambda_2)) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & f(\mathbf{J}_k(\lambda_k)) \end{bmatrix} \quad (2.17)$$

and

$$f(\mathbf{J}_i(\lambda_i)) = \begin{bmatrix} f(\lambda_i) & f'(\lambda_i) & \frac{f''(\lambda_i)}{2!} & \cdots & \frac{f^{(k-1)}(\lambda_i)}{(k-1)!} \\ 0 & f(\lambda_i) & f'(\lambda_i) & \cdots & \frac{f^{(k-2)}(\lambda_i)}{(k-2)!} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & f(\lambda_i) & f'(\lambda_i) \\ 0 & 0 & \cdots & 0 & f(\lambda_i) \end{bmatrix} \quad (2.18)$$

Obviously, both definitions of matrix functions, based on diagonalization and Jordan canonical form, presuppose that the spectral radius of \mathbf{A} , $\rho(\mathbf{A})$, is less than r , the radius of convergence.

2.3 Interpolation-based definition

Interestingly, in this section we will show that for any $\mathbf{A} \in \mathbb{C}^{n \times n}$ and any sufficiently differentiable f , we can find a polynomial p such that $f(\mathbf{A}) = p(\mathbf{A})$. First, let us observe from previous sections that only the eigenvalues of \mathbf{A} are actually important for matrix polynomials. Also recall that every matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$ with spectrum $\{\lambda_1, \dots, \lambda_n\}$ has a minimal polynomial $\phi_{\mathbf{A}}$ given by

$$\phi_{\mathbf{A}}(t) := \prod_{i=1}^k (t - \lambda_k)^{n_i} \quad (2.19)$$

which is the unique monic minimal degree ($\deg(\phi_{\mathbf{A}}) = n_1 + \dots + n_k \leq n$) polynomial such that $\phi_{\mathbf{A}}(\mathbf{A}) = 0$.

Theorem 5. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a matrix with eigenvalues $\{\lambda_1, \dots, \lambda_k\}$, and minimal polynomial given by equation 2.19. Then for any two polynomials p_1, p_2 we have that $p_1(\mathbf{A}) = p_2(\mathbf{A})$ if and only if

$$\forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\} : p_1^{(i)}(\lambda_j) = p_2^{(i)}(\lambda_j).$$

Proof. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$, with spectrum $\{\lambda_1, \dots, \lambda_k\}$, and two polynomials p_1 and p_2 such that $p_1(\mathbf{A}) = p_2(\mathbf{A})$. Let us define q such that

$$q := p_1 - p_2$$

Then, $q(\mathbf{A}) = 0$, and is thus divisible by $\phi_{\mathbf{A}}$, meaning that

$$\forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\}, q(\lambda_j) = 0 \Rightarrow p_1^{(i)}(\lambda_j) = p_1^{(i)}(\lambda_i)$$

Similarly, consider two polynomials p_1 and p_2 such that

$$\forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\}, p_1^{(i)}(\lambda_j) = p_2^{(i)}(\lambda_i)$$

For $i = 0$, $q := p_1 - p_2 = 0$ on the spectrum of \mathbf{A} , and is then divisible by $\phi_{\mathbf{A}}$. Then

$$q = K\phi_{\mathbf{A}}$$

with K a polynomial. Then, $q(\mathbf{A}) = K(\mathbf{A})\phi_{\mathbf{A}}(\mathbf{A}) = 0$ since by definition, $\phi_{\mathbf{A}}(\mathbf{A}) = 0$. And thus, $p_1(\mathbf{A}) = p_2(\mathbf{A})$.

From this reasoning, we conclude that

$$\begin{aligned} p_1(\mathbf{A}) &= p_2(\mathbf{A}) \\ \Leftrightarrow \forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\} : p_1^{(i)}(\lambda_j) &= p_2^{(i)}(\lambda_j) \end{aligned}$$

□

In theorem 5, the conditions involve the evaluation of the polynomials p_1 and p_2 and their derivatives up to order $n_k - 1$ at the eigenvalues of A .

However, when the spectrum of A is simple, each eigenvalue λ_j is of multiplicity $n_j = 1$. This means that there are no higher order terms corresponding to these eigenvalues in the minimal polynomial, or in other words, there are no repeated roots. Consequently, there is no need to consider the derivatives of the polynomials p_1 and p_2 because there are no repeated roots for the polynomials to “match up” with. Therefore, in this simpler case, we only need to check that the polynomials p_1 and p_2 agree at the eigenvalues of A . In formal terms, the condition becomes:

Corollary 5.1. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a matrix with eigenvalues $\{\lambda_1, \dots, \lambda_k\}$, and minimal polynomial given by equation 2.19. If \mathbf{A} has simple spectrum, then for any two polynomials p_1, p_2 we have that $p_1(\mathbf{A}) = p_2(\mathbf{A})$ if and only if*

$$\forall j \in \{1, \dots, k\} : p_1(\lambda_j) = p_2(\lambda_j).$$

From this corollary, and from theorem 5, we can confirm our earlier statement : only the spectrum of \mathbf{A} is important for matrix polynomials. More importantly, we observe that $p(\mathbf{A})$ is uniquely defined by its values on the spectrum of \mathbf{A} . It seems then natural to extend this definition to any function f .

Definition 1. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a matrix with minimal polynomial given as in equation 2.19, and let f be a function that is at least $\max_k \{n_k - 1\}$ times differentiable. Say p is its (n_1, \dots, n_k) -Hermite interpolant i.e. the polynomial satisfying*

$$\forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\} : p^{(i)}(\lambda_j) = f^{(i)}(\lambda_j)$$

of minimal degree. Then we define $f(\mathbf{A}) = p(\mathbf{A})$.

3 The matrix-vector product $f(\mathbf{A})\mathbf{b}$

3.1 Introduction

To motivate the need for a matrix-vector product, we will consider a matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$. Let us consider the matrix exponential, such as :

$$e^{\mathbf{A}} = \sum_{k=0}^{\infty} \frac{\mathbf{A}^k}{k!} \quad (3.1)$$

Not only does this computation is very heavy (see section 5.1 for computation strategies), but it also affects the structure of the matrix. Say for example, we have the following laplacian matrix:

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 0 & 0 & \dots & \dots & 0 \\ -1 & 2 & -1 & 0 & 0 & \dots & 0 \\ 0 & -1 & 2 & -1 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & -1 & 2 & -1 \\ 0 & 0 & \dots & 0 & 0 & -1 & 2 \end{bmatrix}$$

Obviously, \mathbf{A} is a sparse matrix, and also a rank-structured one : it is a tridiagonal matrix. However, by computing $e^{\mathbf{A}}$, one will get a dense matrix. Storing a dense matrix often becomes challenging as the dimension of the problem grows. Besides storing, computing such a dense matrix is also an issue. However, the matrix-vector product $f(\mathbf{A})\mathbf{b}$ can be stored and computed efficiently.

3.2 The Method

As motivated by 3.1, when encountering specific structure in \mathbf{A} , such as sparsity, there is a lot to gain if we can store $f(\mathbf{A})\mathbf{b}$. In this section we will work towards a way to evaluate $f(\mathbf{A})\mathbf{b}$ in an intuitive way, thanks to Arnoldi method.

3.2.1 Formal Definitions

Firstly, we will define the notion of $\phi_{\mathbf{A},\mathbf{b}}$, i.e. the minimal polynomial of \mathbf{A} with respect to the vector \mathbf{b} . This is simply the polynomial

$$\phi_{\mathbf{A},\mathbf{b}}(t) := \prod_{i=1}^k (t - \lambda_i)^{m_i} \quad (3.2)$$

of minimal degree such that $\phi_{\mathbf{A},\mathbf{b}}(\mathbf{A})\mathbf{b} = \mathbf{0}$. Here $\lambda_1, \dots, \lambda_k$ are again the eigenvalues of \mathbf{A} .

Lemma 2. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a matrix with eigenvalues $\{\lambda_1, \dots, \lambda_k\}$, and minimal polynomial given by equation 2.19. Then for any two polynomials p_1, p_2 we have that $p_1(\mathbf{A})\mathbf{b} = p_2(\mathbf{A})\mathbf{b}$ if and only if*

$$\forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\} : p_1^{(i)}(\lambda_j) = p_2^{(i)}(\lambda_j).$$

Proof. From theorem 5, we know that

$$\begin{aligned} p_1(\mathbf{A}) &= p_2(\mathbf{A}) \\ \Leftrightarrow \forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\} : p_1^{(i)}(\lambda_j) &= p_2^{(i)}(\lambda_j) \end{aligned}$$

We also know that

$$p_1(\mathbf{A}) = p_2(\mathbf{A}) \Leftrightarrow p_1(\mathbf{A})\mathbf{b} = p_2(\mathbf{A})\mathbf{b}$$

Thus, we conclude that

$$\begin{aligned} p_1(\mathbf{A})\mathbf{b} &= p_2(\mathbf{A})\mathbf{b} \\ \Leftrightarrow \forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, n_k - 1\} : p_1^{(i)}(\lambda_j) &= p_2^{(i)}(\lambda_j) \end{aligned}$$

□

Theorem 6. *Let f be a sufficiently differentiable function that has no singularities on the spectrum of a given matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$. Then, with p the unique Hermite interpolating polynomial of \mathbf{A} w.r.t. \mathbf{b} i.e.*

$$\forall j \in \{1, \dots, k\} : \forall i \in \{0, \dots, m_k - 1\} : p^{(i)}(\lambda_j) = f^{(i)}(\lambda_j)$$

we have that $f(\mathbf{A})\mathbf{b} = p(\mathbf{A})\mathbf{b}$.

Proof. Let f be a function that is at least $\max_k \{m_k - 1\}$ times differentiable, and let p_1 be its (m_1, \dots, m_k) -Hermite interpolant. Then, by definition 1, we have that

$$f(\mathbf{A}) = p_1(\mathbf{A})$$

Then, by lemma 2, let us define p_2 such that

$$p_2(\mathbf{A})\mathbf{b} = p_1(\mathbf{A})\mathbf{b}$$

Then, we have that

$$p_1(\mathbf{A})\mathbf{b} = p_2(\mathbf{A})\mathbf{b} = f(\mathbf{A})\mathbf{b}$$

□

3.2.2 The Arnoldi Method

The Arnoldi method is a reduction method that allows low-rank approximation of a given matrix \mathbf{A} . It is based on the Hessenberg reduction of a matrix \mathbf{A} . More formally

Definition 2. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$, the Arnoldi method approximates its Hessenberg reduction given by*

$$\mathbf{V}^* \mathbf{A} \mathbf{V} = \mathbf{H} \tag{3.3}$$

where $\mathbf{V} \in \mathbb{C}^{n \times n}$ is unitary, and $\mathbf{H} \in \mathbb{C}^{n \times n}$ is upper Hessenberg. As it is a (low-rank) approximation, Arnoldi will compute the following decomposition

$$\mathbf{V}_k^* \mathbf{A} \mathbf{V}_k = \mathbf{H}_k \tag{3.4}$$

with $\mathbf{V}_k \in \mathbb{C}^{n \times k}$ unitary, and $\mathbf{H}_k \in \mathbb{C}^{k \times k}$ upper Hessenberg. That means that \mathbf{H}_k is the orthogonal projection of \mathbf{A} onto the k th Krylov subspace $\mathcal{K}_k(\mathbf{A}, \mathbf{v}_1)$, with \mathbf{v}_1 the first column of \mathbf{V} .

The Arnoldi method is an iterative method, following this algorithm:

Here the orthogonalization process (lines 3-6) is a modified Gram-Schmidt process. Also, when implementing, we will consider an ℓ_2 norm. Numerical conditions will often imply loss of orthogonality. To avoid this, it will be necessary to reorthogonalize, i.e. to reapply the Gram-Schmidt process.

Algorithm 1: Arnoldi Iteration

Data: $A \in \mathbb{C}^{n \times n}$, $v_1 \in \mathbb{C}^n$ a unit vector in the chosen norm (line 7)

Result: $H_n \in \mathbb{C}^{n \times n}$

```
1 for  $k = 1$  to  $n$  do
2    $\mathbf{w} \leftarrow A\mathbf{v}_k$ ;
3   for  $i = 1$  to  $k$  do
4      $h_{ik} \leftarrow \mathbf{v}_i^* \mathbf{w}$ ;
5      $\mathbf{w} \leftarrow \mathbf{w} - h_{ik} \mathbf{v}_i$ ;
6   end
7    $h_{k+1,k} \leftarrow \|\mathbf{w}\|$ ;
8   if  $h_{k+1,k} = 0$  then
9     break;
10  end
11   $\mathbf{v}_{k+1} \leftarrow \mathbf{w} / h_{k+1,k}$ 
12 end
```

3.2.3 The matrix-vector product $f(\mathbf{A})\mathbf{b}$

Recall the problem is to approximate the matrix-vector product $f(\mathbf{A})\mathbf{b}$. Now we have all the tools to achieve that efficiently.

Lemma 3. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ and $\mathbf{b} \in \mathbb{C}^n$. Then, the matrix vector product $f(\mathbf{A})\mathbf{b}$ can be approximated by*

$$f(\mathbf{A})\mathbf{b} \approx \|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{e}_1 \quad (3.5)$$

Where \mathbf{V}_k and \mathbf{H}_k are the matrices computed by the Arnoldi method after k iterations. \mathbf{H}_k is upper Hessenberg and $\text{span}(\mathbf{V}_k) = \mathcal{K}_k(\mathbf{A}, \mathbf{b}) = \text{span}(\mathbf{b}, \mathbf{A}\mathbf{b}, \mathbf{A}^2\mathbf{b}, \dots, \mathbf{A}^{k-1}\mathbf{b})$. \mathbf{e}_1 is the first column of the identity matrix.

Proof. Consider we want to approximate the matrix-vector product $f(\mathbf{A})\mathbf{b}$. We will use the Arnoldi method (algorithm 12). For initial unit vector we choose $\mathbf{v}_1 = \mathbf{b} / \|\mathbf{b}\|_2$ which is indeed an ℓ_2 unit vector. According to the Hessenberg reduction of \mathbf{A} , we have that

$$f(\mathbf{A})\mathbf{b} \approx f(\mathbf{V}_k \mathbf{H}_k \mathbf{V}_k^*) \mathbf{b}$$

Since \mathbf{V}_k is unitary, we have

$$f(\mathbf{A})\mathbf{b} \approx \mathbf{V}_k f(\mathbf{H}_k) \mathbf{V}_k^* \mathbf{b}$$

Furthermore, as we chose $\mathbf{v}_1 = \mathbf{b} / \|\mathbf{b}\|_2$, we have that

$$\mathbf{b} = \|\mathbf{b}\|_2 \mathbf{v}_1 = \|\mathbf{b}\|_2 \mathbf{V}_k \mathbf{e}_1$$

Thus, we have that

$$f(\mathbf{A})\mathbf{b} \approx \|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{V}_k^* \mathbf{V}_k \mathbf{e}_1 = \|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{e}_1$$

□

Lemma 4. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ and let $\mathbf{V}_k, \mathbf{H}_k$ be the matrices computed by the Arnoldi method after k iterations. Then, for any polynomial p_j of degree $j \leq k-1$, we have that*

$$p_j(\mathbf{A})\mathbf{b} = \mathbf{V}_k p_j(\mathbf{H}_k) \mathbf{e}_1 \quad (3.6)$$

Lemma 5. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ and let $\mathbf{V}_k, \mathbf{H}_k$ be the matrices computed by the Arnoldi method after k iterations. Then, given p_{k-1} the unique Hermite interpolant of \mathbf{A} w.r.t. \mathbf{b} of degree $k-1$, we have that

$$\|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{e}_1 = p_{k-1}(\mathbf{A}) \mathbf{b} \quad (3.7)$$

Proof. Since p_{k-1} is the unique Hermite interpolant of \mathbf{A} w.r.t. \mathbf{b} of degree $k-1$, we have that

$$\|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{e}_1 = \|\mathbf{b}\|_2 \mathbf{V}_k p_{k-1}(\mathbf{H}_k) \mathbf{e}_1$$

And since

$$\|\mathbf{b}\|_2 \mathbf{V}_k p_{k-1}(\mathbf{H}_k) \mathbf{e}_1 = \|\mathbf{b}\|_2 p_{k-1}(\mathbf{A}) \mathbf{q}_1 = p_{k-1}(\mathbf{A}) \mathbf{b}$$

We indeed have

$$\|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{e}_1 = p_{k-1}(\mathbf{A}) \mathbf{b}$$

□

Theorem 7. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ and let $m = \deg(\phi_{\mathbf{A}, \mathbf{b}})$ then

$$f(\mathbf{A}) \mathbf{b} = \|\mathbf{b}\|_2 \mathbf{V}_m f(\mathbf{H}_m) \mathbf{e}_1 \quad (3.8)$$

Proof. Following theorem 6, we have that

$$f(\mathbf{A}) \mathbf{b} = p_{m-1}(\mathbf{A}) \mathbf{b}$$

With p_{m-1} the unique Hermite interpolant of \mathbf{A} w.r.t. \mathbf{b} of degree $m-1$ with $\deg(\phi_{\mathbf{A}, \mathbf{b}}) = m$. Then, by lemma 3, 4 and 5, we have that

$$f(\mathbf{A}) \mathbf{b} = \|\mathbf{b}\|_2 \mathbf{V}_m f(\mathbf{H}_m) \mathbf{e}_1$$

□

4 Algorithms

In this section, we will implement basic algorithms. From the previous section, we will distinguish two cases :

- the dense case, where $f(\mathbf{A})$ is to be computed
- and the case where only the matrix-vector product $f(\mathbf{A}) \mathbf{b}$ is to be computed

4.1 Dense case

First, let us consider the computation of $f(\mathbf{A})$. Meaning we put aside the matrix-vector product (for now). More specifically, our approach will be to use definition 1 to compute $f(\mathbf{A})$. All implementations are done in `Matlab 2023a`.

4.1.1 Dependencies

We are provided with a function `hess_and_phi()` that computed the Hessenberg reduction of a given matrix and its associated minimal polynomial. It takes as input a matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$. It then follows these simple steps:

1. Compute the Hessenberg reduction of \mathbf{A} , i.e. $\mathbf{V}^* \mathbf{A} \mathbf{V} = \mathbf{H}$
2. It computes its Jordan Canonical Form from \mathbf{H}
3. It computes the minimal polynomial of \mathbf{A} , $\phi_{\mathbf{A}}$ through its Jordan Canonical Form
4. It returns \mathbf{V} , \mathbf{H} , \mathbf{J} , λ_j (the eigenvalues of \mathbf{A}) and n_i (the multiplicity of λ_i in $\phi_{\mathbf{A}}$)

Note that step 2 makes perfect sense.

Theorem 8. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ with a Hessenberg reduction defined by $\mathbf{V}^* \mathbf{A} \mathbf{V} = \mathbf{H}$. Then \mathbf{A} and \mathbf{H} share the same Jordan Canonical Form.*

Proof. \mathbf{V} is unitary, hence \mathbf{A} and \mathbf{H} are similar matrices, meaning they have the same eigenvalues. Thus, they share the same Jordan Canonical Form. \square

To recover the minimal polynomial of \mathbf{A} , one simply retrieves the λ_j and the n_i from the previously described steps, and then compute the polynomial this way:

$$\phi_{\mathbf{A}}(t) = \prod_{i=1}^k (t - \lambda_k)^{n_i}$$

In the supplementary materials, you will find an implementation of this, called `construct_minimal_polynomial()`. One quick way to assess if the minimal polynomial is constructed correctly, is to evaluate it at \mathbf{A} . Taking the ℓ_2 norm of the result should be close to zero. Here, with `test1.mat` (a 5 by 5 matrix), we obtain $\phi_{\mathbf{A}}(\mathbf{A}) = 3.3523e - 12$. This is indeed numerically close to zero. For the Hermite interpolation, several options are possible. First, a routine `hermite_interp()` is provided in the supplementary materials, it computed the Hermite interpolation by divided differences. However, I had more precise results with an alternative method proposed by Or Werner, BGU, Israel, based on the Hermite method. In the following, I used the latter approach, the routine is provided in the supplementary material aswell.

4.1.2 Implementation

From the previous elements, we can construct a routine called `matrix_function()` that takes as input a matrix $\mathbf{A} \in \mathbb{C}^{n \times n}$, and a function f . It has extremely simple steps:

1. Retrieve λ_j and n_i from `hess_and_phi()`
2. Construct the matrix \mathbf{FdF} which is f and its derivatives evaluated at the eigenvalues of \mathbf{A}
3. Construct the Hermite interpolation of \mathbf{A} with `hermite_interp()`
4. Evaluate the Hermite interpolation at \mathbf{A} , and return the result

The routine is provided in the supplementary materials.




Figure 1: Measure of performance of the algorithm to compute $f(\mathbf{A})$ according to definition 1. We assume that Davies and Higham 2003 (Schur Parlett) algorithm is correct, and we measure the relative error. We compare the performance of our algorithm on several types of matrices : diagonal (- -), symmetric (-*) and dense (-). Matrix coefficients are randomly filled such that for all non-zero entry: $a_{ij} \sim \mathcal{U}(0, 1)$, this ensures \mathbf{A} is full rank. We test it on three different functions : Sine (a), Cosine (b) and Exponential (c). We note that starting from a certain matrix size, the algorithm lose considerably in performance.

4.1.3 Results

In this part, we will compare the performance of our routine `matrix_function()` with the built-in `funm()` function from `Matlab`. The built-in Matlab function uses a Schur-Parlett algorithm from Davies and Higham 2003. To compare both methods, we generate three types of matrices of different sizes. More precisely, we will investigate both algorithm's performances on symmetric, diagonal and dense matrices, on the following functions: cosine, sine and exponential.

Figure 1 illustrates the result of the measurement of performance of the algorithm. The relative error between Davies and Higham 2003 and our results is computed:

$$\text{relative error} = \frac{\|f(\mathbf{A}) - \text{funm}(\mathbf{A}, f)\|_2}{\|\text{funm}(\mathbf{A}, f)\|_2}$$

We note that for small matrices ($n \leq 10$), the hermite interpolation approach is precise and has residual error inferior to 10^{-11} , which is somewhat close to numerical precision (around 10^{-16} as we work with double precision). However, starting from a certain matrix size, the algorithm loses considerably in performance: both in precision and in computation time. Interestingly, this behavior is independant to the matrix structure : diagonal, symmetric or dense. It also seem to be independant to the function we are evaluating (though we has slightly lower relative error for the exponential function as depicted in figure 1c).

We conclude that this algorithm, while being elegant, is quickly inefficient, and lead to both numerical and computational issues, even at small matrix sizes.

4.2 Matrix-vector product

4.2.1 Implementation

In this section, we will implement the matrix-vector product $f(\mathbf{A})\mathbf{b}$, as described in section 3.1. We will use the Arnoldi method to achieve this. The implementation is done in `Matlab 2023a`. As the theory has been well described in section 3.1, the implementation is straight-forward :

Algorithm 2: Matrix-Vector Product

Data: $\mathbf{A} \in \mathbb{C}^{n \times n}$, $\mathbf{b} \in \mathbb{C}^n$, f a function and $k \in \mathbb{N}$ the dimension of the Krylov subspace

Result: $f(\mathbf{A})\mathbf{b}$

- 1 $\mathbf{v}_1 \leftarrow \mathbf{b}/\|\mathbf{b}\|_2$;
 - 2 $[\mathbf{V}, \mathbf{H}] \leftarrow \text{arnoldi}(\mathbf{A}, \mathbf{v}_1, k)$;
 - 3 $\mathbf{e}_1 \leftarrow$ first column of the identity matrix;
 - 4 $f \leftarrow \|\mathbf{b}\|_2 \mathbf{V} f(\mathbf{H}) \mathbf{e}_1$;
-

Where `arnoldi()` is the Arnoldi iteration described in algorithm 12. This routine is provided in the supplementary materials.

4.2.2 Results

To evaluate the performance of this Matrix-vector product routine, we will test it versus the naive way of doing it in `Matlab`, meaning computing $f(\mathbf{A})$ explicetely, then multiplying it by \mathbf{b} . In our first experiment (figure 2), where \mathbf{A} is a large sparse matrix. More specifically it is a graded L-shape pattern from George and Liu 1978. The vector \mathbf{b} is filled with uniformly distributed random coefficients such that $b_i \sim \mathcal{U}(0, 1)$. The figure 2b shows that for such a setup, the proposed algorithm reaches a relative error equals to numerical precision few iterations after 20. This means

$$\frac{\|\|\mathbf{b}\|_2 \mathbf{V}_k f(\mathbf{H}_k) \mathbf{e}_1 - f(\mathbf{A})\mathbf{b}\|_2}{\|f(\mathbf{A})\mathbf{b}\|_2} = \epsilon$$

for $k > 20$, and where ϵ is the machine precision. Note that this result is very specific to the problem depicted in figure 2, and the threshold for k will differ in function of the considered problem. Another interpretation is that computing $f(\mathbf{A})\mathbf{b}$ on the smaller Krylov subspace $\mathcal{K}_k(\mathbf{A}, \mathbf{b})$ gives extremely good results, even when $k \ll n$.

Obviously, this very interesting result lets us think that there is a lot of potential computational gain in evaluating $f(\mathbf{A})\mathbf{b}$ on this smaller Krylov subspace. The computational gain for the problem solved in figure 2 is described in Table 1. We note that for this problem, our method considerably outperforms the naive approach of evaluating $f(\mathbf{A})$ seperately, regardless of the function (though the biggest gain seem to come from the matrix exponential).

However, this does not give us any insight about how those two algorithms scale up or down. To investigate this, we will use the BCSPWR matrix collection. This collection contains matrices of different sizes, and different sparsity patterns. We will test our method against the naive approach on this collection. The results are depicted in Table 2. We notice that for very low dimension ($n < 300$) the naive approach slightly outperforms ours for trigonometric functions. However, for larger matrices, and for the exponential function, working in the Kryloc subspace is the way to go. For the largest matrix, where $n = 5300$, the naive approach takes half a minute, while our method takes less than a tenth of a second. This is a considerable gain in performance, and could be crucial in some applications. As this margin increases with the matrix size, we can expect our method to be even more efficient for larger matrices.




Figure 2: Evaluation of the performance of the matrix-vector product routine. We compare the performance of our method against the naive way of doing it in `Matlab`. We test it on three different functions : Sine (---), Cosine (-*) and Exponential (—) on figure (b). Those functions are applied to a large matrix \mathbf{A} with sparsity pattern given in (a). Our method allows for precise approximation of $f(\mathbf{A})\mathbf{b}$ even when working on a very low rank Hessenberg reduction (b). The vector \mathbf{b} is a random vector whose coefficients are uniformly distributed such that $b_i \sim \mathcal{U}(0, 1)$.

Table 1: Comparison of computational performance of both methods for evaluating $f(\mathbf{A}\mathbf{b})$ with \mathbf{A} the matrix in 2a. Performance was measured in ms. Optimal k was determined when the relative error went below $1e - 14$, as we worked on double precision. Tests were run on an Intel Core i7-1185G7 @ 3.00GHz with 16GB RAM. We note that for this problem, our method is outperforming by a significant margin the naive approach.

fun	Naive way (ms)	Krylov method (ms)	Optimal k
exp()	151.2	6.3	21
cos()	121.8	11.2	23
sin()	112.6	8.1	23

Table 2: Comparison of the naive approach and the Krylov approach on BCSPWR matrix collection. Performance was measured in ms. Optimal k was determined when the relative error went below $1e - 14$.

Matrix	n	exp()		cos()		sin()	
		Naive	Krylov	Naive	Krylov	Naive	Krylov
BCSPWR01	39	1.0	1.3	0.55	1.5	0.38	5.3
BCSPWR02	49	3.2	1.9	0.66	3.7	0.48	1.6
BCSPWR03	118	16.6	2.3	1.3	3.0	0.87	3.1
BCSPWR04	274	6.5	1.6	5.2	8.7	5.9	6.4
BCSPWR05	443	26.4	2.7	21.8	4.6	21.5	3.6
BCSPWR06	1454	436.6	11.9	299.6	10.3	399.6	12.1
BCSPWR07	1612	484.0	11.9	392.0	14.7	393.2	14.7
BCSPWR08	1624	504.4	13.4	414.5	12.4	433.8	14.6
BCSPWR09	1723	628.1	11.7	593.7	10.3	531.8	10.1
BCSPWR10	5300	32868	98.4	27447	96.7	23948	89.7

5 Applications

5.1 Matrix Exponential

Consider the simple system of ODEs

$$\frac{d\mathbf{x}}{dt} = \mathbf{A}\mathbf{x} \quad (5.1)$$

with initial condition $\mathbf{x}(0) = \mathbf{x}_0 \in \mathbb{R}^n$. Then we know the solution to be given by $\mathbf{x}(t) = e^{\mathbf{A}t}\mathbf{x}_0$. However, for all but the stablest systems, this is not a good method, due to issues such as stability and stiffness. Here we consider for instance the 2D convection-diffusion equation for the flow $\mathbf{u}(x, y)$:

$$\frac{d\mathbf{u}}{dt} = \epsilon \Delta \mathbf{u} + \alpha \cdot \nabla \mathbf{u}$$

with Dirichlet boundary conditions and $\epsilon \in \mathbb{R}_0^+$ and $\alpha \in \mathbb{R}^2$. Simple time-stepping methods are known to be unstable at large time-steps, and our exponential scheme suffers from similar problems, i.e. t cannot be taken too large.

5.2 The sign function

In control theory we are often interested in the eigenvalues λ of system matrices with $\text{Re}(\lambda) > 0$, since they correspond to unstable poles. In the design of controllers it is therefore interesting to have an efficient way to count the number of eigenvalues of a matrix in the right half-plane $\text{Re}(z) > 0$. Here we will build such a method.

Theorem 9. *Let $\mathbf{A} \in \mathbb{C}^{n \times n}$ be a matrix with k_- eigenvalues in the left plane, k_+ eigenvalues in the right plane and none on the imaginary axis, counting multiplicity. Let $\text{sgn} : \mathbb{C} \mapsto \{1, -1\}$ be defined by*

$$\text{sgn}(z) = \begin{cases} 1 & \text{Re}(z) \geq 0 \\ -1 & \text{Re}(z) < 0. \end{cases}$$

Then $\text{trace}(\text{sgn}(\mathbf{A})) = k_+ - k_-$.

Proof. Let $\mathbf{A} \in \mathbb{C}^{n \times n}$. To stay in a very general scenario, and consider cases where \mathbf{A} , let us consider its Jordan Canonical Form :

$$\mathbf{A} = \mathbf{V}\mathbf{J}\mathbf{V}^{-1}$$

Recall that by theorem 4, consider f a function, then

$$f(\mathbf{A}) = \mathbf{V}f(\mathbf{J})\mathbf{V}^{-1}$$

Thus let us consider the decomposition where

$$\mathbf{J} = \begin{pmatrix} \mathbf{J}_{k_+} & 0 \\ 0 & \mathbf{J}_{k_-} \end{pmatrix}$$

such that \mathbf{J}_{k_+} has k_+ eigenvalues in the right plane, and \mathbf{J}_{k_-} has k_- eigenvalues in the left plane. Then, we have that

$$\text{sgn}(\mathbf{J}) = \begin{pmatrix} \mathbf{I}_{k_+} & 0 \\ 0 & -\mathbf{I}_{k_-} \end{pmatrix}$$

where \mathbf{I}_n is the identity matrix of size n . Then, we have that

$$\text{trace}(\text{sgn}(\mathbf{J})) = k_+ - k_-$$

And thus, we have that

$$\text{trace}(\text{sgn}(\mathbf{A})) = \text{trace}(\text{sgn}(\mathbf{V}\mathbf{J}\mathbf{V}^{-1})) = \mathbf{V}\text{trace}(\text{sgn}(\mathbf{J}))\mathbf{V}^{-1} = (k_+ - k_-)\mathbf{V}\mathbf{V}^{-1} = k_+ - k_-$$

□

References

- Davies, Philip I and Nicholas J Higham (2003). “A Schur-Parlett algorithm for computing matrix functions”. In: *SIAM Journal on Matrix Analysis and Applications* 25.2, pp. 464–485.
- Frommer, Andreas and Valeria Simoncini (2008). “Matrix functions”. In: *Model order reduction: theory, research aspects and applications*. Springer, pp. 275–303.
- George, Alan and Joseph WH Liu (1978). “An automatic nested dissection algorithm for irregular finite element problems”. In: *SIAM Journal on Numerical Analysis* 15.5, pp. 1053–1069.