


Observation Synthesis for Games with Imperfect Information

Nathan Lhote

Joint work with Paulin Fournier


Wydział Matematyki, Informatyki i Mechaniki




Games with Imperfect Information



Games with Imperfect Information




Games with Imperfect Information



- Finite arena:


$$V = \{v_1, v_2, v_3, v_4, v_5\}$$

Games with Imperfect Information




- ▶ Finite arena:
 $V = \{v_1, v_2, v_3, v_4, v_5\}$
- ▶ Players:
 $V = V_1 \uplus V_2 \uplus \dots \uplus V_n$

Games with Imperfect Information




- ▶ Finite arena:
 $V = \{v_1, v_2, v_3, v_4, v_5\}$
- ▶ Players:
 $V = V_1 \uplus V_2 \uplus \dots V_n$
- ▶ Histories: V^* , Plays: V^ω

Games with Imperfect Information



- ▶ Finite arena:
 $V = \{v_1, v_2, v_3, v_4, v_5\}$
- ▶ Players:
 $V = V_1 \uplus V_2 \uplus \dots \uplus V_n$
- ▶ Histories: V^* , Plays: V^ω
- ▶ Winning condition:
 $\mathcal{W} : V^\omega \rightarrow \mathcal{S}$
($\{0, 1\}, \mathbb{N}, \mathbb{R}^d$, etc)


Games with Imperfect Information



- ▶ Finite arena:
 $V = \{v_1, v_2, v_3, v_4, v_5\}$
- ▶ Players:
 $V = V_1 \uplus V_2 \uplus \dots \uplus V_n$
- ▶ Histories: V^* , Plays: V^ω
- ▶ Winning condition:
 $\mathcal{W} : V^\omega \rightarrow \mathcal{S}$
($\{0, 1\}, \mathbb{N}, \mathbb{R}^d$, etc)

Indistinguishable histories (for Player 1):

Games with Imperfect Information




- ▶ Finite arena:
 $V = \{v_1, v_2, v_3, v_4, v_5\}$
- ▶ Players:
 $V = V_1 \uplus V_2 \uplus \dots \uplus V_n$
- ▶ Histories: V^* , Plays: V^ω
- ▶ Winning condition:
 $\mathcal{W} : V^\omega \rightarrow \mathcal{S}$
($\{0, 1\}, \mathbb{N}, \mathbb{R}^d$, etc)

Indistinguishable histories (for Player 1):

Equivalence relation $R \subseteq V^* \times V^*$

Games with Imperfect Information




- ▶ Finite arena:
 $V = \{v_1, v_2, v_3, v_4, v_5\}$
- ▶ Players:
 $V = V_1 \uplus V_2 \uplus \dots \uplus V_n$
- ▶ Histories: V^* , Plays: V^ω
- ▶ Winning condition:
 $\mathcal{W} : V^\omega \rightarrow \mathcal{S}$
($\{0, 1\}, \mathbb{N}, \mathbb{R}^d$, etc)

Indistinguishable histories (for Player 1):

Equivalence relation $R \subseteq V^* \times V^*$

Given as a transducer

Observation functions



Observation functions

Imperfect information

Observation functions

Imperfect information

- Finite arena V

Observation functions

Imperfect information

- ▶ Finite arena V
- ▶ $f : V^* \rightarrow O^*$, from histories to *observations*

Observation functions

Imperfect information

- ▶ Finite arena V
- ▶ $f : V^* \rightarrow O^*$, from histories to *observations*
Given as a transducer

Observation functions

Imperfect information

- ▶ Finite arena V
- ▶ $f : V^* \rightarrow O^*$, from histories to *observations*
Given as a transducer

Remark

$\ker f := \{(u, v) \in (V^*)^2 \mid f(u) = f(v)\} = f^{-1} \circ f$
 \rightarrow indistinguishability relation

Observation functions


Imperfect information

- ▶ Finite arena V
- ▶ $f : V^* \rightarrow O^*$, from histories to *observations*
Given as a transducer


Remark

$\ker f := \{(u, v) \in (V^*)^2 \mid f(u) = f(v)\} = f^{-1} \circ f$
 \rightarrow indistinguishability relation


Ex:




Arena:





Observation function:




Arena:





Observation function:



Arena:



Observation function:



Problem:

Problem:

- ▶ Input: Arena V , equivalence relation $R \subseteq V^* \times V^*$

Problem:

- ▶ Input: Arena V , equivalence relation $R \subseteq V^* \times V^*$
- ▶ Question: Can one compute an observation function $f : V^* \rightarrow O^*$ such that $\ker f = R$

Problem:

- ▶ Input: Arena V , equivalence relation $R \subseteq V^* \times V^*$
- ▶ Question: Can one compute an observation function $f : V^* \rightarrow O^*$ such that $\ker f = R$

$R \setminus f$	Functional	Sequential	Mealy
Arbitrary	?		X
Deterministic	Yes		X
Letter-to-letter	Yes		

Mealy observation function

Arena V , indistinguishability relation R .

Is there f given by a *sequential* and *letter-to-letter* transducer, such that $\ker f = R$?

Mealy observation function

Arena V , indistinguishability relation R .

Is there f given by a *sequential* and *letter-to-letter* transducer, such that $\ker f = R$?

Necessary conditions

Mealy observation function

Arena V , indistinguishability relation R .

Is there f given by a *sequential* and *letter-to-letter* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter

Mealy observation function

Arena V , indistinguishability relation R .

Is there f given by a *sequential* and *letter-to-letter* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter
- ▶ 2) R is prefix closed

If uRv then $\forall w \leq_{\text{pref}} u, z \leq_{\text{pref}} v$ with $|w| = |z|$, wRz

Mealy observation function

Arena V , indistinguishability relation R .

Is there f given by a *sequential* and *letter-to-letter* transducer, such that $\ker f = R$?

Necessary conditions


► 1) R is letter-to-letter

► 2) R is prefix closed


If uRv then $\forall w \leq_{\text{pref}} u, z \leq_{\text{pref}} v$ with $|w| = |z|$, wRz

► 3) R satisfies condition 3)

Co-example




Co-example



Syntactic congruence of R

uSv if $\forall w \ uwRvw$

Co-example




Syntactic congruence of R

uSv if $\forall w \ uwRvw$

Condition 3):

- S has finite index with respect to R

Co-example



Syntactic congruence of R


uSv if $\forall w \ uwRvw$

Condition 3):

- S has finite index with respect to R

First result:

Co-example



Syntactic congruence of R

uSv if $\forall w \ uwRvw$


Condition 3):

- S has finite index with respect to R

First result:

- 1)-3) are decidable

Co-example



Syntactic congruence of R

uSv if $\forall w \ uwRvw$

Condition 3):

- ▶ S has finite index with respect to R

First result:

- ▶ 1)-3) are decidable
- ▶ 1)-3) are sufficient

Sequential observation function

Arena V, indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Sequential observation function

Arena V, indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Necessary conditions

Sequential observation function

Arena V, indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter

Sequential observation function

Arena V, indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter
- ▶ 2) S has finite index with respect to $TC(PC(R))$
(hard to show)

Sequential observation function

Arena V, indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter
- ▶ 2) S has finite index with respect to $TC(PC(R))$
(hard to show)

Second result

Sequential observation function

Arena V , indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter
- ▶ 2) S has finite index with respect to $TC(PC(R))$
(hard to show)

Second result

- ▶ 1)-2) are sufficient (same kind of proof)

Sequential observation function

Arena V, indistinguishability relation R letter-to-letter.

Is there f given by a *sequential* transducer, such that $\ker f = R$?

Necessary conditions

- ▶ 1) R is letter-to-letter
- ▶ 2) S has finite index with respect to $TC(PC(R))$
(hard to show)

Second result

- ▶ 1)-2) are sufficient (same kind of proof)
- ▶ 2) is undecidable (thanks Bruno Guillon)

Sum up *etc*

$R \setminus f$	Functional	Sequential	Mealy
Arbitrary	?	U	X
Deterministic	Yes	U	X
Letter-to-letter	Yes	U	D

Sum up *etc*

$R \setminus f$	Functional	Sequential	Mealy
Arbitrary	?	U	X
Deterministic	Yes	U	X
Letter-to-letter	Yes	U	D

Qs

Sum up *etc*

$R \setminus f$	Functional	Sequential	Mealy
Arbitrary	?	U	X
Deterministic	Yes	U	X
Letter-to-letter	Yes	U	D

Qs

- Find interesting classes of observation functions
(*i.e.* deterministic, increasing)

Sum up *etc*

$R \setminus f$	Functional	Sequential	Mealy
Arbitrary	?	U	X
Deterministic	Yes	U	X
Letter-to-letter	Yes	U	D

Qs

- ▶ Find interesting classes of observation functions (*i.e.* deterministic, increasing)
- ▶ Solve games!

Thanks !

