Richard Goldberg Solutions

Nathanael Seen

July 2, 2020

Preface

In this book, I would be providing solutions to selected exercises in Richard R. Goldberg's Real Analysis book.

Disclaimer: The solutions I provide are not official solutions, just my own solutions to the exercises.

This book is only available as an electronic copy, and not available in print.

If you spot any mistakes in this book, have suggestions on how to improve this book, or have any other queries, you may reach me at my email.

 \sim Nathanael Seen

Contents

1	Sets and Functions			
	1.1	Exercise 1.7 Solutions		
2	Seq	uences of Real Numbers		
	2.1	Exercise 2.1 Solutions		
	2.2	Exercise 2.2 Solutions		
	2.3	Exercise 2.3 Solutions		
	2.4	Exercise 2.4 Solutions		
	2.5	Exercise 2.5 Solutions		
	2.6	Exercise 2.6 Solutions		
	2.7	Exercise 2.7 Solutions		

Chapter 1

Sets and Functions

1.1 Exercise 1.7 Solutions

Q1.

 $\underline{\mathbf{Ans.}}$

- (a) 7
- (b) $\pi + 1$
- (c) π

Chapter 2

Sequences of Real Numbers

2.1 Exercise 2.1 Solutions

Q5.

Proof.

Let S be a Sequence in the set A.

We note that it is given by the function; $f: \mathbb{N} \to A$.

Now, consider an (arbitary) subsequence S' of S, which has the form $f \circ g$, where $h : \mathbb{N} \to \mathbb{N}$, satisfies h(n) < h(n+1), for all n.

(Claim:
$$(g \circ h)(n) < (g \circ h)(n+1), \forall n \in \mathbb{N}$$
)

We note that h(n) < h(n+1), and g(n) < g(n+1).

Now, let
$$b = h(n)$$
, and $c = h(n+1)$.

Also,
$$b < c$$
 since $h(n) < h(n+1)$.

Thus,

$$b < b+1 \le c$$

=> $g(b) < g(b+1) \le g(c)$
=> $g(b) < g(c)$
=> $g(h(n)) < g(h(n+1)).$

- 2.2 Exercise 2.2 Solutions
- 2.3 Exercise 2.3 Solutions
- 2.4 Exercise 2.4 Solutions
- 2.5 Exercise 2.5 Solutions
- 2.6 Exercise 2.6 Solutions
- 2.7 Exercise 2.7 Solutions