

EIGENBASES

NATHAN GRIGG

Let A be an $n \times n$ matrix. An **eigenbasis** corresponding to A is a basis for R^n consisting entirely of eigenvectors for A . Such a basis only exists if A is not defective. To find an eigenbasis, you find a basis for each eigenspace of A . The collection of all these basis vectors for an eigenbasis for A .

Example. The matrix $A = \begin{bmatrix} -83 & 4 & 35 \\ -348 & 20 & 140 \\ -174 & 8 & 74 \end{bmatrix}$ has eigenvalues 3, 4, 4. The eigenspace E_3 has a basis $\left\{ \begin{bmatrix} 35 \\ 0 \\ 87 \end{bmatrix}, \begin{bmatrix} 4 \\ 87 \\ 0 \end{bmatrix} \right\}$ and the eigenspace E_4 has a basis $\left\{ \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix} \right\}$. So an eigenbasis corresponding to A is $\left\{ \begin{bmatrix} 35 \\ 0 \\ 87 \end{bmatrix}, \begin{bmatrix} 4 \\ 87 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix} \right\}$.

Exercises.

- (1) Find an eigenbasis corresponding to $A = \begin{bmatrix} 3 & 0 & -3 \\ 0 & 3 & -3 \\ 0 & 0 & 0 \end{bmatrix}$. Write $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ as a linear combination of the basis and use this to find $A^{10}\mathbf{x}$.
- (2) Find an eigenbasis corresponding to $A = \begin{bmatrix} 0 & -1 & 1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$. Write $\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ as a linear combination of the basis and use this to find a general formula for $A^t\mathbf{x}$.