

Chapter 1: Introduction

Math 307i – Winter 2011 – January 3

What is a differential equation?

$$(1 - x^2)y'' - xy' + 3y = 0$$

$$\frac{dy}{dx} = \frac{3x^2 + 4 + 2}{2(y - 1)}$$

An equation that involves a derivative

Mathematical Modeling

or: finally, we can factor in air resistance (kind of)


Determine the how the velocity of a falling object changes over time.

Ignoring air resistance

- Newton's first law: $F = ma$
- In diff eq form: $v' = F/m$
- Force from gravity: $F = 9.8m$
- Final equation: $v' = 9.8$
- Solution: $v(t) = 9.8t + c$

Air resistance

- Assume the drag force is proportional to the velocity squared.



$$\begin{aligned} v' &= \frac{9.8m - \gamma v^2}{m} \\ &= 9.8 - \frac{\gamma v^2}{m} \end{aligned}$$


Analyze it

- Suppose an object has mass 60 kg and γ is 0.2 kg/m:


$$v' = 9.8 - v^2/300$$

- What is happening?
- How can you tell if the speed is increasing or decreasing?
- What is terminal velocity?

Direction Fields



Direction Fields



Direction Field Plotter

dfield

<http://math.rice.edu/~dfield/dfpp.html>

also linked on the class website

Is this a good model?

- Does it agree with experimental results?
- What assumptions does our model make?
- Where does the model break down?

Classifying Differential Equations

- If it looks like [----] then do [----].
- Learn to recognize patterns.
- Call the equations by their names.


Ordinary or Partial?



Harder

- Ordinary differential equation (ODE): involves a function of **one variable**.
- Partial differential equation (PDE): involves a function of 2+ variables (and **partial derivatives**).
- In this class, we will study ODEs only.

Single equation or system of equations?



- We will study a single function at a time.
- You might have several functions that depend on one another.

First order, second order, or higher?



Harder

- Order refers to the highest derivative that appears in the equation.

$$(1 - x^2)y'' - xy' + 3y = 0$$

$$\frac{dy}{dx} = \frac{3x^2 + 4 + 2}{2(y - 1)}$$

Linear or nonlinear?



Harder

- Linear:

$$t^2 y'' + y' + (\sin t)y = \sqrt{t}$$

- Nonlinear:

$$y' = \sin y$$

$$\sqrt{y'} + y = 0$$

$$(y')^2 = t$$

$$yy' = t^2$$

Homogeneous or non-homogeneous



Harder

- This is a way to classify linear equations

- Homogeneous

$$ty'' + (\sin t)y' + y = 0$$

- Non-homogeneous

$$ty'' + (\sin t)y' + y = t$$

Classify it!

$$ty' + y = 1$$

Ordinary, linear, first order, non-homogeneous, single equation

$$yy'' = 0$$

Ordinary, nonlinear, 2nd order, single equation

$$\begin{cases} y_1' + y_2' = \cos t \\ y_1 + y_2 = \sin t \end{cases}$$

Ordinary, first order, linear, non-homogeneous, system of equations