

Data-Driven Neuroimaging Analysis

Nathaniel E. Helwig

Assistant Professor of Psychology and Statistics
University of Minnesota (Twin Cities)



UMN VA Brain Sciences Center — June 10, 2015

Outline of Talk

1) Component Analysis Methods:

- Bilinear Models
- Multiway Models
- Some analyses and results

2) Functional Modeling:

- Parametric Regression
- Nonparametric Regression
- Some analyses and results

Component Analysis Methods

Bilinear Model Form

Let $\mathbf{X} = \{x_{ij}\}_{I \times J}$ denote a data matrix.

- I locations (voxels or channels) by J time points

A **bilinear model** with $R \leq \min(I, J)$ components has the form

$$\mathbf{X} = \mathbf{AB}' + \mathbf{E} \quad \Longleftrightarrow \quad x_{ij} = \sum_{r=1}^R a_{ir} b_{jr} + e_{ij}$$

where

- $\mathbf{A} = \{a_{ir}\}_{I \times R}$ are the voxel weights (spatial maps)
- $\mathbf{B} = \{b_{jr}\}_{J \times R}$ are the time weights (time maps)
- $\mathbf{E} = \{e_{ij}\}_{I \times J}$ are the error terms

Rotational Indeterminacy Problem

Suppose that $\mathbf{R}$ is an $R \times R$ orthogonal rotation matrix.

- $\mathbf{R}'\mathbf{R} = \mathbf{R}\mathbf{R}' = \mathbf{I}$ (identity matrix)

Rotational indeterminacy problem of the bilinear model:

$$\mathbf{AB}' = \tilde{\mathbf{A}}\tilde{\mathbf{B}}'$$

where $\tilde{\mathbf{A}} = \mathbf{AR}$ and $\tilde{\mathbf{B}} = \mathbf{BR}$.

Need to make some (non-data-driven) assumptions to solve the rotational indeterminacy.

Principal Components and Independent Components

Principal Component Analysis Assumptions:

- Columns of **A** are orthogonal; $d_1 \geq \dots \geq d_R$ where $\mathbf{D} = \text{diag}(\mathbf{A}'\mathbf{A})$
- Columns of **B** are orthonormal
- See Pearson (1901) and Hotelling (1933)

Independent Component Analysis Assumptions:

- Columns of **A** are orthogonal and statistically independent
- Columns of **B** are orthonormal
- See Comon (1994) and De Lathauwer et al. (2000)

Historical Development of Multiway Models

Cattell's (1944) **Principle of Parallel Proportional Profiles:**

- Attempted to resolve rotational indeterminacy in bilinear model
- “Real” components should have an invariant structure across different conditions, and should vary proportionally depending on the relative importance of each component in each condition

Some popular multiway models:

- Tucker's (1966) 3-way factor model (Tucker3 Model)
- Harshman's (1970) Parallel Factors (PARAFAC) Model
- Carroll and Chang's (1970) Canonical Decomposition Approach
- Harshman's (1972b) Parallel Factors 2 (PARAFAC2) Model

PARAFAC/CANDECOMP Model Form

Let $\mathbf{X} = \{x_{ijk}\}_{I \times J \times K}$ denote a three-way data array.

- x_{ijk} is signal from i -th voxel at j -th time point of k -th subject

A **trilinear model** has the form


$$x_{ijk} = \sum_{r=1}^R a_{ir} b_{jr} c_{kr} + e_{ijk}$$

where


- $\mathbf{A} = \{a_{ir}\}_{I \times R}$ are the Mode A (voxel) weights
- $\mathbf{B} = \{b_{jr}\}_{J \times R}$ are the Mode B (time) weights
- $\mathbf{C} = \{c_{kr}\}_{K \times R}$ are the Mode C (subject) weights
- $\mathbf{E} = \{e_{ijk}\}_{I \times J \times K}$ is the 3-way error array

Visualization of PCA and Parafac (Helwig et al., 2012)

$$\text{a) PCA: } \mathbf{X} = \sum_{r=1}^R \mathbf{a}_r \circ \mathbf{b}_r + \mathbf{E}$$



$$\text{b) Parafac: } \underline{\mathbf{X}} = \sum_{r=1}^R \mathbf{a}_r \circ \mathbf{b}_r \circ \mathbf{c}_r + \underline{\mathbf{E}}$$



Simultaneous Component Analysis Framework

Let $\mathbf{X}_k = \{x_{ij(k)}\}_{I \times J_k}$ denote a two-way data

- $x_{ij(k)}$ is signal from i -th voxel at j -th time point of k -th subject

The **Simultaneous Component Analysis** framework assumes

$$\mathbf{X}_k = \mathbf{A}\mathbf{D}'_k + \mathbf{E}_k$$

where

- $\mathbf{A} = \{a_{ir}\}_{I \times R}$ are the common Mode A (voxel) weights
- $\mathbf{D}_k = \{d_{jr(k)}\}_{J_k \times R}$ are the subject-specific Mode B (time) weights
- $\mathbf{E}_k = \{e_{ij(k)}\}_{I \times J_k}$ is the 2-way error array

SCA Covariance Structures

Four different SCA models assume different covariance structures:

$$\text{SCA-P: } J_k^{-1} \mathbf{D}_k' \mathbf{D}_k = \Phi_k$$

$$\text{SCA-PF2: } J_k^{-1} \mathbf{D}_k' \mathbf{D}_k = \mathbf{C}_k \Phi \mathbf{C}_k$$

$$\text{SCA-IND: } J_k^{-1} \mathbf{D}_k' \mathbf{D}_k = \mathbf{C}_k^2$$

$$\text{SCA-ECP: } J_k^{-1} \mathbf{D}_k' \mathbf{D}_k = \Phi$$

where

- Φ_k is cross-product matrix specific to k -th subject
- Φ is cross-product matrix common to all K subjects
- $\mathbf{C}_k = \text{diag}(\mathbf{c}_k)$ where $\mathbf{c}_k$ is k -th row of $\mathbf{C} = \{c_{kr}\}_{K \times R}$

Uniqueness of Multiway Solutions

Multiway models can have essentially unique solutions

- No rotational indeterminacy!


The following models have an **intrinsic axis property**:

- Parafac, see Harshman (1970, 1972a) and Kruskal (1977)
- SCA-PF2, see Harshman (1972b); Harshman and Lundy (1996)
- SCA-IND, see Timmerman & Kiers (2003)

The following models have a rotational indeterminacy

- Tucker3 model
- SCA-P model
- SCA-ECP model

Some Simulated fMRI Data (Helwig & Hong, 2013)



Spatial pattern of activity differs across four patterns:

- (a) No overlapping activity
- (b) 10% overlapping activity
- (c) 30% overlapping activity
- (d) 50% overlapping activity

Parafac versus TPICA for fMRI Data Analysis

TPICA: Tensor Probabilistic Independent Component Analysis

- Combines ideas of ICA and Parafac
- Assumes spatial maps are independent


Comparing TPICA and Parafac for fMRI:

- If independence assumption is correct. . .
 - TPICA outperforms Parafac at small SNR
 - Orthogonal-Parafac performs as well as TPICA at small SNR
- If independence assumption is incorrect. . .
 - TPICA produces noticeable bias for all SNR
 - Orthogonal-Parafac produces smaller bias than TPICA

See Helwig & Hong (2013) for thorough critique of TPICA

Parafac versus TPICA (SNR=0.10)

Figure 4 from Helwig and Hong (2013):



Visual Stimulus ERP Study

Data from UCI Machine Learning Repository; originally collected by **Henri Begleiter** at the Neurodynamics Laboratory at SUNY Brooklyn.

Experiment Details:

- Data from 120 subjects
 - 76 alcoholics
 - 44 controls
- 64-channel EEG cap
 - Ground: Nz
 - Reference: Cz
- Snodgrass & Vanderwart
 - Set of 260 pictures
- 10 trials sampled at 256 Hz for one second after stimulus





Figure: The 64-channel EEG cap.

SCA-PF2 Solution ($R^2 = 0.68$)



Functional Modeling

Multiple Linear Regression Model

The **multiple linear regression** model has the form

$$y_i = b_0 + \sum_{j=1}^p b_j x_{ij} + e_i \quad \Longleftrightarrow \quad \mathbf{y} = \mathbf{X}\mathbf{b} + \mathbf{e}$$

for $i \in \{1, \dots, n\}$ where

- $y_i \in \mathbb{R}$ is the real-valued **response** for the i -th observation
- $b_0 \in \mathbb{R}$ is the regression **intercept**
- $b_j \in \mathbb{R}$ is the j -th predictor's regression **slope**
- $x_{ij} \in \mathbb{R}$ is the j -th **predictor** for the i -th observation
- $e_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, \sigma^2)$ is Gaussian **measurement error**

Ordinary Least Squares and Inference

Typically minimize the ordinary least squares (OLS) problem

$$\sum_{i=1}^n (y_i - \hat{y}_i)^2$$

where $\hat{y}_i = \hat{b}_0 + \sum_{j=1}^p \hat{b}_j x_{ij}$ is the fitted value.

$\hat{\mathbf{b}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ and from assumptions $\hat{\mathbf{b}} \sim N(\mathbf{b}, \sigma^2(\mathbf{X}'\mathbf{X})^{-1})$.

- Inference follows from the normality assumption

Limitations of Parametric Regression

MLR parameters have conditional interpretation.

- b_j gives effect of X_j holding other X 's constant
- Hard to interpret if X_j variables are correlated

MLR assumes effects have known parametric (e.g., linear) relationship.

- With real data, one rarely knows the parametric form
- Typically we collect data because we want to discover the form of the relationships between variables

Nonparametric Regression Model

Assume observed data follow a **nonparametric regression** model

$$y_i = \eta(\mathbf{x}_i) + e_i$$

where

- $y_i \in \mathbb{R}$ is observed response for the i -th observation
- $\mathbf{x}_i \equiv (x_{i1}, \dots, x_{ip})'$ is the i -th observation's known predictor vector
- η is an unknown smooth function
- $e_i \stackrel{\text{iid}}{\sim} N(0, \sigma^2)$ is unobserved measurement error

Goal is to estimate the unknown function η from the noisy data.

Model Building and Penalized Least Squares

You can include any combination of effects in the model. For example, with two predictors $\mathbf{x} = (x_1, x_2)$ we could fit two possible models:

- Additive: $\eta(\mathbf{x}) = \eta_0 + \eta_1(x_1) + \eta_2(x_2)$
- Interaction: $\eta(\mathbf{x}) = \eta_0 + \eta_1(x_1) + \eta_2(x_2) + \eta_{12}(x_1, x_2)$

Once η is specified, minimize penalized least squares functional

$$\frac{1}{n} \sum_{i=1}^n (y_i - \eta(\mathbf{x}_i))^2 + \lambda J(\eta)$$

where

- $J(\eta) > 0$ is a **penalty functional** quantifying roughness of η
- $\lambda > 0$ is a **smoothing parameter** controlling influence of J

Smoothing Spline Analysis of Variance

Estimates η in tensor product reproducing kernel Hilbert space $\mathcal{H}$.

Two-way SSANOVA useful for neuroimaging data analysis:

$$\eta(\mathbf{x}) = \eta_0 + \eta_1(\text{space}) + \eta_2(\text{time}) + \eta_{12}(\text{space, time})$$

Three-way SSANOVA useful for neuroimaging data analysis:

$$\begin{aligned}\eta(\mathbf{x}) = & \eta_0 + \eta_1(\text{space}) + \eta_2(\text{time}) + \eta_3(\text{group}) \\ & + \eta_{12}(\text{space, time}) + \eta_{13}(\text{space, group}) \\ & + \eta_{23}(\text{time, group}) + \eta_{123}(\text{space, time, group})\end{aligned}$$

Visual Stimulus ERP Study

Data from UCI Machine Learning Repository; originally collected by **Henri Begleiter** at the Neurodynamics Laboratory at SUNY Brooklyn.

Experiment Details:

- Data from 20 subjects
 - 10 alcoholics
 - 10 controls
- 64-channel EEG cap
 - Ground: Nz
 - Reference: Cz
- Snodgrass & Vanderwart
 - Set of 260 pictures
- 10 trials sampled at 256 Hz for one second after stimulus





Figure: The 64-channel EEG cap.

Data

For the multi-electrode (spatial) analysis, I focus on EEG activity at all 61 electrodes from controls and alcoholics

- Electrode: $x_{i1} \in \{1, 2, 3, 4, \dots, 61\} \rightarrow \mathbb{R}^3$
- Time: $x_{i2} \in \{0, 1, 2, 3, \dots, 255\} \rightarrow [0, 1]$
- Group: $x_{i3} \in \{\text{control, alcoholic}\} \rightarrow \{1, 2\}$

Map each electrode to a 3-dimensional coordinate in Euclidean space.



Analyses

Three-way SSANOVA on $\mathcal{X} \equiv \mathbb{R}^3 \times [0, 1] \times \{1, 2\}$


- Thin-plate spline for electrode effect
- Cubic spline for time effect
- Nominal spline for group effect

I analyzed data from all subjects and electrodes, so that $n = 3123200$ (20 subjects $\times$ 256 time points $\times$ 61 electrodes $\times$ 10 trials).

Examined the solution using $q = 1520$ bin-sampled knots.


Predicted ERPs

Can predict function in 3D space with dense cap (633 electrodes), and then project results to 2D space for visualization.



Useful for visualizing **predicted ERPs and ERP differences**.

Predicted ERPs



Some R Packages

Some R packages for multiway modeling:

- `ThreeWay`: Three-way component analysis. (Ferraro et al., 2014)
- `multiway`: Component Models for Multiway Data. (Helwig, 2015b)

Some R packages for nonparametric regression:

- `gss`: General Smoothing Splines. (Gu, 2014)
- `mgcv`: Mixed GAM Computation Vehicle. (Wood, 2015)
- `bigsplines`: Smoothing Splines for Large Samples. (Helwig, 2015a)

References I

- Carroll, J. D. and J.-J. Chang (1970). Analysis of individual differences in multidimensional scaling via an n-way generalization of "eckart-young" decomposition. *Psychometrika* 35, 283–319.
- Cattell, R. B. (1944). "Parallel Proportional Profiles" and other principles for determining the choice of factors by rotation. *Psychometrika* 9, 267–283.
- Comon, P. (1994). Independent component analysis, a new concept? *Signal Processing* 36, 287–314.
- De Lathauwer, L., B. De Moor, and J. Vandewalle (2000). An introduction to independent component analysis. *Journal of Chemometrics* 14, 123–149.
- Ferraro, M. A. D., H. A. L. Kiers, and P. Giordani (2014). *Component analysis for three-way data arrays by means of Candecomp/Parafac, Tucker3, Tucker2 and Tucker1 models*. R package version 1.1.2.
- Gu, C. (2014). *gss: General Smoothing Splines*. R package version 2.1-4.
- Harshman, R. A. (1970). Foundations of the PARAFAC procedure: Models and conditions for an "explanatory" multimodal factor analysis. *UCLA Working Papers in Phonetics* 16, 1–84.
- Harshman, R. A. (1972a). Determination and proof of minimum uniqueness conditions for parafac1. *UCLA Working Papers in Phonetics* 22, 111–117.
- Harshman, R. A. (1972b). Parafac2: Mathematical and technical notes. *UCLA Working Papers in Phonetics* 22, 30–44.
- Harshman, R. A. and M. E. Lundy (1996). Uniqueness proof for a family of models sharing features of Tucker's three-mode factor analysis and PARAFAC/CANDECOMP. *Psychometrika* 61, 133–154.
- Helwig, N. E. (2015a). *bigsplines: Smoothing Splines for Large Samples*. R package version 1.0-6.
- Helwig, N. E. (2015b). *multiway: Component Models for Multi-Way Data*. R package version 1.0-0.
- Helwig, N. E. and S. Hong (2013). A critique of Tensor Probabilistic Independent Component Analysis: Implications and recommendations for multi-subject fMRI data analysis. *Journal of Neuroscience Methods* 213, 263–273.
- Helwig, N. E., S. Hong, and J. D. Polk (2012). Parallel Factor Analysis of gait waveform data: A multimode extension of Principal Component Analysis. *Human Movement Science* 31, 630–648.

References II

- Hotelling, H. (1933). Analysis of a complex of statistical variables into principal components. *Journal of Educational Psychology* 24, 417–441, 498–520.
- Kruskal, J. B. (1977). Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics. *Linear Algebra and Its Applications* 18, 95–138.
- Pearson, K. (1901). One lines and planes of closest fit to systems of points in space. *Philosophical Magazine* 2, 559–572.
- Timmerman, M. E. and H. A. L. Kiers (2003). Four simultaneous component analysis models for the analysis of multivariate time series from more than one subject to mode intraindividual and interindividual differences. *Psychometrika* 68, 105–121.
- Tucker, L. R. (1966). Some mathematical notes on three-mode factor analysis. *Psychometrika* 31, 279–311.
- Wood, S. N. (2015). *mgcv: Mixed GAM Computation Vehicle with GCV/AIC/REML smoothness estimation and GAMMs by REML/PQL*. R package version 1.8-5.