

Linear Algebra Primer

Nathaniel E. Helwig

Associate Professor of Psychology and Statistics
University of Minnesota



January 12, 2024

Copyright © 2024 by Nathaniel E. Helwig

Table of Contents

1. Basic Definitions

- Vector and Matrix
- Transpose and Trace
- Symmetric and Diagonal
- Special Matrices

2. Basic Calculations

- Equality
- Addition/Subtraction
- Vector Products
- Matrix Products

3. Matrix Decompositions

- Eigenvalue Decomposition
- Cholesky Decomposition
- Singular Value Decomp
- QR Decomposition

4. Miscellaneous Topics

- Definiteness
- Determinants
- Inverses
- R Code

Content adapted from:

Johnson, R. A., & Wichern, D. W. (2007). Applied Multivariate Statistical Analysis (6th ed).

Table of Contents

1. Basic Definitions

Vector and Matrix
Transpose and Trace
Symmetric and Diagonal
Special Matrices

2. Basic Calculations

Equality
Addition/Subtraction
Vector Products
Matrix Products

3. Matrix Decompositions

Eigenvalue Decomposition
Cholesky Decomposition
Singular Value Decomp
QR Decomposition

4. Miscellaneous Topics

Definiteness
Determinants
Inverses
R Code

Vectors and Matrices

A **vector** is a one-dimensional array: $\mathbf{a} = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}_{n \times 1}$

A **matrix** is a two-dimensional array: $\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1p} \\ a_{21} & a_{22} & \cdots & a_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{np} \end{bmatrix}_{n \times p}$

The **order** of a matrix refers to the number of rows and columns:

- $\mathbf{a}$ has order n -by-1
- $\mathbf{A}$ has order n -by- p

Rank of a Matrix

The **rank** of $\mathbf{A}$ is the number of linearly independent rows/columns.

- **column rank** of $\mathbf{A}$ is number of linearly independent columns
- **row rank** of $\mathbf{A}$ is number of linearly independent rows

We say that $\mathbf{A}$ is **full rank** if $\text{rank}(\mathbf{A}) = \min(n, p)$.

- If $n < p$, **full rank** implies **full row rank**, i.e., $\text{rank}(\mathbf{A}) = n$
- If $n > p$, **full rank** implies **full column rank**, i.e., $\text{rank}(\mathbf{A}) = p$

Rank Example

The matrix $\mathbf{A}$ is NOT full rank

$$\mathbf{A} = \begin{bmatrix} 1 & 3 \\ 2 & 6 \\ 5 & 15 \end{bmatrix}$$

because we have $3\mathbf{a}_1 = \mathbf{a}_2$ where $\mathbf{a}_j$ denotes the j -th column of $\mathbf{A}$.

In contrast, the matrix $\mathbf{A}$ is full rank

$$\mathbf{A} = \begin{bmatrix} 1 & 3 \\ 2 & 6 \\ 4 & 15 \end{bmatrix}$$

because we cannot write $\sum_{j=1} b_j \mathbf{a}_j = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$ unless we set $b_j = 0 \ \forall j$.

Matrix Transpose: Definition

We will denote the **transpose** with a $^\top$ symbol.

The **transpose** of a vector turns a column vector into a row vector:

$$\mathbf{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}_{n \times 1} \iff \mathbf{a}^\top = [a_1 \quad a_2 \quad \cdots \quad a_n]_{1 \times n}$$

The **transpose** of a matrix exchanges rows and columns, such as

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1p} \\ a_{21} & a_{22} & \cdots & a_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{np} \end{bmatrix}_{n \times p} \iff \mathbf{A}^\top = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{n1} \\ a_{12} & a_{22} & \cdots & a_{n2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1p} & a_{2p} & \cdots & a_{np} \end{bmatrix}_{p \times n}$$

Matrix Transpose: Example

The transpose of $\mathbf{a} = \begin{bmatrix} 1 \\ 7 \\ 5 \\ 9 \end{bmatrix}_{4 \times 1}$ is given by $\mathbf{a}^\top = [1 \ 7 \ 5 \ 9]_{1 \times 4}$

The transpose of $\mathbf{A} = \begin{bmatrix} 1 & 3 \\ 7 & 2 \\ 5 & 7 \\ 9 & 4 \end{bmatrix}_{4 \times 2}$ is given by $\mathbf{A}^\top = \begin{bmatrix} 1 & 7 & 5 & 9 \\ 3 & 2 & 7 & 4 \end{bmatrix}_{2 \times 4}$

Matrix Transpose: Properties

Some useful properties of matrix transposes include:

- $(\mathbf{A}^\top)^\top = \mathbf{A}$
- $(\mathbf{A} + \mathbf{B})^\top = \mathbf{A}^\top + \mathbf{B}^\top$ (where $\mathbf{A} + \mathbf{B}$ is matrix addition)
- $(b\mathbf{A})^\top = b\mathbf{A}^\top$ (where $b\mathbf{A}$ is scalar multiplication)
- $(\mathbf{AB})^\top = \mathbf{B}^\top \mathbf{A}^\top$ (where $\mathbf{AB}$ is matrix multiplication)
- $(\mathbf{A}^{-1})^\top = (\mathbf{A}^\top)^{-1}$ (where $\mathbf{A}^{-1}$ is matrix inverse)

Matrix Trace: Definition

The **trace** of a square matrix $\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1p} \\ a_{21} & a_{22} & \cdots & a_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ a_{p1} & a_{p2} & \cdots & a_{pp} \end{bmatrix}_{p \times p}$ is

$$\text{tr}(\mathbf{A}) = \sum_{j=1}^p a_{jj} \quad (1)$$

which is the sum of the diagonal elements.

Matrix Trace: Example

The trace of the matrix $\mathbf{A} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \\ 5 & 9 & 4 & 3 \end{bmatrix}$ is

$$\begin{aligned}\operatorname{tr}(\mathbf{A}) &= 1 + 8 + 6 + 3 \\ &= 18\end{aligned}$$

Matrix Trace: Properties

Some useful properties of matrix traces include:

- $\text{tr}(\mathbf{A}) = \text{tr}(\mathbf{A}^\top)$
- $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}(\mathbf{A}) + \text{tr}(\mathbf{B})$
- $\text{tr}(b\mathbf{A}) = b\text{tr}(\mathbf{A})$
- $\text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{BA})$ if both products are defined
- If $\mathbf{A}$ is symmetric, $\text{tr}(\mathbf{A}) = \sum_{j=1}^p \lambda_j$ where λ_j is j -th eigenvalue

Symmetric Matrix: Definition

A **symmetric** matrix is square and symmetric along the main diagonal:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}_{n \times n} \quad (2)$$

with $a_{ij} = a_{ji}$ for all $i \neq j$.

Note that $\mathbf{A} = \mathbf{A}^\top$ for all symmetric matrices (by definition).

Symmetric Matrix: Example

The matrix $\mathbf{A} = \begin{bmatrix} 9 & 1 & 0 & 4 \\ 1 & 4 & 2 & 1 \\ 0 & 2 & 5 & 6 \\ 4 & 1 & 6 & 8 \end{bmatrix}$ is a symmetric 4×4 matrix.

The matrix $\mathbf{A} = \begin{bmatrix} 9 & 1 & 0 & 4 \\ 1 & 4 & 2 & 1 \\ 0 & 2 & 5 & 6 \\ 3 & 1 & 6 & 8 \end{bmatrix}$ is NOT a symmetric 4×4 matrix.

Diagonal Matrix

A **diagonal** matrix is a square matrix that has zeros in the off-diagonals:

$$\mathbf{D} = \begin{bmatrix} d_1 & 0 & 0 & \cdots & 0 \\ 0 & d_2 & 0 & \cdots & 0 \\ 0 & 0 & d_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & d_p \end{bmatrix}_{p \times p} \quad (3)$$

We often write $\mathbf{D} = \text{diag}(d_1, \dots, d_p)$ to define a diagonal matrix.

Identity Matrix

The **identity matrix** of order p is a $p \times p$ matrix that has ones along the main diagonal and zeros in the off-diagonals:

$$\mathbf{I}_p = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}_{p \times p} \quad (4)$$

Note that $\mathbf{I}_p$ is a special type of diagonal matrix.

Zero and One Matrices

A vector or matrix of all zeros will be denoted using the notation:

$$\mathbf{0}_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{n \times 1}$$

$$\mathbf{0}_{n \times p} = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}_{n \times p}$$

A vector or matrix of all ones will be denoted using the notation:

$$\mathbf{1}_n = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}_{n \times 1}$$

$$\mathbf{1}_{n \times p} = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 1 \end{bmatrix}_{n \times p}$$

Table of Contents

1. Basic Definitions

Vector and Matrix

Transpose and Trace

Symmetric and Diagonal

Special Matrices

2. Basic Calculations

Equality

Addition/Subtraction

Vector Products

Matrix Products

3. Matrix Decompositions

Eigenvalue Decomposition

Cholesky Decomposition

Singular Value Decomp

QR Decomposition

4. Miscellaneous Topics

Definiteness

Determinants

Inverses

R Code

Matrix Equality

Given two matrices of the same order $\mathbf{A} = \{a_{ij}\}_{n \times p}$ and $\mathbf{B} = \{b_{ij}\}_{n \times p}$, $\mathbf{A}$ is equal to $\mathbf{B}$ (written $\mathbf{A} = \mathbf{B}$) if and only if $a_{ij} = b_{ij} \forall i, j$.

$$\text{If } \mathbf{A} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \end{bmatrix}, \text{ then } \mathbf{A} = \mathbf{B}.$$

$$\text{If } \mathbf{A} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \end{bmatrix} \text{ and } \mathbf{B} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 0 \end{bmatrix}, \text{ then } \mathbf{A} \neq \mathbf{B}.$$

Matrix Addition and Subtraction: Definition

Given two matrices of the same order $\mathbf{A} = \{a_{ij}\}_{n \times p}$ and $\mathbf{B} = \{b_{ij}\}_{n \times p}$, the addition $\mathbf{A} + \mathbf{B}$ produces $\mathbf{C} = \{c_{ij}\}_{n \times p}$ such that $c_{ij} = a_{ij} + b_{ij}$.

Given two matrices of the same order $\mathbf{A} = \{a_{ij}\}_{n \times p}$ and $\mathbf{B} = \{b_{ij}\}_{n \times p}$, the subtraction $\mathbf{A} - \mathbf{B}$ produces $\mathbf{C} = \{c_{ij}\}_{n \times p}$ such that $c_{ij} = a_{ij} - b_{ij}$.

Note: matrix addition and subtraction is only defined for two matrices of the same order.

Matrix Addition and Subtraction: Example

Given $\mathbf{A} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} 5 & 6 & 1 & 7 \\ 1 & 3 & 0 & 2 \\ 2 & 5 & 3 & 5 \end{bmatrix}$, we have that

$$\mathbf{A} + \mathbf{B} = \begin{bmatrix} 1+5 & 4+6 & 8+1 & 13+7 \\ 2+1 & 8+3 & 11+0 & 2+2 \\ 7+2 & 2+5 & 6+3 & 9+5 \end{bmatrix} = \begin{bmatrix} 6 & 10 & 9 & 20 \\ 3 & 11 & 11 & 4 \\ 9 & 7 & 9 & 14 \end{bmatrix}$$

$$\mathbf{A} - \mathbf{B} = \begin{bmatrix} 1-5 & 4-6 & 8-1 & 13-7 \\ 2-1 & 8-3 & 11-0 & 2-2 \\ 7-2 & 2-5 & 6-3 & 9-5 \end{bmatrix} = \begin{bmatrix} -4 & -2 & 7 & 6 \\ 1 & 5 & 11 & 0 \\ 5 & -3 & 3 & 4 \end{bmatrix}$$

Vector Inner Products: Definition

The **inner product** of $\mathbf{x} = (x_1, \dots, x_n)^\top$ and $\mathbf{y} = (y_1, \dots, y_n)^\top$ is

$$\begin{aligned}\mathbf{x}^\top \mathbf{y} &= [x_1 \cdots x_n] \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix} \\ &= [\sum_{i=1}^n x_i y_i]_{1 \times 1}\end{aligned}\tag{5}$$

Note that $\mathbf{x}$ and $\mathbf{y}$ must have the same length (i.e., n).

Vector Inner Products: Example

Given $\mathbf{x} = (3, 9, -2, 5)^\top$ and $\mathbf{y} = (2, 0, 2, 1)^\top$, we have that

$$\begin{aligned}\mathbf{x}^\top \mathbf{y} &= \begin{bmatrix} 3 & 9 & -2 & 5 \end{bmatrix} \begin{bmatrix} 2 \\ 0 \\ 2 \\ 1 \end{bmatrix} \\ &= 3(2) + 9(0) - 2(2) + 5(1) \\ &= 7\end{aligned}$$

Vector Outer Products: Definition

The **outer product** of $\mathbf{x} = (x_1, \dots, x_m)^\top$ and $\mathbf{y} = (y_1, \dots, y_n)^\top$ is

$$\begin{aligned}\mathbf{xy}^\top &= \begin{bmatrix} x_1 \\ \vdots \\ x_m \end{bmatrix} [y_1 \cdots y_n] \\ &= \begin{bmatrix} x_1 y_1 & x_1 y_2 & \cdots & x_1 y_n \\ x_2 y_1 & x_2 y_2 & \cdots & x_2 y_n \\ \vdots & \vdots & \ddots & \vdots \\ x_m y_1 & x_m y_2 & \cdots & x_m y_n \end{bmatrix}_{m \times n}\end{aligned}\tag{6}$$

Note that $\mathbf{x}$ and $\mathbf{y}$ can have different lengths (i.e., m and n).

Vector Outer Products: Example

Given $\mathbf{x} = (3, 9, -2, 5)^\top$ and $\mathbf{y} = (2, 0, 2, 1)^\top$, we have that

$$\begin{aligned}\mathbf{xy}^\top &= \begin{pmatrix} 3 \\ 9 \\ -2 \\ 5 \end{pmatrix} \begin{bmatrix} 2 & 0 & 2 & 1 \end{bmatrix} \\ &= \begin{pmatrix} 6 & 0 & 6 & 3 \\ 18 & 0 & 18 & 9 \\ -4 & 0 & -4 & -2 \\ 10 & 0 & 10 & 5 \end{pmatrix}\end{aligned}$$

Matrix-Scalar Products: Definition

The **matrix-scalar product** of $\mathbf{A} = \{a_{ij}\}_{n \times p}$ and $b \in \mathbb{R}$ is

$$\mathbf{A}b = b\mathbf{A} = \begin{bmatrix} ba_{11} & ba_{12} & \cdots & ba_{1p} \\ ba_{21} & ba_{22} & \cdots & ba_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ ba_{n1} & ba_{n2} & \cdots & ba_{np} \end{bmatrix}_{n \times p} \quad (7)$$

which is the matrix $\mathbf{C} = \{c_{ij}\}_{n \times p}$ such that $c_{ij} = ba_{ij}$.

Matrix-Scalar Products: Example

Given $\mathbf{A} = \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \end{bmatrix}$ and $b = 2$, we have that

$$\begin{aligned} b\mathbf{A} &= \begin{bmatrix} 1 & 4 & 8 & 13 \\ 2 & 8 & 11 & 2 \\ 7 & 2 & 6 & 9 \end{bmatrix} 2 \\ &= \begin{bmatrix} 2 & 8 & 16 & 26 \\ 4 & 16 & 22 & 4 \\ 14 & 4 & 12 & 18 \end{bmatrix} \end{aligned}$$

Matrix-Vector Products: Definition

The **matrix-vector product** of $\mathbf{A} = \begin{bmatrix} a_{11} & \cdots & a_{1p} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{np} \end{bmatrix}$ and $\mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_p \end{bmatrix}$ is

$$\begin{aligned} \mathbf{Ax} &= \begin{bmatrix} a_{11} & \cdots & a_{1p} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{np} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_p \end{bmatrix} \\ &= \begin{bmatrix} \sum_{j=1}^p a_{1j}x_j \\ \vdots \\ \sum_{j=1}^p a_{nj}x_j \end{bmatrix}_{n \times 1} \end{aligned} \tag{8}$$

Note that length of $\mathbf{x}$ must match number of columns of $\mathbf{A}$ (i.e., p).

Matrix-Vector Products: Example

Given $\mathbf{A} = \begin{bmatrix} 3 & 4 & 1 \\ 4 & 7 & 5 \end{bmatrix}$ and $\mathbf{x} = \begin{bmatrix} 1 \\ 6 \\ 3 \end{bmatrix}$, we have that

$$\begin{aligned}\mathbf{Ax} &= \begin{bmatrix} 3 & 4 & 1 \\ 4 & 7 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 6 \\ 3 \end{bmatrix} \\ &= \begin{bmatrix} 3(1) + 4(6) + 1(3) \\ 4(1) + 7(6) + 5(3) \end{bmatrix} \\ &= \begin{bmatrix} 30 \\ 61 \end{bmatrix}\end{aligned}$$

Matrix-Matrix Products: Definition

The **matrix-matrix product** of $\mathbf{A} = \{a_{ij}\}_{m \times n}$ and $\mathbf{B} = \{b_{jk}\}_{n \times p}$ is

$$\begin{aligned} \mathbf{AB} &= \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & \cdots & b_{1p} \\ \vdots & \ddots & \vdots \\ b_{n1} & \cdots & b_{np} \end{bmatrix} \\ &= \begin{bmatrix} \sum_{j=1}^n a_{1j}b_{j1} & \sum_{j=1}^n a_{1j}b_{j2} & \cdots & \sum_{j=1}^n a_{1j}b_{jp} \\ \sum_{j=1}^n a_{2j}b_{j1} & \sum_{j=1}^n a_{2j}b_{j2} & \cdots & \sum_{j=1}^n a_{2j}b_{jp} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{j=1}^n a_{mj}b_{j1} & \sum_{j=1}^n a_{mj}b_{j2} & \cdots & \sum_{j=1}^n a_{mj}b_{jp} \end{bmatrix}_{m \times p} \end{aligned} \quad (9)$$

Note that $\#$ of rows of $\mathbf{B}$ must match $\#$ of columns of $\mathbf{A}$ (i.e., n), and note that $\mathbf{AB} \neq \mathbf{BA}$ even if both products are defined.

Matrix-Matrix Products: Example

Given $\mathbf{A} = \begin{bmatrix} 3 & 4 & 1 \\ 4 & 7 & 5 \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} 1 & 2 \\ 6 & 1 \\ 3 & 4 \end{bmatrix}$, we have that

$$\begin{aligned} \mathbf{AB} &= \begin{bmatrix} 3 & 4 & 1 \\ 4 & 7 & 5 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 6 & 1 \\ 3 & 4 \end{bmatrix} \\ &= \begin{bmatrix} 3(1) + 4(6) + 1(3) & 3(2) + 4(1) + 1(4) \\ 4(1) + 7(6) + 5(3) & 4(2) + 7(1) + 5(4) \end{bmatrix} \\ &= \begin{bmatrix} 30 & 14 \\ 61 & 35 \end{bmatrix} \end{aligned}$$

Multiplying by Identity Matrix

Given $\mathbf{A} = \{a_{ij}\}_{m \times n}$, pre-multiplying by the identity matrix returns $\mathbf{A}$

$$\mathbf{I}_m \mathbf{A} = \mathbf{A}$$

and post-multiplying by the identity matrix returns $\mathbf{A}$

$$\mathbf{A} \mathbf{I}_n = \mathbf{A}$$

This is the reason we call $\mathbf{I}_m$ and $\mathbf{I}_n$ “identity” matrices.

Table of Contents

1. Basic Definitions

- Vector and Matrix
- Transpose and Trace
- Symmetric and Diagonal
- Special Matrices

2. Basic Calculations

- Equality
- Addition/Subtraction
- Vector Products
- Matrix Products

3. Matrix Decompositions

- Eigenvalue Decomposition
- Cholesky Decomposition
- Singular Value Decomp
- QR Decomposition

4. Miscellaneous Topics

- Definiteness
- Determinants
- Inverses
- R Code

Overview of Matrix Decompositions

A **matrix decomposition** decomposes (i.e., separates) a given matrix into a matrix multiplication of two (or more) simpler matrices.

Matrix decompositions are useful for many things:

- Solving systems of equations
- Obtaining low-rank approximations
- Finding important features of data

We will briefly discuss four matrix decompositions:

- Eigenvalue Decomposition
- Cholesky Decomposition
- Singular Value Decomposition
- QR Decomposition

Eigenvalue (Spectral) Decomposition

The **eigenvalue decomposition** (EVD) decomposes a symmetric¹ matrix $\mathbf{A} = \{a_{ij}\}_{n \times n}$ into a product of three matrices:

$$\mathbf{A} = \mathbf{\Gamma} \mathbf{\Lambda} \mathbf{\Gamma}^\top \quad (10)$$

such that

- $\mathbf{\Gamma} = (\gamma_1 \cdots \gamma_n)_{n \times n}$ where $\gamma_j = (\gamma_{1j}, \dots, \gamma_{nj})^\top$ is j -th **eigenvector**
- $\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_n)$ where λ_j is j -th **eigenvalue**
- Eigenvalues/vectors are ordered such that $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$

Note that $\mathbf{\Gamma}$ is an **orthogonal matrix**: $\mathbf{\Gamma} \mathbf{\Gamma}^\top = \mathbf{\Gamma}^\top \mathbf{\Gamma} = \mathbf{I}_n$

¹EVD is defined for asymmetric matrices, but we will consider symmetric case.

Cholesky Decomposition

The **Cholesky decomposition** (CD) decomposes a positive definite matrix $\mathbf{A} = \{a_{ij}\}_{n \times n}$ into a product of a two matrices:

$$\mathbf{A} = \mathbf{L}\mathbf{L}^\top \quad (11)$$

where

• $\mathbf{L} = \begin{bmatrix} l_{11} & 0 & 0 & \cdots & 0 \\ l_{21} & l_{22} & 0 & \cdots & 0 \\ l_{31} & l_{32} & l_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ l_{n1} & l_{n2} & l_{n3} & \cdots & l_{nn} \end{bmatrix}$ is a **lower (left) triangular** matrix

Singular Value Decomposition

The **singular value decomposition** (SVD) decomposes any matrix $\mathbf{A} = \{a_{ij}\}_{n \times p}$ into a product of three matrices:

$$\mathbf{A} = \mathbf{U}\mathbf{S}\mathbf{V}^\top \quad (12)$$

such that

- $\mathbf{U} = (\mathbf{u}_1 \cdots \mathbf{u}_r)_{n \times r}$ and $\mathbf{u}_k = \{u_{ik}\}_{n \times 1}$ is k -th **left singular vector**
- $\mathbf{S} = \text{diag}(s_1, \dots, s_r)$ and $s_k > 0$ is k -th **singular value**
- $\mathbf{V} = (\mathbf{v}_1 \cdots \mathbf{v}_r)_{p \times r}$ and $\mathbf{v}_k = \{v_{jk}\}_{p \times 1}$ is k -th **right singular vector**
- $r \leq \min(m, n)$ and $r = \min(m, n)$ if $\mathbf{A}$ is full-rank

Note that $\mathbf{U}$ and $\mathbf{V}$ are **columnwise orthogonal**: $\mathbf{U}^\top \mathbf{U} = \mathbf{V}^\top \mathbf{V} = \mathbf{I}_r$

QR Decomposition

The **QR decomposition** (QRD) decomposes any long (i.e., $n \geq p$) matrix $\mathbf{A} = \{a_{ij}\}_{n \times p}$ into a product of two matrices:

$$\begin{aligned}\mathbf{A} &= \mathbf{Q}\mathbf{R} \\ &= [\mathbf{Q}_1 \quad \mathbf{Q}_2] \begin{bmatrix} \mathbf{R}_1 \\ \mathbf{0}_{(n-p) \times p} \end{bmatrix} \\ &= \mathbf{Q}_1 \mathbf{R}_1\end{aligned}\tag{13}$$

such that

- $\mathbf{Q}$ is an orthogonal matrix

- $\mathbf{R}_1 = \begin{bmatrix} r_{11} & r_{12} & r_{13} & \cdots & r_{1p} \\ 0 & r_{22} & r_{23} & \cdots & r_{2p} \\ 0 & 0 & r_{33} & \cdots & r_{3p} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & r_{pp} \end{bmatrix}$ is upper (right) triangular matrix

Table of Contents

1. Basic Definitions

- Vector and Matrix
- Transpose and Trace
- Symmetric and Diagonal
- Special Matrices

2. Basic Calculations

- Equality
- Addition/Subtraction
- Vector Products
- Matrix Products

3. Matrix Decompositions

- Eigenvalue Decomposition
- Cholesky Decomposition
- Singular Value Decomp
- QR Decomposition

4. Miscellaneous Topics

- Definiteness
- Determinants
- Inverses
- R Code

Quadratic Forms

The **quadratic form** of a symmetric matrix $\mathbf{A} = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix}$ is

$$\begin{aligned} \mathbf{x}^\top \mathbf{A} \mathbf{x} &= \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix} \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \left[\sum_{i=1}^n \sum_{j=1}^n x_i x_j a_{ij} \right]_{1 \times 1} \end{aligned} \quad (14)$$

where $\mathbf{x} = \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix}^\top$ is any arbitrary vector of length n .

Positive, Negative, and Semi-Definite Matrices

A symmetric matrix $\mathbf{A} = \{a_{ij}\}_{n \times n}$ is said to be

- **positive definite** if $\mathbf{x}^\top \mathbf{A} \mathbf{x} > 0$ for every $\mathbf{x} \neq \mathbf{0}_n$
- **positive semi-definite** if $\mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0$ for every $\mathbf{x} \neq \mathbf{0}_n$
- **negative definite** if $\mathbf{x}^\top \mathbf{A} \mathbf{x} < 0$ for every $\mathbf{x} \neq \mathbf{0}_n$
- **negative semi-definite** if $\mathbf{x}^\top \mathbf{A} \mathbf{x} \leq 0$ for every $\mathbf{x} \neq \mathbf{0}_n$

Note if $\mathbf{x}^\top \mathbf{A} \mathbf{x} \geq 0$ for some $\mathbf{x}$ and $\mathbf{x}^\top \mathbf{A} \mathbf{x} < 0$ for other $\mathbf{x}$, then $\mathbf{A}$ is said to be an **indefinite** matrix.

Matrix Definiteness: Example

The matrix $\mathbf{A} = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ is positive definite:

$$\begin{aligned}\mathbf{x}^\top \mathbf{A} \mathbf{x} &= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \\ &= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 2x_1 - x_2 \\ -x_1 + 2x_2 \end{bmatrix} \\ &= 2x_1^2 - 2x_1x_2 + 2x_2^2 \\ &= x_1^2 + x_2^2 + (x_1 - x_2)^2 \\ &\geq 0\end{aligned}$$

with the equality holding only when $x_1 = x_2 = 0$.

Matrix Definiteness: Properties

Let λ_j denote the j -th eigenvalue of $\mathbf{A}$ for $j \in \{1, \dots, n\}$.

Some useful properties of matrix definiteness include:

- If $\mathbf{A}$ is positive definite, then $\lambda_j > 0 \forall j$
- If $\mathbf{A}$ is positive semi-definite, then $\lambda_j \geq 0 \forall j$
- If $\mathbf{A}$ is negative definite, then $\lambda_j < 0 \forall j$
- If $\mathbf{A}$ is negative semi-definite, then $\lambda_j \leq 0 \forall j$
- If $\mathbf{A}$ is indefinite, then $\lambda_i > 0$ and $\lambda_j < 0$ for some $i \neq j$

Matrix Determinant: Definition

The **determinant** of a square matrix $\mathbf{A} \in \mathbb{R}^{p \times p}$ is a real-valued function from $\mathbb{R}^{p \times p} \rightarrow \mathbb{R}$, and is typically denoted by $|\mathbf{A}|$ or $\det(\mathbf{A})$.

Determinants provide information about systems of linear equations:

- Suppose that $\mathbf{A} \in \mathbb{R}^{p \times p}$, $\mathbf{x} \in \mathbb{R}^{p \times 1}$, and $\mathbf{b} \in \mathbb{R}^{p \times 1}$
- System $\mathbf{Ax} = \mathbf{b}$ has a unique solution if and only if $|\mathbf{A}| \neq 0$

Determinants provide information about linear transformations:

- Magnitude of $|\mathbf{A}|$ is the transformation's **scale factor**
- Sign of $|\mathbf{A}|$ is the transformation's **orientation**

Matrix Determinant: Calculation

- For 1×1 matrix $\mathbf{A} = (a)$, we have
$$|\mathbf{A}| = a$$
- For 2×2 matrix $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we have
$$|\mathbf{A}| = ad - bc$$
- For 3×3 matrix $\mathbf{A} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$, we have
$$|\mathbf{A}| = aei + bfg + cdh - (ceg + bdi + afh)$$

Matrix Determinant: Calculation (continued)

For $p \times p$ matrix $\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1p} \\ a_{21} & a_{22} & \cdots & a_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ a_{p1} & a_{p2} & \cdots & a_{pp} \end{bmatrix}$, we have

$$|\mathbf{A}| = \sum_{j=1}^p (-1)^{i+j} a_{ij} M_{ij} = \sum_{i=1}^p (-1)^{i+j} a_{ij} M_{ij}$$

where

- $M_{ij} = |\mathbf{A}_{-ij}|$ is the **minor** corresponding to cell (i, j) of $\mathbf{A}$
- $(-1)^{i+j} M_{ij}$ is the **cofactor** corresponding to cell (i, j) of $\mathbf{A}$
- $\mathbf{A}_{-ij}$ is the $(p-1) \times (p-1)$ matrix formed by deleting the i -th row and j -th column of $\mathbf{A}$

Note: can use any column (or row) to define the determinant of $\mathbf{A}$.

Properties of Matrix Determinants

Some useful properties of matrix determinants include:

- $|\mathbf{A}| = |\mathbf{A}^\top|$
- $|\mathbf{A}^{-1}| = |\mathbf{A}|^{-1}$ (where $\mathbf{A}^{-1}$ is defined on the next slide)
- $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$ (if $\mathbf{A}$ and $\mathbf{B}$ are both square)
- $|b\mathbf{A}| = b^p|\mathbf{A}|$ (if $b \in \mathbb{R}$ and $\mathbf{A}$ is $p \times p$)
- If $\mathbf{A}$ is symmetric, $|\mathbf{A}| = \prod_{j=1}^p \lambda_j$ where λ_j is j -th eigenvalue of $\mathbf{A}$.

Matrix Inverses: Definition

A square (not necessarily symmetric) matrix $\mathbf{A} = \{a_{ij}\}_{n \times n}$ is **invertible** (or **nonsingular**) if there exists another matrix $\mathbf{B} = \{b_{ij}\}_{n \times n}$ such that

$$\mathbf{AB} = \mathbf{I}_n \tag{15}$$

where $\mathbf{I}_n$ is the $n \times n$ **identity matrix**.

If $\mathbf{B}$ exists, the matrix $\mathbf{B}$ is called the **inverse** of the matrix $\mathbf{A}$ and is denoted by $\mathbf{A}^{-1}$ (so that $\mathbf{AA}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}_n$).

Matrix Inverses: Calculation for 2×2 Case

Claim:

For 2×2 matrix $\mathbf{A} = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we have $\mathbf{A}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$

Proof:

$$\begin{aligned} \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} &= \frac{1}{ad-bc} \begin{bmatrix} da-bc & db-bd \\ -ca+ac & -cb+ad \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

Matrix Inverses: Example

Given $\mathbf{A} = \begin{bmatrix} 1 & 3 \\ 2 & 1 \end{bmatrix}$, the inverse is $\mathbf{A}^{-1} = \begin{bmatrix} -1/5 & 3/5 \\ 2/5 & -1/5 \end{bmatrix}$:

$$\mathbf{A}\mathbf{A}^{-1} = \begin{bmatrix} 1 & 3 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} -1/5 & 3/5 \\ 2/5 & -1/5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\mathbf{A}^{-1}\mathbf{A} = \begin{bmatrix} -1/5 & 3/5 \\ 2/5 & -1/5 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Matrix Inverses: Properties

Some useful properties of matrix inverses include:

- $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$
- $(b\mathbf{A})^{-1} = b^{-1}\mathbf{A}^{-1}$
- $(\mathbf{A}^{-1})^{\top} = (\mathbf{A}^{\top})^{-1}$
- $\mathbf{A}^{-1} = \mathbf{A}^{\top}$ if and only if $\mathbf{A}$ is orthogonal
- $|\mathbf{A}^{-1}| = |\mathbf{A}|^{-1}$
- $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$ if both $\mathbf{A}^{-1}$ and $\mathbf{B}^{-1}$ exist
- $\mathbf{A}^{-1}$ exists only if $|\mathbf{A}| \neq 0$
- If $\mathbf{A}$ is positive definite, then $\mathbf{A}^{-1} = \mathbf{\Gamma}\mathbf{\Lambda}^{-1}\mathbf{\Gamma}^{\top} = (\mathbf{L}^{-1})^{\top}\mathbf{L}^{-1}$, where $\mathbf{\Gamma}\mathbf{\Lambda}\mathbf{\Gamma}^{\top}$ and $\mathbf{L}\mathbf{L}^{\top}$ denote the EVD and CD of $\mathbf{A}$, respectively

Matrix Function: Overview

To create a matrix in R, we use the `matrix` function.

The relevant inputs of the `matrix` function include

- `data`: the data that will be arranged into a matrix
- `nrow`: the number of rows of the matrix
- `ncol`: the number of columns of the matrix
- `byrow`: logical indicating if the data should be read-in by rows (default reads in data by columns)

Matrix Function: Example

```
> x <- 1:9
```

```
> x  
[1] 1 2 3 4 5 6 7 8 9
```

```
> matrix(x, nrow = 3, ncol = 3)  
      [,1] [,2] [,3]  
[1,]    1    4    7  
[2,]    2    5    8  
[3,]    3    6    9
```

```
> matrix(x, nrow = 3, ncol = 3, byrow = TRUE)  
      [,1] [,2] [,3]  
[1,]    1    2    3  
[2,]    4    5    6  
[3,]    7    8    9
```

Matrix Function: Warning

R recycles numbers if the dimensions do not conform:

```
> x <- 1:9
```

```
> x  
[1] 1 2 3 4 5 6 7 8 9
```

```
> matrix(x, nrow = 3, ncol = 4)  
      [,1] [,2] [,3] [,4]  
[1,]    1    4    7    1  
[2,]    2    5    8    2  
[3,]    3    6    9    3
```

Warning message:

```
In matrix(x, nrow = 3, ncol = 4) :
```

```
  data length [9] is not a sub-multiple or multiple of the  
number of columns [4]
```

R Matrix Calculations: Overview

Remember: scalar multiplication is performed using:

`*`

In contrast, matrix multiplication is performed using:

`%*%`

Note: the matrix multiplication symbol is really three symbols in a row:

- percent sign
- asterisk
- percent sign

R Matrix Calculations: Example

```
> x <- 1:9
> y <- 9:1
> X <- matrix(x, 3, 3)
> Y <- matrix(y, 3, 3)
> X
```

```
      [,1] [,2] [,3]
[1,]     1     4     7
[2,]     2     5     8
[3,]     3     6     9
```

```
> Y
      [,1] [,2] [,3]
[1,]     9     6     3
[2,]     8     5     2
[3,]     7     4     1
```

```
> X * Y
      [,1] [,2] [,3]
[1,]     9    24    21
[2,]    16    25    16
[3,]    21    24     9
```

```
> X %*% Y
      [,1] [,2] [,3]
[1,]    90    54    18
[2,]   114    69    24
[3,]   138    84    30
```


R Matrix Calculations: Error Messages

```
> x <- 1:6  
> y <- 6:1  
> X <- matrix(x, 2, 3)  
> Y <- matrix(y, 3, 2)  
> X
```

```
      [,1] [,2] [,3]  
[1,]     1     3     5  
[2,]     2     4     6
```

```
> Y  
      [,1] [,2]  
[1,]     6     3  
[2,]     5     2  
[3,]     4     1
```

```
> X * Y  
Error in X * Y :  
non-conformable arrays
```

```
> X %*% Y  
      [,1] [,2]  
[1,]    41    14  
[2,]    56    20
```

R Matrix Calculations: Error Messages (continued)

```
> x <- 1:6
> y <- 6:1
> X <- matrix(x, 2, 3)
> Y <- matrix(y, 2, 3)
> X
```

```
      [,1] [,2] [,3]
[1,]     1     3     5
[2,]     2     4     6
```

```
> Y
```

```
      [,1] [,2] [,3]
[1,]     6     4     2
[2,]     5     3     1
```

```
> X * Y
```

```
      [,1] [,2] [,3]
[1,]      6    12    10
[2,]     10    12     6
```

```
> X %*% Y
```

```
Error in X %*% Y :
non-conformable arguments
```

Transpose Function

To obtain the transpose of a matrix in R, we use the `t` function.

```
> X <- matrix(1:6,2,3)
```

```
> X
```

```
      [,1] [,2] [,3]
[1,]     1     3     5
[2,]     2     4     6
```

```
> t(X)
```

```
      [,1] [,2]
[1,]     1     2
[2,]     3     4
[3,]     5     6
```

Dimension Function

To obtain the dimensions of a matrix in R, we use the `dim` function.

```
> X <- matrix(1:6, 2, 3)
```

```
> X
```

```
      [,1] [,2] [,3]  
[1,]     1     3     5  
[2,]     2     4     6
```

```
> dim(X)
```

```
[1] 2 3
```

```
> dim(t(X))
```

```
[1] 3 2
```

Crossproduct Function

Given $\mathbf{X} = \{x_{ij}\}_{n \times p}$ and $\mathbf{Y} = \{y_{ik}\}_{n \times q}$, we can obtain the crossproduct $\mathbf{X}^\top \mathbf{Y}$ using the `crossprod` function.

```
> X <- matrix(1:6, 3, 2)
> Y <- matrix(1:9, 3, 3)
> crossprod(X, Y)
      [,1] [,2] [,3]
[1,]   14   32   50
[2,]   32   77  122
> t(X) %*% Y
      [,1] [,2] [,3]
[1,]   14   32   50
[2,]   32   77  122
```

Note that `crossprod` produces same result as using transpose and matrix multiplication symbol.

However, you should prefer `crossprod` because it is faster.

Transpose-Crossproduct Function

Given $\mathbf{X} = \{x_{ij}\}_{n \times p}$ and $\mathbf{Y} = \{y_{hj}\}_{m \times p}$, we can obtain the transpose-crossproduct $\mathbf{XY}^\top$ using the `tcrossprod` function.

```
> X <- matrix(1:6, 2, 3)
> Y <- matrix(1:9, 3, 3)
> tcrossprod(X, Y)
      [,1] [,2] [,3]
[1,]   48   57   66
[2,]   60   72   84
> X %*% t(Y)
      [,1] [,2] [,3]
[1,]   48   57   66
[2,]   60   72   84
```

Note that `tcrossprod` produces same result as using transpose and matrix multiplication symbol.

However, you should prefer `tcrossprod` because it is faster.

Row and Column Summation Functions

We can obtain rowwise and columnwise summations using the `rowSums` and `colSums` functions.

```
> X <- matrix(1:6, 2, 3)
> X
      [,1] [,2] [,3]
[1,]     1     3     5
[2,]     2     4     6
> rowSums(X)
[1]  9 12
> colSums(X)
[1]  3  7 11
```

Row and Column Mean Functions

We can obtain rowwise and columnwise means using the `rowMeans` and `colMeans` functions.

```
> X <- matrix(1:6, 2, 3)
> X
      [,1] [,2] [,3]
[1,]    1    3    5
[2,]    2    4    6
> rowMeans(X)
[1] 3 4
> colMeans(X)
[1] 1.5 3.5 5.5
```


Diagonal Function

The `diag` function has multiple purposes:

- If you input a square matrix, `diag` returns the diagonal elements
- If you input a vector, `diag` creates a diagonal matrix
- If you input a scalar, `diag` creates an identity matrix

```
> X <- matrix(1:4, 2, 2)
```

```
> X
```

```
      [,1] [,2]
[1,]    1    3
[2,]    2    4
```

```
> diag(X)
```

```
[1] 1 4
```

```
> diag(1:3)
```

```
      [,1] [,2] [,3]
[1,]    1    0    0
[2,]    0    2    0
[3,]    0    0    3
```

```
> diag(2)
```

```
      [,1] [,2]
[1,]    1    0
[2,]    0    1
```

Functions for Matrix Decompositions

R has built-in functions for popular matrix decompositions:

- Eigenvalue Decomposition: `eigen`
- Cholesky Decomposition: `chol`
- Singular Value Decomposition: `svd`
- QR Decomposition: `qr`

We will not directly use these functions, but some of the methods we will use call these functions internally.

Eigenvalue Decomposition

```
> X <- matrix(1:9, 3, 3)
> X <- crossprod(X)
> x eig <- eigen(X, symmetric = TRUE)
> x eig$values
[1] 2.838586e+02 1.141413e+00 3.283090e-14
> x eig$ vectors
      [,1]      [,2]      [,3]
[1,] -0.2148372  0.8872307  0.4082483
[2,] -0.5205874  0.2496440 -0.8164966
[3,] -0.8263375 -0.3879428  0.4082483
> Xhat = x eig$ vec %*% diag(x eig$val) %*% t(x eig$ vec)
> max(abs(X - Xhat))
[1] 1.421085e-13
```

Cholesky Decomposition

```
> set.seed(1)
> X <- matrix(runif(9), 3, 3)
> X <- crossprod(X)
> xchol <- chol(X)
> t(xchol)
           [,1]      [,2]      [,3]
[1,] 0.7328929 0.0000000 0.0000000
[2,] 1.1336353 0.6224886 0.0000000
[3,] 1.1694863 0.3705306 0.4688907
> Xhat <- crossprod(xchol)
> max(abs(X - Xhat))
[1] 1.110223e-16
```

Singular Value Decomposition

```
> X <- matrix(1:6, 3, 2)
> xsvd <- svd(X)
> xsvd$d
[1] 9.5080320 0.7728696
> xsvd$u
      [,1]      [,2]
[1,] -0.4286671  0.8059639
[2,] -0.5663069  0.1123824
[3,] -0.7039467 -0.5811991
> xsvd$v
      [,1]      [,2]
[1,] -0.3863177 -0.9223658
[2,] -0.9223658  0.3863177
> Xhat <- xsvd$u %*% diag(xsvd$d) %*% t(xsvd$v)
> max(abs(X - Xhat))
[1] 9.992007e-16
```

QR Decomposition

```
> X <- matrix(1:6, 3, 2)
> xqr <- qr(X)
> Q <- qr.Q(xqr)
> Q
      [,1]      [,2]
[1,] -0.2672612  0.8728716
[2,] -0.5345225  0.2182179
[3,] -0.8017837 -0.4364358
> R <- qr.R(xqr)
> R
      [,1]      [,2]
[1,] -3.741657 -8.552360
[2,]  0.000000  1.963961
> Xhat <- Q %*% R[,sort(xqr$pivot,index=TRUE)$ix]
> max(abs(X - Xhat))
[1] 8.881784e-16
```