Quiz 2 Solutions

MATH 103A August 2, 2018

(1) (Q) Determine whether the following functions are entire (holomorphic on all finite points of \mathbb{C}):

a)
$$f(x,y) = y^2 - ix^2$$

b)
$$f(x,y) = \sin(x)\cosh(y) + i\cos(x)\sinh(y)$$

(A)

(a) For the Cauchy-Riemann equations we identify $u(x,y) = y^2$ and $v(x,y) = -x^2$ to see that:

$$u_x = 0 = 0 = v_y$$
$$u_y = 2y \neq 2x = -v_x$$

Since the equations are not satisfied for all of \mathbb{C} it follows that f(x,y) is not an entire function.

(b) For the Cauchy-Riemann equations we identify $u(x,y) = \sin(x)\cosh(y)$ and $v(x,y) = \cos(x)\sinh(y)$ to see that:

$$u_x = \cos(x)\cosh(y) = v_y$$

 $u_y = \sin(x)\sinh(y) = -v_x$

Since the equations are satisfied for all of $\mathbb C$ with continuous partial derivatives it follows that f(x,y) is an entire function.

(2) (Q) Does the function:

$$f(z) = \begin{cases} \frac{\overline{z}^2}{z} & z \neq 0\\ 0 & z = 0 \end{cases}$$

satisfy the Cauchy-Riemann equations at z = 0? Is f'(0) well-defined?

(A) It helps to rewrite:

$$\frac{\overline{z}^2}{z} = \frac{\overline{z}^3}{\|z\|^2} = \frac{(x - iy)^3}{x^2 + y^2} = \frac{x^3 - 3xy^2}{x^2 + y^2} + \frac{y^3 - 3x^2y}{x^2 + y^2}i = u(x, y) + iv(x, y)$$

For the Cauchy-Riemann equations we need the following calculations:

$$u_x(0) = \lim_{h \to 0} \frac{u(0+h,0) - u(0,0)}{h} = \lim_{h \to 0} \frac{\frac{h^3}{h^2}}{h} = 1$$

$$u_y(0) = \lim_{h \to 0} \frac{u(0,0+h) - u(0,0)}{h} = \lim_{h \to 0} \frac{0}{h} = 0$$

$$v_x(0) = \lim_{h \to 0} \frac{v(0+h,0) - v(0,0)}{h} = \lim_{h \to 0} \frac{0}{h} = 0$$

$$v_y(0) = \lim_{h \to 0} \frac{v(0,0+h) - v(0,0)}{h} = \lim_{h \to 0} \frac{\frac{h^3}{h^2}}{h} = 1$$

It follows that the Cauchy-Riemann equations are satisfied at z=0. Now for the derivative behavior, we calculate directly:

$$f'(0) = \lim_{h \to 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \to 0} \frac{\frac{\overline{h}^2}{h}}{h} = \left(\lim_{h \to 0} \frac{h}{\overline{h}}\right)^{-2} = \text{DNE}$$