

Answer Key to Group Exercises

MATH 11A - Discussion Sections C & F

- (1) For each function transformation graph the result when acted on $f(x)$ and write down its corresponding formula given $f(x) = e^x$:
- (a) Shift down by 3 $[g(x) = f(x) - 3 = e^x - 3]$
 - (b) Shift right by 7 $[g(x) = f(x - 7) = e^{x-7}]$
 - (c) Reflect about the x -axis $[g(x) = -f(x) = -e^x]$
 - (d) Reflect about the y -axis $[g(x) = f(-x) = e^{-x}]$
 - (e) Reflect about the origin $[g(x) = -f(-x) = -e^{-x}]$
- (2) Find the domain of the following functions:
- (a) $f(x) = \frac{7-x}{e^x-1}$ $[x \in (-\infty, 0) \cup (0, \infty)]$
 - (b) $g(x) = \frac{e^x}{x^2-x-1}$ $\left[x \in \left(-\infty, \frac{1-\sqrt{5}}{2}\right) \cup \left(\frac{1-\sqrt{5}}{2}, \frac{1+\sqrt{5}}{2}\right) \cup \left(\frac{1+\sqrt{5}}{2}, \infty\right)\right]$
 - (c) $h(x) = \frac{x^\pi - x + 1}{x^3 + x^2 - x}$ $\left[x \in \left(-\infty, \frac{-1-\sqrt{5}}{2}\right) \cup \left(\frac{-1-\sqrt{5}}{2}, 0\right) \cup \left(0, \frac{-1+\sqrt{5}}{2}\right) \cup \left(\frac{-1+\sqrt{5}}{2}, \infty\right)\right]$
 - (d) $j(x) = \frac{1}{e^{x^2+2x-1}}$ $[x \in \mathbb{R}]$
 - (e) $m(x) = \frac{x+\pi}{e^{\sin(x)}}$ $[x \in \mathbb{R}]$
- (3) The half-life of Uranium-238 is 4.51×10^9 years. Given that we start with a sample of 500mg, answer the following:
- (a) Write down a formula that models how much of the sample is left after t years. $\left[P = 500\left(\frac{1}{2}\right)^{\frac{t}{4.51 \times 10^9}}\right]$
 - (b) How much of the sample remains after $t = 10^{20}$ years? $\left[P(10^{20}) = 500\left(\frac{1}{2}\right)^{\frac{10^{20}}{4.51}}\right]$
 - (c) At what value of t does the sample have 200mg left? $\left[t = (4.51 \times 10^9) \log_{\frac{1}{2}}\left(\frac{2}{5}\right)\right]$
- (4) The half-life of Polonium-210 is 138.376 days. Given that we start with a sample of 500mg, answer the following:
- (a) Write down a formula that models how much of the sample is left after t days. $\left[P = 500\left(\frac{1}{2}\right)^{\frac{t}{138.376}}\right]$
 - (b) How much of the sample remains after 2 years? $\left[P(730) = 500\left(\frac{1}{2}\right)^{\frac{730}{138.376}}\right]$
 - (c) At what value of t does the sample have 200mg left? $\left[t = 138.376 \log_{\frac{1}{2}}\left(\frac{2}{5}\right)\right]$
- (5) Determine whether the following functions are one-to-one (injective) on the provided domains:
- (a) $f(x) = \cos(x)$ where $x \in \mathbb{R}$ [No]
 - (b) $g(x) = \cos(x)$ where $x \in [0, \pi]$ [Yes]
 - (c) $h(x) = \frac{x^2-x+7}{x-1}$ where $x \in (-\infty, 1) \cup (1, \infty)$ [No]
 - (d) $j(x) = \ln(x)$ where $x \in (0, \infty)$ [Yes]
 - (e) $m(x) = \ln|x|$ where $x \in \mathbb{R}$ [No]
 - (f) $n(x) = \frac{x}{|x|}$ where $x \in (-\infty, 0) \cup (0, \infty)$ [No]
- (6) Determine a formula for the inverse, if it exists:
- (a) $f(x) = \frac{2x-1}{5x+3}$ where $x \in \left(-\infty, -\frac{3}{5}\right) \cup \left(-\frac{3}{5}, \infty\right)$ $\left[f^{-1}(x) = -\frac{3x+1}{5x-2}\right]$
 - (b) $g(x) = e^{x-3}$ where $x \in \mathbb{R}$ $[g^{-1}(x) = \ln(x) + 3]$
 - (c) $h(x) = e^{(7-x)^3+10}$ where $x \in \mathbb{R}$ $\left[h^{-1}(x) = 7 - \sqrt[3]{\ln(x) - 10}\right]$
 - (d) $j(x) = \ln(x^3 + 4)$ where $x \in \mathbb{R}$ [DNE]
 - (e) $m(x) = \ln(x^2)$ where $x \in \mathbb{R}$ [DNE]
 - (f) $n(x) = x^2$ where $x \in \mathbb{R}$ [DNE]
 - (g) $p(x) = x^2$ where $x \in [0, \infty)$ $[p^{-1}(x) = \sqrt{x}]$

(7) Graph the following piecewise functions and determine if they are continuous:

$$\text{a) } f(x) = \begin{cases} |x|, & x < 0 \\ x^2, & x \geq 0 \end{cases} \quad [\text{Yes}] \quad \text{b) } g(x) = \begin{cases} x+1, & x < 0 \\ x+10, & x \geq 0 \end{cases} \quad [\text{No}] \quad \text{c) } h(x) = \begin{cases} x+1, & x < 0 \\ 0, & x = 0 \\ 1-x, & x > 0 \end{cases} \quad [\text{No}]$$

(8) Simplify the following expressions in terms of a single logarithm:

$$\begin{aligned} \text{(a) } & 7 \log_{12}(x) + 21 \log_{12}(y) \quad \left[\log_{12}(x^7 y^{21}) \right] \\ \text{(b) } & 3 \log(x) - 6 \log(y) \quad \left[\log(x^3 y^{-6}) \right] \\ \text{(c) } & 5 \ln(x+y) - 21 \ln(x) - 8 \ln(y) \quad \left[\ln((x+y)^5 x^{-21} y^{-8}) \right] \\ \text{(d) } & \log_7(x) + \ln(y) \quad \left[\ln(x^{\frac{1}{\ln(7)}} y) \right] \\ \text{(e) } & \ln(y^2) - \ln(x) + 2 \log(y) \quad \left[\ln\left(y^{2+\frac{2}{\ln(10)}} x^{-1}\right) \right] \end{aligned}$$

(9) Determine whether the following are true or false:

- (a) If $x \in \mathbb{R}$, then $\sqrt{x^2} = x$. [False]
- (b) If f and g are two continuous functions, then $f \circ g = g \circ f$. [False]
- (c) For any given function f we have $f(x+y) = f(x) + f(y)$. [False]
- (d) Given that f is a linear function, then we must have $f(\alpha x + \beta y) = \alpha f(x) + \beta f(y)$. [False]
- (e) Given that f is a linear function that passes through the origin, then we must have $f(\alpha x + \beta y) = \alpha f(x) + \beta f(y)$. [True]
- (f) If $x < y$ and f is a strictly decreasing function, then $f(x) > f(y)$. [True]
- (g) If $\alpha, x \in \mathbb{R}$, then $\ln(x^\alpha) = \alpha \ln(x)$. [False]

(10) Determine a formula for the general term a_n given the following sequences:

- (a) $\{1, -1, 1, -1, 1, -1, \dots\}$ $\left[a_n = (-1)^n \text{ where } n \geq 0 \right]$
- (b) $\{2, 3, 2, 3, 2, 3, \dots\}$ $\left[a_n = \frac{5}{2} + \frac{(-1)^n}{2} \text{ where } n \geq 1 \right]$
- (c) $\{1, \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \frac{1}{9}, \dots\}$ $\left[a_n = \frac{1}{2n+1} \text{ where } n \geq 0 \right]$
- (d) $\{1, -\frac{1}{3}, \frac{1}{9}, -\frac{1}{27}, \frac{1}{81}, \dots\}$ $\left[a_n = \left(-\frac{1}{3}\right)^n \text{ where } n \geq 0 \right]$
- (e) $\{1, 1.9, 1.99, 1.999, 1.9999, 1.99999, \dots\}$ $\left[a_n = 2 - \frac{1}{10^n} \text{ where } n \geq 0 \right]$
- (f) $\{1, r, r^2, r^3, r^4, \dots\}$ $\left[a_n = r^n \text{ where } n \geq 0 \right]$
- (g) $\{\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \dots\}$ $\left[a_n = 2^{\frac{2^n - 1}{2^n}} \text{ where } n \geq 0 \right]$

(11) Determine a formula for each situation:

- (a) Linear function that passes through $(-2, 4)$ and $(2, 0)$. $[f(x) = -x + 2]$
- (b) Exponential function that passes through $(-1, 10)$ and $(2, 30)$. $\left[f(x) = 30(3)^{\frac{x-2}{3}} \right]$
- (c) Linear function that has slope $m = 5$ and passes through $(0, 10)$. $[f(x) = 5x + 10]$
- (d) Quadratic function that has roots at $x = -3$ and $x = 3$ while passing through $(10, 10)$. $\left[f(x) = \frac{10}{91}(x-3)(x+3) \right]$
- (e) Exponential function that has a horizontal asymptote of $y = -1$ and passes through $(1, 1)$ and $(5, 10)$. $\left[f(x) = 2\left(\frac{11}{2}\right)^{\frac{x-1}{4}} - 1 \right]$

(12) Find the exact value of each expression without the use of a calculator:

(a) $\log_3 \left(\frac{1}{27} \right)$ $[-3]$

(b) $\ln \left(\frac{1}{e^k} \right)$ where $k \in \mathbb{R}$ $[-k]$

(c) $e^{-2 \ln(5)}$ $[5^{-2}]$

(d) $\ln \left(\ln \left(e^{e^{10}} \right) \right)$ $[10]$

(e) $2 \log_2(6) - \log_2(15) + \log_2(20)$ $[\log_2(48)]$

(13) Solve the following for x exactly:

(a) $e^{7-4x} = 6$ $\left[x = \frac{7-\ln(6)}{4} \right]$

(b) $\ln(3x - 10) = 2$ $\left[x = \frac{e^2+10}{3} \right]$

(c) $\ln(x^2 - 1) = 3$ $\left[x = \pm \sqrt{e^3 + 1} \right]$

(d) $e^{2x} - 3e^x + 2 = 0$ $[x = 0, \ln(2)]$

(e) $\ln(\ln(x)) = 1$ $[x = e^e]$

(14) Solve the following inequalities for x :

(a) $\ln(x^2 - 1) > e$ $[|x| > \sqrt{e^e + 1}]$

(b) $e^{(x-1)^2} > 5$ $\left[x < 1 - \sqrt{\ln(5)} \text{ or } x > 1 + \sqrt{\ln(5)} \right]$

(c) $1 - 5 \ln(x) < 7$ $\left[x > e^{-\frac{6}{5}} \right]$

(d) $1 < e^{3x-1} < 2$ $\left[\frac{1}{3} < x < \frac{1+\ln(x)}{3} \right]$

(e) $\ln|x - 3| \geq 5$ $[x \geq 3 + e^5 \text{ or } x \leq 3 - e^5]$

(15) Determine the inverse function of the following, if it exists:

(a) $f(x) = 1 + \sqrt{2 + 3x}$ where $x \in \left[-\frac{2}{3}, \infty \right)$ $\left[f^{-1}(x) = \frac{x^2 - 2x - 1}{3} \right]$

(b) $g(x) = \frac{4x-1}{2x+3}$ where $x \in \left(-\infty, -\frac{3}{2} \right) \cup \left(-\frac{3}{2}, \infty \right)$ $\left[g^{-1}(x) = -\frac{3x+1}{2x-4} \right]$

(c) $h(x) = e^{2x-1}$ where $x \in \mathbb{R}$ $\left[h^{-1}(x) = \frac{\ln(x)+1}{2} \right]$

(d) $j(x) = \frac{e^x}{1+2e^x}$ where $x \in \mathbb{R}$ $\left[j^{-1}(x) = \ln \left(\frac{x}{1-2x} \right) \right]$

(e) $m(x) = \begin{cases} \sqrt{|x|}, & x \leq 0 \\ -x, & x > 0 \end{cases}$ $\left[m^{-1}(x) = \begin{cases} -x^2, & x > 0 \\ -x, & x \leq 0 \end{cases} \right]$

(16) Find the domain and range of the following functions:

(a) $f(x) = \frac{2}{3x-1}$ $\left[\text{D:} \left(-\infty, \frac{1}{3} \right) \cup \left(\frac{1}{3}, \infty \right) \text{ and R:} (-\infty, 0) \cup (0, \infty) \right]$

(b) $g(x) = \sqrt{16 - x^4}$ $\left[\text{D:} [-2, 2] \text{ and R:} [0, 4] \right]$

(c) $h(x) = \ln(x^2 - 1)$ $\left[\text{D:} (-\infty, -1) \cup (1, \infty) \text{ and R:} \mathbb{R} \right]$

(d) $j(x) = 3 + \cos(2x)$ $\left[\text{D:} \mathbb{R} \text{ and R:} [2, 4] \right]$

(e) $m(x) = \left| \frac{x+3}{x^2-1} \right|$ $\left[\text{D:} (-\infty, -1) \cup (-1, 1) \cup (1, \infty) \text{ and R:} \mathbb{R}_{\geq 0} \right]$

- (17) Determine whether the following sequences converge or diverge. If they converge determine the exact value they converge to.
- (a) $a_n = \frac{1}{3n^4}$ [Convergent since $a_n \rightarrow 0$]
 - (b) $a_n = \frac{n^3-1}{n^3+1}$ [Convergent since $a_n \rightarrow 1$]
 - (c) $a_n = \frac{10^n}{1+9^n}$ [Divergent]
 - (d) $a_n = \ln(2n^2 + 1) - \ln(n^2 + 1)$ [Convergent since $a_n \rightarrow \ln(2)$]
 - (e) $a_n = \frac{\sqrt[3]{n}}{\sqrt[3]{n} + \sqrt{n}}$ [Convergent since $a_n \rightarrow 0$]
- (18) For the following assume that $a_n \rightarrow \mathcal{L}$ and determine the value of \mathcal{L} exactly:
- (a) $a_{n+1} = \frac{1}{2}a_n + 1$ where $a_1 = 1$ [$\mathcal{L} = 2$]
 - (b) $a_{n+1} = 2a_n - 1$ where $a_1 = 2$ [$\mathcal{L} = 1$]
 - (c) $a_{n+1} = \sqrt{5a_n}$ where $a_1 = 1$ [$\mathcal{L} = 5$]
 - (d) $a_{n+1} = \frac{6}{1+a_n}$ where $a_1 = 1$ [$\mathcal{L} = 2$]
 - (e) $a_{n+1} = \frac{1}{2}\left(a_n + \frac{25}{a_n}\right)$ where $a_1 = 100$ [$\mathcal{L} = 5$]
- (19) Calculate the following:
- (a) $\lim_{n \rightarrow \infty} \frac{\sqrt{n+n^2}}{2n-n^2}$ [0]
 - (b) $\lim_{x \rightarrow \infty} \frac{(2x^2+1)^2}{(x-1)^2(x^2+x)}$ [4]
 - (c) $\lim_{t \rightarrow \infty} (\sqrt{9t^2 + t} - 3t)$ $\left[\frac{1}{6}\right]$
 - (d) $\lim_{x \rightarrow \infty} [\ln(x^2) - \ln(x^2 + 1)]$ [0]
 - (e) $\lim_{x \rightarrow \infty} \frac{e^{3x} - e^{-3x}}{e^{3x} + e^{-3x}}$ [1]
- (20) Evaluate the following limits, if they exist:
- (a) $\lim_{x \rightarrow 5} \frac{x^2-6x+5}{x-5}$ [4]
 - (b) $\lim_{h \rightarrow 0} \frac{(4+h)^2-16}{h}$ [8]
 - (c) $\lim_{x \rightarrow 0} \left(\frac{1}{x} - \frac{1}{x^2+x}\right)$ [1]
 - (d) $\lim_{x \rightarrow -2} \frac{x+2}{x^3+8}$ $\left[\frac{1}{12}\right]$
 - (e) $\lim_{x \rightarrow 0} \frac{\sqrt{1+x}-1}{x}$ $\left[\frac{1}{2}\right]$
- (21) In the context of *Classical Mechanics* as first described by Sir Isaac Newton, *momentum* is defined as $p = mv$ where m and v are the mass and velocity respectively of the object of interest. In the context of *Special Relativity*, momentum is defined as:

$$p^* = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}}$$

where c is the speed of light. What happens as $v \rightarrow c^-$? [$p \rightarrow mc$ but $p^* \rightarrow \infty$]

(22) Evaluate the following limits:

- (a) $\lim_{n \rightarrow \infty} (2^{-n} + 3^{-n})$ [0]
- (b) $\lim_{n \rightarrow \infty} \frac{6n+5}{n-7}$ [6]
- (c) $\lim_{n \rightarrow \infty} \frac{\sqrt{n}+n^2}{100n-3n^2}$ $\left[-\frac{1}{3}\right]$
- (d) $\lim_{n \rightarrow \infty} \frac{e^{3n}-e^{-3n}}{10e^{3n}+e^{-2n}}$ $\left[\frac{1}{10}\right]$
- (e) $\lim_{n \rightarrow \infty} \frac{n^4-8n^2+n}{n^3-n+90}$ $[\infty]$

(23) Determine whether the following approach negative or positive infinity:

- (a) $\lim_{n \rightarrow 3^+} \frac{e^n}{(n-3)^5}$ [+]
- (b) $\lim_{n \rightarrow -4^-} \frac{n+3}{n+4}$ [+]
- (c) $\lim_{n \rightarrow 5^+} \frac{n^2-5n}{n^2-10n+25}$ [+]
- (d) $\lim_{n \rightarrow 3^+} \ln(n^2 - 9)$ [-]
- (e) $\lim_{x \rightarrow (2\pi)^-} x \csc(x)$ [-]

(24) Evaluate the following limits:

- (a) $\lim_{h \rightarrow 0} \frac{(4+h)^2-16}{h}$ [8]
- (b) $\lim_{h \rightarrow 0} \frac{(2+h)^3-8}{h}$ [12]
- (c) $\lim_{h \rightarrow 0} \frac{\sqrt{1+h}-1}{h}$ $\left[\frac{1}{2}\right]$
- (d) $\lim_{h \rightarrow 0} \left(\frac{1}{h} - \frac{1}{h^2+h}\right)$ [1]
- (e) $\lim_{x \rightarrow 16} \frac{4-\sqrt{x}}{16x-x^2}$ $\left[\frac{1}{128}\right]$

(25) Evaluate the following limits:

- (a) $\lim_{x \rightarrow 3} (2x - |x-3|)$ [6]
- (b) $\lim_{x \rightarrow -6} \frac{2x+12}{|x+6|}$ [DNE]
- (c) $\lim_{x \rightarrow 0^-} \left(\frac{1}{x} - \frac{1}{|x|}\right)$ $[-\infty]$
- (d) $\lim_{x \rightarrow -2} \frac{2-|x|}{2+x}$ [1]
- (e) $\lim_{x \rightarrow 0} \frac{|2x-1|-|2x+1|}{x}$ [-4]

(26) Is it true that $\frac{x^2+x-6}{x-2} = x+3$ for all $x \in \mathbb{R}$? If not, explain why. Next determine whether $\lim_{x \rightarrow 2} \frac{x^2+x-6}{x-2} = \lim_{x \rightarrow 2} (x+3)$ is true or not. [False and True respectively]

(27) Assuming $a_i, b_j \in \mathbb{R}$ for $i \leq n$ and $j \leq m$, prove the following:

$$\lim_{x \rightarrow \infty} \frac{a_0 + a_1x + a_2x^2 + \cdots + a_nx^n}{b_0 + b_1x + b_2x^2 + \cdots + b_mx^m} = \begin{cases} 0 & n < m \\ \frac{a_n}{b_n} & n = m \\ \infty & n > m \end{cases}$$

[For each case divide by the highest power of x and compute the limit accordingly]

(28) Evaluate the following using an analytical approach:

$$\lim_{n \rightarrow \infty} n^{\frac{1}{n}} = 1$$

(29) Find numbers $a, b \in \mathbb{R}$ s.t.:

$$\lim_{x \rightarrow 0} \frac{\sqrt{ax+b}-2}{x} = 1 \quad [a = b = 4]$$

(30) Evaluate the following using an analytical approach:

$$\lim_{x \rightarrow \infty} \frac{\sin(x)}{x} = 0$$

(31) The gravitational force exerted by the planet Earth on a unit mass at a distance r from the center of the planet is given by:

$$F(r) = \begin{cases} \frac{GM}{r^2} & r > R \\ \frac{GM}{R^3} r & r \leq R \end{cases}$$

where M is the mass of Earth, R is its radius, and G is the gravitational constant. Is F a continuous function of r ?
[Yes]