

# Sensitivity in relative binding free energies from incremental ligand changes within a simple model binding site using modern free energy calculation protocols and force fields

Nathan M. Lim,<sup>\*,†</sup> Lingle Wang,<sup>‡</sup> Robert Abel,<sup>‡</sup> and David L. Mobley<sup>†</sup>

*Department of Pharmaceutical Sciences, University of California—Irvine, Irvine, California 92697, United States, and Schrödinger, Inc., 120 West 45th Street, New York, New York 10036, United States*

E-mail: limn1@uci.edu

## Abstract

Tremendous recent improvements in sampling techniques for molecular dynamics (MD), classical force fields, and computer hardware optimization, are now allowing protein ligand binding free energy calculations to be routinely used to advance drug discovery project. However, despite these recent innovations, clear need remains to further improve sampling algorithms to more completely and reliably sample protein motion relevant to protein-ligand binding. Evidence of such clear and remaining need has recently been found in the well studied model binding sites like the apolar cavity of T4 Lysozyme L99A. In this study, we model recent experimental results that show

---

<sup>\*</sup>To whom correspondence should be addressed

<sup>†</sup>University of California—Irvine

<sup>‡</sup>Schrödinger, Inc.

the progressive opening of the binding pocket in response to a series of homologous ligands.<sup>1</sup> Even while using enhanced sampling techniques, we demonstrate that the predicted relative binding free energies (RBFE) are still highly sensitive to the initial protein conformational state. Particularly, we highlight the importance of sufficient sampling of protein conformational changes and possible techniques for addressing the issue.

# 1 Introduction

Proteins play a central role in many biological processes by modulating many key signaling pathways. It is unsurprising to find that proteins make up the vast majority of pharmaceutical drug targets. Many small-molecules on the market today induce their therapeutic effect by modulating the proteins' biological activity through binding.<sup>2-4</sup> Thus, optimization of protein-ligand binding affinities is a central goal in early pharmaceutical drug design projects.

Recent advancements in technology and computational chemistry have led to increasing use of computer-aided drug design techniques to assist in lead optimization in pharmaceutical drug discovery. Since accuracy and reliability of the approach being used is critical for success, it is here where the most rigorous of methods like free energy calculations can be applied in the prediction of protein-ligand binding affinities. Using methods like alchemical free energy perturbation (FEP), thermodynamic integration, and  $\lambda$ -dynamics on molecular dynamics simulations, the difference in binding free energies between two ligands can be computed in a robust and accurate manner. By computing relative binding free energies, much of the computational cost and difficulties of absolute binding free energy calculations is avoided.<sup>5-11</sup> Other advancements in forcefields, sampling algorithms, and emergence of GPU computing have considerably improved the accuracy and robustness of FEP calculations. The recent development of FEP+, a fully automated FEP protocol with implemented modern methodologies, reduces overall workload and the potential for human error in setup and analysis of these types of calculations. Using this modern FEP protocol, previous studies<sup>12</sup> have demonstrated it to yield highly accurate free energy predictions in a wide range of pharmaceutically relevant protein targets and ligands. The accuracy and reliability of FEP+ make it a very powerful tool in the hands of medicinal chemists for efficient optimization of lead compounds.

Although the previous report on FEP+<sup>12</sup> demonstrates its robustness—like much of the FEP literature—the primary concern focuses on the size of the ligand perturbation or the

initial ligand pose and how these factors impact the accuracy of calculated free energies.<sup>13,14</sup> Consideration of effects arising from protein conformational changes have generally received less attention due to the difficulty in addressing the sampling challenges encountered. Some computational studies have shown even small simple changes in the protein—like a side-chain rotamer flipping—can cause large errors in affinity predictions if not sampled<sup>15,16</sup> and thus protein conformational changes should not be ignored. In the previous FEP+ study,<sup>12</sup> many of the protein targets used were either extremely rigid (i.e. tyk2 and thrombin) or ligand pairs appear to be selected so issues concerning protein conformational changes did not have to be addressed. Potentially, this could mean the domain of applicability for the FEP+ protocol is restricted to either extremely rigid proteins or ligand pairs that do not induce a significant protein conformational change. Considering it is extremely common to find that ligand binding causes the protein to adopt a discrete conformation, this limits the potential impact FEP calculations can have in industrial drug discovery projects despite being shown to be highly accurate. Furthermore, protein-ligand binding can be more complex than this, where several studies have found a protein can adopt multiple conformations even for the same ligand.<sup>1,17,18</sup> A thorough investigation on how protein conformational changes and the initial protein configuration affects the accuracy of free energy predictions has yet to be performed, to which we turn to now.

Here, we apply the now-standard FEP protocol used in ,<sup>12</sup> which utilizes replica exchange with solute tempering (FEP/REST),<sup>19</sup> to a very simple model binding site in an engineered mutant of T4 lysozyme (L99A). In this mutant, the L99A mutation creates a small apolar binding site that has been studied extensively experimentally<sup>20–23</sup> and computationally by docking<sup>1,24–26</sup> and free energy methods.<sup>16,19,27–32</sup> Recent studies on this binding site have found that the protein adopts three discrete conformations in response to ligand binding.<sup>1</sup> Through a series of eight congeneric ligands(Fig 1f, each growing by addition of a single methyl, the protein responds by a single helix rearrangement to accommodate the growing ligand (Fig 1). As the ligand grows, the binding cavity was observed to incrementally

reorganize into three discrete conformations which we will refer to as the closed(Fig 1c), intermediate(Fig 1d), and open states(Fig 1e). Using the FEP/REST protocol and the aforementioned homologous ligand series, we calculate relative protein-ligand binding affinities between ligands that occupy different discrete protein conformational states. In this study—while using the implemented default FEP/REST protocol—we demonstrate how the kinetically distinct protein states and structural rearrangement affects the accuracy and reliability of our predicted relative binding affinities. Further, we illustrate the importance of sufficient sampling of protein conformational changes and how modification of the REST region can potentially address this issue.

## 2 Methods

### Protein/Ligand preparation

All proteins were prepared and aligned in Maestro<sup>33</sup> using the ‘Protein Preparation Wizard’<sup>34–38</sup> tool and with the following settings enabled (as they appear in the Maestro GUI menu):

- Preprocess: Assign bond orders, Add hydrogens, Create zero-order bonds to metals, Create disulfide bonds, Cap termini, Delete waters beyond 5Åfrom het groups
- Refine: Sample water orientations, Use PROPKA pH: 7.0, Remove waters with less than 3 H-bonds to non-waters, and restrained minimization.

Protein structures were taken from PDBs: 4W52, 4W53, 4W54, 4W55, 4W56, 4W57, 4W58, and 4W59 corresponding to protein-ligand bound structures of benzene, toluene, ethylbenzene, n-propylbenzene, sec-butylbenzene, n-butylbenzene, n-pentylbenzene, and n-hexylbenzene, respectively.<sup>1</sup> Each simulation starts from either the protein closed state (PDB:4W52) or the open state (PDF:4W59). Using LigandFEP, our system preparation follows a similar workflow to the tutorial.<sup>39</sup> Generally, two options were taken:

(1a) If the simulation starts from the protein closed state, the benzene crystal position was used as a reference for fragment building (PDB:4W52).

(1b) The corresponding ligand in the transformation was built by duplicating benzene in place and adding methyl groups.

(2a) If the simulation starts from the protein open state, the n-hexylbenzene crystal position was used as reference for fragment building (PDF:4W59).

(2b) The corresponding ligand in the transformation was built by duplicating n-hexylbenzene in place and deleting methyl groups.

Ligand tail fragments were added using the Build/Fragments toolbar in Maestro and were not overlaid or docked. As the ligand tails were built, bonds were manually rotated so that the tail was oriented in a similar manner as in their corresponding crystal structure. Following, the newly added atoms in the tail were locally minimized while leaving the core in its initial position. This was done in an attempt to correct bond angles and minimize the core RMSD, which LigandFEP uses to determine the core atoms between the two ligands.

## Classification of alchemical transformations and color coding

Here, we classified ligands based on the primary protein conformation (closed, intermediate, or open) the ligand occupies from the experimental studies (Table 1). To be explicit, the set of closed ligands refers to benzene, toluene, ethylbenzene, and n-propylbenzene; n-/sec-butylbenzene for intermediate; and n-pentyl/n-hexylbenzene for open ligands. The various protein states and ligands are then assigned a color accordingly: purple for closed, cyan for intermediate, and green for the open state.

Our sets of alchemical transformations consisted of 3 groups: ‘closed-intermediate’, ‘closed-open’, and an ‘experimental’ ligand set. The first two sets were classified based on the expected conformational change that would result from the alchemical transformation. For example, the alchemical transformation of benzene to n-butylbenzene falls into closed-intermediate while benzene to n-hexylbenzene is a closed-open transformation. In our experimental set,

we perform all possible combinations of transformations for ligands with available experimental binding affinities (Table 1). Ligands with available experimental binding affinities consisted of benzene, toluene, ethylbenzene, n-propylbenzene, and n-butylbenzene. This gives a total of 26 alchemical transformations in this study, 8 from ‘closed-intermediate’, 8 from ‘closed-open’, and 10 from the experimental set.

## FEP protocols

Using the Schrödinger application suite (release 2015-3),<sup>40</sup> we utilize a similar protocol to FEP+<sup>12</sup> called LigandFEP.<sup>39</sup> FEP+ is a fully automated work flow that plans perturbation pathways based off the LOMAP<sup>41</sup> mapping algorithm which uses the maximum common substructure (MCS) between any pair of compounds. LigandFEP is an academic toolkit that generates the configuration files to perform the free energy calculation but is limited in the sense that the user must plan each perturbation path instead. Both LigandFEP and FEP+ use the same Desmond relaxation protocol and the FEP/REST methodology.<sup>19,32,42,43</sup> By utilizing LigandFEP, we demonstrate LigandFEP can be a powerful tool for academics as it strictly differs only in the level of automation.

## Simulation details

Desmond<sup>44–47</sup> simulation protocols have been described previously in the supporting information<sup>12</sup> or can be found in greater detail in the Desmond User Manual.<sup>48</sup> The relaxation protocol begins with a simulation where solute molecules are restrained to their initial positions while minimizing using a Brownian dynamics NVT integrator for 100ps, followed by 12ps simulations at 10K with a NVT ensemble and then a NPT ensemble using the Langevin method.<sup>49</sup> Next is a 24ps simulation followed by a final 240ps simulation with solute molecules unrestrained, both are carried at room temperature with a NPT ensemble using Langevin. Production simulations with the default REST region were ran for the default setting of 5ns. Initial ‘pREST’ production simulations were also simulated for 5ns and then were carried

out to a length of up to 55ns for all closed-open transformations. For closed-intermediate transformations, we extended ‘pREST’ production simulations up to 25ns, only for cases that were far from convergence with the 5ns simulation time. Here, we use the final 15ns for closed-open simulations and the final 10ns for closed-intermediate to calculate our final free energies, discarding the initial time as additional equilibration time. FEP/REST simulations were run on four GeForce GTX Titan Black GPUs using the Desmond/GPU engine with the recently developed OPLS3<sup>50</sup> forcefield parameters.

## REST region selection

In this study, by default, only heavy atoms in the ligand were included in the REST region unless specified otherwise. Further details on the temperature profile and how the REST region is normally selected can be found in previous studies<sup>12,19</sup> in the supporting information. Simulations that included protein heavy atoms in the REST region are referred to with the ‘pREST’ label, where selection of the particular residues is described as follows.

Based on visual inspection of our molecular dynamics simulations and considering the F-helix spans residues 107-115, we selected residues Glu108, Val111, and Gly113 to include into the REST region (Fig 2a). Glu108 sits near the start of the helix which appears as a hinge point for the opening and closing of the binding cavity (Fig 2b). Following, Val111 appears in the middle of the helix and was observed to undergo the largest motion during protein conformational changes (Fig 2c). Gly113 was included in order to collectively have hot regions approximately at the start, middle and end points of the helix.

## Calculation of free energies and measurement of inconsistency

Throughout this study, we measure the inconsistency ( $\Delta\Delta G_{\varepsilon_n}$ ) between the final calculated free energies between simulations that start from the protein closed state ( $\Delta\Delta G_{C_n}$ ) versus

the protein open state ( $\Delta\Delta G_{\text{O}_n}$ ) by simply taking the difference.

$$\Delta\Delta G_{\varepsilon_n} = |\Delta\Delta G_{\text{C}_n} - \Delta\Delta G_{\text{O}_n}| \quad (1)$$

Then we compute the overall inconsistency—referred to as the 'Root-Mean-Square-Inconsistency' (RMSI)—for each set of alchemical transformations. The RMSI is calculated by using the differences ( $\Delta\Delta G_{\varepsilon_n}$ ) obtained from the comparisons between protein open and closed simulations.

$$\text{RMSI} = \sqrt{\frac{\sum_n (\Delta\Delta G_{\varepsilon_n})^2}{n}} \quad (2)$$

Similarly, we compute the 'Root-Mean-Square-Error' (RMSE) when comparing with experimental free energies for both simulations starting from the protein closed and open state.

$$\text{RMSE}^{\text{O}} = \sqrt{\frac{\sum_n (\Delta\Delta G_{\text{O}_n} - \Delta\Delta G_{\text{exp}_n})^2}{n}} \quad \text{RMSE}^{\text{C}} = \sqrt{\frac{\sum_n (\Delta\Delta G_{\text{C}_n} - \Delta\Delta G_{\text{exp}_n})^2}{n}} \quad (3)$$

Calculated free energies were determined using the Bennett acceptance ratio<sup>51</sup> (BAR) with error estimations using both bootstrapping and BAR analytical error prediction.<sup>52</sup> Hysteresis around closed thermodynamic cycles and best estimates of the free energies with their errors were calculated using the cycle closure algorithm discussed in a previous publication.<sup>19</sup>

## Determining the protein conformation state using RMSD

In this study, we determine the state of the protein by computing the 'Root-Mean-Square-Deviation' (RMSD) of the protein backbone atoms spanning the F-helix relative to their positions found in the closed (PDB:4W53), intermediate (PDB:4W57), and open (PDB:4W59) crystal structures (Fig 1b). The set of RMSDs—that is, the RMSD relative to the closed, intermediate, and open states—is computed at each frame over the course of the entire simulation. Then, we use the protein conformational state with the lowest RMSD to correspondingly color each time point in our analyses of 'RMSD/time' and 'Color maps'. Again,

we use purple to denote the protein closed state, cyan for the intermediate state, and green for the open state. Here, we use VMD<sup>53,54</sup> to align and compute the RMSD of our Desmond trajectories relative to crystal structures. Further details on the procedure and the scripts used for these analyses are provided in the supplementary info.

For the ‘RMSD/time’ analysis, see Figure 3b) for reference. Here, we plot the RMSD to the closed helix, represented by the black line, where each time point is colored according to the lowest RMSD state. We apply the RMSD/time analysis only to the simulation corresponding to the end state ligand of interest ( $\lambda_{11}$ ). By tracking the RMSD relative to the closed helix, we can monitor if the protein opens (by high RMSD with green points) or closes (by low RMSD with purple points). Additionally, we gain some insight on the time required to capture the opening or closing of the binding cavity.

It is important to note that by limiting our view to only the end-state replica, we encounter time points that may appear contradictory, at a glance. For example in figure 3b, between 3-4ns there are two frames where the protein is closed and the RMSD is slightly greater than 2.0Å. Following, there is one frame where the protein is open but the RMSD is lower than 2.0Å. This appears to be an artifact of coordinate swapping during replica exchanges from other intermediate lambda windows, resulting in apparently contradictory time points.

We address this limitation by viewing all replicas in what we call ‘Color maps’, see Figure 6b for an example. Essentially, our ‘Color map’ analysis is the same as our ‘RMSD/time’ plots but without the RMSD line plot. In other words, we color time points according to the protein state of lowest RMSD and do this for all replicas but do not track the RMSD relative to the closed state. Through a collective view of all replicas, we gain a better perspective of the overall protein conformational sampling and the states they occupy over each replica’s separate trajectories. Using color maps, it becomes visually easy to see if intermediate—higher temperature—lambda windows are able to sample, say the open state, and if this leads to an enhancement in sampling at the end states via replica exchange.

### 3 Results

#### Calculated free energies depend strongly on starting protein conformation

Using the default FEP/REST methodology,<sup>19</sup> we find calculated free energies significantly depend on the protein starting conformation, especially for large perturbations (i.e. opening the cavity from the closed state). To illustrate this, we begin our molecular dynamics simulations both from the protein closed and open conformations then perform alchemical transformations to ligands that occupy another protein conformational state. For example, in the alchemical transformation of benzene to n-hexylbenzene—starting from the protein closed state—we expect to see opening of the binding cavity when the ligand is in the fully interacting n-hexylbenzene state. In this study, we demonstrate using the default FEP protocol settings of a 5ns simulation time and REST region selection does not generate adequate sampling of the motion in the F-helix and does not eliminate the dependence on the initial protein state.

#### Closed-Open Ligand Transformations

An examination of the largest alchemical transformation, benzene to n-hexylbenzene, clearly highlights the sampling challenges faced when using the default FEP/REST protocol. From experimental data of ligand occupancies (Table 1), we expect in our simulations of n-hexylbenzene to see the protein primarily in the open state over the closed state. Instead, we find the protein remains trapped in its initial conformational state whether we start from closed (Fig 3a) or open (Fig 3b) over the course of the 5ns simulation. From the protein closed simulation, the protein only briefly samples the intermediate state around 3ns but never enters the open conformation. As the protein tries to accommodate n-hexylbenzene and enter its preferred open state, protein-ligand strain results, yielding a positive value for  $\Delta\Delta G_{calc}$ (+4.13 kcal/mol). On the other hand, in the protein open simulations, the protein

already begins in its preferred state for n-hexylbenzene and stays only in this open state. As expected, the  $\Delta\Delta G_{calc}$  is negative(-0.61 kcal/mol) as there is no occurrence of large protein-ligand strain in order to open the cavity. By remaining trapped in the initial state, we under-sample the open state if we begin from the closed state or over-sample it if we begin from the open state. Ultimately, we arrive at two very different relative free energies values, where the inconsistency is as large as +4.74 kcal/mol for the same transformation of benzene to n-hexylbenzene.

In the overall set, we similarly observe protein closed simulations to yield positive free energies and negative for protein open simulations. In turn, we find the overall inconsistency to be very high with a RMSI of +4 kcal/mol (Fig 7a, Table 5). Clearly, despite the use of implemented default FEP/REST protocol, we are unable to get sufficient sampling in the protein within the standard 5ns time frame. This is admittedly unsurprising since in the default calculation, only the perturbed ligand R-group is added to the enhanced sampling region. Instead, we encounter sampling problems as the protein remains in its initial conformational state throughout the simulation. As a result, our calculated free energies exhibit high dependence on the initial protein configuration which is reflected by the large RMSI.

### Closed-Intermediate Ligand Transformations

In the case of closed-intermediate alchemical transformations, we find that the calculated free energies still have some (albeit much smaller) dependence on the initial protein conformation, using the default protocol. For this set of alchemical transformations, we find the RMSI to be +0.60 kcal/mol (Fig 7c, Table 2). Considering this set involves a smaller protein conformational change and smaller perturbations to the ligand, it is unsurprising to find the RMSI to be much smaller than our closed-open transformation set.

Although, the collective RMSI for closed-intermediate transformations falls below the acceptable range of less than 1 kcal/mol, we can still see a dependence on the initial protein

configuration by viewing transformations involving n-butylbenzene. For these cases in particular, we observe the same pattern of protein closed simulations yielding positive free energies and negative for protein open simulations. We do not see this pattern for transformations with sec-butylbenzene as it does not partially occupy the open state, unlike n-butylbenzene (Table 1). Through this observation, we demonstrate further that with the default protocol does not eliminate the free energy dependence on the protein starting conformation, even for smaller perturbations. Although, the level of discrepancy is much more comparable to the typically expected sampling error for routine scoring work.

## Experimental Ligand Transformations

Now, when we compare  $\Delta\Delta G_{calc}$  against  $\Delta\Delta G_{exp}$ , we find that simulations starting from the protein closed conformation are further from converging to  $\Delta\Delta G_{exp}$  than when starting from the protein open conformation. Here, we calculate the RMS-'Error' with experiment and find the RMSE for protein closed simulations to be +1.0 kcal/mol and +0.58 kcal/mol with protein open (Fig 8a,Table 8). Interesting, the accuracy versus experiment of using either starting structure with the default protocol is well within the earlier published estimates, and exceptional accuracy is obtained using the open structure, comparable to the experimental accuracy of most common experimental binding assays. However, the fact that protein open simulations are much closer to  $\Delta\Delta G_{exp}$ , once again does demonstrates our calculated free energies depend on the initial protein state. Unsurprisingly, the large RMSE seen for protein closed simulations primarily comes from transformations involving n-butylbenzene. Evidently, we find the simulations involving n-butylbenzene remain trapped in their respective starting conformations, resulting in inadequate sampling in the protein closed simulations (Fig. 4a) versus the protein open simulations (Fig. 4b). Despite performing much smaller alchemical transformations, this shows we still encounter sampling problems that result in  $\Delta\Delta G_{calc}$  that depend on the initial protein conformation, evident when comparing to  $\Delta\Delta G_{exp}$ .

## Including protein residues into the REST region (pREST) improves sampling

Primarily, we encounter major sampling problems when we begin our simulations from the protein closed state and attempt a mutation which should result in the binding cavity opening. In order to facilitate protein motion, thereby enhancing protein sampling, we included 3 key residues spanning the F-helix region into the REST region, which we will denote simulations using this with ‘pREST’ (Fig 2a). By expanding the REST region, we are able to drive the F-helix out its initial state trap by locally heating up key regions and thereby reduce our sampling problem.

To demonstrate the REST improvement over the default protocol, we return to the case of benzene to n-hexylbenzene. Here, we show the facilitation of the helix motion by first referring to Figure 3a which shows that there is no sampling of the open state for the default protocol. Now with pREST, we see a few open state points around 3ns and even a single open point before closing again after our initial step (Fig 5a). Alternatively, we can further illustrate the enhancement of protein sampling by viewing all replicas collectively, using the color maps. In reference to Figure 6a and Figure 6b, we illustrate that there is far less sampling of the intermediate or open protein states in default simulations versus the pREST simulations.

Collectively, we find only some minor improvements in the RMSI for all our closed-open and closed-intermediate transformations while using pREST. For closed-open transformations (Table 6), the RMSI reduces to +2.82 kcal/mol (previously +4 kcal/mol). On the other hand, for closed-intermediate, the RMSI raises slightly to +0.79 kcal/mol (previously +0.60 kcal/mol), which may be due to statistical noise (Table 3). Generally, simulations starting from the closed state had  $\Delta\Delta G_{calc}$  values that moved towards favorability (i.e. more negative  $\Delta\Delta G_{calc}$ ), while for protein open simulations  $\Delta\Delta G_{calc}$  values tended towards unfavorability (i.e. more positive  $\Delta\Delta G_{calc}$ ). This is indicative of the fact that pREST is indeed improving sampling, but it is evident that our  $\Delta\Delta G_{calc}$  are still far from convergence,

given the RMSI is still large, especially for closed/open transformations.

## Long simulations enhances protein conformational sampling from more exchanges

Although we see improvements in sampling with pREST, the standard implemented time frame of 5ns clearly is not long enough to gain adequate sampling, particularly if we start from the protein closed state. By running longer, we allow our simulations to perform more exchanges across replicas and thereby allow for better sampling of all conformational states at the relevant end state replicas.

Returning to our most extreme transformation, benzene to n-hexylbenzene, we have shown pREST alone does not facilitate adequate sampling of the open state (Fig 5a). Now, when we run much longer we see far more sampling of the open protein conformational state in the final 10ns window (Fig 5b).<sup>1</sup> In viewing all the replicas (Fig 6c), we illustrate the dramatic increase in protein conformational sampling in stark contrast to our previous 5ns simulations (Fig 6b).

By simulating longer with pREST we dramatically increase our sampling of the intermediate and open protein states and almost entirely eliminate the dependence on the initial protein conformational state. For the set of closed-open transformations the RMSI dramatically falls to +0.57 kcal/mol (Fig. 7b,Table 7) and a RMSI of +0.43 kcal/mol for the closed-intermediate (Fig. 7d,Table 4). Similarly, for experimental ligand transformations, our RMSE for protein closed simulations falls to +0.54 kcal/mol (Fig. 8b,Table 9). Now, all our inconsistencies in the final calculated free energies and error from experiment fall within a much more reasonable range of less than +1 kcal/mol. It should be noted that this accuracy is comparable to, or possibly better than, the accuracy of many assays used in drug discovery campaigns by pharmaceutical companies.

---

<sup>1</sup>We remind the reader that for closed-open transformations we use the final 15ns to compute the final free energies. We only show the final 10ns in our RMSD/time analysis to avoid overcrowding data points.

## 4 Discussion

In this study, we find for this system that relative free energy calculations can suffer from substantial convergence problems resulting from relatively modest protein conformational changes. Although, the protein conformational changes in T4 lysozyme (L99A) are extremely localized to a rearrangement of a single helix (Fig 1), we still encounter challenges in sampling, likely due to the high energy barrier associated with the reorganization of protein secondary structure. These problems have profound implications for the accuracy of computed relative free energies in these cases. Particularly, we find that calculated relative free energies can depend on the initial protein conformational state by up to 4 kcal/mol.

By looking at alchemical transformations that involve a conformation change in the protein, we show the  $\Delta\Delta G_{s_{calc}}$  are sensitive to the initial protein conformational state when utilizing the default implemented FEP protocol. This sensitivity occurs primarily when the alchemical transformation involves mutating ligands that mainly occupy the closed state (i.e. benzene to n-propyl) into ligands that occupy the intermediate state (i.e. sec-/n-butyl) or, especially, the open state (n-pentyl/n-hexyl) (Table 1). By our RMSD analyses, we show the protein remains trapped in its initial state throughout the simulation when using the implemented default protocol. Because it remains trapped, we are unable to adequately sample the necessary protein conformational states and thereby obtain inconsistent  $\Delta\Delta G_{s_{calc}}$  depending on whether we start simulations from the open or closed protein configuration. Interestingly, despite these slow degrees of freedom being present in the system, the accuracy of the default protocol is still in line with earlier reported work, and exceptionally accurate if the more open crystal structure is initially used.

Without prior knowledge of preferred protein conformational states on ligand binding, we can arrive at very different binding affinity predictions based on the initial protein state being used in the simulation. For this system, by starting from the protein close state and growing the ligand we obtain  $\Delta\Delta G_{calc}$  values that appear overly positive or unfavorable due to high protein-ligand energy strain and inability to sample the open state. However, when

we begin with the protein open state our  $\Delta\Delta G_{calc}$  values have deeply surprising agreement with experiment, despite our inability inability to sample the closed state. If we only had the crystal structure of the closed protein-ligand complexes, we would blindly conclude that the much larger, open-ligands bind to T4 lysozyme worse than smaller ones. On the other hand, if only the open protein-ligand complexes were available, the opposite would be concluded in that larger ligands are better binders than smaller ligands. We speculatively note, this may be problematic for fragment based drug discovery campaigns, but is somewhat unlikely for high-throuput screening campaigns, where most of the screened molecules will be highly elaborated.

By including key residues into the REST region and simulating longer, we reduce the  $\Delta\Delta G_{calc}$  dependence on the initial protein configuration to a more reasonable range of less than 1 kcal/mol. Through expanding the REST region, intermediate lambda windows are able to more easily access the intermediate and open conformations by effectively heating key residues that facilitated protein motion, illustrated in Fig 6b. Further, by simulating longer we allow for more exchanges between replicas, which in turn enhances sampling at our physically relevant end state replica (Fig 6c). With these modifications to the default protocol, we almost completely converge our  $\Delta\Delta G_{calc}$  to the same value regardless of the starting protein conformation.

Generally, our brute-force approach of simulating longer and multiple trials with varied protein structures is not a desirable or feasible approach, especially in early drug discovery phases. At the industrial level, ligand libraries can be large—driving computational cost exponentially if we simulate longer—or experimental structures can be sparse for new therapeutic protein targets. For future studies, approaches using Markov State Models (MSMs)<sup>55</sup> can potentially be of great use for identifying discrete protein conformations. MSMs build a representation of the conformational space from batches of short molecular simulations, whereby the discrete states and transition rates between them can be determined in an efficient manner. Utilizing MSMs can thereby provide useful insight on the various protein

conformational states before running free energy predictions.

## 5 Conclusions

Overall, we have shown that the presence of kinetically distinct protein conformational states can dramatically impact the accuracy free energy calculations. In a worse-case scenario for this study, if only the apo protein structure was available and with no prior knowledge of the discrete states, identifying the bias in the free energies would have been challenging, if at all possible. Remarkably, the absolute agreement with experiment of the default protocol ranges from within the previously reported accuracy of 1.0 kcal/mol, to exceptionally accurate at 0.58 kcal/mol, despite such a slow degree of freedom being present in the system. It would have been especially challenging as there would essentially be no indicators that the final free energies were sensitive to the initial protein configuration. Only from prior knowledge of the discrete states and by our tedious systematic trials were we able to identify and address the bias in our final calculated free energies. It may be conforming that even without this prior knowledge, good to very good agreement with experiment is obtained for the majority of ligands.

Although FEP calculations have shown tremendous recent successes on a variety of protein targets,<sup>12</sup> we demonstrate challenges in protein sampling remain. Using T4 lysozyme (L99A) as our simple model system, we highlight sampling problems even from a relatively small ( 1-3.5Å) and localized single helix rearrangement in response to a series of growing ligands. Through this study, we show using a typical 5ns simulation with only ligand atoms in the REST region, yields free energies that are sensitive to the initial protein conformation. Only by longer simulation times and expansion of the REST region to include key protein residues were we able to eliminate the dependence on the starting conformation. This study demonstrates that special attention and care should be exercised when performing FEP calculations where regions of flexibility surround the binding site. More importantly, prior to

performing binding free energy calculations, we present strong evidence on the importance of identifying the occurrence of protein conformational changes upon ligand binding,

## Acknowledgement

N.M.L. thanks Dmitry Lupyan and Joseph Goose for helpful discussions and technical support. Financial support for N.M.L. was provided by Schrödinger and the National Science Foundation Graduate Research Fellowship (DGE-1321846). D.L.M. appreciates financial support from the National Institutes of Health (1R01GM108889-01). L.W. and R.A. are supported by Schrödinger.

## Figures





Figure 1: In Fig.1a are overlaid cartoon representations of T4 lysozyme using the protein-ligand bound crystal structures of the toluene(4W53), n-butylbenzene(4W57), and n-hexylbenzene(4W59) complexes where the F-helix region is highlighted in yellow. Fig.1b illustrates a closer view of strictly the F-helix in the closed(purple), intermediate(cyan), and open(green) conformational states and their corresponding ligands as found in the crystal structure. Below are the molecular surface representations of the binding cavity in the closed(1c), intermediate(1d), and open(1e) states with the ligands represented in an orange space-filling model. In Fig.1f is the full series of congeneric ligands used in this study; their protein state occupancies can be found in Table 1. Images were created using Maestro.<sup>33</sup>

(a) Selected residues in pREST




(b) Residue Glu108

(c) Residue Val111

Figure 2: Fig. 2a highlights the 3 residues selected to be included into the REST region for ‘pREST’ simulations. Fig. 2b and Fig. 2c illustrates the motion of the C<sub>α</sub>’s during the transition between the protein closed and open states, where the C<sub>α</sub> in Glu108 undergoes a motion of approximately 2 Å, while in Val11 the C<sub>α</sub> move approximately 3.5 Å.



(a) Protein closed simulation



(b) Protein open simulation

Figure 3: Plotted is the RMSD( $\text{\AA}$ ) relative to the closed helix conformation (black line), where each time point is colored according to the protein state of lowest RMSD to the trajectory. Here, using the default protocol in the transformation of benzene to n-hexylbenzene, RMSD/time plots correspond to the final end state of n-hexylbenzene ( $\lambda_{11}$ ). Fig. 3a corresponds to the simulation that began from the protein closed state and Fig. 3b is from the simulation started from the protein open state. The legends indicate the percentage of sampling of each protein conformational state from the trajectory.


(a) Protein closed simulation



(b) Protein open simulation

Figure 4: RMSD/time plots correspond to the final end state of n-butylbenzene ( $\lambda_{11}$ ) while using the default protocol for transformation of benzene to n-butylbenzene. Fig. 4a and Fig. 4b correspond to simulations that were started from the protein closed or open state, respectively. The legends indicate the percentage of sampling of each protein conformational state from the trajectory.

(a) Protein closed simulation, pREST



(b) Protein closed simulation, pREST extended

Figure 5: Using the modified REST region (pREST) in the transformation of benzene to n-hexylbenzene, we plot the RMSD/time corresponding to the n-hexylbenzene state ( $\lambda_{11}$ ) with the simulation starting from protein closed for the first 5ns (Fig. 5a) and the final 10ns from an extended run up to 55ns (Fig. 5b).




Figure 6: Color maps of simulations starting from the protein closed state for the benzene to n-hexylbenzene alchemical transformation. Lines at every frame are colored accordingly to the protein state of lowest RMSD (purple-closed, cyan-intermediate, green-open). Fig. 6a corresponds to simulations using the default protocol. Fig. 6b represents simulations using the modified REST region protocol (pREST). Fig. 6c illustrates the enhancement in protein conformational sampling through extending the pREST simulation time up to 55ns.




Figure 7:  $\Delta\Delta G_{calc}$  from MD simulations beginning from the protein closed versus open state. Fig. 7a plots relative free energies obtained using the default protocol from the ‘closed-open’ alchemical transformations set, yielding a RMSI of 4.0 kcal/mol. Fig. 7b are the final computed free energies with simulations carried out to 55ns using pREST, giving RMSI of 0.57 kcal/mol. Fig. 7c plots relative free energies from the ‘closed-intermediate’ set using the default protocol which gives an RMSI of 0.6 kcal/mol. Fig. 7d are free energies with simulations carried out up to 25ns using pREST, giving RMSI of 0.43 kcal/mol. Numerical data for each plot can be found in Tables 2, 4, 5, and 7.




Figure 8:  $\Delta\Delta G_{calc}$  from MD simulations beginning from the protein closed (purple) and open (green) state compared against  $\Delta\Delta G_{exp}$ . Fig. 8a plots free energies obtained using the default protocol which gives a 1.0 kcal/mol and 0.58 kcal/mol RMSE for protein closed and open simulations, respectively. Fig. 8b plots the final free energies after applying pREST, yielding an RMSE of 0.54 for protein closed and 0.50 kcal/mol for protein closed simulations. Numerical data for each plot can be found in Tables 8 and 9.

## Tables

Table 1: Experimental protein-ligand occupancies and affinities<sup>a</sup>

PDB	Ligand	C	I	O	$\Delta G_{exp}$	$\sigma_{exp}$	Set
4W52	benzene	90	-	-	-5.19	0.16	C
4W53	toluene	80	20	-	-5.52	0.04	C
4W54	ethylbenzene	50	50	-	-5.76	0.07	C
4W55	n-propylbenzene	60	40	-	-6.55	0.02	C
4W56	sec-butylbenzene	40	60	-	N/A	-	I
4W57	n-butylbenzene	10	60	30	-6.70	0.02	I
4W58	n-pentylbenzene	30	-	70	N/A	-	O
4W59	n-hexylbenzene	30	-	70	N/A	-	O

<sup>a</sup> Listed are the PDB codes of the corresponding protein-ligand complex crystal structures and the ligand occupancies<sup>1</sup> for the closed(C), intermediate(I), and open(O) protein conformations by percentage. Under the ‘Set’ column, shaded in grey, is each ligand’s primary state of occupancy which we use to categorize our alchemical transformation sets.

Binding affinities for n-pentylbenzene and n-hexylbenzene were inaccessible due to solubility limits.<sup>1</sup> Experimental binding free energies and their uncertainties are in units of kcal/mol.<sup>22</sup>

Table 2: Closed-Intermediate Transformations<sup>a</sup>

Ligand 1	Ligand 2	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$	$\Delta\Delta G_\epsilon$
benzene	n-butylbenzene	0.58	0.07	-0.59	0.08	1.17
toluene	n-butylbenzene	-0.28	0.06	-1.27	0.09	0.99
ethylbenzene	n-butylbenzene	0.24	0.06	-0.23	0.07	0.47
n-propylbenzene	n-butylbenzene	0.99	0.06	0.63	0.04	0.36
benzene	sec-butylbenzene	2.36	0.09	2.14	0.11	0.22
toluene	sec-butylbenzene	1.47	0.07	1.14	0.08	0.33
ethylbenzene	sec-butylbenzene	1.90	0.08	1.77	0.08	0.13
n-propylbenzene	sec-butylbenzene	2.86	0.06	2.67	0.05	0.19

<sup>a</sup> Calculated free energies and their uncertainties in ‘closed-intermediate’ alchemical transformations from simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies are in units of kcal/mol. The inconsistency( $\Delta\Delta G_\epsilon$ ) is obtained from the difference in the relative free energies between protein closed and open simulations. Inconsistency is highlighted red or green if above or below 1 kcal/mol, respectively. RMSI is 0.6 kcal/mol. Table is represented in Fig 7c.

Table 3: Closed-Intermediate Transformations pREST<sup>a</sup>

Ligand 1	Ligand 2	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$	$\Delta\Delta G_\epsilon$
benzene	n-butylbenzene	-0.07	0.10	-0.72	0.12	0.65
toluene	n-butylbenzene	0.89	0.10	-0.42	0.08	1.31
ethylbenzene	n-butylbenzene	-0.20	0.09	-0.43	0.08	0.23
n-propylbenzene	n-butylbenzene	1.02	0.06	0.49	0.06	0.53
benzene	sec-butylbenzene	0.45	0.08	1.34	0.12	0.89
toluene	sec-butylbenzene	0.60	0.12	0.60	0.10	0.00
ethylbenzene	sec-butylbenzene	1.09	0.10	1.69	0.09	0.60
n-propylbenzene	sec-butylbenzene	1.88	0.07	3.05	0.07	1.17

<sup>a</sup> Calculated free energies and their uncertainties in ‘closed-intermediate’ alchemical transformations from ‘pREST’ simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies were calculated using frames from 0 to 5ns. Free energies are in units of kcal/mol. The inconsistency( $\Delta\Delta G_\epsilon$ ) is obtained from the difference in the relative free energies between protein closed and open simulations. Inconsistency is highlighted red or green if above or below 1 kcal/mol, respectively. RMSI is 0.79 kcal/mol. Table is represented in Fig 7d.

Table 4: Closed-Intermediate Transformations pREST Extended<sup>a, b</sup>

Ligand 1	Ligand 2	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$	$\Delta\Delta G_\epsilon$
benzene	n-butylbenzene	-1.14 <sup>b</sup>	0.09	-0.72	0.12	0.42
toluene	n-butylbenzene	-0.62 <sup>b</sup>	0.09	-0.96 <sup>a</sup>	0.09	0.34
ethylbenzene	n-butylbenzene	-0.20	0.09	-0.43	0.08	0.23
n-propylbenzene	n-butylbenzene	0.51 <sup>b</sup>	0.06	0.49	0.06	0.02
benzene	sec-butylbenzene	0.45	0.08	1.34	0.12	0.89
toluene	sec-butylbenzene	0.60	0.12	0.60	0.10	0.00
ethylbenzene	sec-butylbenzene	1.09	0.10	1.69	0.09	0.60
n-propylbenzene	sec-butylbenzene	1.88	0.07	1.84 <sup>b</sup>	0.06	0.04

<sup>a</sup> Calculated free energies and their uncertainties in ‘closed-intermediate’ alchemical transformations from ‘pREST’ simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies were calculated using frames from 15 to 25ns. Free energies are in units of kcal/mol. The inconsistency( $\Delta\Delta G_\epsilon$ ) is obtained from the difference in the relative free energies between protein closed and open simulations. Inconsistency is highlighted red or green if above or below 1 kcal/mol, respectively. RMSI is 0.43 kcal/mol. Table is represented in Fig 7d.

<sup>b</sup> Only these alchemical transformations were extended up to 25ns of simulation time in order to lower the inconsistency and error with experiment.

Table 5: Closed-Open Transformations<sup>a</sup>

Ligand 1	Ligand 2	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$	$\Delta\Delta G_\epsilon$
benzene	n-pentylbenzene	2.36	0.12	-1.33	0.10	3.69
toluene	n-pentylbenzene	1.77	0.09	0.34	0.10	1.43
ethylbenzene	n-pentylbenzene	2.45	0.08	0.46	0.09	1.99
n-propylbenzene	n-pentylbenzene	3.46	0.08	-0.22	0.07	3.68
benzene	n-hexylbenzene	4.13	0.16	-0.61	0.15	4.74
toluene	n-hexylbenzene	2.90	0.13	-1.63	0.08	4.53
ethylbenzene	n-hexylbenzene	3.63	0.11	-0.76	0.09	4.39
n-propylbenzene	n-hexylbenzene	5.85	0.10	0.13	0.06	5.72

<sup>a</sup> Calculated free energies and their uncertainties in ‘closed-open’ alchemical transformations from simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies were calculated using frames from 0 to 5ns. Free energies are in units of kcal/mol. The inconsistency( $\Delta\Delta G_\epsilon$ ) is obtained from the difference in the relative free energies between protein closed and open simulations. Inconsistency is highlighted red or green if above or below 1 kcal/mol, respectively. RMSI is 4.0 kcal/mol. Table is represented in Fig 7a.

Table 6: Closed-Open Transformations pREST<sup>a</sup>

Ligand 1	Ligand 2	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$	$\Delta\Delta G_\epsilon$
benzene	n-pentylbenzene	1.49	0.13	0.07	0.10	1.42
toluene	n-pentylbenzene	1.41	0.12	0.79	0.13	0.62
ethylbenzene	n-pentylbenzene	2.90	0.10	1.23	0.09	1.67
n-propylbenzene	n-pentylbenzene	4.25	0.09	1.01	0.07	3.24
benzene	n-hexylbenzene	2.75	0.14	1.25	0.11	1.50
toluene	n-hexylbenzene	3.22	0.11	-1.15	0.11	4.37
ethylbenzene	n-hexylbenzene	3.34	0.12	-0.22	0.11	3.56
n-propylbenzene	n-hexylbenzene	4.93	0.12	1.21	0.11	3.72

<sup>a</sup> Calculated free energies and their uncertainties in ‘closed-open’ alchemical transformations from ‘pREST’ simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies are in units of kcal/mol. The inconsistency( $\Delta\Delta G_\epsilon$ ) is obtained from the difference in the relative free energies between protein closed and open simulations. The inconsistency is highlighted red or green if above or below 1 kcal/mol, respectively. RMSI is 2.82 kcal/mol.

Table 7: Closed-Open Transformations pREST Extended<sup>a</sup>

Ligand 1	Ligand 2	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$	$\Delta\Delta G_\epsilon$
benzene	n-pentylbenzene	1.37	0.08	1.41	0.08	0.04
toluene	n-pentylbenzene	1.08	0.08	0.81	0.07	0.27
ethylbenzene	n-pentylbenzene	1.71	0.08	1.50	0.07	0.21
n-propylbenzene	n-pentylbenzene	3.20	0.06	2.34	0.05	0.86
benzene	n-hexylbenzene	2.15	0.09	1.14	0.08	1.01
toluene	n-hexylbenzene	0.32	0.10	1.10	0.07	0.78
ethylbenzene	n-hexylbenzene	1.96	0.08	2.17	0.07	0.21
n-propylbenzene	n-hexylbenzene	3.38	0.07	3.12	0.06	0.26

<sup>a</sup> Calculated free energies and their uncertainties in ‘closed-open’ alchemical transformations from extended ‘pREST’ simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies are in units of kcal/mol. Free energies were calculated using frames from 40 to 55ns. The inconsistency( $\Delta\Delta G_\epsilon$ ) is obtained from the difference in the relative free energies between protein closed and open simulations. Inconsistency is highlighted red or green if above or below 1 kcal/mol, respectively. RMSI is 0.57 kcal/mol. Table is represented in Fig 7b.

Table 8: Experimental Ligand Transformations<sup>a</sup>

Ligand 1	Ligand 2	$\Delta G_{exp}$	$\sigma_{exp}$	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$
benzene	toluene	-0.33	0.16	0.71	0.04	0.59	0.05
benzene	ethylbenzene	-0.19	0.17	0.15	0.05	-0.09	0.06
benzene	n-propylbenzene	-1.36	0.16	-0.82	0.06	-0.69	0.08
benzene	n-butylbenzene	-1.51	0.16	0.58	0.07	-0.59	0.08
toluene	ethylbenzene	-0.24	0.08	-0.59	0.04	-0.40	0.05
toluene	n-propylbenzene	-1.03	0.04	-2.21	0.05	-0.89	0.05
toluene	n-butylbenzene	-1.18	0.04	-0.28	0.06	-1.27	0.09
ethylbenzene	n-propylbenzene	-0.79	0.07	-0.86	0.05	-0.80	0.03
ethylbenzene	n-butylbenzene	-0.94	0.07	0.24	0.06	-0.23	0.07
n-propylbenzene	n-butylbenzene	-0.15	0.03	0.99	0.06	0.63	0.04

<sup>a</sup> Experimental and calculated free energies with their uncertainties in alchemical transformations involving ligands with available experimental affinities. Calculated free energies are from simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies are in units of kcal/mol. Here, we compare  $\Delta\Delta G_C$  and  $\Delta\Delta G_O$  against  $\Delta G_{exp}$  to compute the error with experiment. RMSE is 1.0 kcal/mol and 0.58 kcal/mol for closed and open simulations, respectively.

Table is represented in Fig 8a.

Table 9: Experimental Ligand Transformations pREST<sup>a</sup>

Ligand 1	Ligand 2	$\Delta G_{exp}$	$\sigma_{exp}$	$\Delta\Delta G_C$	$\sigma_C$	$\Delta\Delta G_O$	$\sigma_O$
benzene	toluene	-0.33	0.16	0.10	0.08	0.31	0.06
benzene	ethylbenzene	-0.19	0.17	-0.20	0.08	-0.62	0.08
benzene	n-propylbenzene	-1.36	0.16	-1.57	0.09	-1.26	0.09
benzene	n-butylbenzene	-1.51	0.16	-1.14	0.11	-0.72	0.12
toluene	ethylbenzene	-0.24	0.08	-0.84	0.07	-0.83	0.07
toluene	n-propylbenzene	-1.03	0.04	-1.91	0.09	-1.44	0.08
toluene	n-butylbenzene	-1.18	0.04	-0.62	0.09	-0.96	0.09
ethylbenzene	n-propylbenzene	-0.79	0.07	-1.08	0.07	-0.80	0.06
ethylbenzene	n-butylbenzene	-0.94	0.07	-0.20	0.09	-0.43	0.08
n-propylbenzene	n-butylbenzene	-0.15	0.03	0.51	0.06	0.49	0.06

<sup>a</sup> Experimental and calculated free energies with their uncertainties in alchemical transformations involving ligands with available experimental affinities. Calculated free energies are from ‘pREST’ simulations that were started from the protein closed state( $\Delta\Delta G_C$ ) versus the protein open state( $\Delta\Delta G_O$ ). Free energies are in units of kcal/mol. Here, we compare  $\Delta\Delta G_C$  and  $\Delta\Delta G_O$  against  $\Delta G_{exp}$  to compute the error with experiment. RMSE is 0.54 kcal/mol and 0.50 kcal/mol for closed and open simulations, respectively. Table is represented in Fig 8b.

## **Supporting Information Available**

The following files are provided in the Supporting Information:

### **Prepared protein-ligand crystal structures: `xtal_preppe.tar.gz`**

Contains the reference protein-ligand crystal structures that have been aligned and prepared through the ‘Protein Preparation Wizard’ tool as `.mae` files.

### **RMSD analysis: `scripts.tar.gz`**

Contains the scripts `RMSD-analysis.tcl` and `plot-RMSD.py` used for RMSD analysis. A `README.txt` file has been provided which details how to generate the RMSD/time and Colormap plots found in the paper.

### **FEP data: `C-I_default.tar.gz`, `C-I_pPREST.tar.gz`, `C-O_default.tar.gz`, `C-O_pREST.tar.gz`, `EXP_default.tar.gz`, and `EXP_pREST.tar.gz`**

Contains archived data from the FEP calculations for each alchemical transformation set, where within each archived set are directories for individual FEP calculations. The input `.mae` files containing the protein/ligand structures, the simulation configuration `.msj` files, energy files, and a `README.txt` that summarizes information about each FEP calculation have been included. Trajectories have been neglected due to file size limitations.

Full set of Supporting Information files can be downloaded at <http://n2t.net/ark:/b7280/d1js3b> (doi:10.7280/D1JS3B). This material is available free of charge via the Internet at <http://pubs.acs.org/>.

## References

- (1) Merski, M.; Fischer, M.; Balias, T. E.; Eidam, O.; Shoichet, B. K. *Proceedings of the National Academy of Sciences* **2015**, *112*, 5039–5044.
- (2) Overington, J. P.; Al-Lazikani, B.; Hopkins, A. L. *Nature reviews Drug discovery* **2006**, *5*, 993–996.
- (3) Landry, Y.; Gies, J.-P. *Fundamental & Clinical Pharmacology* **2008**, *22*, 1–18.
- (4) Lundstrom, K. In *G Protein-Coupled Receptors in Drug Discovery*; Leifert, R. W., Ed.; Humana Press: Totowa, NJ, 2009; Chapter An Overview on GPCRs and Drug Discovery: Structure-Based Drug Design and Structural Biology on GPCRs, pp 51–66.
- (5) Homeyer, N.; Stoll, F.; Hillisch, A.; Gohlke, H. *Journal of Chemical Theory and Computation* **2014**, *10*, 3331–3344, PMID: 26588302.
- (6) Chipot, C.; Pohorille, A. *Free energy calculations*; Springer, 2007.
- (7) Chodera, J. D.; Mobley, D. L.; Shirts, M. R.; Dixon, R. W.; Branson, K.; Pande, V. S. *Current opinion in structural biology* **2011**, *21*, 150–160.
- (8) Knight, J. L.; Brooks, C. L. *Journal of computational chemistry* **2009**, *30*, 1692–1700.
- (9) Zheng, L.; Chen, M.; Yang, W. *Proceedings of the National Academy of Sciences* **2008**, *105*, 20227–20232.
- (10) Gallicchio, E.; Levy, R. M. *Current opinion in structural biology* **2011**, *21*, 161–166.
- (11) Hansen, N.; van Gunsteren, W. F. *Journal of Chemical Theory and Computation* **2014**, *10*, 2632–2647, PMID: 26586503.
- (12) Wang, L.; Wu, Y.; Deng, Y.; Kim, B.; Pierce, L.; Krilov, G.; Lupyan, D.; Robinson, S.; Dahlgren, M. K.; Greenwood, J.; Romero, D. L.; Masse, C.; Knight, J. L.;

- Steinbrecher, T.; Beuming, T.; Damm, W.; Harder, E.; Sherman, W.; Brewer, M.; Wester, R.; Murcko, M.; Frye, L.; Farid, R.; Lin, T.; Mobley, D. L.; Jorgensen, W. L.; Berne, B. J.; Friesner, R. A.; Abel, R. *Journal of the American Chemical Society* **2015**, *137*, 2695–2703, PMID: 25625324.
- (13) Mobley, D. L.; Klimovich, P. V. *The Journal of chemical physics* **2012**, *137*, 230901.
- (14) Kaus, J. W.; Harder, E.; Lin, T.; Abel, R.; McCammon, J. A.; Wang, L. *Journal of Chemical Theory and Computation* **2015**, *11*, 2670–2679.
- (15) Mobley, D. L.; Dill, K. A. *Structure* **2009**, *17*, 489 – 498.
- (16) Mobley, D. L.; Graves, A. P.; Chodera, J. D.; McReynolds, A. C.; Shoichet, B. K.; Dill, K. A. *Journal of Molecular Biology* **2007**, *371*, 1118 – 1134.
- (17) Jonas BostrÃum, â., \*; Anders Hogner, â.; ; SchmittâAq, S. *Journal of Medicinal Chemistry* **2006**, *49*, 6716–6725, PMID: 17154502.
- (18) Gutteridge, A.; Thornton, J. *Journal of Molecular Biology* **2005**, *346*, 21 – 28.
- (19) Wang, L.; Berne, B. J.; Friesner, R. A. *Proceedings of the National Academy of Sciences* **2012**, *109*, 1937–1942.
- (20) Eriksson, A. E.; Baase, W. A.; Zhang, X.-J.; Heinz, D. W.; Blaber, M.; Baldwin, E. P.; Matthews, B. W. *Science* **1992**, *255*, 178–183.
- (21) Eriksson, A.; Baase, W. A.; Matthews, B. W. *Journal of molecular biology* **1993**, *229*, 747–769.
- (22) Morton, A.; Baase, W. A.; Matthews, B. W. *Biochemistry* **1995**, *34*, 8564–8575, PMID: 7612598.
- (23) Morton, A.; Matthews, B. W. *Biochemistry* **1995**, *34*, 8576–8588, PMID: 7612599.

- (24) Wei, B. Q.; Baase, W. A.; Weaver, L. H.; Matthews, B. W.; Shoichet, B. K. *Journal of molecular biology* **2002**, *322*, 339–355.
- (25) Wei, B. Q.; Weaver, L. H.; Ferrari, A. M.; Matthews, B. W.; Shoichet, B. K. *Journal of molecular biology* **2004**, *337*, 1161–1182.
- (26) Graves, A. P.; Brenk, R.; Shoichet, B. K. *Journal of medicinal chemistry* **2005**, *48*, 3714–3728.
- (27) Hermans, J.; Wang, L. *Journal of the American Chemical Society* **1997**, *119*, 2707–2714.
- (28) Boresch, S.; Tettinger, F.; Leitgeb, M.; Karplus, M. *The Journal of Physical Chemistry B* **2003**, *107*, 9535–9551.
- (29) Deng, Y.; Roux, B. *Journal of Chemical Theory and Computation* **2006**, *2*, 1255–1273.
- (30) Mann, G.; Hermans, J. *Journal of molecular biology* **2000**, *302*, 979–989.
- (31) Boyce, S. E.; Mobley, D. L.; Rocklin, G. J.; Graves, A. P.; Dill, K. A.; Shoichet, B. K. *Journal of Molecular Biology* **2009**, *394*, 747 – 763.
- (32) Wang, L.; Deng, Y.; Knight, J. L.; Wu, Y.; Kim, B.; Sherman, W.; Shelley, J. C.; Lin, T.; Abel, R. *Journal of Chemical Theory and Computation* **2013**, *9*, 1282–1293, PMID: 26588769.
- (33) Schrödinger, LLC, Maestro, version 10.3. 2015.
- (34) Schrödinger, LLC, Schrödinger Suite 2015-3 Protein Preparation Wizard. 2015.
- (35) Schrödinger, LLC, Epik, version 3.3. 2015.
- (36) Schrödinger, LLC, Impact, version 6.8. 2015.
- (37) Schrödinger, LLC, Prime, version 4.1. 2015.

- (38) Madhavi Sastry, G.; Adzhigirey, M.; Day, T.; Annabhimoju, R.; Sherman, W. *Journal of Computer-Aided Molecular Design* **2013**, *27*, 221–234.
- (39) Lim, N. M. Schrödinger Academy Molecular Dynamics Ligand FEP Tutorial. Schrödinger: New York, NY, 2015.
- (40) Schrödinger, LLC, Maestro-Desmond Interoperability Tools, version 4.3. 2015.
- (41) Liu, S.; Wu, Y.; Lin, T.; Abel, R.; Redmann, J. P.; Summa, C. M.; Jaber, V. R.; Lim, N. M.; Mobley, D. L. *Journal of Computer-Aided Molecular Design* **2013**, *27*, 755–770.
- (42) Liu, P.; Kim, B.; Friesner, R. A.; Berne, B. J. *Proceedings of the National Academy of Sciences of the United States of America* **2005**, *102*, 13749–13754.
- (43) Wang, L.; Friesner, R. A.; Berne, B. J. *The Journal of Physical Chemistry B* **2011**, *115*, 9431–9438, PMID: 21714551.
- (44) D.E. Shaw Research, Desmond Molecular Dynamics System, version 4.3. 2015.
- (45) Bowers, K. J.; Chow, E.; Xu, H.; Dror, R. O.; Eastwood, M. P.; Gregersen, B. A.; Klepeis, J. L.; Kolossvary, I.; Moraes, M. A.; Sacerdoti, F. D.; Salmon, J. K.; Shan, Y.; Shaw, D. E. Scalable Algorithms for Molecular Dynamics Simulations on Commodity Clusters. Proceedings of the 2006 ACM/IEEE Conference on Supercomputing. New York, NY, USA, 2006.
- (46) Shrivakumar, D.; Williams, J.; Wu, Y.; Damm, W.; Shelley, J.; Sherman, W. *Journal of Chemical Theory and Computation* **2010**, *6*, 1509–1519, PMID: 26615687.
- (47) Guo, Z.; Mohanty, U.; Noehre, J.; Sawyer, T. K.; Sherman, W.; Krilov, G. *Chemical Biology & Drug Design* **2010**, *75*, 348–359.
- (48) D.E. Shaw Research, Desmond Users Guide. version 3.6.1.1/0.8.

- (49) Feller, S. E.; Zhang, Y.; Pastor, R. W.; Brooks, B. R. *The Journal of Chemical Physics* **1995**, *103*, 4613–4621.
- (50) Harder, E.; Damm, W.; Maple, J.; Wu, C.; Reboul, M.; Xiang, J. Y.; Wang, L.; Lupyan, D.; Dahlgren, M. K.; Knight, J. L.; Kaus, J. W.; Cerutti, D. S.; Krilov, G.; Jorgensen, W. L.; Abel, R.; Friesner, R. A. *Journal of Chemical Theory and Computation* **2016**, *12*, 281–296, PMID: 26584231.
- (51) Bennett, C. H. *Journal of Computational Physics* **1976**, *22*, 245–268.
- (52) Hahn, A. M.; Then, H. *Physical Review E* **2009**, *80*, 031111.
- (53) Humphrey, W.; Dalke, A.; Schulten, K. *Journal of Molecular Graphics* **1996**, *14*, 33–38.
- (54) Eargle, J.; Wright, D.; Luthey-Schulten, Z. *Bioinformatics* **2006**, *22*, 504–506.
- (55) Bowman, G. R.; Pande, V. S.; Noé, F. *An introduction to markov state models and their application to long timescale molecular simulation*; Springer Science & Business Media, 2013; Vol. 797.