

Basic R: Matrices

Nathan Mokhtarzadeh

January 25, 2018

Matrix problems

1. Suppose

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}$$

- (a) Check that $A^3 = \mathbf{0}$
- (b) Replace the third column of A by the sum of the second and third columns

First, produce A

```
A <- matrix(c(1,1,3,5,2,6,-2,-1,-3), nrow = 3, byrow = TRUE)
A
##      [,1] [,2] [,3]
## [1,]    1    1    3
## [2,]    5    2    6
## [3,]   -2   -1   -3
```

Then, add the columns 2 and 3 and assign the sum to the third column

```
A[,3] <- A[,2] + A[,3]
A
##      [,1] [,2] [,3]
## [1,]    1    1    4
## [2,]    5    2    8
## [3,]   -2   -1   -4
```

2. Create the following matrix B with 15 rows

$$B = \begin{bmatrix} 10 & -10 & 10 \\ 10 & -10 & 10 \\ \dots & \dots & \dots \\ 10 & -10 & 10 \end{bmatrix}$$

Calculate the 3x3 matrix $B^T B$. You can make this calculation with the function `crossprod()`. See the documentaion.

```
B <- matrix(c(10, -10, 10), ncol = 3, nrow = 15)
crossprod(B)

##      [,1] [,2] [,3]
## [1,] 1500 1500 1500
## [2,] 1500 1500 1500
## [3,] 1500 1500 1500
```

3. Create a 6 x 6 matrix `matE` with every element equal to 0. check what the functions `row()` and `col()` return when applied to `matE`.

Now, create the 6 x 6 matix:

$$\begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}$$

Here is `matE`, a 6x6 matrix of 0's followed by `row(matE)` and `col(matE)`

```
matE <- matrix(rep(0,36), nrow = 6, byrow = TRUE)

# Note what the functions row() and col() do
row(matE)

##      [,1] [,2] [,3] [,4] [,5] [,6]
## [1,]    1    1    1    1    1    1
## [2,]    2    2    2    2    2    2
## [3,]    3    3    3    3    3    3
## [4,]    4    4    4    4    4    4
## [5,]    5    5    5    5    5    5
## [6,]    6    6    6    6    6    6

col(matE)

##      [,1] [,2] [,3] [,4] [,5] [,6]
## [1,]    1    2    3    4    5    6
## [2,]    1    2    3    4    5    6
## [3,]    1    2    3    4    5    6
## [4,]    1    2    3    4    5    6
## [5,]    1    2    3    4    5    6
## [6,]    1    2    3    4    5    6
```

```

# With a little experimentation you would see
# that the specified pattern is in the |1|'s
row(matE)-col(matE)

##      [,1] [,2] [,3] [,4] [,5] [,6]
## [1,]    0   -1   -2   -3   -4   -5
## [2,]    1    0   -1   -2   -3   -4
## [3,]    2    1    0   -1   -2   -3
## [4,]    3    2    1    0   -1   -2
## [5,]    4    3    2    1    0   -1
## [6,]    5    4    3    2    1    0

# so you use the locations of the 1's to modify matE
matE[abs(row(matE)-col(matE))==1] <- 1
matE

##      [,1] [,2] [,3] [,4] [,5] [,6]
## [1,]    0    1    0    0    0    0
## [2,]    1    0    1    0    0    0
## [3,]    0    1    0    1    0    0
## [4,]    0    0    1    0    1    0
## [5,]    0    0    0    1    0    1
## [6,]    0    0    0    0    1    0

```

4. Look at the help for the function `outer()`. Now, create the following patterned matrix:

$$\begin{bmatrix} 0 & 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 & 5 \\ 2 & 3 & 4 & 5 & 6 \\ 3 & 4 & 5 & 6 & 7 \\ 4 & 5 & 6 & 7 & 8 \end{bmatrix}$$

```

a <- 0:4
A <- outer(a,a,"+")
A

##      [,1] [,2] [,3] [,4] [,5]
## [1,]    0    1    2    3    4
## [2,]    1    2    3    4    5
## [3,]    2    3    4    5    6
## [4,]    3    4    5    6    7
## [5,]    4    5    6    7    8

```

Use `outer()` a little more to make sure you get it.

```

B <- outer(a,a, "*")
B

```

```
##      [,1] [,2] [,3] [,4] [,5]
## [1,]    0    0    0    0    0
## [2,]    0    1    2    3    4
## [3,]    0    2    4    6    8
## [4,]    0    3    6    9   12
## [5,]    0    4    8   12   16
```

and

```
b <- 5:10
```

```
C <- outer(a,b,"+")
```

C

```
##      [,1] [,2] [,3] [,4] [,5] [,6]
## [1,]    5    6    7    8    9   10
## [2,]    6    7    8    9   10   11
## [3,]    7    8    9   10   11   12
## [4,]    8    9   10   11   12   13
## [5,]    9   10   11   12   13   14
```

and finally -- make sure you check the values.

```
D <- outer(b,a, "%%")
```

D

```
##      [,1] [,2] [,3] [,4] [,5]
## [1,]   NA    0    1    2    1
## [2,]   NA    0    0    0    2
## [3,]   NA    0    1    1    3
## [4,]   NA    0    0    2    0
## [5,]   NA    0    1    0    1
## [6,]   NA    0    0    1    2
```

5. Create the following patterned matrices. Your solutions should be generalizable to enable creating larger matrices with the same structure.

$$\begin{bmatrix} 0 & 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 & 0 \\ 2 & 3 & 4 & 0 & 1 \\ 3 & 4 & 0 & 1 & 2 \\ 4 & 0 & 1 & 2 & 3 \end{bmatrix}$$

```
a <- outer(0:4, 0:4, '+')%5
```

a

```
##      [,1] [,2] [,3] [,4] [,5]
## [1,]    0    1    2    3    4
## [2,]    1    2    3    4    0
## [3,]    2    3    4    0    1
## [4,]    3    4    0    1    2
## [5,]    4    0    1    2    3
```

$$\begin{bmatrix} 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 8 & 9 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 9 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \end{bmatrix}$$

```
b <- outer(0:9,0:9,'+')%%10
```

```
b
```

```
##      [,1] [,2] [,3] [,4] [,5] [,6] [,7] [,8] [,9] [,10]
## [1,]  0    1    2    3    4    5    6    7    8    9
## [2,]  1    2    3    4    5    6    7    8    9    0
## [3,]  2    3    4    5    6    7    8    9    0    1
## [4,]  3    4    5    6    7    8    9    0    1    2
## [5,]  4    5    6    7    8    9    0    1    2    3
## [6,]  5    6    7    8    9    0    1    2    3    4
## [7,]  6    7    8    9    0    1    2    3    4    5
## [8,]  7    8    9    0    1    2    3    4    5    6
## [9,]  8    9    0    1    2    3    4    5    6    7
## [10,] 9    0    1    2    3    4    5    6    7    8
```

$$\begin{bmatrix} 0 & 8 & 7 & 6 & 5 & 4 & 3 & 2 & 1 \\ 1 & 0 & 8 & 7 & 6 & 5 & 4 & 3 & 2 \\ 2 & 1 & 0 & 8 & 7 & 6 & 5 & 4 & 3 \\ 3 & 2 & 1 & 0 & 8 & 7 & 6 & 5 & 4 \\ 4 & 3 & 2 & 1 & 0 & 8 & 7 & 6 & 5 \\ 5 & 4 & 3 & 2 & 1 & 0 & 8 & 7 & 6 \\ 6 & 5 & 4 & 3 & 2 & 1 & 0 & 8 & 7 \\ 7 & 6 & 5 & 4 & 3 & 2 & 1 & 0 & 8 \\ 8 & 7 & 6 & 5 & 4 & 3 & 2 & 1 & 0 \end{bmatrix}$$

```
c <- outer(0:8, 0:8, '-')%%9
```

```
c
```

```
##      [,1] [,2] [,3] [,4] [,5] [,6] [,7] [,8] [,9]
## [1,]  0    8    7    6    5    4    3    2    1
## [2,]  1    0    8    7    6    5    4    3    2
## [3,]  2    1    0    8    7    6    5    4    3
## [4,]  3    2    1    0    8    7    6    5    4
## [5,]  4    3    2    1    0    8    7    6    5
## [6,]  5    4    3    2    1    0    8    7    6
## [7,]  6    5    4    3    2    1    0    8    7
## [8,]  7    6    5    4    3    2    1    0    8
## [9,]  8    7    6    5    4    3    2    1    0
```

6. Solve the following system of linear equations by setting up and solving the matrix equation $Ax = y$.

$$x_1 + 2x_2 + 3x_3 + 4x_4 + 5x_5 = 7$$

$$2x_1 + x_2 + 2x_3 + 3x_4 + 4x_5 = -1$$

$$\begin{aligned}
3x_1 + 2x_2 + x_3 + 2x_4 + 3x_5 &= -3 \\
4x_1 + 3x_2 + 2x_3 + x_4 + 2x_5 &= 5 \\
5x_1 + 4x_2 + 3x_3 + 2x_4 + x_5 &= 17
\end{aligned}$$

```

y <- c(7, -1, -3, 5, 17)
mat <- matrix(0, nrow = 5, ncol = 5)
mat <- abs(col(mat) - row(mat)) + 1
solve(mat)%*%y

##      [,1]
## [1,]   -2
## [2,]    3
## [3,]    5
## [4,]    2
## [5,]   -4

```

7. Create a 6 x 10 matrix of random integers chosen from 1,2,...,10 by executing the following two lines of code:

```

set.seed(75)
aMat <- matrix(sample(10, size=60, replace=TRUE), nr=6)

```

```

set.seed(75)
aMat <- matrix(sample(10, size=60, replace=TRUE), nr=6)

```

Use the matrix you have created to answer these questions:

(a) Find the number of entries in each row which are greater than 4.

```

temp <- function(num) {
  sum(num > 4)
}
apply(aMat, 1, temp)

## [1] 4 7 6 2 6 7

```

(b) Which rows contain exactly two occurrences of the number seven?

```

temp2 <- function(num) {
  sum(num == 7) == 2
}
which(apply(aMat, 1, temp2))

## [1] 5

```

(c) Find those pairs of columns whose total (over both columns) is greater than 75. The answer should be a matrix with two columns; so, for example, the row (1,2) in the output matrix means that the sum of columns 1 and 2 in the original matrix is greater than 75. Repeating a column is permitted; so, for example, the final output matrix could contain the rows (1,2), (2,1), and (2,2).

```

pairs <- colSums(aMat)
cbind(rep(1:10, rep(10,10)), rep(1:10,10))[outer(pairs,pairs,'+') > 75,]

```

```
##      [,1] [,2]
## [1,]    2    2
## [2,]    2    6
## [3,]    2    8
## [4,]    6    2
## [5,]    6    8
## [6,]    8    2
## [7,]    8    6
## [8,]    8    8
```

What if repetitions are not permitted? Then only (1,2) from (1,2),(2,1) and (2,2) would be permitted.

```
pairs <- colSums(aMat)
x <- (outer(pairs,pairs,'+')>75)
x[lower.tri(x)] <- F
which(x,arr.ind = T)
```

```
##      row col
## [1,]    2    2
## [2,]    2    6
## [3,]    2    8
## [4,]    6    8
## [5,]    8    8
```

8. Calculate

(a) $\sum_{i=1}^{20} \sum_{j=1}^5 \frac{i^4}{(3+j)}$

```
sum((1:20)^4 * sum(1/(3+(1:5))))
```

```
## [1] 639215.3
```

or

```
sum(outer((1:20)^4, (3+(1:5)), "/"))
```

```
## [1] 639215.3
```

(b) $\sum_{i=1}^{20} \sum_{j=1}^5 \frac{i^4}{(3+ij)}$

```
sum((1:20)^4 / (3 + outer(1:20,1:5,'*')))
```

```
## [1] 89912.02
```

(c) $\sum_{i=1}^{10} \sum_{j=1}^i \frac{i^4}{(3+ij)}$

```
temp <- function(x, y) {
  (x >= y) * x^4 / (3 + x * y)
```

```
}  
sum(outer(1:10,1:10,temp))  
## [1] 6944.743
```