

Math 110BH Homework 6

Nathan Solomon

March 19, 2024

1

Let M be a (left) R -module generated by one element. Prove that M is isomorphic to the factor module R/I where I is a (left) ideal of R .

2

Let R be a commutative ring. Show that for every two R modules M and N , the group $\text{Hom}_R(M, N)$ has a structure of an R -module.

3

Let M be a (left) R -module, $N \subset M$ a submodule. Prove that if N and M/N are finitely generated, then so is M .

4

Prove that for any (left) R -module M , the groups $\text{Hom}_R(R, M)$ and M are isomorphic.

5

Let $f : R^n \rightarrow R^m$ be a ring R -module homomorphism. Show that there is a unique $m \times n$ matrix A such that $F(x) = A \cdot x$ for any $x \in R^n$.

6

Let R be a commutative ring and $I \subset R$ an ideal. Prove that if I is a free R -module, then I is a principal ideal.

7

Show that \mathbb{Q} is not a free abelian group (\mathbb{Z} -module).

Suppose it has a basis and then try to find a contradiction?

8

Prove that a free finitely generated (left) R -module has a finite basis.

9

Let M be a (left) R -module, $I \subset R$ an ideal. Denote by IM the submodule of M generated by the products am for all $a \in I$ and $m \in M$.

- (a) Assume that $IM = 0$. Show that M admits a structure of a (left) module over the factor ring R/I .
- (b) Show that M/IM admits a structure of a (left) module over the factor ring R/I .
- (c) Prove that if M is a free R -module, then M/IM is a free R/I -module. Hint: Show that if S is a basis for M then the set of cosets $\{s + IM, s \in S\}$ is a basis for M/IM .
- (d) Let R be a nonzero commutative ring. Prove that if (left) R -module R^n and R^m are isomorphic, then $n = m$. Deduce that every two bases for a free finitely generated R -module have the same number of elements. Hint: Consider modules over the factor ring R/I where I is a maximal ideal of R .

- (a)
- (b)
- (c)
- (d)

10

Let A be an abelian group, $f \in \text{End}(A)$. Show that A admits a $\mathbb{Z}[x]$ -module structure such that $x \cdot a = f(a)$ for all $a \in A$.