

M 431: Assignment 1

Nathan Stouffer

Page 7 — Problem 14

Problem. If C is a finite set, let $m(C)$ denote the number of elements in C . If A, B are finite sets, prove that

$$m(A \cup B) = m(A) + m(B) - m(A \cap B)$$

Proof. We take for granted the fact that $m(A \cup B) = m(A) + m(B)$ if A, B are disjoint sets. Now let's establish some notation. Let $A' = A \setminus (A \cap B)$ and $B' = B \setminus (A \cap B)$.

We now show a quick proof of the fact that $m(A \cap B)$ is finite. By definition of intersection, $A \cap B \subset A$. But then A cannot contain fewer elements than $A \cap B$, so we must have $m(A) \geq m(A \cap B)$. Then since A is finite, $A \cap B$ is also finite.

We also show that $A' \cap B' = \emptyset$. $A' = A \setminus (A \cap B)$ and $B' = B \setminus (A \cap B)$ so

$$\begin{aligned} A' \cap B' &= \{x \mid x \in A' \text{ and } x \in B'\} \\ &= \{x \mid x \in A \setminus (A \cap B) \text{ and } x \in B \setminus (A \cap B)\} \\ &= \{x \mid (x \in A \text{ and } x \notin A \cap B) \text{ and } (x \in B \text{ and } x \notin A \cap B)\} \\ &= \{x \mid x \in A \text{ and } x \in B \text{ and } x \notin A \cap B\} \\ &= \{x \mid x \in A \cap B \text{ and } x \notin A \cap B\} \\ &= \emptyset \end{aligned}$$

So $A' \cap B' = \emptyset$.

We now show, via induction, that given $k \in \{0, 1, 2, \dots, m(A \cap B)\}$, every subset $C \subset A \cap B$ such that $m(C) = k$ satisfies $m(A' \cup B' \cup C) = m(A' \cup C) + m(B' \cup C) - m(C)$. We will denote a subset of $A \cap B$ containing k elements as C_k . Consider the base case where $k = 0$. Then there is only one C_k and it must be $C_k = \emptyset$. So we must show that $m(A' \cup B' \cup \emptyset) = m(A' \cup \emptyset) + m(B' \cup \emptyset) - m(\emptyset)$. But then

$$\begin{aligned} m(A' \cup B' \cup \emptyset) &= m(A' \cup \emptyset) + m(B' \cup \emptyset) - m(\emptyset) \\ m(A' \cup B') &= m(A') + m(B') - 0 \\ m(A' \cup B') &= m(A') + m(B') \end{aligned}$$

which we already know to be true since A' and B' are disjoint sets. So the base case is proved.

Now suppose that for some $i \in \{0, 1, 2, \dots, m(A \cap B) - 1\}$, that we have $m(A' \cup B' \cup C_i) = m(A' \cup C_i) + m(B' \cup C_i) - m(C_i)$ for all $C_i \subset A \cap B$. We must now show that $m(A' \cup B' \cup C_{i+1}) = m(A' \cup C_{i+1}) + m(B' \cup C_{i+1}) - m(C_{i+1})$ for all $C_{i+1} \subset A \cap B$.

For every i , we C_{i+1} contains one more element than C_i so we must have $m(C_{i+1}) = m(C_i) + 1 \iff m(C_i) = m(C_{i+1}) - 1$. Furthermore, we constructed A', B' so that for any k , $A' \cap C_k = B' \cap C_k = (A' \cup B') \cap C_k = \emptyset$. Then we can use our fact about disjoint sets to say that

$$\begin{aligned} m(A' \cup B' \cup C_i) &= m(A' \cup C_i) + m(B' \cup C_i) - m(C_i) \\ m(A' \cup B') + m(C_i) &= m(A') + m(C_i) + m(B') + m(C_i) - m(C_i) \\ m(A' \cup B') + m(C_{i+1}) - 1 &= m(A') + m(C_{i+1}) - 1 + m(B') + m(C_{i+1}) - 1 - (m(C_{i+1}) - 1) \\ m(A' \cup B' \cup C_{i+1}) - 1 &= m(A' \cup C_{i+1}) - 1 + m(B' \cup C_{i+1}) - 1 - m(C_{i+1}) + 1 \end{aligned}$$

Then the ones cancel, leaving $m(A' \cup B' \cup C_{i+1}) = m(A' \cup C_{i+1}) + m(B' \cup C_{i+1}) - m(C_{i+1})$, which completes the induction.

So now we know that given $k \in \{0, 1, 2, \dots, m(A \cap B)\}$, every $C_k \subset A \cap B$ must satisfy $m(A' \cup B' \cup C_k) = m(A' \cup C_k) + m(B' \cup C_k) - m(C_k)$. Take $k = m(A \cap B)$, then there is only one C_k and it must be $C_k = A \cap B$. So we must have $m(A' \cup B' \cup (A \cap B)) = m(A' \cup (A \cap B)) + m(B' \cup (A \cap B)) - m(A \cap B)$. But $A' \cup B' \cup (A \cap B) = A \cup B$, $A' \cup (A \cap B) = A$, and $B' \cup (A \cap B) = B$, so we have $m(A \cup B) = m(A) + m(B) - m(A \cap B)$, which is what needed to be shown.

□

Page 7 — Problem 20

Problem. Show, for finite sets A, B , that $m(A \times B) = m(A)m(B)$.

Proof.

□

Page 13 — Problem 6

Problem. If $f : S \longrightarrow T$ is onto and $g : T \longrightarrow U$ and $h : T \longrightarrow U$ are such that $g \circ f = h \circ f$, then $g = h$.

Proof. To show that $g = h$, we must show that $g(t) = h(t)$ for all $t \in T$. Since $g \circ f = h \circ f$, we can say that $(g \circ f)(s) = g(f(s)) = (h \circ f)(s) = h(f(s))$ for all $s \in S$. Then since f is onto, for every $t \in T$, there exists $s_t \in S$ such that $f(s_t) = t$. Then $f(s)$ can be replaced with t to give $g(f(s)) = g(t) = h(f(s)) = h(t)$, which proves that $g = h$.

□

Page 20 — Problem 11

Problem. Can you find a positive integer m such that $f^m = i$ for all $f \in S_4$?

Proof.

□

Page 20 — Problem 13

Problem. Show that there is a positive integer t such that $f^t = i$ for all $f \in S_n$.

Proof.

□