

M 383: Assignment 3

Nathan Stouffer

Exercises 2.2.4 — Problem 3

Problem. If x is a real number, show that there exists a Cauchy sequence of rationals x_1, x_2, \dots representing x such that $x_n < x$ for all n .

Proof.

Exercises 2.2.4 — Problem 7

Problem. Prove that $|x - y| \geq |x| - |y|$ for any real numbers x and y .

Proof. We must show that $|x - y| \geq |x| - |y|$ holds for all real numbers x and y . We take $|x + y| \leq |x| + |y|$ (the triangle inequality) to be true. Then let x and y be any real numbers. Certainly it is true that $|x| = |x - y + y|$ since \mathbb{R} is an ordered field. Also, by the triangle inequality, $|x - y + y| \leq |x - y| + |y|$. Since $|x - y + y| = |x|$, we can say that $|x - y| + |y| \geq |x|$ which we can reorder to be $|x - y| \geq |x| - |y|$. So we have shown the desired inequality for all real numbers x and y .

Exercises 2.3.3 — Problem 1

Problem. Write out a proof that $\lim_{k \rightarrow \infty} (x_k + y_k) = x + y$ if $\lim_{k \rightarrow \infty} x_k = x$ and $\lim_{k \rightarrow \infty} y_k = y$ for sequences of real numbers.

Proof. We want to show (for sequences of real numbers) that the limit of a sum is the same as the sum of the limits. Given that $\lim_{k \rightarrow \infty} x_k = x$ and $\lim_{k \rightarrow \infty} y_k = y$, we want to show that $\{x_k + y_k\}$ satisfies the Cauchy criterion and converges to $x + y$.

Saying that $\{x_k\}$ and $\{y_k\}$ have limits is equivalent to saying $\{x_k\}$ and $\{y_k\}$ are Cauchy. We also know the sum of two Cauchy sequences to be Cauchy, so certainly $\{x_k + y_k\}$ is Cauchy.

But what is the limit of $\{x_k + y_k\}$? Let's begin by noting that $\lim_{k \rightarrow \infty} x_k = x \implies \forall n, \exists m_a$ such that $|x_a - x| \leq 1/2n$ for all $a \geq m_a$. Similarly $\lim_{k \rightarrow \infty} y_k = y \implies \forall n, \exists m_b$ such that $|y_b - y| \leq 1/2n$ for all $b \geq m_b$. Choose $k = \max\{a, b\}$. Then $|(x_k + y_k) - (x + y)| = |(x_k - x) + (y_k - y)| \leq 1/2n + 1/2n = 1/n$ which satisfies the definition of a limit. Thus the limit of a sum is equal the sum of the limits.

Exercises 2.3.3 — Problem 3

Problem. Let x_1, x_2, \dots be a sequence of real numbers such that $|x_n| \leq 1/2^n$, and set $y_n = x_1 + x_2 + \dots + x_n$. Show that the sequence y_1, y_2, \dots converges.

Proof. We must show that y_1, y_2, \dots converges. To this end, we introduce the following result: $1/2^n = \sum_{k=n+1}^{\infty} 1/2^k$. Let's now prove this result. First note that for some natural number a , $1/2^{a+1} = 1/2 * 1/2^a$. Then for any a , we have

$$1/2^a + 1/2^{a+1} + 1/2^{a+2} + \dots = 1/2^a + 1/2 * 1/2^a + 1/2 * 1/2^{a+1} + \dots = 1/2^a + 1/2 * (1/2^a + 1/2^{a+1} + \dots)$$

which implies that

$$1/2^a = 1/2 * (1/2^a + 1/2^{a+1} + \dots) = 1/2 * 1/2^a + 1/2 * 1/2^{a+1} + \dots = 1/2^{a+1} + 1/2^{a+2} + \dots = \sum_{k=a+1}^{\infty} 1/2^k$$

Now to show that y_1, y_2, \dots converges, we will show that y_1, y_2, \dots is Cauchy. So, given a natural number n , we must show the existence of an index m such that $|y_j - y_k| \leq 1/n$ for all $j, k \geq m$. For any index m , we can maximize $|y_j - y_k|$ by choosing the largest possible y_j and the smallest possible y_k . Because $1/2^n = \sum_{k=n+1}^{\infty} 1/2^k$ and $x_n \leq 1/2^n$, we can say that the largest y_j could be is $y_m + 1/2^m$ and the smallest y_k could be is $y_m - 1/2^m$. Then

$$|y_j - y_k| \leq |(y_m + 1/2^m) - (y_m - 1/2^m)| = |2/2^m| = 1/2^{m-1}$$

Choosing $m = n$, we can say that $1/2^{m-1} = 1/2^{n-1} \leq 1/n$ for all n . So there exists an index m such that $|y_j - y_k| \leq 1/n$ for all $j, k \geq m$. Thus y_1, y_2, \dots satisfies the Cauchy criterion and must also converge.