

Probability and Statistics by DeGroot Notes

Nathan Ueda

February 12, 2024

Contents

1	Introduction to Probability	3
1.1	The History of Probability	3
1.2	Interpretations of Probability	3
1.3	Experiments and Events	3
1.4	Set Theory	3
1.5	The Definition of Probability	4
1.6	Finite Sample Spaces	4
1.7	Counting Methods	4
1.8	Combinatorial Methods	5
1.9	Multinomial Coefficients	5
1.10	The Probability of a Union of Events	5
2	Conditional Probability	6
2.1	The Definition of Conditional Probability	6

1 Introduction to Probability

1.1 The History of Probability

1.2 Interpretations of Probability

Probability Interpretations

- Frequency: If an experiment is carried out many times, the frequency with which a particular outcome occurred would define its probability.
- Classical: If an outcome of some experiment must be one of n different, equally likely outcomes, the probability of each outcome is $\frac{1}{n}$.
- Subjective: An entity assigns probabilities to each possible outcome.

Probability theory does not depend on interpretation.

1.3 Experiments and Events

Probability allows us to quantify how likely an outcome is to occur.

Experiments: Any process in which the possible outcomes can be identified ahead of time.

Events: A well defined set of possible outcomes of the experiment.

Although there is controversy in regard to the proper meaning and interpretation of some of the probabilities that are assigned to the outcomes of many experiments, once these probabilities are assigned, there is complete agreement upon the mathematical theory of probability.

Almost all work in the mathematical theory of probability is related to:

- Methods for determining probabilities of certain events from given probabilities for each possible outcome in an experiment.
- Methods for revising probabilities of events when additional relevant information is obtained.

1.4 Set Theory

Sample Space: The collection of all possible outcomes of an experiment.

Empty Set: Subset of S containing no elements, denoted \emptyset , representing any events that cannot occur.

Complement: For some set A , its complement, denoted A^c , is the set containing all elements of S not in A .

Union: For n sets A_1, \dots, A_n , their union, denoted $A_1 \cup \dots \cup A_n$ or $\bigcup_{i=1}^n A_i$, is defined as the set containing all outcomes that belong to at least one of these n sets.

Intersection: For n sets A_1, \dots, A_n , their intersection, denoted $A_1 \cap \dots \cap A_n$ or $\bigcap_{i=1}^n A_i$, is defined as the set containing the elements common to all these n sets.

Disjoint/Mutually Exclusive: Two sets A and B are disjoint/mutually exclusive if they have no outcomes in common, that is, if $A \cap B = \emptyset$, representing that both A and B cannot occur.

1.5 The Definition of Probability

Axioms of Probability:

1. For every event A , $P(A) \geq 0$
2. $P(S) = 1$
3. $P(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$

Basic Theorems:

1. $P(\emptyset) = 0$
2. For every finite sequence of n disjoint events, A_1, \dots, A_n , $P(\bigcup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i)$
3. For every event A , $P(A^c) = 1 - P(A)$
4. If $A \subset B$, then $P(A) \leq P(B)$
5. For every event A , $0 \leq P(A) \leq 1$
6. For every two events A and B , $P(A \cap B^c) = P(A) - P(A \cap B)$
7. For every two events A and B , $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
8. Bonferroni Inequality: For all events A_1, \dots, A_n , $P(\bigcap_{i=1}^n A_i) \geq 1 - \sum_{i=1}^n P(A_i^c)$

1.6 Finite Sample Spaces

Simple Sample Space

- Has a finite number (n) of possible outcomes
- Each outcome has an equal probability ($\frac{1}{n}$)
- If an event A has m outcomes, then $P(A) = \frac{m}{n}$

1.7 Counting Methods

Multiplication Rule: An experiment with k parts where the i th part has n_i possible outcomes (regardless of which specific outcomes have occurred in the other parts) has a sample space $S = n_1 n_2 \dots n_k$

Permutations ($P_{n,k}$):

- Number of ways to arrange a set (order matters)
- Sampling considering n different items and making k choices from them
 - Sampling with replacement: n^k
 - Sampling without replacement: $n(n-1) \dots (n-k+1)$
 - * n options for first choice, $n-1$ options for second choice, $n-k+1$ options for k th choice

- The number of permutations of n different items is $P_{n,n} = n!$
- The number of permutations of n different items making k choices ($0 \leq k \leq n$) is

$$P_{n,k} = n(n-1) \dots (n-k+1)$$

$$P_{n,k} = n(n-1) \dots (n-k+1) \left(\frac{1}{1}\right)$$

$$P_{n,k} = n(n-1) \dots (n-k+1) \left(\frac{(n-k)(n-k-1) \dots 1}{(n-k)(n-k-1) \dots 1}\right)$$

$$P_{n,k} = \frac{n(n-1) \dots (n-k+1)(n-k)(n-k-1) \dots 1}{(n-k)(n-k-1) \dots 1}$$

$$P_{n,k} = \frac{n!}{(n-k)!}$$

1.8 Combinatorial Methods

Combinations ($C_{n,k}$):

- Number of subsets (order does not matter)
- Permutations may be thought of as combinations of size k chosen out of n , multiplied by the number of ways to arrange the size k subsets, $k!$. More formally, this says

$$P_{n,k} = C_{n,k} k!$$

- Combinations (binomial coefficient) are the number of distinct subsets of size k that can be chosen from a set of size n :

$$C_{n,k} = \binom{n}{k} = \frac{P_{n,k}}{k!} = \frac{\frac{n!}{(n-k)!}}{k!} = \frac{n!}{(n-k)!k!}$$

- Combinations without replacement: $\binom{n+k-1}{k}$

1.9 Multinomial Coefficients

The total number of different ways of dividing n elements into k groups is

$$\binom{n}{n_1, n_2, \dots, n_k} = \binom{n}{n_1} \binom{n-n_1}{n_2} \binom{n-n_1-n_2}{n_3} \dots \binom{n-n_1-\dots-n_{k-2}}{n_{k-1}} = \frac{n!}{n_1! n_2! \dots n_k!}$$

1.10 The Probability of a Union of Events

- Union of two events A_1 and A_2 :

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2)$$

- Union of three events A_1 , A_2 , and A_3 :

$$P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3) - [P(A_1 \cap A_2) + P(A_1 \cap A_3) + P(A_2 \cap A_3)] \\ + P(A_1 \cap A_2 \cap A_3)$$

- Union of n events A_1, \dots, A_n :

$$P\left(\bigcup_{i=1}^n A_i\right) = \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) - \sum_{i < j < k < l} P(A_i \cap A_j \cap A_k \cap A_l) + \dots + (-1)^{n+1} P(A_1 \cap A_2 \cap \dots \cap A_n)$$

2 Conditional Probability

2.1 The Definition of Conditional Probability

- Conditional probability is the updating of probabilities when certain events are observed
- The updated probability of event A after we learn that event B has occurred is the conditional probability of A given B , denoted $P(A|B)$
- When we go from $P(A)$ to $P(A|B)$, we say we are conditioning on B
- For $P(B) > 0$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

- Intuitively, this is saying the probability of A occurring, given B has occurred is equal to the outcomes where both A and B occurred (which makes sense since we know B has occurred and we want to find the probability of A also occurring) divided by the probability of B occurring (which makes sense since we know B occurred, we can renormalize the sample space to only contain outcomes where B occurred).

Multiplication Rule for Conditional Probabilities

- For 2 events A, B

– If $P(B) > 0$

$$P(A \cap B) = P(B)P(A|B)$$

– If $P(A) > 0$

$$P(A \cap B) = P(A)P(B|A)$$

- For n events A_1, \dots, A_n such that $P(A_1 \cap \dots \cap A_{n-1}) > 0$

$$P(A_1 \cap \dots \cap A_n) = P(A_1)P(A_2|A_1)P(A_3|A_1 \cap A_2) \dots P(A_n|A_1 \cap \dots \cap A_{n-1})$$

- For n events A_1, \dots, A_n, B such that $P(B) > 0$ and $P(A_1 \cap \dots \cap A_{n-1}) > 0$

$$P(A_1 \cap \dots \cap A_n|B) = P(A_1|B)P(A_2|A_1 \cap B)P(A_3|A_1 \cap A_2 \cap B) \dots P(A_n|A_1 \cap \dots \cap A_{n-1} \cap B)$$

Law of Total Probability

- Tells us that to get the unconditional probability of A , we can divide the sample space into disjoint slices B_j , find the conditional probability of A within each of these slices, then take a weighted sum of the conditional probabilities, where the weights are the probabilities $P(B_i)$

- Often used in tandem with Bayes' Rule
- Partition: Let S denote the sample space and consider k events B_1, \dots, B_k in S such that B_1, \dots, B_k are disjoint and $\bigcup_{i=1}^k B_i = S$. Then events B_1, \dots, B_k form a partition in S . In other words, only one of these events can occur and combined they fill the entire sample space
- Suppose events B_1, \dots, B_k form a partition of the space S and $P(B_j) > 0$ for $j = 1, \dots, k$. Then

$$P(A) = \sum_{j=1}^k P(B_j \cap A) = \sum_{j=1}^k P(B_j)P(A|B_j)$$

- Conditional LOTP

$$P(A|C) = \sum_{j=1}^k P(B_j|C)P(A|B_j \cap C)$$