


The image features a light beige background with a subtle, mottled texture. In the four corners, there are decorative geometric patterns. The top-left and bottom-right corners have patterns made of overlapping lines in teal, orange, and magenta. The top-right and bottom-left corners have patterns featuring yellow stars on a magenta and black background. The central focus is the Basmala in a highly stylized, calligraphic script. The word 'Bismillah' is written in magenta, 'ar-Rahman' in brown, and 'ar-Rahim' in orange and yellow. The script is fluid and elegant, with thick black outlines for the letters.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ
Bismillah al rahman al rahim


A scenic view of a university campus. A paved path leads through a row of cherry blossom trees in full bloom, their pink petals falling onto the path. To the left is a green metal fence with stone pillars. A person is walking in the distance on the path. The background shows more trees and a building under a clear sky.

دانشگاه علوم کشاورزی و منابع طبیعی گرگان
دانشکده صنایع غذایی

شیمی مواد غذایی ۱

WATER ACTIVITY (a_w)

- Water quantity report :
 - Wet – weight basis, Moisture content, M
 - Dry – weight basis, Moisture value, m



$$M = \frac{\text{water mass}}{\text{sample mass}} \times 100 = \frac{\text{water mass}}{\text{water mass} + \text{solid mass}} \times 100$$

$$m = \frac{\text{water mass}}{\text{solid mass}}$$

↓
“bone dry” or “oven dry” mass

$$M = \frac{100 m}{1 + m}$$

WATER ACTIVITY (a_w)


- **Practice :**

- 250 kg tomato, $M = 94\%$
 - Freeze drying \rightarrow 25 kg
 - $M = ?$
 - $m = ?$
-
- $0.94 \times 250 = 235$ kg water (15 kg dry matter)
 - $M = [(25-15)/25] \times 100 = 40\%$
 - $m = (25-15)/15 = 0.67$

MOISTURE SORPTION ISOTHERM (MSI)

- **Zone I**

- Most strongly sorbed and least mobile
- Accessible polar sites by water-ion or water-dipole interactions
- Unfreezable at -40°C
- has no ability to dissolve solutes
- No Plastisizing effect
- Boundary of zones I and II : BET monolayer water (Brunauer, Emmett, Teller)
- $< 1\%$ in high moisture food




MOISTURE SORPTION ISOTHERM (MSI)

- **Zone II**

- Slightly less mobile than bulk water
- Incipient swelling of the solid matrix
- significant plasticizing action on solutes
- Slows their glass transition temperatures
- < 5% in high moisture food
- Boundary of zones II and III : full hydration (a true monolayer hydration shell)

- **Zone III**


- Bulk-phase water
- Physically entrapped in gels and cells
- > 95% in high moisture foods



MOISTURE SORPTION ISOTHERM (MSI)

- The types of water in food

- BET monolayer water
 - Constitutional water ($<0.03\%$)
 - Vicinal water ($0.5\pm0.4\%$)
 - A monolayer coverage
- True monolayer water ($3\pm2\%$)
 - A multilayer coverage
- Bulk-phase water ($\sim 95\%$)
 - Entrapped water
 - Free water



Types of water in food: some properties

All foods contain at least some water

- *Free water*: can be extracted easily from foods by **squeezing, cutting or pressing**.
 - free flowing
 - great solvent for food components
 - can be removed by pressure
- *Bound water*: cannot be extracted easily.
 - It is not free to act as a solvent for salts and sugars
 - It can be frozen only at very low temperatures (below the freezing point of water).
 - It exhibits essentially no vapor pressure
 - Its density is greater than that of free water

An example: water present in cacti



Types of bound water:


All water found in tissue is bound water

Structural water:


- Small portion of total bound water
- Found **inside protein macromolecule**
- Not available for chemical reactions.
- **Bound directly to protein molecule** by hydrogen bonding
- Stabilize the native conformation of the protein molecule; determine the three-dimensional conformation of the protein.
- Structural water molecules may act as prosthetic groups indispensable for proper protein function

prosthetic group is a tightly bound, specific non-polypeptide unit required for the biological function of some proteins..

- Participate in the formation of enzyme active site
(http://www1.lsbu.ac.uk/water/protein_hydration.html)



Monolayer water:




- Bound to the surface of the protein molecules by **hydrogen bonding or dipole interactions**.
- It represents 4-9% of the water associated with the protein.
- It has kinetic and thermodynamic properties which are different from that of the pure water.
- Not available as solvent.
- May be available for certain reactions.
- Hard to remove from food.

Multilayer Water:

- Additional layer of water around food particle.
- Not as hard to remove as the monolayer.

Sorption Isotherm of Water

- **Hysteresis**
- Resorption or adsorption
- Desorption
- Superimposability
- Hysteresis loop




- ▶ **Relative humidity** can be simply defined as the ratio of the actual amount of water in the air compared to how much the air can hold at a specific temperature
- ▶ When the water content of food and relative humidity of the environment are not equal, the food gains or loses water.
- ▶ **Desorption:** Lowering water content of moist foods to reach equilibrium with its surroundings
- ▶ **Adsorption:** Increasing water content of dry foods to reach equilibrium with its surroundings
- ▶ The water content of food and environment are same at equilibrium stage. After the equilibrium is reached, the water content of food does not change.

MOISTURE SORPTION ISOTHERM (MSI)

- **Hysteresis**

- Ink bottle theory



Capillary rise of a liquid that wets the walls of a tube.

$$\Delta P = -\frac{2\gamma}{r} \cos \theta \quad \text{Laplace equation}$$

$$\Delta P = \Delta \rho \cdot g \cdot h \quad \ln a_w = \ln \frac{P}{P_0} = -\frac{2\gamma}{r} \cos \theta \cdot \frac{V_m}{RT}$$

$r \text{ (}\mu\text{m)}$	a_w
0.0005	0.116
0.001	0.34
0.002	0.583
0.005	0.806
0.01	0.898
0.02	0.948
0.05	0.979
0.1	0.989
1.0	0.999
10	0.9999
100	0.99999

Kelvin equation

MOISTURE SORPTION ISOTHERM (MSI)



- **Hysteresis**


- Ink bottle theory
 - Resorption or adsorption

$$\ln a_w = -\frac{2\gamma}{r_2} \cos \theta \cdot \frac{V_m}{RT}$$

- Desorption

$$\ln a_w = -\frac{2\gamma}{r_1} \cos \theta \cdot \frac{V_m}{RT}$$





WATER ACTIVITY AND FOOD STABILITY

- Microbial growth
- Chemical reactions
- Food texture
- Food flavor
- IMF : intermediate moisture foods

Relative Vapor Pressure and Growth of Microorganisms in Food

Range of p/p_0	Microorganisms generally inhibited by lowest p/p_0 in this range	Foods generally within this range
1.00–0.95	<i>Pseudomonas</i> , <i>Escherichia</i> , <i>Proteus</i> , <i>Shigella</i> , <i>Klebsiella</i> , <i>Bacillus</i> , <i>Clostridium perfringens</i> , some yeasts	Highly perishable (fresh) foods and canned fruits, vegetables, meat, fish, and milk; cooked sausages and breads; foods containing up to approximately 40% (w/w) sucrose or 7% sodium chloride
0.95–0.91	<i>Salmonella</i> , <i>Vibrio parahaemolyticus</i> , <i>C. botulinum</i> , <i>Serratia</i> , <i>Lactobacillus</i> , <i>Pediococcus</i> , some molds, yeasts (<i>Rhodotorula</i> , <i>Pichia</i>)	Some cheeses (Cheddar, Swiss, Muenster, Provolone), cured meat (ham), some fruit juice concentrates; foods containing up to 55% (w/w) sucrose or 12% sodium chloride
0.91–0.87	Many yeasts (<i>Candida</i> , <i>Torulopsis</i> , <i>Hansenula</i>), <i>Micrococcus</i>	Fermented sausage (salami), sponge cakes, dry cheeses, margarine; foods containing up to 65% (w/w) sucrose (saturated) or 15% sodium chloride
0.87–0.80	Most molds (mycotoxigenic penicillia), <i>Staphylococcus aureus</i> , most <i>Saccharomyces (bailii)</i> spp., <i>Debaryomyces</i>	Most fruit juice concentrates, sweetened condensed milk, chocolate syrup, maple and fruit syrups; flour, rice, pulses containing 15–17% moisture; fruit cake; country-style ham, fondants, high-ratio cakes
0.80–0.75	Most halophilic bacteria, mycotoxigenic aspergilli	Jam, marmalade, marzipan, glacé fruits, some marshmallows
0.75–0.65	Xerophilic molds (<i>Aspergillus chevalieri</i> , <i>A. candidus</i> , <i>Wallemia sebi</i>), <i>Saccharomyces bisporus</i>	Rolled oats containing approximately 10% moisture; grained nougats, fudge, marshmallows, jelly, molasses, raw cane sugar, some dried fruits, nuts
0.65–0.60	Osmophilic yeasts (<i>Saccharomyces rouxii</i>), few molds (<i>Aspergillus echinulatus</i> , <i>Monascus bisporus</i>)	Dried fruits containing 15–20% moisture; some toffees and caramels; honey
0.50	No microbial proliferation	Pasta containing approximately 12% moisture; spices containing approximately 10% moisture
0.40	No microbial proliferation	Whole egg powder containing approximately 5% moisture
0.30	No microbial proliferation	Cookies, crackers, bread crusts, etc. containing 3–5% moisture
0.20	No microbial proliferation	Whole milk powder containing 2–3% moisture; dried vegetables containing approximately 5% moisture; corn flakes containing approximately 5% moisture; country style cookies, crackers

Table 2 Water Activity and Growth of Microorganisms in Food*

Range of a_w	Microorganisms Generally Inhibited by Lowest a_w in This Range	Foods Generally within This Range
1.00 – 0.95	<i>Pseudomonas</i> , <i>Escherichia</i> , <i>Proteus</i> , <i>Shigella</i> , <i>Klebsiella</i> , <i>Bacillus</i> , <i>Clostridium perfringens</i> , some yeasts	Highly perishable (fresh) foods and canned fruits, vegetables, meat, fish, and milk
0.95 – 0.91	<i>Salmonella</i> , <i>Vibrio parahaemolyticus</i> , <i>C. botulinum</i> , <i>Serratia</i> , <i>Lactobacillus</i> , <i>Pediococcus</i> , some molds, yeasts (<i>Rhodotorula</i> , <i>Pichia</i>)	Some cheeses (Cheddar, Swiss, Muenster, Provolone), cured meat (ham)
0.91 – 0.87	Many yeasts (<i>Candida</i> , <i>Torulopsis</i> , <i>Hansenula</i>), <i>Micrococcus</i>	Fermented sausage (salami), sponge cakes, dry cheeses, margarine
0.87 – 0.80	Most molds (mycotoxigenic penicillia), <i>Staphylococcus aureus</i> , most <i>Saccharomyces</i> (<i>bailii</i>) spp., <i>Debaryomyces</i>	Most fruit juice concentrates, sweetened condensed milk, syrups
0.80 – 0.75	Most halophilic bacteria, mycotoxigenic aspergilli	Jam, marmalade, marzipan, glacé fruits
0.75 – 0.65	Xerophilic molds (<i>Aspergillus chevalieri</i> , <i>A. candidus</i> , <i>Wallemia sebi</i>), <i>Saccharomyces bisporus</i>	Jelly, molasses, raw cane sugar, some dried fruits, nuts
0.65 – 0.60	Osmophilic yeasts (<i>Saccharomyces rouxii</i>), few molds (<i>Aspergillus echinulatus</i> , <i>Monascus bisporus</i>)	Dried fruits containing 15-20% moisture; some toffees and caramels; honey
0.60 – 0.50	No microbial proliferation	Dry pasta, spices
0.50 - 0.40	No microbial proliferation	Whole egg powder
0.40 - 0.30	No microbial proliferation	Cookies, crackers, bread crusts
0.30 - 0.20	No microbial proliferation	Whole milk powder; dried vegetables

* Adapted from Beuchat (1981).




Figure 1. Water Activity – Stability Map (adapted from Labuza, (1970))