

Turbulence

J. Nättälä¹

Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden. e-mail: joonas.nattila@su.se

Received XXX / Accepted XXX

ABSTRACT

Blaa blaa.

Key words. Plasmas – Turbulence – Methods: numerical

1. Theory

(Sobacchi & Lyubarsky 2019) paragraphs:

The most important property of MHD turbulence is its strong anisotropy, with the turbulent eddies becoming strongly elongated in the direction of the background magnetic field at small scales. As first proposed by Goldreich & Sridhar (1995), in the course of the turbulent cascade the ratio of the longitudinal scale of the eddies, λ_{\parallel} , to the Alfvén velocity remains equal to the ratio of the, perpendicular scale, λ_{\perp} , to the turbulent velocity. This condition is known as critical balance. From this condition one finds

$$\frac{\lambda_{\perp}}{\lambda_{\parallel}} \sim \sqrt{\frac{\lambda_{\parallel}}{R}} \quad (1)$$

The statement that in collisionless plasmas the anisotropic MHD turbulence decays by heating/accelerating particles along the background magnetic field is general. Indeed, the dissipation occurs at the wave-particle resonances

$$\omega - \mathbf{k} \cdot \mathbf{v} = n\omega_B \quad (2)$$

where ω and \mathbf{k} are the frequency and the wavenumber, \mathbf{v} is the particle velocity, ω_B is the Larmor frequency, and n is an integer. The cyclotron resonance condition $n \neq 0$ is satisfied when the longitudinal scale of the wave packet, λ_{\parallel} is of the order of the particle Larmor radius, $r_L = c/\omega_B$. Since $r_L \ll R$, due to the strong anisotropy of the MHD turbulence a particle crosses many wave packets during one Larmor orbit. It was first noticed by Gruzinov (1998), Quataert (1998), and Quataert & Gruzinov (1999) that in this case the dissipation occurs at the Landau resonance $n = 0$. Since the two physical mechanisms of wave-particle interaction at the $n = 0$ resonance are due to i) the longitudinal electric field of the wave, and ii) the interaction between the effective particle's magnetic moment and the longitudinal magnetic perturbation, one is led to the conclusion that the turbulent energy is primarily dissipated on to the longitudinal particle motion.

It has also been found that the turbulent fluctuations tend to align with one another forming small-scale current sheets (e.g. Beresnyak & Lazarian 2006; Boldyrev

2006; Mason, Cattaneo & Boldyrev 2006), which could be disrupted via magnetic reconnection thus providing an additional dissipation mechanism (e.g. Boldyrev & Loureiro 2017; Loureiro & Boldyrev 2017; Mallet, Schekochihin & Chandran 2017a,b). Note that the background magnetic field, which is much larger than the reconnecting field and lies in the same plane of the current sheet, plays the role of a guide field. Since the magnetic energy is transferred to the plasma particles at the Landau resonance between the particles and the tearing mode that disrupts the current sheet, also in this case one would expect the particles to be heated in the longitudinal direction.

Even if the perpendicular heating is negligible, in a weakly magnetized plasma the fire-hose instability quickly erases any momentum anisotropy (e.g. Parker 1958; Lerche 1966). However, since the fire-hose instability develops once $P_{\parallel} - P_{\perp} > B^2/4\pi$, one immediately sees that this instability is not effective if the magnetic-to-plasma energy ratio exceeds $\frac{1}{2}$.

2. Literature

2.1. Non-thermal particles from turbulence

Melrose 1980 Petrosian 2012 Lazarian 2012

2.2. Turbulence in astrophysics

turbulence in stellar coronae Matthaeus 1999, Cranmer 2007

ISM Armstrong 1995, Lithwick & Goldreich 2001
SNRs Weiler & Sramek 1988, Roy 2009
PWN Porth 2014, Lyutikov 2019
BH disks Balbus & Hawley 1998, Brandenburg & Subramanian 2005
jets from AGNs Marscher 2008, MacDonald & Marscher 2018
radio lobes Vogt & Ensslin 2005, O'Sullivan 2009
GRBs Piran 2004, Kumar & Narayan 2009
Galaxy clusters Zweibel & Heiles 1997, Subramanian 2006

2.3. Magnetically dominated turbulence

Sustained relativistic turbulence (force-free) (Thompson & Blaes 1998): extension of Goldreich & Sridhar 1995 to extreme relativistic limit (no plasma inertia; force-free MHD). Anisotropic cascade is formed, dissipation occurs at the scale of current starvation (when not enough charge carriers in plasma to maintain currents from Alfvén waves).

(Cho 2005) Inoue 2011 (Cho & Lazarian 2014) (Zrake & East 2016)

Relativistic MHD (Zrake & MacFadyen 2012) (Zrake 2014)

2.4. Bright non-thermal synchrotron and inverse Compton signatures

pulsar magnetospheres and winds Buhler & Blandford 2014
jets from AGNs Begelman 1984
coronae of accretion disks Yuan & Narayan 2014

2.5. Kinetic turbulence

Kinetic turbulence: (Zhdankin et al. 2017b) Letter
(Zhdankin et al. 2017a) Paper
(Zhdankin et al. 2018) System size convergence
(Zhdankin et al. 2019a) electron-proton plasma
(Zhdankin et al. 2019b) radiative turbulence
(Comisso & Sironi 2018) acceleration
(Wong et al. 2019) acceleration
(Nättilä 2019) Runko and turbulence
(Comisso & Sironi 2019) acceleration

2.6. Radiative turbulence

Analytic work on radiative turbulence (Uzdensky 2018)
(Zrake et al. 2018); GRBs (Sobacchi & Lyubarsky 2019)

PIC simulations: (Zhdankin et al. 2019b) radiative turbulence

Fokker-Planck equation in momentum space with radiative cooling term (Schlickeiser 1984, 1985); not in original list (Schlickeiser 1989) (Stawarz & Petrosian 2008)

2.7. radiative PIC simulations

Reconnection: (Jaroschek & Hoshino 2009) (Cerutti et al. 2013, 2014b,a) (Kagan et al. 2016b,a) (Hakobyan et al. 2019) (Werner et al. 2019) (Schoeffler et al. 2019)

decay of magnetostatic equilibria (Yuan et al. 2016) (Nalewajko et al. 2018)

pulsar wind (Cerutti & Philippov 2017)
pulsar magnetospheres (Cerutti et al. 2016) (Philippov & Spitkovsky 2018)

Synchrotron and jitter radiative signatures of collisionless shocks (Medvedev & Spitkovsky 2009) (Sironi & Spitkovsky 2009) (Kirk & Reville 2010) (Nishikawa et al. 2011)

Radiative turbulence (Zhdankin et al. 2019c)

Cerutti, B., Philippov, A. A., & Spitkovsky, A. 2016, MNRAS, 457, 2401
Cerutti, B., Werner, G. R., Uzdensky, D. A., & Begelman, M. C. 2013, ApJ, 770, 147
Cerutti, B., Werner, G. R., Uzdensky, D. A., & Begelman, M. C. 2014a, Physics of Plasmas, 21, 056501
Cerutti, B., Werner, G. R., Uzdensky, D. A., & Begelman, M. C. 2014b, ApJ, 782, 104
Cho, J. 2005, ApJ, 621, 324
Cho, J. & Lazarian, A. 2014, ApJ, 780, 30
Comisso, L. & Sironi, L. 2018, Physical Review Letters, 121
Comisso, L. & Sironi, L. 2019, arXiv e-prints, arXiv:1909.01420
Hakobyan, H., Philippov, A., & Spitkovsky, A. 2019, ApJ, 877, 53
Jaroschek, C. H. & Hoshino, M. 2009, Phys. Rev. Lett., 103, 075002
Kagan, D., Nakar, E., & Piran, T. 2016a, ApJ, 826, 221
Kagan, D., Nakar, E., & Piran, T. 2016b, ApJ, 833, 155
Kirk, J. G. & Reville, B. 2010, ApJ, 710, L16
Medvedev, M. V. & Spitkovsky, A. 2009, ApJ, 700, 956
Nalewajko, K., Yuan, Y., & Chruslińska, M. 2018, Journal of Plasma Physics, 84, 755840301
Nättilä, J. 2019, arXiv e-prints
Nishikawa, K. I., Niemiec, J., Medvedev, M., et al. 2011, Advances in Space Research, 47, 1434
Philippov, A. A. & Spitkovsky, A. 2018, ApJ, 855, 94
Schlickeiser, R. 1984, A&A, 136, 227
Schlickeiser, R. 1985, A&A, 143, 431
Schlickeiser, R. 1989, ApJ, 336, 243
Schoeffler, K. M., Grismayer, T., Uzdensky, D., Fonseca, R. A., & Silva, L. O. 2019, ApJ, 870, 49
Sironi, L. & Spitkovsky, A. 2009, ApJ, 707, L92
Sobacchi, E. & Lyubarsky, Y. E. 2019, MNRAS, 484, 1192
Stawarz, L. & Petrosian, V. 2008, ApJ, 681, 1725
Thompson, C. & Blaes, O. 1998, Phys. Rev. D, 57, 3219
Uzdensky, D. A. 2018, Monthly Notices of the Royal Astronomical Society, 477, 2849
Werner, G. R., Philippov, A. A., & Uzdensky, D. A. 2019, MNRAS, 482, L60
Wong, K., Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2019
Yuan, Y., Nalewajko, K., Zrake, J., East, W. E., & Blandford, R. D. 2016, ApJ, 828, 92
Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2017a, Monthly Notices of the Royal Astronomical Society, 474, 2514
Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2018, The Astrophysical Journal, 867, L18
Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2019a, Phys. Rev. Lett., 122, 055101
Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2019b, arXiv e-prints, arXiv:1908.08032
Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2019c, arXiv e-prints, arXiv:1908.08032
Zhdankin, V., Werner, G. R., Uzdensky, D. A., & Begelman, M. C. 2017b, Physical Review Letters, 118
Zrake, J. 2014, ApJ, 794, L26
Zrake, J., Beloborodov, A. M., & Lundman, C. 2018
Zrake, J. & East, W. E. 2016, ApJ, 817, 89
Zrake, J. & MacFadyen, A. I. 2012, ApJ, 744, 32

References

Cerutti, B. & Philippov, A. A. 2017, A&A, 607, A134