

Journal  
of the Fisheries  
Research  
Board  
of Canada

Journal  
de l'office  
des recherches  
sur les pêcheries  
du Canada

Volume 34, No. 3, March 1977

Volume 34, n° 3, mars 1977

---

**Statistical Assessment of the Age–Length Key**

DANIEL K. KIMURA

*Washington Department of Fisheries, Olympia, Wash. 98504, USA*

KIMURA, D. K. 1977. Statistical assessment of the age–length key. *J. Fish. Res. Board Can.* 34: 317–324.

Since 1934 when Fridriksson originated the age–length key, it has been widely used by fisheries biologists to estimate age distributions of populations. In recent years, there has been a general recognition that often the key has little value, or even worse, gives biased results. The analysis presented here indicates why the age–length key is so susceptible to bias. More importantly, a criterion is presented for determining whether the age–length key should be used in a particular situation. If the key is to be used, results from examples indicate that random age subsamples (i.e. the number of specimens aged from each length category proportional to the number in each length category) are superior to fixed age subsamples (i.e. a constant number of specimens aged from each length category). Generally, small increases in the age sample will likely increase the accuracy of an age–distribution determination more effectively than relatively large increases in the length sample.

KIMURA, D. K. 1977. Statistical assessment of the age–length key. *J. Fish. Res. Board Can.* 34: 317–324.

Introduite en 1934 par Fridriksson, la clé âge–longueur a été depuis largement utilisée par les biologistes des pêches pour estimer la distribution des âges dans une population. Depuis quelques années toutefois, on admet généralement que la clé a souvent peu de valeur ou, ce qui est pire, donne des résultats biaisés. L'analyse que nous présentons ici explique pourquoi la clé âge–longueur est si susceptible de biais. Ce qui est plus important, nous donnons un critère permettant de décider si la clé âge–longueur doit être utilisée dans une situation particulière. Dans les cas où la clé doit être utilisée, des exemples indiquent que les sous-échantillons d'âges pris au hasard (i.e. le nombre de spécimens dont l'âge est déterminé dans chaque catégorie de longueur étant proportionnel au nombre de poissons dans chaque catégorie de longueur) sont supérieurs à des sous-échantillons d'âges fixes (i.e. un nombre uniforme de spécimens dont l'âge est déterminé dans chaque catégorie de longueur). Généralement, des petites augmentations dans l'échantillon d'âges résulteront probablement en une détermination plus précise de la répartition des âges que ne le feraient des augmentations relativement grandes de l'échantillon de longueurs.

Received September 7, 1976  
Accepted December 2, 1976

Reçu le 7 septembre 1976  
Accepté le 2 décembre 1976

Printed in Canada (J4544)  
Imprimé au Canada (J4544)

ONE of the fundamental problems of fisheries biologists is to estimate the age distribution of a population. Sampling for ages may be accomplished by either taking a simple random sample, or by using Fridriksson's (1934) age-length key. The biologist must decide if the gains made by using the age-length key justify its use. There appears to be little in the literature to help him make this decision. In this paper, a criterion is presented for determining whether the key should be used in a particular situation.

If the age-length key is to be used, the biologist must decide whether the age subsample should be random (i.e. the number of specimens aged from each length category proportional to the number in each length category) or fixed (i.e. a constant number of specimens aged from each length category). In the literature, one can find proponents of random and fixed age subsamples. Ketchen (1949), in an early paper, felt that fixed age subsamples were superior. Southward (1976) found that random age subsamples were more efficient. The analyses and examples presented in the following sections show that from a statistical point of view random age subsamples are more efficient.

### Age-Length Key

Suppose the proportions of a population of age  $i = 1, \dots, n_a$  are  $p_1, \dots, p_{n_a}$ . Suppose also that the distribution of lengths at each age is given by  $q_{ij}$ ,  $j = 1, \dots, n_l$ .<sup>1</sup> Here,

$$\sum_{j=1}^{n_l} q_{ij} = 1$$

for all  $i$ . The proportion of length  $j$  is then

$$l_j = \sum_{i=1}^{n_a} p_i q_{ij}.$$

Of the  $j$  length fish, the proportion of fish of age  $i$  is  $q_{ij}' = (p_i q_{ij})/l_j$ . Therefore,

$$\sum_{i=1}^{n_a} q_{ij}' = 1$$

for all  $j$ . The quantities directly estimable from sample data are  $l_j$  and  $q_{ij}'$ . If a random sample of  $N_l$  length frequencies is taken and  $n_j$  are of length  $j$ , and if  $n_j^*$  of the  $n_j$  fish are aged, of which  $m_{ij}$  are of age  $i$ , then  $\hat{l}_j = n_j/N_l$  and  $\hat{q}_{ij}' = m_{ij}/n_j^*$  are unbiased estimates of  $l_j$  and  $q_{ij}'$ . Since

$$p_i = \sum_{j=1}^{n_l} l_j q_{ij}',$$

it follows that

$$\hat{p}_i = \sum_{j=1}^{n_l} \hat{l}_j \hat{q}_{ij}'$$

provides an unbiased estimate of  $p_i$ . This is the usual application of the age-length key. It should be noted that  $q_{ij}'$  is dependent on the  $p_i$ 's. This means that the age-length key will give biased results if applied to a population where the age composition differs from that of the population from which the age-length key was drawn.

The value of this key can be assessed directly by calculating the variance of the estimates of the various  $p_i$ 's. Any gains from increasing the length-frequency sample should be reflected in the variance of the  $\hat{p}_i$ 's. Since we are concerned with the overall variance of the  $\hat{p}_i$ 's, the total variance

$$\text{Vartot} = E \sum_{i=1}^{n_a} (\hat{p}_i - p_i)^2 = \sum_{i=1}^{n_a} \text{var}(\hat{p}_i)$$

<sup>1</sup>It should be noted that  $n_a$  and  $n_l$  actually refer to the number of categories. The ages and lengths are assumed to be coded so that the smallest values are one.

seems an appropriate error index. Using this error index, the efficiency of the age-length key may be assessed, as compared with increasing the age sample. Also, it may be determined whether the age subsample should be random or fixed.

#### DERIVATION OF THE VARIANCE OF THE $\hat{p}_i$ 's

The variance for the  $\hat{p}_i$ 's estimated from the age-length key may be derived by straightforward methods. However, for simplification, the following assumptions are made concerning  $n_j^*$ , the subsample from length category  $j$  which is aged. For the case of fixed age subsample sizes, the variance is conditioned on  $n_j^* = n_s/n_l$ , where  $n_s$  is the total sample to be aged and  $n_l$  is the number of length categories. The weakness of this approach is that  $n_j$ , the number of specimens of length  $j$ , may be actually less than  $n_j^*$ . When  $N_l$  is large compared with  $n_s$ , this should not be a problem. For the case of random age subsamples, it is assumed that  $n_j^* = sn_j$ , where  $s = n_s/N_l$ . Hence, this variance is not conditioned on  $n_j^*$ .

The variance of  $\hat{p}_i$ ,

$$\begin{aligned}\text{Var}(\hat{p}_i) &= E(\hat{p}_i - p_i)^2 \\ &= E \sum_{j=1}^{n_l} (\hat{l}_j \hat{q}'_{ij} - l_j q'_{ij}) \sum_{j'=1}^{n_l} (\hat{l}_{j'} \hat{q}'_{ij'} - l_{j'} q'_{ij'}) \\ &= \sum_{j=1}^{n_l} \sum_{j'=1}^{n_l} E(\hat{l}_j \hat{q}'_{ij} \hat{l}_{j'} \hat{q}'_{ij'} - l_j q'_{ij} l_{j'} q'_{ij'})\end{aligned}\quad (1)$$

The expectations can most easily be evaluated by breaking them into the cases  $j = j'$  and  $j \neq j'$ . When  $j = j'$  the expectation gives a different result under fixed and random subsampling.

Case 1a.  $j = j'$  subsample fixed

$$\begin{aligned}E(\hat{l}_j \hat{q}'_{ij} \hat{l}_j \hat{q}'_{ij}) &= E\left(\frac{n_j}{N_l} \frac{m_{ij}}{n_j^*} \frac{n_j}{N_l} \frac{m_{ij}}{n_j^*}\right) \\ &= E_{n_j} \frac{n_j}{N_l} \frac{n_j}{N_l} \left(\frac{q'_{ij}(1 - q'_{ij})}{n_j^*} + q'^2_{ij}\right) \\ &= \left[\frac{l_j(1 - l_j)}{N_l} + l_j^2\right] \left[\frac{q'_{ij}(1 - q'_{ij})}{n_j^*} + q'^2_{ij}\right]\end{aligned}$$

Case 1b.  $j = j'$  subsample random

$$\begin{aligned}E(\hat{l}_j \hat{q}'_{ij} \hat{l}_j \hat{q}'_{ij}) &= E\left(\frac{n_j}{N_l} \frac{m_{ij}}{sn_j} \frac{n_j}{N_l} \frac{m_{ij}}{sn_j}\right) \\ &= E_{n_j} \frac{n_j}{N_l} \frac{n_j}{N_l} \left[\frac{q'_{ij}(1 - q'_{ij})}{sn_j} + q'^2_{ij}\right] \\ &= \frac{l_j q'_{ij}(1 - q'_{ij})}{n_s} + q'^2_{ij} \left[\frac{l_j(1 - l_j)}{N_l} + l_j^2\right]\end{aligned}$$

Case 2.  $j \neq j'$  subsample fixed or random

$$\begin{aligned}E(\hat{l}_j \hat{q}'_{ij} \hat{l}_{j'} \hat{q}'_{ij'}) &= E\left(\frac{n_j}{N_l} \frac{m_{ij}}{n_j^*} \frac{n_{j'}}{N_l} \frac{m_{ij'}}{n_{j'}^*}\right) \\ &= E_{n_j, n_{j'}} \left[\frac{n_j}{N_l} \frac{n_{j'}}{N_l} q'_{ij} q'_{ij'}\right] \\ &= q'_{ij} q'_{ij'} \left[\frac{-l_j l_{j'}}{N_l} + l_j l_{j'}\right]\end{aligned}$$

Substituting the appropriate expectations back into (1), and simplifying, yields the following variance formulas:

$$\text{a) Var } (\hat{p}_i) = \frac{p_i(1-p_i)}{n_s} \quad \text{(random sample)} \\ \text{ages only}$$

$$\text{b) Var } (\hat{p}_i) = \sum_{j=1}^{n_l} \left[ \frac{l_j(1-l_j)}{N_l} \frac{q'_{ij}(1-q'_{ij})}{n_j^*} + \frac{l_j^2 q'_{ij}(1-q'_{ij})}{n_j^*} + \frac{l_j q'^2_{ij}}{N_l} \right] - \frac{p_i^2}{N_l} \\ \text{age subsample fixed}$$

$$\text{c) Var } (\hat{p}_i) = \sum_{j=1}^{n_l} \left[ \frac{l_j q'_{ij}(1-q'_{ij})}{n_s} + \frac{l_j q'^2_{ij}}{N_l} \right] - \frac{p_i^2}{N_l} \\ \text{age subsample random}$$

Note that  $\text{Var } (\hat{p}_i)$  and  $\text{Vartot} = \sum_{i=1}^{n_a} \text{Var } (\hat{p}_i)$  have the following forms:

$$\text{a) Vartot} = \frac{a_1}{n_s} \\ \text{ages only}$$

$$\text{b) Vartot} = \frac{b_1}{N_l n_j^*} + \frac{b_2}{n_j^*} + \frac{b_3}{N_l} \\ \text{age subsample fixed}$$

$$\text{c) Vartot} = \frac{c_1}{n_s} + \frac{c_2}{N_l} \\ \text{age subsample random}$$

The form of these Vartot expressions allow us to easily calculate the asymptotic value of Vartot as  $N_l$  goes to infinity. This value of Vartot would be that given by an age-length key when an infinite length sample was taken. We can then calculate the asymptotic percentage reduction in Vartot (ages only), which would then represent an upper bound to the value of using the key. An important property of this percentage reduction is that it is independent of the actual value of  $n_s$ . In other words, the gain in using the age-length key is the same when  $n_s = 100$ , or  $n_s = 1000$ , when measured in this way.

The asymptotic variances can be written as

$$\lim_{N_l \rightarrow \infty} \frac{\text{Vartot}}{\text{age subsample fixed}} = \frac{b_2}{n_j^*} = n_l \frac{b_2}{n_s}$$

and

$$\lim_{N_l \rightarrow \infty} \frac{\text{Vartot}}{\text{age subsample random}} = \frac{c_1}{n_s}$$

The percentage reduction in Vartot (ages only) that can be achieved when age subsamples are fixed is then

$$\gamma_F = \frac{\frac{a_1}{n_s} - n_l \frac{b_2}{n_s}}{\frac{a_1}{n_s}} \times 100 = \frac{a_1 - n_l b_2}{a_1} \times 100$$

Since  $p_i = \sum_{j=1}^{n_i} l_j q'_{ij}$  implies  $a_1 = c_1 + c_2$ , it may be deduced that

Vartot ages only	=	Vartot age subsample random
------------------------	---	-----------------------------------

when  $N_i = n_s$  (i.e. when the length observations are taken from specimens that are aged). Therefore, the percentage reduction in Vartot (ages only) that can be achieved when age subsamples are random is

$$\gamma_R = \frac{\frac{a_1}{n_s} - \frac{c_1}{n_s}}{\frac{a_1}{n_s}} \times 100 = \frac{c_2}{a_1} \times 100$$

The age-length key, when age subsamples are random, cannot do worse than a random sample of ages without lengths. However, when age subsamples are fixed, the age-length key may do worse than a random sample of ages even if an infinite length sample were taken. In the following section, an example is given in which this is the case.

### Examples

As examples of how the preceding analyses can be used, age-length data were compiled from biological market samples for the following two species:

- a) Species: Pacific ocean perch  
(*Sebastes alutus*)  
Location: Queen Charlotte Sound  
Time: May-June-July 1975  
Sex: male  
Ages: 9-20 yr  
Lengths: 32-43 cm

- b) Species: Pacific cod  
(*Gadus macrocephalus*)  
Location: miscellaneous  
Time: February-March 1974  
Sex: male  
Ages: 3-5 yr  
Lengths: 53-77 cm

From the data presented in Tables 1 and 2, estimates of  $p_i$  and  $q_{ij}$  were made which were then assumed to be the true parameters of the population. The coefficients for Vartot were then calculated from the formulas given in the preceding section. The calculated coefficients are as follows:

Pacific ocean perch	Pacific cod
$a_1 = .85135$	$a_1 = .54270$
$b_1 = .66564$	$b_1 = .25866$
$b_2 = .09106$	$b_2 = .01798$
$b_3 = .09465$	$b_3 = .26605$
$c_1 = .75670$	$c_1 = .27665$
$c_2 = .09465$	$c_2 = .26605$
$\gamma_F = -28\%$	$\gamma_F = +17\%$
$\gamma_R = +11\%$	$\gamma_R = +49\%$

From the  $\gamma_F$  value, it is seen that for Pacific ocean perch, Vartot (ages only) increases 28% when an age-length key is used with fixed age subsamples. This means that the age determination is worse

TABLE 1. Age-length data for Pacific ocean perch (*Sebastes alutus*).

Length (cm)	Age (yr)											
	9	10	11	12	13	14	15	16	17	18	19	20
32	10	5										
33	15	13	13	1								
34	14	27	27	19	4	1						
35	26	36	59	32	3	2						
36	5	48	74	67	18	3	1					
37	3	23	51	37	20	7		1				
38		6	23	40	27	12	2		2			
39			5	10	13	23	9	6		2		1
40			1	1	12	13	14	18	6	2	1	
41				1	5	9	11	18	11	3	6	1
42						5	7	7	7	9	6	5
43						1	2	1	2	1	1	4

TABLE 2. Age-length data for Pacific cod (*Gadus macrocephalus*).

Length (cm)	Age (yr)		
	3	4	5
53	5		
54	3		
55	6		
56	13	3	
57	11		
58	14	3	
59	18	4	
60	20	2	
61	18	2	
62	15	10	
63	13	1	
64	8	9	1
65	6	14	
66	3	13	
67	1	13	
68	1	8	2
69		13	4
70	1	7	
71		4	2
72			2
73			2
74		1	
75			2
76			1
77			2

than a simple random sample of ages even if an infinite length sample is taken. From the  $\gamma_R$  value, it is seen that Vartot (ages only) decreases 11% when an age-length key is used with random age subsamples. The age-length key is apparently of marginal value in this case.

For Pacific cod,  $\gamma_F = 17\%$  and  $\gamma_R = 49\%$ . The age-length key may be of significant value. Again, random age subsamples are favored over fixed age subsamples. Tables 3 and 4 give values of Vartot at various sampling levels. For consistency, sample sizes were rounded to integers. From these tables, it appears that random samples are generally favored over fixed samples. For populations

TABLE 3. Values of Vartot for Pacific ocean perch. F and R refer to fixed and random age subsamples.

Aged sample	Length sample = aged sample	Total length sample											
		100		200		300		400		500		1000	
		F	R	F	R	F	R	F	R	F	R	F	R
50	.01703	.02538	.01608	.02407	.01561	.02364	.01545	.02342	.01537	.02329	.01532	.02303	.01523
100	.00851	.01316	.00851	.01227	.00804	.01198	.00788	.01183	.00780	.01174	.00776	.01156	.00766
150	.00568			.00773	.00552	.00749	.00536	.00737	.00528	.00730	.00523	.00715	.00514
200	.00426			.00603	.00426	.00580	.00410	.00569	.00402	.00562	.00397	.00549	.00388
250	.00341					.00476	.00334	.00465	.00326	.00459	.00322	.00446	.00312
300	.00284					.00405	.00284	.00395	.00276	.00389	.00271	.00376	.00262
350	.00243							.00343	.00240	.00338	.00235	.00326	.00226
400	.00213							.00305	.00213	.00299	.00208	.00287	.00199
450	.00189									.00262	.00187	.00251	.00178
500	.00170									.00239	.00170	.00228	.00161

TABLE 4. Values of Vartot for Pacific cod. F and R refer to fixed and random age subsamples.

Aged sample	Length sample = aged sample	Total length sample											
		100		200		300		400		500		1000	
		F	R	F	R	F	R	F	R	F	R	F	R
50	.01085	.01295	.00819	.01097	.00686	.01031	.00642	.00998	.00620	.00978	.00607	.00939	.00580
100	.00543	.00780	.00543	.00615	.00410	.00560	.00365	.00532	.00343	.00516	.00330	.00483	.00303
150	.00362			.00454	.00317	.00403	.00273	.00377	.00251	.00362	.00238	.00331	.00211
200	.00271			.00374	.00271	.00324	.00227	.00299	.00205	.00284	.00192	.00255	.00165
250	.00217					.00277	.00199	.00253	.00177	.00238	.00164	.00209	.00137
300	.00181					.00246	.00181	.00222	.00159	.00207	.00145	.00179	.00119
350	.00155							.00200	.00146	.00185	.00132	.00157	.00106
400	.00136							.00183	.00136	.00169	.00122	.00141	.00096
450	.00121									.00156	.00115	.00128	.00088
500	.00109									.00146	.00109	.00118	.00082

of different age structures, the analysis could change significantly. However, these changes can be examined by the method presented here.

In addition to the analysis presented here, the variance formulas can be used to examine each  $\hat{p}_i$  on an individual basis. For example, the coefficient of variation of each  $\hat{p}_i$  may be calculated. This type of analysis tends to be voluminous when many sampling levels are being examined.

Finally, the variance of the  $\hat{p}_i$ 's estimated from the actual use of an age-length key may be estimated by simply substituting the appropriate parameter estimates into the  $\text{var}(\hat{p}_i)$  formula for the case where the age subsample is fixed. The derivation of this formula only required that the  $n_j^*$ 's were fixed constants, not necessarily equal.

### Acknowledgments

I thank Dr D. Gunderson of the Northwest Fisheries Center of the National Marine Fisheries Service who suggested this analysis and was kind enough to review a draft.

- FRIDRIKSSON, A. 1934. On the calculation of age-distribution within a stock of cod by means of relatively few age determinations as a key to measurements on a large scale. *Rapp. P.V. Reun., Cons. Int. Explor. Mer.* 86: 1-14.
- KETCHEN, K. S. 1949. Stratified subsampling for determining age distributions. *Trans. Am. Fish. Soc.* 79: 205-212.
- SOUTHWARD, G. M. 1976. Sampling landings of halibut for age composition. *Int. Pac. Halibut Comm. Sci. Rep.* 58: 31 p.



# This article has been cited by:

1. Thomas Carruthers, Laurence Kell, Carlos Palma. Accounting for uncertainty due to data processing in virtual population analysis using Bayesian multiple imputation. *Canadian Journal of Fisheries and Aquatic Sciences*, ahead of print1-14. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
2. E. M. Schemmel, M. K. Donovan, C. Wiggins, M. Anzivino, A. M. Friedlander. 2016. Reproductive life history of the introduced peacock grouper *Cephalopholis argus* in Hawaii. *Journal of Fish Biology* **89**:2, 1271-1284. [[Crossref](#)]
3. R.I.C. Chris Francis. 2016. Growth in age-structured stock assessment models. *Fisheries Research* **180**, 77-86. [[Crossref](#)]
4. . References 283-304. [[Crossref](#)]
5. Sondre Aanes, Jon Helge Vølstad. 2015. Efficient statistical estimators and sampling strategies for estimating the age composition of fish. *Canadian Journal of Fisheries and Aquatic Sciences* **72**:6, 938-953. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
6. J Giddens, AM Friedlander, E Conklin, C Wiggins, K Stamoulis, MK Donovan. 2014. Experimental removal of the invasive peacock hind (roi) *Cephalopholis argus*, in Puakō, Hawai'i: methods for assessing and managing marine invasive species. *Marine Ecology Progress Series* **511**, 209-221. [[Crossref](#)]
7. Yingbin Wang, Ji Zheng, Cungen Yu. 2013. Stock assessment of chub mackerel (*Scomber japonicus*) in the central East China Sea based on length data. *Journal of the Marine Biological Association of the United Kingdom* 1-7. [[Crossref](#)]
8. Alec D. MacCall, Steven L.H. Teo. 2013. A hybrid stock synthesis—Virtual population analysis model of Pacific bluefin tuna. *Fisheries Research* **142**, 22-26. [[Crossref](#)]
9. Lewis G. Coggins, Daniel C. Gwinn, Micheal S. Allen. 2013. Evaluation of Age–Length Key Sample Sizes Required to Estimate Fish Total Mortality and Growth. *Transactions of the American Fisheries Society* **142**:3, 832-840. [[Crossref](#)]
10. J. Stewart, W. Sumpton, M. Lockett, J. M. Hughes. 2013. Age-based demographics of the pearl perch *Glaucosoma scapulare*. *Journal of Applied Ichthyology* n/a-n/a. [[Crossref](#)]
11. Caroline Bouchard, Louis Fortier. 2011. Circum-arctic comparison of the hatching season of polar cod *Boreogadus saida*: A test of the freshwater winter refuge hypothesis. *Progress in Oceanography* **90**:1-4, 105-116. [[Crossref](#)]
12. B. W. Kendall, C. A. Gray, D. Bucher. 2009. Age validation and variation in growth, mortality and population structure of *Liza argentea* and *Myxus elongatus* (Mugilidae) in two temperate Australian estuaries. *Journal of Fish Biology* **75**:10, 2788-2804. [[Crossref](#)]
13. J. Horppila, L. Nurminen. 2009. Food niche segregation between two herbivorous cyprinid species in a turbid lake. *Journal of Fish Biology* **75**:6, 1230-1243. [[Crossref](#)]
14. Ching-Ping Chih. 2009. Evaluation of the Sampling Efficiency of Three Otolith Sampling Methods for Commercial King Mackerel Fisheries. *Transactions of the American Fisheries Society* **138**:5, 990-999. [[Crossref](#)]
15. A. G. Murta, C. Vendrell. 2009. Using the EM algorithm to age fish eggs. *ICES Journal of Marine Science* **66**:4, 607-616. [[Crossref](#)]
16. Eliza C. Heery, Jim Berkson. 2009. Systematic Errors in Length Frequency Data and their Effect on Age-Structured Stock Assessment Models and Management. *Transactions of the American Fisheries Society* **138**:1, 218-232. [[Crossref](#)]
17. P.Doering-ArjesP. Doering-Arjes, M.CardinaleM. Cardinale, H.MosegaardH. Mosegaard. 2008. Estimating population age structure using otolith morphometrics: a test with known-age Atlantic

- cod (*Gadus morhua*) individuals. *Canadian Journal of Fisheries and Aquatic Sciences* **65**:11, 2342-2350. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
18. Hugo Robotham, Zaida I. Young, Juan C. Saavedra-Nievas. 2008. Jackknife method for estimating the variance of the age composition using two-phase sampling with an application to commercial catches of swordfish (*Xiphias gladius*). *Fisheries Research* **93**:1-2, 135-139. [[Crossref](#)]
19. Debra A. Prince, Lyle W. Konigsberg. 2008. New Formulae for Estimating Age-at-Death in the Balkans Utilizing Lamendin's Dental Technique and Bayesian Analysis. *Journal of Forensic Sciences* **53**:3, 578-587. [[Crossref](#)]
20. Faith A. Ochwada, James P. Scandol, Charles A. Gray. 2008. Predicting the age of fish using general and generalized linear models of biometric data: A case study of two estuarine finfish from New South Wales, Australia. *Fisheries Research* **90**:1-3, 187-197. [[Crossref](#)]
21. A J R Cotter, G M Pilling. 2007. Landings, logbooks and observer surveys: improving the protocols for sampling commercial fisheries. *Fish and Fisheries* **8**:2, 123-152. [[Crossref](#)]
22. Keith L. Hurley. 2007. Effects of a reduced subsampling protocol on calculation of growth and body condition indices. *Fisheries Research* **83**:2-3, 360-363. [[Crossref](#)]
23. Timothy J. Miller, John R. Skalski, James N. Ianelli. 2007. Optimizing a stratified sampling design when faced with multiple objectives. *ICES Journal of Marine Science* **64**:1, 97-109. [[Crossref](#)]
24. Laurence Challier, Pia Orr, Jean-Paul Robin. 2006. Introducing inter-individual growth variability in the assessment of a cephalopod population: application to the English Channel squid *Loligo forbesi*. *Oecologia* **150**:1, 17-28. [[Crossref](#)]
25. EILEEN E. HOFMANN, JOHN M. KLINCK, JOHN N. KRAEUTER, ERIC N. POWELL, RAY E. GRIZZLE, STUART C. BUCKNER, V. MONICA BRICELJ. 2006. A POPULATION DYNAMICS MODEL OF THE HARD CLAM, *MERCENARIA MERCENARIA*: DEVELOPMENT OF THE AGE- AND LENGTH-FREQUENCY STRUCTURE OF THE POPULATION. *Journal of Shellfish Research* **25**:2, 417-444. [[Crossref](#)]
26. Louis Fortier, Pascal Sirois, Josée Michaud, David Barber. 2006. Survival of Arctic cod larvae (*Boreogadus saida*) in relation to sea ice and temperature in the Northeast Water Polynya (Greenland Sea). *Canadian Journal of Fisheries and Aquatic Sciences* **63**:7, 1608-1616. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
27. T J Miller, J R Skalski. 2006. Integrating design- and model-based inference to estimate length and age composition in North Pacific longline catches. *Canadian Journal of Fisheries and Aquatic Sciences* **63**:5, 1092-1114. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
28. Nancy A. Nate, Mary T. Bremigan. 2005. Comparison of Mean Length at Age and Growth Parameters of Bluegills, Largemouth Bass, and Yellow Perch from Length-Stratified Subsamples and Samples in Michigan Lakes. *North American Journal of Fisheries Management* **25**:4, 1486-1492. [[Crossref](#)]
29. V. S. Troynikov, S. G. Robertson. 2005. Estimating age composition using the Fredholm first-kind equation. *Marine and Freshwater Research* **56**:5, 745. [[Crossref](#)]
30. A. K. Morison, J. Burnett, W. J. McCurdy, E. Moksness. 2005. Quality issues in the use of otoliths for fish age estimation. *Marine and Freshwater Research* **56**:5, 773. [[Crossref](#)]
31. R. I. C. Chris Francis, Shelton J. Harley, Steven E. Campana, Peer Doering-Arjes. 2005. Use of otolith weight in length-mediated estimation of proportions at age. *Marine and Freshwater Research* **56**:5, 735. [[Crossref](#)]
32. A J R Cotter, L Burt, C G M Paxton, C Fernandez, S T Buckland, J-X Pan. 2004. Are stock assessment methods too complicated?. *Fish and Fisheries* **5**:3, 235-254. [[Crossref](#)]
33. RIC Chris Francis, Steven E Campana. 2004. Inferring age from otolith measurements: a review and a new approach. *Canadian Journal of Fisheries and Aquatic Sciences* **61**:7, 1269-1284. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]

34. John Stewart, Douglas J. Ferrell, Bryan van der Walt. 2004. Sizes and ages in commercial landings with estimates of growth, mortality and yield per recruit of yellowtail kingfish ( *Seriola lalandi* ) from New South Wales, Australia. *Marine and Freshwater Research* **55**:5, 489. [[Crossref](#)]
35. Hans-Georg Müller, Brad Love, Robert D. Hoppa. 2002. Semiparametric method for estimating paleodemographic profiles from age indicator data. *American Journal of Physical Anthropology* **117**:1, 1-14. [[Crossref](#)]
36. Anna Rindorf, Peter Lewy. 2001. Analyses of length and age distributions using continuation-ratio logits. *Canadian Journal of Fisheries and Aquatic Sciences* **58**:6, 1141-1152. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
37. Phillip W. Bettoli, Leandro E. Miranda. 2001. Cautionary Note about Estimating Mean Length at Age with Subsampled Data. *North American Journal of Fisheries Management* **21**:2, 425-428. [[Crossref](#)]
38. David G Reddin, Kevin D Friedland. 1999. A history of identification to continent of origin of Atlantic salmon (*Salmo salar* L.) at west Greenland, 1969-1997. *Fisheries Research* **43**:1-3, 221-235. [[Crossref](#)]
39. Rosangela Lessa, Francisco Marcante Santana, Renato Paglerani. 1999. Age, growth and stock structure of the oceanic whitetip shark, *Carcharhinus longimanus*, from the southwestern equatorial Atlantic. *Fisheries Research* **42**:1-2, 21-30. [[Crossref](#)]
40. A JR Cotter. 1998. Method for estimating variability due to sampling of catches on a trawl survey. *Canadian Journal of Fisheries and Aquatic Sciences* **55**:7, 1607-1617. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
41. KEVIN M. BAILEY, SUSAN J. PICQUELLE, STELLA M. SPRING. 1996. Mortality of larval walleye pollock *Theragra chalcogramma* in the western Gulf of Alaska, 1988-91. *Fisheries Oceanography* **5**:s1, 124-136. [[Crossref](#)]
42. D. G. Worthington, A. J. Fowler, P. J. Doherty. 1995. Determining the most efficient method of age determination for estimating the age structure of a fish population. *Canadian Journal of Fisheries and Aquatic Sciences* **52**:11, 2320-2326. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
43. D. J. Jellyman. 1995. Longevity of longfinned eels *Anguilla dieffenbachii* in a New Zealand high country lake. *Ecology of Freshwater Fish* **4**:3, 106-112. [[Crossref](#)]
44. M. J. Unwin, D. H. Lucas. 1993. Scale Characteristics of Wild and Hatchery Chinook Salmon (*Oncorhynchus tshawytscha*) in the Rakaia River, New Zealand, and Their Use in Stock Identification. *Canadian Journal of Fisheries and Aquatic Sciences* **50**:11, 2475-2484. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
45. Thomas P. Quinn, Martin J. Unwin. 1993. Variation in Life History Patterns among New Zealand Chinook Salmon (*Oncorhynchus tshawytscha*) Populations. *Canadian Journal of Fisheries and Aquatic Sciences* **50**:7, 1414-1421. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
46. Lyle W. Konigsberg, Susan R. Frankenberg. 1992. Estimation of age structure in anthropological demography. *American Journal of Physical Anthropology* **89**:2, 235-256. [[Crossref](#)]
47. Jukka Horppila, Heikki Peltonen. 1992. Optimizing Sampling from Trawl Catches: Contemporaneous Multistage Sampling for Age and Length structures. *Canadian Journal of Fisheries and Aquatic Sciences* **49**:8, 1555-1559. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
48. Barry D. Smith, Gordon A. McFarlane, Mark W. Saunders. 1992. Inferring the Summer Distribution of Migratory Pacific Hake (*Merluccius productus*) from Latitudinal Variation in Mean Lengths-at-Age and Length Frequency Distributions. *Canadian Journal of Fisheries and Aquatic Sciences* **49**:4, 708-721. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
49. Dominique Pelletier, Philippe Gros. 1991. Assessing the impact of Sampling Error on Model-Based Management Advice: Comparison of Equilibrium Yield per Recruit Variance Estimators. *Canadian Journal of Fisheries and Aquatic Sciences* **48**:11, 2129-2139. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
50. Philip J. Smith. 1989. Is Two-Phase Sampling Really Better for Estimating Age Composition?. *Journal of the American Statistical Association* **84**:408, 916-921. [[Crossref](#)]

51. John M. Hoenig, Dennis M. Heisey, William D. Lawing, Dennis H. Schupp. 1987. An Indirect Rapid Methods Approach to Assessment. *Canadian Journal of Fisheries and Aquatic Sciences* **44**:S2, s324-s338. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]
52. 1984. LETTERS AND COMMENTS/LETTRES ET COMMENTAIRES. *Canadian Journal of Fisheries and Aquatic Sciences* **41**:5, 825-834. [[Citation](#)] [[PDF](#)] [[PDF Plus](#)]