

## Planarity

**Problem:** Let  $G$  be a graph. Can  $G$  be drawn in such a manner so that no two edges intersect?

### Example 1. *Example Graph here*

Any drawing of  $K_5$  or  $K_{3,3}$  have (at lease) 2 edges which cross (*Proof to come*)

**Definition 1.** A graph  $G$  is said to be planer if it can be drawn in  $\mathbb{R}^2$  so that no two edges cross. Such a drawing is called a plane drawing. The graph associated with a plane drawing is usually refered to as a plane graph.

**Remark.** 1. Any subgraph of a planer graph is planer.


2. Every plane graph is planar.

*graph here* is a planer but not plane.

Our aim is to determine condition which ensures that a graph  $G$  is planar. Let  $G$  be a plane graph. Consider the set  $S$  obtained from  $\mathbb{R}^2$  by deleting (the vertices and edges) of  $G$ . We observe that  $S$  is the disjoint union of finitely many subsets  $F_1, F_2, \dots, F_l$  of  $\mathbb{R}^2$  having the following two properties:

1. Any two points of  $F_i$  can be joined by a curve not crossing  $G$ .
2. Any curve in  $\mathbb{R}^2$  which joins a point of  $F_i$  to one of  $F_j, i \neq j$ , must cross  $G$

**Definition 2.** Let  $G$  be a plane graph. The sets  $F_1, \dots, F_l$  described above are called the faces of  $G$ .



### Example 2.

**Remark.** One face is always unbounded, with the remaining faces all bounded.

**Definition 3.** Let  $F$  be a face of a plane graph  $G$ . The boundary of  $F$  consists of a finite # of vertices and edges of  $G$

The length of a closed walk around the boundary of  $F$  is called the degree of  $F$ , usually denoted  $\deg F$

### Example 2. (Cont.)

1.  $\deg F_1 = \deg F_2 = 5$

2.  $\deg F_2 = 2$  the closed walk is  $x \rightarrow y \rightarrow x$

3.  $\deg F_1 = \deg F_2 = 3$   
 $\deg F_3 = 6$


**Theorem 1.** (Euler's Formula, 1780) Let  $G$  be a connected plane graph. If  $G$  has  $n$  nodes and  $f$  faces then

$$n - m + f = 2$$

proof of **Theorem 1** requires a couple of preliminary lemmas.


**Lemma.** Let  $G$  be a plane graph.  $G$  contains a cycle if and only if the number of faces of  $G \geq 2$ .

*Proof.*  $\rightarrow$  Let  $C$  be a cycle of  $G$ . Let  $x$  and  $y$  be in the interior and exterior of  $C$ , respectively, then any curve in  $\mathbb{R}^2$  connecting  $x$  and  $y$  must cross  $C$ , have cross  $G$



Choosing  $x, y$  not lying on  $G$   $x$  and  $y$  belong to different faces of  $G$ .

$\leftarrow$  The absence of a cycle,  $G$  is a forest. On induction on the number of components of  $G$  shows  $G$  has only one (unbounded) face.




□


**Lemma.** Let  $e$  be an edge of a plane graph  $G$ .

1. If  $e$  is a bridge then it lies on boundary of exactly one face.
2. if  $e$  is not a bridge then it lies on the boundary of exactly two faces of  $G$

*Proof.* 1. Let  $H_1$  and  $H_2$  be the components of  $G \setminus e$ . The edge  $e$  lies in face  $F_1$  of  $H_1$ , as well as a face  $F_2$  of  $H_2$ . The intersection  $F_1 \cap F_2$  contains a unique face  $F_o$  of  $G$ .  $e$  lies on the boundary of  $F_o$  and is the unique face of  $G$ .



2. Let  $C$  be a cycle containing  $e$ . The edge  $e$  lies on the boundary of one face lying in the interior of  $C$  and one face lying in the exterior of  $C$ . Thus,  $e$  lies on the boundary of at least 2 faces of  $G$ .



The fact faces are disjoint can be used to show that  $e$  lies on the boundary of at most 1 face lying in the interior of  $C$ , and one face lying in the exterior of  $C$ . Then  $e$  lies in the boundary of at most 2 faces.

□

**Corollary.** (*Handshaking Lemma for Planar Graphs*) If  $G$  is a plane graph of size  $m$  then

$$2m = \sum_{\text{faces } F} \deg F$$

*Proof.* Each edge  $e$  of  $G$  contributes 2 to the sum on the right:

If  $e$  is a bridge lying on the face  $F_o$  then it contributes 2 to  $\deg F_o$  and 0 to the remaining  $\deg F$ .

If  $e$  is not a bridge then it contributes 1 to the degree of two distinct faces of  $G$ , and 0 to the remaining. □


*Proof.* Proof of Euler's Formula

We proved by induction on # of faces of  $G$  on the case  $f = 1$ . **Lemma 2** asserts  $G$  is a forest, where  $G$  is a tree. Thus,  $m = n - 1$ , and hence  $n - m + f = n - (n - 1) + 1 = 2$

Assume the result is true  $f$  connected plane graph  $H$  having  $f$  faces, and let  $G$  have  $f + 1$  faces, say  $F_1, F_2, \dots, F_l, F_{l+1}$ . If  $n = |G|, m = e(G)$  then we are required to show

$$n - m + (f + 1) = 2$$

Since  $f + 1 \geq 2$ ,  $G$  contains a cycle  $C$  (**Lemma 2**). Fix an edge  $e$  lying on  $C$ . Assume  $e$  is not a bridge of  $G$ . **Lemma 2** ensures that  $e$  lies on the boundary of 2 distinct faces of  $G$ , say  $F_l$  and  $F_{l+1}$ .



Construct the subgraph  $H = G - e$ . We note  $H$  is a connected plane graph of order  $n$  and size  $m - 1$ . Furthermore, the number of faces of  $H$  is  $f$ . denoted each of the faces  $F_1, f_2, \dots, F_{l-1}$  of  $G$  occur on faces of  $H$ . Assume  $e$  does not appear in the boundary of any of these faces.

The remaining faces of  $H$  is obtained by joining  $F_l$  and  $F_{l+1}$  along the edge  $e$ .

$$\hat{F}_l = F_l \cup F_{l+1} \cup e$$

Since  $H$  has  $f$  faces, the induction hypotheses allows in  $G$  to conclude that

$$2 = n - (m - 1) + f = n - m + (f + 1)$$

as required

□

**Corollary.** Let  $G$  be a connected planer graph. Each plane drawing of  $G$  has the same number of faces, namely  $2 + m - n$

Euler's formula can be used to obtain a necessary condition for a simple connected graph to be planar.

**Definition 4.** Let  $G$  be a graph, If  $G$  contains a cycle then the **girth**  $gr(G)$  is defined as the length of the smallest cycle in  $G$ . If  $G$  is a forest then we set  $gr(G) = \infty$

a.)  has girth 3

b.) Peterson has girth 5

c.)  $K_5$  has girth 3,  $K_{3,3}$  has girth 4

**Example 3.**

**Remark.** If  $G$  is simple then  $gr(G) \geq 3$

**Theorem 4.** Let  $G$  be a connected simple planar graph. If  $G$  has order  $n$  and size  $m$  then

$$m \leq \begin{cases} n - 1 & \text{if } gr(G) = \infty \\ \frac{gr(G)}{gr(G)-2}(n - 2) & \text{if } gr(G) \text{ is finite} \end{cases} \quad (1)$$

*Proof.* If  $gr(G) = \infty$  then  $G$  is a tree, hence  $m = n - 1$  if  $gr(G)$  is finite then  $G$  has  $\geq 2$  faces. In this case, the boundary of each face of  $G$  contains a cycle, hence

$$\deg(F) \geq gr(G)$$

If each face  $F$  of  $G$ . therefore, if  $f = \#$  of faces of  $G$  then the handshaking lemma of planer graphs

$$2m = \sum_{\text{faces } F} \deg F \geq \sum gr(G) = gr(G)f$$

From Euler's formula,  $f = 2 + m - n$ , substitutes  $f_i$  in the preceding yields

$$2m \geq gr(G)(2 + m - n)$$

□

**Example 4.** 1. Does there exist a simple connected planar graph  $G$  of order 12 and size 40?

**Solution 1.** Suppose such a graph  $G$  exists, Note that  $G$  cannot be a tree, hence  $gr(G)$  is finite. Thus, **Theorem 3** asserts

$$40 \geq \frac{gr(G)}{gr(G) - 2}(12 - 2)$$

Solving for  $gr(G)$

$$gr(G) \leq \frac{8}{3} < 3$$

This contradicts that  $gr(G) \geq 3$  No such  $G$  exists.

2. Let  $G$  be a planar graph of size and girth 5. What can one say about  $n = |G|$

**Solution 2.** From **Theorem 2**

$$14 \leq \frac{5}{5 - 2}(n - 2)$$

solving for  $n$  yields  $n \geq 52/5$  some  $n$  is an integer  $n \geq 11$ .

**Corollary.**  $K_5$  and  $K_{3,3}$  are non-planar

**Proof here**

**Exercise Show that Peterson is non-planar**

**Corollary.** Let  $G$  be a simple connected planar graph.

1. if  $G$  has order  $n \geq 3$  then  $e(G) \leq 3n - 6$
2. Furthermore, if  $G$  contains no triangles then  $e(G) \leq 2n - 4$

**proof here**


**Remark.** The preceding two results are often used as tools for non-planarity, they are weaker than the full theorem 4.

**Corollary.** Every simple planar graph  $G$  contains a vertex of degree at most 5 ( $\delta(G) \leq 5$ )

**proof here**

**Example 5.**  $K_n$  is non-planar of  $n \geq 7$ . Each vertex of  $K_n$  has degree  $n - 1 \geq 6$

**Example 6.** The condition of theorem 4 (as well as its corollaries) is only necessary for planarity, not sufficient. For example, the following graph  $G$  is non-planar, as it contains a copy of  $K_{3,3}$



On the other hand,  $|G| = 8, e(G) = 12$  and  $gr(G) = 4$

$$\frac{4}{4 - 2}(8 - 2) = 12$$


**Remark.** Euler's formula can be extended to disconnected plane graphs using induction on the # of components. If  $G$  is a planar graph of order  $n$ , size  $m$ , with  $f$  faces and  $k$  components then

$$n - m + f = k + 1$$

There are also analogues for Theorem 3 and its corollaries.


We have already observed that Theorem 3 and its corollaries present many necessary conditions for a graph to be planar.

**Definition 5.** Let  $G$  and  $H$  be graphs.  $H$  is said to be a subdivision of  $G$  if the former graph can be constructed from the latter by introduction of a finite # of new vertices along existing edges.



**Example 7.**

c.) Each cycle graph  $C_n$  is a subdivision of  $C_1$ .  $C_n$  can be obtained from  $C_1$  by the addition of  $n - 1$  new vertices on the single edge of  $C_1$



**Remark.** 1. The process of subdivision only introduces new vertices of degree 2

2. Note that if  $H$  is a subdivision of  $G$  then  $H$  has the same shape as  $G$ .


**Definition 6.** Two graphs  $G$  and  $H$  are said to be homeomorphic if they are both subdivisions of a common graph.

**Example 8.** 1. If  $n, m \geq 1$  then  $C_n$  is homeomorphic to  $C_m$  as both are subdivisions of  $C_1$ .

2. If  $n, m \geq 2$  then  $P_n$  and  $P_m$  are homeomorphic as both are subdivisions of  $P_2$

**Remark.** 1.  $G$  and  $H$  are homeomorphic if the latter can be obtained from the former by a finite sequence of following 2 operations:


- (a) addition of new vertex along an existing edge.
- (b) filling in of an existing vertex of degree 2.



2. homeomorphic graphs have the same shape.

**Theorem 5.** A graph is planar if and only if it contains no subgraph homeomorphic to  $K_5$  or  $K_{3,3}$ .

**Example 9.** The graph




an alternate characterization of non-planar graphs can be obtained using the notion of contractions.

**Definition 7.** Let  $G$  and  $H$  be graphs.  $G$  is said to be contractible to  $H$  if  $H$  can be obtained from  $G$  by successively contracting a finite number of edges.

**Theorem 6.** A graph  $G$  is planar if and only if it contains no subgraph which is contractible to  $K_5$  or  $K_{3,3}$ .


**Example 10.** Peterson graph is contractible to  $K_5$ .



Thus peterson is non-planar.


We have observed that  $K_5$  and  $K_{3,3}$  are non-planar. What happens if we consult other surface?

**Example 11.** Torous (Donut)



**Definition 8.** A non-planar graph  $G$  is said to be toroidal if it can be drawn on the torus so as no two edges cross.

**Example 12.** 1.  $K_5$  is toroidal



2.  $K_{3,3}$  is toroidal

**Question:** Is there analogous of Euler's Formula for the toros?

**Theorem 7.** Let  $G$  be a simple connected torodial graph of order  $n$ , size  $m$  and  $f$  faces, Then

$$n - m + f = 0$$

Furthermore,

$$m \leq \frac{gr(G)}{gr(G) - 2} n$$

**Definition 9.** A graph  $G$  is said to have genus  $g$  if  $G$  can be drawn on a surface of genus  $g$  with no edges crossing, but no drawing on a surface of genus  $g - 1$  exists. (i.e planar = genus 0, torodial = genus 1)



**Theorem 8.** *Let  $G$  be a connected graph of genus  $g$ , order  $n$ , size  $m$ , and face  $f$ . Then*

$$n - m + f = 2 - 2g$$

*Furthermore, if  $G$  is simple of finite girth then*

$$m \leq \frac{gr(G)}{gr(G) - 2}(n + 2g - 2)$$

**Corollary.** *Let  $G$  be a connected simple graph of genus  $g$ , order  $n \geq 3$  and size  $m$  then,*

$$m \leq 3(n + 2g - 2)$$

$$m \leq 2(n + 2g - 2) \text{ if no triangle present}$$

**Corollary.** *Let  $G$  be a connected simple graph of order  $n \geq 4$  and size  $m$ . Then the genus  $g$  satisfies*

$$g \geq \lceil \frac{m - 3n}{6} + 1 \rceil$$

$\lceil x \rceil = \text{least integer greater than or equal to } x$

**Remark.** *Let  $G$  be a graph. The crossing number  $cr(G)$  is the minimum # of crossing that can occur when  $G$  is drawn in the plane.*

**Example 13.** *graph here*


**Theorem 9.** *The genus of the graph  $G$  is  $\leq cr(G)$*

### Dual Graphs

**Definition 10.** *Let  $G$  be a planar graph. The geometric dual  $G^*$  of  $G$  is the graph constructed as follows:*

1. *denote each face of  $G$ , choose a point  $v^*$ . The  $v^*$  forms the vertex set  $V(G^*)$*
2. *For each edge  $e$  of  $G$ , join the vertices  $v^*$  and  $w^*$  in the adjacent face by a curve  $e^*$  that crosses  $e$  and no other edge of  $G$ . The collection of  $e^*$  forms the edge set  $E(G^*)$*

**Example 14.**



### Platonic Graphs

<u>Graphs</u>	<u>Geometric Dual</u>	
tetrahedron	tetrahedron	self-dual
cube	octahedron	
octahedron	cube	
dodecahedron	icosahedron	
icosahedron	dodecahedron	

**Remark.** 1. The construction of  $G^*$  depends on the plane drawing of  $G$ . For example, the graphs  $G$  in example 13 (i) and (ii) are isomorphic, but their geometric duals are not. One has vertex of degree 5 but the other has no such vertex.

2.  $G^*$  is planar and connected.

**Lemma.** Let  $G$  be a connected planar graph of order  $n$ , size  $m$ , and face  $f$ . If  $G^*$  is a geometric dual then the nodes  $n^*$  size  $m^*$  and number of faces  $f^*$  satisfies

$$n^* = f, m^* = m, f^* = f$$

**Theorem 10.** Let  $G$  be a connected planar graph. Then  $G^{**}$  is isomorphic to  $G$ .

**Theorem 11.** Let  $G$  be a connected planar graph and  $G^*$  a geometric dual of  $G$ , A subset of  $E(G)$  forms a cycle of  $G$  if and only if the corresponding subset of  $E(G^*)$  is a cutset of  $G^*$ .

**Remark.** If  $e$  is an edge of  $G$  then there is a unique edge  $e^*$  of  $G^*$  which crosses  $e$ .


**Corollary.** Let  $G$  be a connected planar graph. A set of edges of  $G$  forms a cutset if and only if the corresponding edges of  $G^*$  forms a cycle.

**Theorem 12.** Let  $G$  be a connected planar graph  $G$  is bipartite if and only if  $G^*$  is eulerian.

**Theorem 13.** Let  $G$  be a connected planar graph. If  $G$  is 3-edge connected then  $G^*$  is simple (of order  $\geq 3$ )

**Definition 11.** Let  $G$  be a graph. A graph  $G^*$  is said to be an abstract dual of  $G$  if there exists a one-one correspondence between the edge of  $G$  and the of  $G^*$  with the property that  
a subset of  $E(G)$  forms a cycle  $\leftrightarrow$  correspong subset of  $E(G^*)$  forms a cutset

**Example 15.** If  $G$  is plane then its geometric dual  $G^*$  is an abstract dual (Theorem 11)



**Theorem 14.** If  $G^*$  is an abstract dual of  $G$  then  $G$  is an abstract dual of  $G^*$ .  
Abstract duals provide another characterization of planar graphs.

**Theorem 15.** A graph  $G$  is planar if and only if  $G$  has an abstract dual.