

Two models from the same provider on one question

MedMCQA question

The retrocuspid papilla is found approx 1mm below the free gingiva on the attached gingiva lingual to:

- ☒ A Mandibular canine. keyed answer
- ☐ B Maxillary canine.
- ☐ C Mandibular premolars
- ☐ D Maxillary premolars.

Qwen 3.5 9B

smaller model · answered (B)

Explanation: The retrocuspid papilla is a specific anatomical landmark found in the oral cavity. It is a small, distinct elevation of the attached gingiva located on the lingual aspect of the maxillary canine. It is situated approximately 1 mm below the free gingival margin. This feature is unique to the maxillary canine region and is not typically found on mandibular teeth or premolars. Therefore, the retrocuspid papilla is associated with the maxillary canine.

Answer: (B)

INCORRECT

Qwen 3.7 Plus

larger model · answered (A)

CORRECT

Explanation: The retrocuspid papilla is a small, circumscribed, and elevated area of normal gingival tissue. It is classically located on the attached gingiva, approximately 1 mm below the free gingival margin, specifically on the lingual aspect of the mandibular canine. While it is a normal anatomical landmark, it can occasionally become inflamed or hypertrophic.

Answer: (A)

Both models were given the identical prompt at temperature 0. The smaller model locates the landmark on the maxillary canine; the larger one on the mandibular canine.