

Questions that every model answered incorrectly

Three of the 105 questions missed by all ten models, one per recurring pattern. Green marks the recorded key; red marks the answer the models converged on.

Example 1 · MedMCQA

models agreeing on one non-key answer: 10 of 10

Wave patterns of EEF, ECG and EMG are depicted below. The B pattern belongs to (Figure was not provided in the exam):

- A NREM sleep
- B REM sleep
- C Wakefulness
- D Quiet wakefulness

model consensus

recorded key

The stimulus the question refers to is absent, so the item cannot be answered from text alone. The question itself records that the figure was not provided.

Example 2 · MedMCQA

models agreeing on one non-key answer: 10 of 10

The police has brought an unresponsive patient to you. What is the first thing you will do?

- A Sta chest compressions immediately
- B Check carotid pulse
- C Check for response and call help
- D Sta rescue breaths

recorded key

model consensus

Every model selects the assess-and-summon-help step; the key selects the pulse check. Two choice strings are also truncated ("Sta" for "Start").

Example 3 · MedMCQA

models agreeing on one non-key answer: 8 of 10

In mandibular primary second molar true statement is

- A ML is largest cusp and distobuccal is smallest
- B All buccal cusp are fo same size> all lingual cusp are of same size
- C DB is largest cusp
- D All of the above

model consensus

recorded key

The keyed choice is typographically corrupted ("fo" for "of", a stray ">"), leaving no well-formed correct choice.

We describe these questions as warranting review rather than as confirmed errors: no independent clinician adjudication was obtained.