

A3

## Serum Creatinine Variability as a Novel Biomarker Predictive of Albuminuria Progression in Type 2 Diabetes Mellitus – Discovery and Validation in Two Longitudinal Cohorts

**Dr Kiat Mun Serena Low<sup>1</sup>**, Associate Professor Su Chi Lim<sup>1</sup>, Dr Xiao Zhang<sup>1</sup>, Dr Keven Ang<sup>1</sup>, Associate Professor Tavintharan Subramaniam<sup>1</sup>, Associate Professor Chee Fang Sum<sup>1</sup>

<sup>1</sup>*Khoo Teck Puat Hospital*

AFES Investigator Award, November 10, 2017, 2:15 PM - 3:15 PM

**Objective(s):** The role of stability of renal function on kidney disease progression is unknown. We examined association between serum creatinine (scr) variability and albuminuria progression in Type 2 Diabetes(T2DM) and elucidated mechanism for the association.

**Materials and Methods:** In a discovery cohort, we identified 589 T2DM patients attending Diabetes Centre with  $\geq 6$  scr readings and no macroalbuminuria at baseline in 2002-2014. Scr variability was expressed adjusted-Cr-standard-deviation(CrSD)= $SD/\sqrt{n/(n-1)}$ . We used cox-proportional hazards regression for outcome on progression (worsening across albuminuria stages), adjusting for demographics, medication, blood pressure, urinary-albumin-to-creatinine ratio (uACR), estimated glomerular filtration rate (eGFR), glycated haemoglobin(HbA1c) and cr-intrapersonal-mean (CrMEAN). We validated the association in a prospective cohort comprising 554 T2DM patients attending Diabetes Centre in 2011-2016. The Homeostatic model assessment (HOMA2)-Insulin Resistance (IR) index was obtained from HOMA Calculator. Sobel-Goodman mediation test was performed to examine extent of mediation between IR(defined by HOMA2-IR>1.8) and progression by adjusted-CrSD.

**Results:** Progression occurred in 31.2% of patients in discovery cohort. The progressors had higher median adjusted-CrSD than non-progressors (9.62(5.90-16.93) vs. 6.94(4.64-10.57);  $p<0.001$ ) and poorer baseline eGFR, uACR and HbA1c ( $p<0.05$ ). Log-transformed adjusted-CrSD was significantly associated with progression (adjusted HR 1.46(1.06-2.01);  $p=0.021$ ). Similar findings were noted in validation cohort (adjusted OR 1.51 (1.09-2.09);  $p=0.013$ ). CrMEAN was not associated with progression in both cohorts. Adjusted-CrSD accounted for 26% of association between IR and progression after adjustment in mediation test in validation cohort ( $p=0.015$ ).

**Conclusion:** Higher scr variability predicted albuminuria progression, independently of intrapersonal mean. It is a potential mediator of association between IR and ACR progression.