Next Generation Sequencing-based Discovery of Novel Genetic Markers in the Vitamin D Receptor Gene correlated with Osteoporotic Fracture among Older Filipino Women

Mr Mark Pretzel Zumaraga^{1,2,3}, Dr Paul Julius Medina^{4,5}, Mr Juan Miguel Recto², Dr Edelyn Azurin⁴, Dr Celeste Tanchoco¹, Dr Cecilia Jimeno⁶, Dr Cynthia Saloma^{2,3}

¹Food And Nutrition Research Institute, ²Philippine Genome Center, ³National Institute of Molecular Biology and Biotechnology, ⁴Philippine Orthopedic Center, ⁵Mindanao State University – College of Medicine, ⁶Philippine Society of Endocrinology, Diabetes and Metabolism

AFES Investigator Award, November 10, 2017, 2:15 PM - 3:15 PM

Objectives: This study aimed to discover novel genetic variants in the entire 101 kB vitamin D receptor (VDR) gene for osteoporotic fractures in a group of older Filipino women using next generation sequencing approach in a case-control study design.

Materials and Methods: A total of fifty women with and without osteoporotic fracture seen at the Philippine Orthopedic Center were included. Blood samples were collected for determination of serum vitamin D, calcium, phosphorus, glucose, blood urea nitrogen, creatinine, aspartate aminotransferase, alanine aminotransferase and as primary source for targeted VDR gene sequencing using the Ion Torrent Personal Genome Machine. The variant calling was based on the GATK best practice workflow and annotated using Annovar tool.

Results A total of 1496 unique variants in the whole 101-kb VDR gene were identified. Novel sequence variations not registered in the dbSNP database were found among cases and controls at a rate of 23.1% and 16.6% of total discovered variants, respectively. Noteworthy is the discovery of two disease-associated novel heterozygous frameshift deletions (Pearson chi square p-value. Neither of the two SNPs was found in control patients nor in the dbSNP. Furthermore, two novel exonic variants were predicted to be damaging and may have increased the probability of acquiring osteoporotic fracture.

Conclusion: These findings show the power of Next Generation Sequencing in identifying sequence variations in a very large gene and the surprising results obtained in this study greatly expand the catalog of known VDR sequence variants that may represent an important clue in inter-individual response to vitamin D treatment for osteoporotic fracture.