

CURVE COUNTING IN GENUS ONE: ELLIPTIC SINGULARITIES & RELATIVE GEOMETRY

LUCA BATTISTELLA, NAVID NABIJOU AND DHRUV RANGANATHAN

ABSTRACT. We construct and study the reduced, relative, genus one Gromov–Witten theory of very ample pairs. These invariants form the principal component contribution to relative Gromov–Witten theory in genus one and are relative versions of Zinger’s reduced Gromov–Witten invariants. We relate the relative and absolute theories by degeneration of the tangency conditions, and the resulting formulas generalise a well-known recursive calculation scheme put forward by Gathmann in genus zero. The geometric input is a desingularisation of the principal component of the moduli space of genus one logarithmic stable maps to a very ample pair, using the geometry of elliptic singularities. Our study passes through general techniques for calculating integrals on logarithmic blowups of moduli spaces of stable maps, which may be of independent interest.

CONTENTS

Introduction	1
1. Constructions and logarithmic smoothness	4
2. Stratification and tropicalization	13
3. Degeneration of contact order	15
4. Splitting the boundary	17
5. Quantum Lefschetz algorithm	31
References	39

INTRODUCTION

The Kontsevich moduli space of stable maps from genus zero curves to projective space exhibits remarkable geometry – it is a smooth orbifold with a “self-similar” normal crossings boundary. These facts have combined to produce a very satisfactory understanding of Gromov–Witten theory in genus zero.

The higher genus situation is much more delicate, and substantial theory has been developed in recent years to treat the genus one case, both for its intrinsic geometry, and for higher genus insights. This began over a decade ago with pioneering work of Li, Vakil, and Zinger [Zin09a, VZ08, LZ07, LZ09, Zin08, Zin09b], and was recently reinvigorated using ideas from singularity theory [BCM20, HL10, Smy11a, Vis12] and logarithmic geometry [RSPW19a, RSPW19b, BC20]. The result is a **reduced** Gromov–Witten theory, which removes degenerate contributions from the ordinary theory. While the Atiyah–Bott localisation for the reduced invariants was used to great effect by Zinger in his proof of the BCOV conjectures, two major components of the subject – relative invariants and the degeneration formula – have not been developed in this setting. Our work provides these as new structural tools in reduced genus one Gromov–Witten theory¹. While the essential starting point is the early work on relative Gromov–Witten theory by Gathmann,

¹Further motivation to consider the genus one case specifically comes simply from the fact that for Calabi–Yau geometries in dimension larger than 3, Gromov–Witten theory vanishes in genera greater than one.

modern tools are required in our setting. We establish both the conceptual framework and the formula by combining tropical geometry and logarithmic Gromov–Witten theory, as developed by Abramovich–Chen–Gross–Siebert, with the recent understanding of elliptic singularities and their interactions with logarithmic structures.

0.1. Results. Our first contribution is the construction of a **reduced relative** Gromov–Witten theory associated to elliptic curves in \mathbb{P}^m with prescribed tangency along a hyperplane, and analogous invariants associated to smooth pairs (X, Y) with Y very ample. Our reduced relative theory differs from relative Gromov–Witten theory in a parallel fashion to how Li–Zinger’s reduced Gromov–Witten theory differs from the standard theory.

The geometric content is the following theorem, which provides a smooth and proper moduli space compactifying the space of elliptic curves in projective space with prescribed contact order data. Let $\mathcal{M}_{1,\alpha}^\circ(\mathbb{P}^m|H, d)$ be the moduli space of positive degree maps from smooth curves of genus one to \mathbb{P}^m with dimensionally transverse contact order $\alpha \in \mathbb{Z}_{\geq 0}^n$ along H . A precise version of the following theorem requires a certain amount of tropical and logarithmic language, and is therefore stated precisely in §1.

Theorem A. There exists a logarithmically smooth and proper Deligne–Mumford stack $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ having expected dimension, and containing $\mathcal{M}_{1,\alpha}^\circ(\mathbb{P}^m|H, d)$ as a dense open subset. A point in this moduli space determines commutative diagrams

$$\begin{array}{ccccc} C & \longrightarrow & \overline{C}_1 & \longrightarrow & \overline{C}_2 \\ & & \downarrow & & \downarrow \\ & & \mathbb{P}^m[s] & \longrightarrow & \mathbb{P}^m, \end{array}$$

where C is a nodal genus one curve, \overline{C}_1 and \overline{C}_2 are Gorenstein curves of genus one, the map $C \rightarrow \mathbb{P}^m[s]$ is a logarithmic map to an s -fold expansion of \mathbb{P}^m along H , and each of the vertical maps does not contract the minimal subcurve of genus one.

The information involving singularities adds substantial complexity to the theory, and this motivates our next result, which shows that calculations are nonetheless possible.

Theorem B. Given a smooth pair (X, Y) with Y very ample, there is an explicit recursive algorithm to calculate

- (1) the (restricted) reduced genus one Gromov–Witten invariants of Y ;
- (2) the reduced genus one relative Gromov–Witten invariants of (X, Y) ;
- (3) the (restricted) reduced genus one rubber invariants of $\mathbb{P} = \mathbb{P}_Y(\mathcal{N}_{Y|X} \oplus \mathcal{O}_Y)$

from the absolute genus zero and absolute reduced genus one invariants of X .

This reconstruction result generalises that of Gathmann [Gat02]. A higher genus reconstruction result for the ordinary theory is proved by Maulik and Pandharipande [MP06], but follows a fundamentally different strategy to ours. In particular, localisation is never used in the analysis here, nor in the work of Gathmann.

We pursue a strategy laid out by Vakil, building on work of Caporaso and Harris [CH98, Vak00]. The first step is to express the locus of maps where the contact order degenerates in terms of Chern classes of tautological vector bundles. The second step is to describe that locus in terms of fibre products of moduli spaces with smaller invariants.

Novelties are introduced in both steps. In the first, we describe the locus of degenerate maps as the zero locus of a logarithmic line bundle, arising from a piecewise linear function with a simple moduli theoretic description. The use of tropical techniques substantially simplifies Gathmann’s calculations.

For the second step, we are forced to contend with the interaction of the relative splitting formula and the geometry of elliptic singularities, which is entirely new (§4.3). We express the factorisation condition – central to the constructions in [RSPW19a, RSPW19b] – in terms of tautological classes, leading to Theorem B.

0.2. Context and further discussion. This paper is most directly related to recent work on aligned logarithmic structures, elliptic singularities, and stable maps, as developed in [RSPW19a, RSPW19b]. The moduli spaces we construct lie in between the absolute and toric-boundary-relative geometries studied in the first and second of those papers, respectively. The degeneration algorithm presented here is a first practical implementation of the theory developed in that work.

We pass through general techniques that we expect will be useful in logarithmic Gromov–Witten theory more broadly. We explain how to describe degenerate moduli spaces in terms of simpler spaces of maps. This is substantially more delicate than the standard situation, where the strata are simply fibre products. First, one must understand the interaction of the relative geometry with elliptic singularities, which involves a study of the factorisation condition in terms of tautological bundles. Second, the degenerate strata of moduli spaces are not simply fibre products of spaces with smaller numerical invariants, but compactified torus bundles over these strata, due to the introduction of logarithmic blowups. This geometry is at the heart of recent work on the degeneration formula [Ran19]. The main text provides a first systematic procedure that uses tropical geometry to calculate with these torus bundles.

A key step in our recursion is a description of the locus of maps with higher than prescribed tangency, which was identified by Gathmann. We realise the locus as the vanishing locus of a section of a line bundle that comes from tropical geometry — from a piecewise linear function on the fan/tropicalization. The systematic understanding of logarithmic line bundles arising in this fashion is expected to play an important role in logarithmic enumerative geometry.

Although the geometric ideas in Gathmann’s recursion are clear, they are implemented within the space of ordinary stable maps, and do not formally fit within the formalism of logarithmic Gromov–Witten theory. On the way to our main results, we reinterpret Gathmann’s idea of increasing tangency within logarithmic mapping spaces.

0.3. Future directions. The results of [RSPW19a] also suggest the existence of a reduced higher genus Gromov–Witten theory formed by replacing contracted elliptic components with singularities. We have been informed by Jonathan Wise that his Ph.D. candidate Sebastian Bozlee has constructed these higher genus reduced Gromov–Witten invariants. A manuscript is in preparation. The discussion here is expected to carry over to this setting.

Our analysis leads naturally to the study of the main component of the double ramification cycle for target manifolds in genus one. The virtual geometry of this was studied in recent work of Janda, Pandharipande, Pixton, and Zvonkine [JPPZ18]. Even in genus one, an understanding of the main component contributions of the double ramification cycle for target manifolds seems to be lacking. Our main algorithm provides a method for calculating integrals over this class, but it would be more natural to have a formula for it.

Finally, it would be natural to generalise the method of degenerating tangency conditions to targets with higher rank logarithmic structures. The extreme case when the target is a toric variety with its toric boundary, is intimately tied with the enumerative geometry of well-spaced tropical curves [LR18, RSPW19b, Spe14].

Acknowledgements. We have benefited from fruitful conversations with friends and colleagues, including Pierrick Bousseau, Francesca Carocci, Cristina Manolache, Daves Maulik, Keli Santos-Parker, and Jonathan Wise. The comments of an anonymous referee have proved very valuable. The authors worked on this project during visits to MPIM-Bonn, MIT, University of Cambridge, and the

University of Glasgow, and thank these institutions for ideal working conditions and support. L.B. and N.N. would in addition like to thank Imperial College London, the LSGNT, the Royal Society, 3CinG, and the MSRI program “Enumerative Geometry beyond Numbers”. N.N. was partially supported by EPSRC grant EP/R009325/1. L.B. is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy EXC-2181/1 - 390900948 (the Heidelberg STRUCTURES Cluster of Excellence).

1. CONSTRUCTIONS AND LOGARITHMIC SMOOTHNESS

We briefly review some basic tropical structures in the theory of logarithmic curves and logarithmic stable maps. A detailed treatment is given by [CCUW20, Section 3]; the following paragraphs are meant only as a reminder.

1.1. Curves and tropical curves. A nodal curve C over $\text{Spec } \mathbb{C}$ has an associated dual graph $\Gamma(C)$. Given a toric monoid P , there is a logarithmic point $\text{Spec}(P \rightarrow \mathbb{C})$ which may be thought of as remembering the information of the embedding of the torus-fixed closed point into the affine toric variety $\text{Spec } \mathbb{C}[P]$. For a logarithmically smooth curve C over $\text{Spec}(P \rightarrow \mathbb{C})$, the dual graph $\Gamma(C)$ is enhanced: given an edge e of $\Gamma(C)$, the logarithmic structure keeps track of a generalised “edge length” ℓ_e , which is an element of P . These data can be repackaged into a cone complex \square together with a map to the dual cone σ_P

$$\pi : \square \rightarrow \sigma_P.$$

A fibre of this map π is a metric space enhancing the topological space $\Gamma(C)$, with edges given lengths in $\mathbb{R}_{\geq 0}$. Over the interior of the cone σ_P these edge lengths are nonzero, and the fibres are homoeomorphic to $\Gamma(C)$. Over the faces of σ_P , the fibres can be identified with edge contractions.

The discussion is globalized in [CCUW20], and the construction can be carried out for the universal curve over the moduli space of logarithmic curves. Specifically, there is a stack $\mathfrak{M}_{g,n}^{\text{trop}}$ over the category of cone complexes equipped with a universal curve

$$\square \rightarrow \mathfrak{M}_{g,n}^{\text{trop}}.$$

The reader is encouraged to think of $\mathfrak{M}_{g,n}^{\text{trop}}$ as playing the same role for $\mathfrak{M}_{g,n}$ as a fan plays for a toric variety. In particular, two constructions are relevant:

- (1) A subdivision of $\mathfrak{M}_{g,n}^{\text{trop}}$ gives rise to a birational modification of $\mathfrak{M}_{g,n}$.
- (2) A piecewise linear function on $\mathfrak{M}_{g,n}^{\text{trop}}$ gives rise to a Cartier divisor on $\mathfrak{M}_{g,n}$.

1.2. Maps and tropical maps. The discussion carries over to the setting of logarithmic maps. Specifically, suppose we are given a logarithmic curve C and a logarithmic map to a smooth pair (X, Y) over a logarithmic scheme S . For simplicity, assume that S is a P -logarithmic point, as above. There is an associated **tropicalization**, i.e. a family of tropical curves $\square \rightarrow \sigma_P$, together with a map of cone complexes

$$\square \rightarrow \mathbb{R}_{\geq 0}.$$

Here $\mathbb{R}_{\geq 0}$ is the tropicalization (or “fan”) of the smooth pair (X, Y) . Each fibre over σ_P encodes a metric enhancement of the dual graph, together with a piecewise linear map to $\mathbb{R}_{\geq 0}$. There is once again a cone stack representing the moduli problem of tropical maps; the technical details are only cosmetically different than those presented in [CCUW20], and we leave these details to the reader (see also [GS13]).

1.3. Maps from curves and elliptic singularities. We provide a brief intuitive picture that underlies the work in [RSPW19a, RSPW19b]. If C is a Gorenstein curve of genus one and $f : C \rightarrow \mathbb{P}^m$ is a morphism, the ampleness of the tangent bundle of \mathbb{P}^m ensures that if f is non-constant on the minimal subcurve of C of arithmetic genus one, then the deformations of f are unobstructed; in fact, the deformations *over the stack of curves* are also unobstructed. It is natural to attempt to resolve the moduli space of stable maps, by contracting the genus one subcurve on which the map is constant. Whilst the moduli space of Gorenstein genus one curves is not smooth, a more sophisticated moduli problem, consisting of nodal curves C together with a *chosen contraction* to a Gorenstein curve \bar{C} , might be expected to resolve these singularities. The formation of such a moduli problem requires logarithmic geometry, and is achieved in [RSPW19a, Section 3]. Once constructed, the corresponding moduli of stable maps out of a Gorenstein curve is also resolved. Note that maps from nodal genus one curves may have stable limits that contract genus one subcurves, but the introduction of Gorenstein singularities ensures properness while disallowing contracted genus one components.

When working with logarithmic targets such as (\mathbb{P}^m, H) , as we do here, similar ideas may be expected to apply, but the deformation theory of maps becomes more delicate. Indeed, the logarithmic Euler sequence for this pair shows that the logarithmic tangent bundle is a quotient of $\mathcal{O} \oplus \mathcal{O}(1)^{\oplus m}$, and additional obstructions appear. In simple terms: there exist logarithmic stable maps to the pair (\mathbb{P}^m, H) with obstructed logarithmic deformations, whose underlying stable map to \mathbb{P}^m is unobstructed.

1.4. Singularities and moduli of attachments. Isolated Gorenstein singularities of genus one were classified by Smyth in [Smy11a]: for every fixed number of branches m , there is precisely one isomorphism type of curve germ. For $m = 1$ and 2 these are the cusp (A_2) and tacnode (A_3) respectively; for $m \geq 3$ they are the union of m general lines through the origin of \mathbb{A}^{m-1} – following Smyth, we refer to these as *elliptic m -fold points*.

The problem of constructing an elliptic singularity from its pointed normalisation is more delicate; it was widely studied in van der Wyck’s thesis [vdW10]. Recall that the seminormalisation C^{sn} of a curve C has the same underlying topological space, but its singularities consist of rational m -fold points only, i.e. they are locally isomorphic to the union of the axes in \mathbb{A}^m .

Lemma 1.1. The seminormalisation $\nu : C^{\text{sn}} \rightarrow C$ of an elliptic m -fold point is the collapse of a general tangent vector v at the rational m -fold point.

This is equivalent to [Smy11a, Lemma 2.2], where he describes the seminormalisation of an elliptic singularity as the data of a hyperplane in the Zariski cotangent space not containing the coordinate lines. We call *moduli of attaching data* [Smy11b, §2.2] (or *crimping space*) the extra data needed to produce a singularity from its pointed normalisation.

Corollary 1.2. The moduli of attachments of an elliptic m -fold point is isomorphic to \mathbb{G}_m^m .

One way to compactify it is to allow *sprouting* on the boundary, i.e. replacing a branch R_i of the singularity with a strictly semistable component \mathbb{P}^1 with the singularity at 0 and the branch R_i glued nodally at ∞ .

There are other ways to look at these data; we recall a construction from linear algebra. Let $V = \sum_{i=1}^m L_i$ be a presentation of a vector space as the direct sum of m lines. Then the data of a general line ℓ in V (or a general hyperplane in V^\vee) is equivalent to the data of compatible isomorphisms $\theta_{ij} : L_i \cong L_j$. Indeed, given a general line ℓ , a vector $v \in \ell$ can be written as:

$$v = a_1 e_1 + \dots + a_m e_m, \quad \text{with } a_i \neq 0 \text{ for all } i$$

for a fixed basis $\{e_i \in L_i\}$ of V . We can then define θ_{ij} by

$$\theta_{ij}(e_i) = (a_j / a_i) \cdot e_j$$

and it is easy to see that it does not depend on any of the choices. On the other hand, given the θ_{ij} , the line ℓ can be taken to be the graph of the morphism:

$$(\theta_{12}, \dots, \theta_{1m}): L_1 \rightarrow L_2 \oplus \dots \oplus L_m.$$

Thus the attaching data can be viewed as specifying an identification of tangent spaces of the pointed normalisation at the singularity.

Lemma 1.1 provides a valuable characterisation of maps from an elliptic singularity C , which we will exploit in §4:

Corollary 1.3. A map $C \rightarrow X$ is the same as a map $C^{\text{sn}} \rightarrow X$ such that its differential sends $v \mapsto 0$. Line bundles on an elliptic m -fold point admit a similar description, see e.g. [Liu02, Lemma 7.5.12].

Lemma 1.4. Let C be a curve of arithmetic genus one containing an elliptic m -fold point. Then $\text{Pic}^0(C) \cong \mathbb{G}_a$.

1.5. Absolute geometry: background. Let $\mathfrak{M}_{1,n}$ be the logarithmic algebraic stack of n -pointed prestable curves of genus one. We begin by recalling the logarithmic moduli spaces of genus one curves constructed in [RSPW19a, §2 & §4].

Consider a logarithmically smooth curve C over a logarithmic geometric point S with monoid P and let $\pi: \sqsubset \rightarrow \sigma_P$ denote its tropicalization. Given a point t in σ_P we obtain a metric graph as explained above. Since the curve in question has genus one, there is a minimal genus one subgraph – either a vertex of genus one or a cycle in the graph. We will refer to this as the **core**, denoted by E below.

Given a point s with $\pi(s)$ equal to t , there is a well-defined distance in the metric graph $\pi^{-1}(t)$ from s to the core. In particular, for each vertex v of the dual graph $\Gamma(C)$, i.e. each component of C , there is a well-defined function $\lambda(v): \sigma_P \rightarrow \mathbb{R}_{\geq 0}$, defined on the base of the tropical family. We may as well think of λ as a piecewise linear function $\lambda: \sqsubset \rightarrow \mathbb{R}_{\geq 0} \times \sigma_P$, see Figure 1. Given any other linear function δ on σ_P , say that δ is **comparable** to $\lambda(v)$ if there is a global order relation, either $\lambda(v) \leq \delta$ or $\lambda(v) \geq \delta$, that holds at all points of σ_P .

Piecewise linear functions on \sqsubset are naturally identified with global sections of the characteristic sheaf \overline{M}_C ; linear functions on σ_P are naturally identified with global sections of the characteristic sheaf \overline{M}_S , and can be pulled back to C . Given an arbitrary logarithmically smooth curve C/S , conditions involving the comparability of piecewise linear functions can be imposed at the level of geometric points; they determine subdivisions of σ_P , and, in turn, logarithmic modifications of S .

Definition 1.5. A **centrally aligned curve** is a pair $(C/S, \delta)$, where $\delta \in \overline{M}_S$, such that:


- (1) the section δ is comparable to $\lambda(v)$ for all vertices v of \sqsubset ;
- (2) for any pair of vertices v and w of distance less than δ , the sections $\lambda(v)$ and $\lambda(w)$ are comparable.

In other words, for the tropical curve \sqsubset , all vertices are either inside, outside, or on the circle of radius δ regardless of the point on the base; the collection of vertices inside the circle is totally ordered by their distance to the core.

The moduli stack $\mathfrak{M}_{1,n}^{\text{cen}}$ of centrally aligned curves is a logarithmic algebraic stack in the smooth topology, and $\mathfrak{M}_{1,n}^{\text{cen}} \rightarrow \mathfrak{M}_{1,n}$ is a logarithmic modification. The main construction of [RSPW19a] canonically associates to any centrally aligned family of curves \mathcal{C}_S a partial destabilisation $\tilde{\mathcal{C}}_S$, introducing 2-pointed components, and a contraction

$$\tilde{\mathcal{C}}_S \rightarrow \overline{\mathcal{C}}_S,$$

where $\overline{\mathcal{C}}_S$ may contain a Gorenstein elliptic singularity [Smy11a]. Intuitively, this is achieved by interpreting the exceptional directions of $\mathfrak{M}_{1,n}^{\text{cen}} \rightarrow \mathfrak{M}_{1,n}$ as moduli of attaching data.


 FIGURE 1. The function λ (distance from the core) and its cutoff value δ .

Formally, given a family of tropical curves, the function δ provides a non-negative real number for each point on the base. We may therefore consider the “**circle**” of points on each fibre whose distance from the core is equal to δ . A partial destabilization, or “bubbling”, introduces 2-pointed rational curves into the curve by adding vertices to the dual graph at the points that lie on this circle (the crosses at height δ in Figure 1); we denote this new tropical curve by $\tilde{\mathcal{C}}$. The open disc of radius δ , i.e. the locus where $\lambda < \delta$, is then contracted to a Gorenstein singularity in $\tilde{\mathcal{C}}_S$. The number of branches is equal to the number of vertices of $\tilde{\mathcal{C}}$ which lie on the circle.

The space of stable maps $\widetilde{\mathcal{VZ}}_{1,n}(\mathbb{P}^m, d)$ from the universal nodal curve \mathcal{C} over the stack $\mathfrak{M}_{1,n}^{\text{cen}}$ is Deligne-Mumford and proper, with projective coarse moduli space. A compatibility condition between the alignment and the map is required, namely that the circle of radius δ passes through *at least one component of positive degree*. A stable map in this space is said to satisfy the **factorisation condition** if the map $\tilde{\mathcal{C}} \rightarrow \mathcal{C} \rightarrow \mathbb{P}^m$ factors through the contraction $\tilde{\mathcal{C}} \rightarrow \bar{\mathcal{C}}$. Maps that are not constant on any genus one subcurve of \mathcal{C} automatically satisfy this condition; indeed, the contraction is an isomorphism in this case.

Theorem 1.6 ([RSPW19a, Theorem B]). The substack $\mathcal{VZ}_{1,n}(\mathbb{P}^m, d)$ of $\widetilde{\mathcal{VZ}}_{1,n}(\mathbb{P}^m, d)$ parameterising maps to \mathbb{P}^m that satisfy the factorisation condition is smooth and proper of the expected dimension.

Remark 1.7 (Constant maps). When the degree of the map is 1, maps from smooth genus one curves to \mathbb{P}^m are constant. The interior of the moduli space of stable maps in this case is empty, though the compactification by stable maps is non-empty. The space of maps satisfying the factorisation condition is also empty. We do not indulge on this any further.

The remainder of this section lifts this construction to the category of relative maps to (\mathbb{P}^m, H) . The core result shows that two factorisation conditions remove the obstructions.

1.6. Relative geometry: compactification. Fix a hyperplane $H \subseteq \mathbb{P}^m$. Let α be a partition of the degree $d > 0$. Consider the moduli space $\mathcal{M}_{1,\alpha}^\circ(\mathbb{P}^m|H, d)$ of maps from smooth genus one curves $C \rightarrow \mathbb{P}^m$ that meet H at finitely many marked points with vanishing orders given by the partition α . This is a smooth, but non-proper Deligne–Mumford stack.

We compactify the space described above, and then desingularise it. For the compactification, we begin with Abramovich–Chen–Gross–Siebert’s space of logarithmic stable maps, though we will work with variants [AC14, Che14b, GS13, Kim10].

Let \mathbb{P}^m be endowed with the divisorial logarithmic structure induced by H . The moduli space $\overline{\mathcal{M}}_{1,\alpha}^{\log}(\mathbb{P}^m|H, d)$ is a fibred category over logarithmic schemes, whose fibre over (S, M_S) is the groupoid of logarithmically smooth curves of genus one over (S, M_S) equipped with a logarithmic map to \mathbb{P}^m of degree d and contact order α . It is a fundamental fact in the subject that this

category is representable by a proper algebraic stack with logarithmic structure, parameterising minimal objects [Che14b].

1.7. Relative geometry: expansions. In order to make the obstruction theory more geometric, we expand the target. To elucidate the connection with the static target, consider a logarithmic stable map $[C \rightarrow (\mathbb{P}^m, H)]$ over $\text{Spec}(\mathbb{N} \rightarrow \mathbb{C})$. At the level of tropicalizations, there is a map of cone complexes:

$$\square \rightarrow \mathbb{R}_{\geq 0}.$$

Choose a subdivision of $\mathbb{R}_{\geq 0}$ whose vertices are the images of vertices of \square . Pull this subdivision back to \square by subdividing it along the preimage of the vertices of $\mathbb{R}_{\geq 0}$. Denote the resulting map $\tilde{\square} \rightarrow \tilde{\mathbb{R}}_{\geq 0}$. These subdivisions induce logarithmic modifications:

$$\tilde{C} \rightarrow \mathbb{P}^m[s],$$

see [AW18]. Here the latter is the s -times iterated deformation to the normal cone of H in \mathbb{P}^m . The irreducible components of the target are in bijection with the vertices in $\tilde{\mathbb{R}}_{\geq 0}$. The curve is modified by adding rational components for the newly introduced vertices.

The result is a logarithmic stable map to an expanded target $\mathbb{P}^m[s]$, together with a contraction to the main target component $\mathbb{P}^m[s] \rightarrow \mathbb{P}^m$. Two maps are considered equivalent if they differ by \mathbb{G}_m -scaling the target at higher level. This is the logarithmic enhancement of the classical theory of relative stable maps, due to Li [Li01, Li02].

Globally, Kim constructs a moduli space of logarithmic maps to expanded degenerations, which on logarithmic points, reduces to the above construction [Kim10]. Kim's space is identified with a subcategory of the Abramovich–Chen–Gross–Siebert space (as stacks over logarithmic schemes), and its dual minimal monoids are cones in a subdivision of the dual minimal cones of the unexpanded space, see [Ran19, § 2].

We typically work with expansions for convenience, and use the notation $\overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H, d)$ for Kim's space of logarithmic stable maps.

1.8. Relative geometry: alignment. We consider the space of logarithmic morphisms from the universal curve $\mathcal{C} \rightarrow \mathfrak{M}_{1,n}^{\text{cen}}$ to the logarithmic scheme (\mathbb{P}^m, H) . As in the absolute case, we require the radius to pass through a vertex of positive degree in \mathbb{P}^m and contain only vertices of degree zero in its strict interior. In this way we obtain a logarithmic modification of the Abramovich–Chen–Gross–Siebert space. If we further perform the subdivisions of target and source outlined in the previous section, we obtain a logarithmic modification of Kim's space:

$$\widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H, d) \rightarrow \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H, d)$$

which we refer to as the **moduli space of centrally aligned maps** to (\mathbb{P}^m, H) . This space parametrises transverse logarithmic maps to the expanded target with an alignment which is compatible with the map. Because of this compatibility condition, there is a morphism

$$\widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H, d) \rightarrow \widetilde{\mathcal{VZ}}_{1,n}(\mathbb{P}^m, d)$$

obtained by forgetting the logarithmic structure on the target and collapsing the map.

1.9. Relative geometry: factorisation. Let $[C \rightarrow \mathbb{P}^m[s] \rightarrow \mathbb{P}^m]$ be a centrally aligned logarithmic map. Recall that $\mathbb{P}^m[s]$ consists of a union of \mathbb{P}^m with s copies of the projective bundle $\mathbb{P} = \mathbb{P}(\mathcal{O}_H \oplus \mathcal{O}_H(1))$. We refer to these latter components as **the higher levels**. Thus, we will say that a subcurve $D \subseteq C$ **maps to higher level** if the composite $C \rightarrow \mathbb{P}^m$ maps D into $H \subset \mathbb{P}^m$.

Let $D_F \subseteq C$ be the maximal genus one subcurve which is mapped to a higher level and contracted by the map $C \rightarrow \mathbb{P}^m[s]$. Let δ_F denote the distance from the core to the nearest component of $C \setminus D_F$. Similarly let $D_B \subseteq C$ be the maximal genus one subcurve that is contracted by the collapsed

map to \mathbb{P}^m , and let δ_B be the associated radius. This coincides with the radius of the underlying map to \mathbb{P}^m . Of course, $\delta_F \leq \delta_B$, so (C, δ_F) is centrally aligned as well.

The datum (δ_F, δ_B) determines a destabilisation \widetilde{C} of C together with successive contractions $\widetilde{C} \rightarrow \overline{C}_F \rightarrow \overline{C}_B$. The contractions are constructed by [RSPW19a, Section 3].

Recall that we name *core* the minimal genus one subcurve of a Gorenstein curve of arithmetic genus one, so the latter decomposes into the nodal union of the core and a (possibly empty) rational forest. If the curve contains an elliptic m -fold point, we say a map is non-constant on the core if at least one branch of the singularity is not contracted. The following condition identifies the locus of smoothable maps.

Definition 1.8. The map $f: C \rightarrow \mathbb{P}^m[s]$ **factors completely** if

- (1) f factors through \overline{C}_F and it is non-constant on its core,
- (2) letting $f_B: C \rightarrow \mathbb{P}^m$ denote the composite of f with the collapsing map $\mathbb{P}^m[s] \rightarrow \mathbb{P}^m$, the map f_B factors through \overline{C}_B and it is non-constant on its core.

Remark. The factorisation conditions are understood to be satisfied when $C \rightarrow \mathbb{P}^m[s]$ (resp. $C \rightarrow \mathbb{P}^m$) does not contract a genus one subcurve.

Let

$$\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d) \subseteq \widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H, d)$$

be the stack of maps from centrally aligned curves to expansions that factor completely.

If $[C \rightarrow \mathbb{P}^m[s] \rightarrow \mathbb{P}^m]$ is a family of centrally aligned maps over S that factors completely, there is a moduli map $S \rightarrow \mathcal{VZ}_{1,n}(\mathbb{P}^m, d)$, to the principal component of the space of absolute maps.

Theorem 1.9. The stack $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ is logarithmically smooth and proper of the expected dimension.

As noted in Remark 1.7, maps from elliptic curves to \mathbb{P}^m will always be taken to have degree at least 2, ensuring that the main component is non-empty.

Proof. We examine the geometry of the space of maps to $(\mathbb{P}^m|H)$ by utilising the forgetful morphism

$$\nu: \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d) \rightarrow \mathcal{VZ}_{1,n}(\mathbb{P}^m, d),$$

which remembers only the stabilisation of the collapsed map. To see that such a forgetful morphism exists, we have already observed that, for a map $[C \rightarrow \mathbb{P}^m[s] \rightarrow \mathbb{P}^m]$, the stabilised composite map is certainly centrally aligned; moreover, it satisfies the factorisation condition by assumption.

The moduli space $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ is proper. Indeed, the verification of the valuative criterion follows from Vakil's Lemma [Vak00, Lemma 5.9]. A direct proof that the factorisation property is a closed condition may be found in [RSPW19a, Theorem 4.3].

We come to logarithmic smoothness. If the elliptic curve does not map into higher level, then deformations of the map are unobstructed exactly as in [RSPW19a, Section 4.5].

Assume now that the elliptic curve maps into higher level. Note that the relative logarithmic tangent bundle of the expansion, over the base \mathbb{P}^m is a trivial line bundle of rank 1, as the fibres are toric. Consider an S -family of maps. After replacing the source C by a destabilisation, we have maps

$$\begin{array}{ccccc} C & \longrightarrow & \overline{C}_F & \longrightarrow & \overline{C}_B \\ & & \downarrow & & \downarrow \\ & & \mathbb{P}^m[s] & \longrightarrow & \mathbb{P}^m \end{array}$$

factoring completely. Let f_B and f be the maps from C to the collapsed and expanded target respectively. Examining the morphism ν , we see that there is a map

$$\mathrm{Def}(C, f_B) \rightarrow \mathrm{Obs}(f/(C, f_B)) = H^1(\overline{C}_F, \mathcal{O}_{\overline{C}_F}),$$

where the former is the space of logarithmic deformations of the curve and map to the collapsed target, as a centrally aligned map satisfying simple factorisation.

More precisely, consider a strict square-zero extension of S , and a deformation of the map (C, f_B) satisfying the factorisation condition. Since the deformation is strict, there is a canonically induced deformation of the expansion $\mathbb{P}^m[s]$. The corresponding element in $\mathrm{Obs}(f/(C, f_B))$ is the obstruction to lifting the deformation of (C, f_B) to this canonical deformation of $\mathbb{P}^m[s]$ over \mathbb{P}^m . By comparing the cotangent complexes of the source, target, and fibre of the morphism ν , we see that the cokernel of this map are the absolute logarithmic obstructions for the map. We will show these vanish, proving logarithmic smoothness.

Consider the manner in which $H^1(\overline{C}_F, \mathcal{O}_{\overline{C}_F})$ functions as the obstruction group for the lifting. Near the minimal genus one subcurve D_F of C , in order to lift the map from \mathbb{P}^m to the expansion, we must provide a rational function on D_F with poles bounded by the divisor associated to the tropicalization, i.e. if α is the piecewise linear function giving this tropical map, the lifting is described by a section of the associated bundle $\mathcal{O}_C(-\alpha)$.

Given a strict square-zero extension S' of S , the piecewise linear function α extends uniquely by strictness to any deformation of the curve. The resulting deformation of $\mathcal{O}_C(-\alpha)$ produces an infinitesimal deformation of the trivial bundle in the Picard group, whose class in $H^1(\overline{C}_F, \mathcal{O}_{\overline{C}_F})$, is the obstruction to deformation. We will show that it is possible to choose a deformation of (C, f_B) that cancels out the obstruction to lifting to f , and thus the map to the obstruction space above is surjective.

On restricting to the minimal genus one subcurve D_F , the line bundle $\mathcal{O}_C(-\alpha)$ is equivalent to the divisor $\sum a_i x_i$ where x_i are the points connecting D_F to the rest of the curve, and a_i are the slopes of α along the edges corresponding to the x_i . Infinitesimally moving the point x_i produces a nontrivial deformation of the line bundle $\mathcal{O}_C(-\alpha)$. The moduli space $\mathcal{VZ}_{1,n}(\mathbb{P}^m, d)$ is a logarithmically smooth fibration over the universal Picard stack over the stack of centrally aligned n -pointed genus one curves [RSPW19a, Remark 4.5.3]. As a consequence, it is possible to find a logarithmic deformation of the map (C, f_B) that induces the deformation of the line bundle that is given by infinitesimal motion. The obstruction group is 1-dimensional, and we have produced a logarithmic deformation of the collapsed map that produces a nontrivial deformation of the line bundle $\mathcal{O}_C(-\alpha)$. Therefore, logarithmic deformations surject onto the obstruction space above. The absolute obstructions, which are the cokernel of this map therefore vanish, and the result follows. **luca's rewrite of log smoothness**

The proof is analogous to that of [RSPW19b, Theorem 3.5.1]. There is a forgetful morphism

$$\nu: \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d) \rightarrow \mathcal{VZ}_{1,n}(\mathbb{P}^m, d),$$

remembering only the stabilisation of the composite $f_B: C \rightarrow \mathbb{P}^m[s] \rightarrow \mathbb{P}^m$. A priori, this is in $\overline{\mathcal{M}}_1(\mathbb{P}^m, d)$, but in fact by assumption (C, δ_B) is centrally aligned, and f_B factors through \overline{C}_B , as we have already remarked.

The expanded target $\mathbb{P}^m[s]$ is logarithmically étale over \mathbb{P}^m with the divisorial log structure associated to H , so it is equivalent to study the deformation theory of the unexpanded map to (\mathbb{P}^m, H) . The logarithmic morphism $(\mathbb{P}^m, H) \rightarrow \mathbb{P}^m$ induces an exact sequence of log tangent bundles:

$$0 \rightarrow T_{\mathbb{P}^m}(-\log H) \rightarrow T_{\mathbb{P}^m} \rightarrow \mathcal{O}_H(H) \rightarrow 0,$$

relating the absolute and relative obstruction theories. This shows that ν can be obstructed only when a genus one subcurve is mapped to higher level in such a way that it is contracted by the

collapsed map f_B . In this case, let C_0 denote the core, and let E be the connected component of $f_B^{-1}(f_B(C_0))$ of arithmetic genus one.

Let $f: C \rightarrow \mathbb{P}^m[s]$ over the base S be a possibly obstructed map to the expansion that factors completely, and let S' be a strict square-zero extension of the base. The deformation problem is local, so we may as well suppose that $\overline{M}_S^{\text{gp}}$ is a constant sheaf of monoids with stalk $\overline{M}_{S,s}^{\text{gp}} =: Q$, for a geometric point $s \in S$. Note that sections of M_S such as δ_B and δ_F extend automatically due to strictness. Moreover, it is equivalent to deform f or $f_F: \overline{C}_F \rightarrow \mathbb{P}_S^m$, where \overline{C}_F is given the trivial logarithmic structure around the elliptic singularity.

The map $f_F: \overline{C}_F \rightarrow \mathbb{P}^m[s]$ induces a tropical map $\alpha: \square_F \rightarrow \mathbb{R}_{\geq 0} \times \sigma_Q$, that we want to think of as a section $\alpha \in H^0(\overline{C}_F, \overline{M}_C^{\text{gp}})$. With notation as above, we obtain a trivialisation of the line bundle $\mathcal{O}_{\overline{C}_F}(-\alpha)$ upon restriction to E . By [RSPW19a, Proposition 2.4.1], the latter is isomorphic to $\mathcal{O}_E(\sum \mu_i x_i)$, where μ_i is the slope/expansion factor of α along the edge e_i of \square_F , and x_i the node corresponding to e_i thought of as a smooth marking on E .

The deformation problem for f_F relative to (C', δ_B, f'_B) reduces to the problem of extending α as a trivialisation of $\mathcal{O}_E(\sum \mu_i x_i)$. This gives an interpretation of $\text{Obs}(f/(C, \delta_B, f_B))$ as $H^1(\overline{C}_F, \mathcal{O})$. The latter is a vector space of dimension 1. Associated to the forgetful morphism ν at the level of perfect obstruction theories, we find a map:

$$(1) \quad \text{Def}(C, \delta_B, f_B) \rightarrow \text{Obs}(f/(C, \delta_B, f_B)).$$

The left-hand side contains in particular the space $\text{Def}(E, \mathbf{x}) = \text{Ext}^1(\Omega_E(\mathbf{x}), \mathcal{O}_E)$, since E is contracted by f_B . Since by assumption at least one of the μ_i is non-zero, $\text{Ext}^1(\Omega_E(\mathbf{x}), \mathcal{O}_E)$ surjects onto $H^1(\overline{C}_F, \mathcal{O})$ by [RSPW19b, Corollary 3.5.3]. Looking back at (1), we see that obstructions to deforming f can be killed by an appropriate choice of lift (C', δ_B, f'_B) . \square

The theorem guarantees that the singularities of the space of completely factoring maps to expansions is logarithmically smooth. One can say more in this case.

Corollary 1.10. The logarithmically smooth stack $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ has at worst orbifold singularities, i.e. admits a non-representable cover by a smooth Deligne–Mumford stack.

Luca’s rewrite of the Corollary: Up to a generalised root construction, $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ is a smooth Deligne–Mumford stack.

Proof. Since $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ is logarithmically smooth, it will suffice to show that the cones of its tropicalization are simplicial. To see this, consider a logarithmic stable map to an expansion $C \rightarrow \mathbb{P}^m[s]$ without a central alignment. The tropical moduli cone obtained as the dual of the minimal base monoid can be identified with $\mathbb{R}_{\geq 0}^s$, see for instance [Che14a, Section 2.2]. The alignment procedure is an iterated barycentric subdivision at the level of tropical moduli spaces, as explained in [RSPW19a, Section 4.6], and such subdivisions preserve the property of being simplicial. We conclude from this that the blowup $\widehat{\mathcal{VZ}}_{1,\alpha}^{\text{exp}}(\mathbb{P}^m|H, d)$ obtained by centrally aligning Kim’s spaces has simplicial cones. The morphism $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d) \rightarrow \widehat{\mathcal{VZ}}_{1,\alpha}^{\text{exp}}(\mathbb{P}^m|H, d)$ is strict, so the cones of $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ are simplicial, as claimed. \square

1.10. Rubber variants. We require a “rubber” variant of the space of maps $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$, where the curve is contained in the higher levels.

Let \mathbb{P} denote the projective bundle $\mathbb{P}(\mathcal{O}(1) \oplus \mathcal{O})$ on $H \cong \mathbb{P}^{N-1}$. Equip this space with the logarithmic structure coming from the 0 and ∞ sections of the bundle. Consider the moduli space $\mathcal{VZ}_{1,\alpha}^{\leftarrow}(\mathbb{P}|H_0 + H_\infty, d)$ of logarithmic maps

$$C \rightarrow \mathbb{P}[s] \rightarrow \mathbb{P}^{N-1},$$

which factor completely, and where automorphisms are taken to cover the identity on the map to \mathbb{P}^{N-1} ; two maps that differ by a \mathbb{G}_m translate in the fibre direction are considered equivalent. As previously, the map to the bundle and the map to the base \mathbb{P}^{N-1} both factor through possibly different singularities.

Theorem 1.11. The stack $\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}|H_0+H_\infty, d)$ is logarithmically smooth and proper of the expected dimension.

The main point in the proof is that on the space of non-degenerate maps with smooth domains, the map from $\mathcal{VZ}_{1,\alpha}^{\circ}(\mathbb{P}|H_0+H_\infty, d) \rightarrow \mathcal{VZ}_{1,\alpha}^{\leftrightarrow,\circ}(\mathbb{P}|H_0+H_\infty, d)$ is a \mathbb{G}_m -torsor. One expects that the map on compact moduli spaces is a nodal curve fibration. To prove this, it is convenient to work with the logarithmic multiplicative group and its torsors. Recall that the **logarithmic multiplicative group** \mathbb{G}_{\log} is the functor on logarithmic schemes whose value on a logarithmic scheme S is the group of global sections $H^0(S, M_S^{\text{gp}})$; it is a proper group functor, and it contains \mathbb{G}_m as a subfunctor (from $\mathcal{O}_S^\times \subseteq M_S^{\text{gp}}$). This functor is representable only after a logarithmically étale cover, see for instance [MW20, RW19].

Each fibre of the bundle $\mathbb{P} \rightarrow \mathbb{P}^{N-1}$ admits a natural log étale morphism to \mathbb{G}_{\log} . Replacing these fibres by \mathbb{G}_{\log} , we obtain a non-representable functor on logarithmic schemes \mathbb{P}_{\log} and a logarithmically étale modification

$$\mathbb{P} \rightarrow \mathbb{P}_{\log}.$$

Proof of Theorem 1.11. Consider the stack over logarithmic schemes $\mathcal{VZ}_{1,\alpha}(\mathbb{P}_{\log}, d)$ of stable logarithmic maps that factor completely. Here, stability for the map coincides with stability for the projection to \mathbb{P}^{N-1} . The arguments in the previous section show that maps to \mathbb{P} that factor completely are unobstructed. Since this space is a logarithmically étale cover of $\mathcal{VZ}_{1,\alpha}(\mathbb{P}_{\log}, d)$, logarithmic smoothness of the latter follows.

The logarithmic multiplicative group \mathbb{G}_{\log} acts on $\mathcal{VZ}_{1,\alpha}(\mathbb{P}_{\log}, d)$ by translation. Tautologically, $\mathcal{VZ}_{1,\alpha}(\mathbb{P}_{\log}, d)$ is a \mathbb{G}_{\log} -torsor over the moduli problem of maps up to this \mathbb{G}_{\log} translation. It is representable after a logarithmically étale modification by a family of nodal curves. It follows that the space $\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}_{\log}, d)$ of maps up to \mathbb{G}_{\log} -translation that factor completely is logarithmically smooth.

To conclude the result, we compare $\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}_{\log}, d)$ and $\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}|H_0 + H_\infty, d)$. Since a map to $(\mathbb{P}, H_0 + H_\infty)$ gives rise immediately to a map to \mathbb{P}_{\log} , there is a morphism

$$\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}|H_0 + H_\infty, d) \rightarrow \mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}_{\log}, d).$$

The lifting criterion shows that this map is logarithmically étale, so the result follows. \square

1.11. Hypersurface pairs. Let (X, Y) be a smooth pair with Y very ample. The definition in §1.9 applies in this setting and yields a moduli space $\mathcal{VZ}_{1,\alpha}(X|Y, \beta)$ with a morphism

$$\mathcal{VZ}_{1,\alpha}(X|Y, \beta) \rightarrow \overline{\mathcal{M}}_{1,\alpha}^{\log}(X|Y, \beta)$$

obtained by performing a logarithmic modification and passing to a closed substack where the factorisation property is satisfied. This moduli space will typically be non-equidimensional, but we equip it with a virtual class as follows. The divisor Y defines an embedding $X \hookrightarrow \mathbb{P}^m$, with $Y = X \cap H$ for some hyperplane H .

Lemma 1.12. The following square is cartesian in the category of ordinary stacks:

$$\begin{array}{ccc} \mathcal{VZ}_{1,\alpha}(X|Y, \beta) & \longrightarrow & \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d) \\ \downarrow & \square & \downarrow \\ \mathcal{VZ}_{1,n}(X, \beta) & \xrightarrow{i} & \mathcal{VZ}_{1,n}(\mathbb{P}^m, d). \end{array}$$




FIGURE 2. An example of well-spacedness: condition (2) becomes $\ell_1 = \ell_2$.

Proof. It is clear from the modular description that this square is cartesian in the category of fs logarithmic stacks, and since i is strict, it is also cartesian in ordinary stacks. \square

Since $\mathcal{VZ}_{1,n}(\mathbb{P}^m, d)$ is smooth and $\mathcal{VZ}_{1,n}(X, \beta)$ carries a natural virtual class [RSPW19a, Theorem 4.4.1], the diagonal of $\mathcal{VZ}_{1,n}(\mathbb{P}^m, d)$ defines a pullback [BN20, Appendix A], and therefore the virtual class on the space of maps to (X, Y) :

$$[\mathcal{VZ}_{1,\alpha}(X|Y, \beta)]^{\text{virt}} := i_{\Delta}^! [\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)].$$

Note that this is parallel to the virtual formalism in [Gat02]. As $\mathcal{VZ}_{1,\alpha}(X|Y, \beta)$ is equipped with evaluation maps and cotangent lines on the universal curve, we arrive at a definition of reduced Gromov–Witten invariants for the pair (X, Y) by the standard mechanics.

2. STRATIFICATION AND TROPICALIZATION

The logarithmic smoothness of Theorem 1.9 implies that $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ is an orbifold toroidal embedding. The irreducible components of its boundary stratify the space, as we now describe.

The tropicalizations of logarithmic maps that factor completely are combinatorially constrained by a condition known as **well-spacedness**. The version we use here is essentially identical to [RSPW19b].

Definition 2.1. Let \square be a tropical curve of genus one and let E be its minimal subcurve of genus one. A tropical map $F : \square \rightarrow \mathbb{R}_{\geq 0}$ is said to be **well-spaced** if one of the following two conditions are satisfied:

- (1) no open neighbourhood of E is contracted to a point in $\mathbb{R}_{>0}$, or
- (2) if an open neighbourhood of E is contracted and t_1, \dots, t_k are the flags whose vertex is mapped to $F(E)$, but along which F has nonzero slope; then, the minimum of the distances from E to a vertex supporting t_i occurs for at least two indices i .

Remark. Condition (2) is not relevant as far as rays of the tropical moduli space (i.e. divisors) are concerned, since there are always at least two parameters involved, namely $F(E)$ - or the distance from $F(E)$ to the origin of $\mathbb{R}_{\geq 0}$ - and the distance from E to the first flag of non-zero slope within \square . As we mostly concentrate on divisors starting from §3, it will not play much of a role in the rest of the paper.

Proposition 2.2. Let $[C \rightarrow \mathbb{P}^m]$ be a logarithmic stable map from a centrally aligned curve to an expansion, that factors completely. Then the tropicalization $\square \rightarrow \mathbb{R}_{\geq 0}$ is well-spaced.

Proof. The required result is essentially contained in [RSPW19b, Section 4], so we explain how to deduce it from the one in loc. cit. We may focus on a single component of the expansion $\mathbb{P}^m[s]$ that contains the image of a contracted genus one subcurve, as this is the only relevant case. We may thus replace the target with the projective bundle $\mathbb{P}(\mathcal{O}(1) \oplus \mathcal{O})$ over \mathbb{P}^{N-1} equipped with the divisorial logarithmic structure from the 0 and ∞ sections. Let p be the point to which the genus one subcurve is contracted. Passing to an open neighbourhood of p , the map to the bundle is given by a rational function f on an open curve C° , determined by the subgraph formed by E and the flags t_i described in the definition above. To describe the tropical map to $\mathbb{R}_{>0}$, we observe that E is contracted to a fixed point $q \in \mathbb{R}_{>0}$. The flags at t_i correspond to nodes or markings of C . The pole orders of f at these distinguished points determine the slopes of the tropical map. We are now exactly in the situation considered in [RSPW19b, Second Paragraph of § 4.6], and Lemma 4.6.1 of loc. cit. guarantees the well-spacedness. \square

2.1. The cone complex. To understand the stratification via combinatorial data, we will build the stratification from known objects.

Step 1. Let Σ^{\log} be the tropical moduli space of genus one tropical stable maps to $\mathbb{R}_{\geq 0}$. This is naturally identified with the tropicalization (in the logarithmic sense) of the Abramovich–Chen–Gross–Siebert space of logarithmic stable maps to the pair (\mathbb{P}^m, H) .

Step 2. Given such a tropical stable map, we may subdivide $\mathbb{R}_{\geq 0}$ such that the image of every vertex of \square is a vertex of $\mathbb{R}_{\geq 0}$. Call this subdivision $\widetilde{\mathbb{R}}_{\geq 0}$. The preimages of vertices of the subdivision form a subdivision of \square . After this procedure, the images of vertices of \square which lie in $\mathbb{R}_{>0}$ are totally ordered, in a manner extending the partial order obtained from the map to $\mathbb{R}_{\geq 0}$. The combinatorial types of such **image-ordered** maps produce the cones of a subdivision of Σ^{\log} which we denote Σ^{Kim} . As the notation suggests, Σ^{Kim} is the tropicalization of Kim’s moduli space of logarithmic maps to (\mathbb{P}^m, H) .

Step 3. Given a tropical map $F : \square \rightarrow \mathbb{R}_{\geq 0}$ parameterised by Σ^{Kim} , there is a largest radius δ (possibly equal to 0) such that every vertex strictly contained in the circle of radius δ around the core has degree-marking $d = 0$. Let Σ^{cen} be the subdivision obtained by requiring that the vertices contained within the circle of radius δ around the core of \square are totally ordered. This involves introducing cones along which certain \mathbb{Z} -linear combinations of edge lengths are identified.

Step 4. Let $\Sigma_{1,\alpha}(\mathbb{P}^m|H, d)$ be the subcomplex of Σ^{cen} consisting of well-spaced tropical maps.

Proposition 2.3. The cone complex $\Sigma_{1,\alpha}(\mathbb{P}^m|H, d)$ is the fan of the toroidal embedding $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$. In particular, the codimension k strata of $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ are in inclusion reversing bijection with the dimension k cones in $\Sigma_{1,\alpha}(\mathbb{P}^m|H, d)$.

Proof. The construction above has been given to mimic the construction of the space $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$. Specifically, the fact that Σ^{cen} is the cone complex attached to the logarithmic stack $\widehat{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H, d)$ of centrally aligned maps to expansions, follows from its description as a subcategory of the fibred category (over logarithmic schemes) of $\mathcal{M}_{1,\alpha}^{\log}(\mathbb{P}^m|H, d)$. To complete the result, note that $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ has a strict map to $\widehat{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H, d)$, so its cone complex is a subcomplex of Σ^{cen} . The fact that it is contained in the subcomplex of well-spaced curves is immediate from Proposition 2.2. \square

2.2. Indexing the strata. The dimension- k cones in $\Sigma_{1,\alpha}(\mathbb{P}^m|H,d)$ can be enumerated as follows. First, the cones in $\Sigma = \Sigma^{\text{Kim}}$ are indexed by **combinatorial types** of tropical maps to $\mathbb{R}_{\geq 0}$. Here a combinatorial type encodes all of the data of a tropical map, except for the edge lengths. To be more precise, a combinatorial type Δ consists of:

- (1) a finite graph \square and topological (not metric) subdivision $\widetilde{\mathbb{R}}_{\geq 0}$ of $\mathbb{R}_{\geq 0}$;
- (2) genus, degree and marking assignments on the vertices;
- (3) the data of which vertex of $\widetilde{\mathbb{R}}_{\geq 0}$ each vertex of \square is mapped to;
- (4) integral slopes along the edges (both finite and infinite).

The corresponding cone $\sigma \in \Sigma^{\text{Kim}}$ is then given by the resulting moduli space of tropical maps, given by choices of edge lengths on source and target which produce a continuous tropical map. Given $\sigma \in \Sigma^{\text{Kim}}$ we then produce all the cones in $\Sigma_{1,\alpha}(\mathbb{P}^m|H,d)$ mapping to σ by performing Steps 3–4 outlined above. This amounts to a subdivision of σ . In this process, new cones are created which map into larger-dimensional cones of Σ^{Kim} . The process for enumerating the codimension k strata of $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$ is therefore:

- (1) fix a combinatorial type Δ ;
- (2) perform the subdivision of the resulting cone $\sigma \in \Sigma^{\text{Kim}}$;
- (3) identify the dimension k cones of that subdivision.

In §4 we will perform this analysis in the case $k = 1$.

3. DEGENERATION OF CONTACT ORDER

Consider the moduli space $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$ of relative maps that factor completely.

Definition 3.1. For each marking x_k , consider the locus $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d) \subseteq \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$ where x_k belongs to an **internal component** of the collapsed map, i.e. a component of the curve that is mapped into H .

In this section, we use the logarithmic structure on $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$ to construct a line bundle \mathcal{L}_k and section s_k which vanishes precisely along $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d)$. We use the correspondence with tropical geometry to identify $c_1(\mathcal{L}_k)$ in terms of tautological classes on $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$, and to compute the vanishing order of s_k along the components of $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d)$. Combined with the relative splitting formulas from the next section, this produces a recursion relation inside $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$.

The pair (\mathcal{L}_k, s_k) is most naturally constructed on the moduli space $\overline{\mathcal{M}}_{1,\alpha}^{\log}(\mathbb{P}^m|H,d)$ of non-expanded logarithmic maps; the corresponding pair on $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$ will be obtained via pull-back. Consider therefore the tropicalization Σ^{\log} of $\overline{\mathcal{M}}_{1,\alpha}^{\log}(\mathbb{P}^m|H,d)$, identified as usual with the moduli space of tropical maps to $\mathbb{R}_{\geq 0}$. We have a universal family

$$\begin{array}{ccc} \square & \xrightarrow{f} & \mathbb{R}_{\geq 0} \\ \downarrow \pi & & \\ \Sigma^{\log} & & \end{array} \quad \begin{array}{c} x_k \curvearrowright \end{array}$$

where x_k is the section which for every point $\lambda \in \Sigma^{\log}$ picks out the vertex of \square_λ supporting the leg x_k . The composition $f \circ x_k: \Sigma^{\log} \rightarrow \mathbb{R}_{\geq 0}$ defines a piecewise-linear function on Σ^{\log} whose preimage over the open cone $\mathbb{R}_{>0}$ consists of those tropical maps where x_k belongs to an internal component. This produces a section of the ghost sheaf on $\overline{\mathcal{M}}_{1,\alpha}^{\log}(\mathbb{P}^m|H,d)$, which in the usual way induces a line bundle and section (\mathcal{L}_k, s_k) on the moduli space, and the tropical description above shows that the zero locus of s_k is (set-theoretically) the locus where x_k belongs to an internal component.

We now calculate $c_1(\mathcal{L}_k)$.

Theorem 3.2. $c_1(\mathcal{L}_k) = \alpha_k \psi_k + \text{ev}_k^* H$.

Proof. Choose a family of logarithmic stable maps over S and let $\mu \in \Gamma(S, \overline{\mathcal{M}}_S)$ be the global section of the ghost sheaf constructed in the previous paragraph. This pulls back along π to give a global section $\pi^b(\mu) \in \Gamma(C, \overline{\mathcal{M}}_C)$. Interpreted as a piecewise-linear function on the tropicalization \square with values in $\overline{\mathcal{M}}_S$ [CCUW20, Remark 7.3], this assigns μ to every vertex and has slope zero along every edge. By construction, the line bundle associated to this section is $\pi^* \mathcal{L}_k$. Consider on the other hand the generator $1 \in \mathbb{N} = \Gamma(\mathbb{P}^m, \overline{\mathcal{M}}_{\mathbb{P}^m})$ with associated line bundle $\mathcal{O}(H)$. The section $f^b(1) \in \Gamma(C, \overline{\mathcal{M}}_C)$ has associated line bundle $f^* \mathcal{O}(H)$. If we let v denote the vertex containing x_k , then by construction $f^b(1)$ assigns μ to v and has slope α_k along the leg x_k . Thus if we consider the difference $f^b(1) - \pi^b(\mu)$ then this assigns 0 to v and still has slope α_k along x_k . Thus by [RSPW19a, Proposition 2.4.1] the corresponding line bundle restricted to C_v is given by

$$\mathcal{O}_{C_v} \left(\alpha_k x_k + \sum_e \mu_e x_e \right)$$

where the sum is over the edges e adjacent to v and distinct from x_k . Thus we have:

$$(f^* \mathcal{O}(H) \otimes \pi^* \mathcal{L}_k^{-1})|_{C_v} = \mathcal{O}_{C_v} \left(\alpha_k x_k + \sum_e \mu_e x_e \right).$$

Since x_k factors through C_v we may pull back along x_k to obtain

$$\mathcal{L}_k = x_k^* \pi^* \mathcal{L}_k = x_k^* \mathcal{O}_{C_k}(-\alpha_k x_k) \otimes x_k^* f^* \mathcal{O}(H) = x_k^* \mathcal{O}_{C_k}(-\alpha_k x_k) \otimes \text{ev}_k^* \mathcal{O}(H)$$

and taking Chern classes gives

$$c_1(\mathcal{L}_k) = \alpha_k \psi_k + \text{ev}_k^* H,$$

as required. \square

This produces (\mathcal{L}_k, s_k) and calculates $c_1(\mathcal{L}_k)$ on $\overline{\mathcal{M}}_{1,\alpha}^{\log}(\mathbb{P}^m|H, d)$; this bundle is pulled back to $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$. The relevant piecewise-linear function is the composition:

$$\Sigma_{1,\alpha}(\mathbb{P}^m|H, d) \rightarrow \Sigma^{\log} \rightarrow \mathbb{R}_{\geq 0}.$$

Note in particular that ψ_k should be interpreted as a **collapsed class** on $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$, i.e. the cotangent class from the source curve of the collapsed and stabilised map. We arrive at the recursion.

Theorem 3.3. We have the following relation in the Chow ring of $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$:

$$(2) \quad (\alpha_k \psi_k + \text{ev}_k^* H) \cap [\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)] = \sum_{\mathcal{D}} \lambda_{\mathcal{D}} [\mathcal{D}].$$

The sum is over irreducible components of the divisor $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d) \subseteq \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$, and $\lambda_{\mathcal{D}}$ is the vanishing order of s_k along this component.

Remark 3.4. This construction gives the natural logarithmic analogue of Gathmann's line bundle and section [Gat02, Construction 2.1]. A benefit of the logarithmic approach is to make the computation of vanishing orders entirely combinatorial (see §4.2.4 below), circumventing the difficult technical calculation given by Gathmann.

4. SPLITTING THE BOUNDARY

A basic phenomenon is responsible for the nature of the forthcoming analysis. In contrast with nodal singularities, the functions on an elliptic singularity are not simply collections of functions on the normalisation that agree at the gluing point. Rather, the contraction map $C \rightarrow \overline{C}$ corresponds to a particular linear combination of tangent vectors at the pointed normalisation, whose kernel consists of those functions descending to the elliptic singularity (see §1.4). In order to compute integrals, it is necessary to give a tautological description of this additional condition.

We provide an explicit description of the terms appearing in the right-hand side of (2). For each term, we provide a combinatorial formula for the vanishing order $\lambda_{\mathcal{D}}$, and a recursive description of the stratum \mathcal{D} in terms of fibre products of moduli spaces with smaller numerical data. The tautological description of the factorisation condition then leads to a recursive structure for calculating integrals. This step is the heart of the analysis.

Remark 4.1. For the reduced relative Gromov–Witten theory of (\mathbb{P}^m, H) with only primary insertions, i.e. where all cohomological insertions in the invariants are pulled back along evaluation maps, computations can be carried out without an analysis of the factorisation condition; this follows from dimension considerations and leads to the main results in [Vak00]. However if we wish to include descendant insertions, work with more general pairs, or produce cycle-level statements, an analysis of the tautological classes arising from the factorisation condition is crucial.

4.1. Irreducible components of the degenerate locus. We explain the tropical procedure to identify irreducible components of $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d)$.

Lemma 4.2. Every irreducible component of $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d)$ is a codimension 1 logarithmic stratum.

Proof. The locus where the logarithmic structure is trivial is the locus where the source curve is smooth and not mapped inside H . Since $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$ is toroidal this locus is open and dense. By definition $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d)$ is contained in the complement of this locus and can be expressed as a union of strata of positive codimension. Since $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d)$ is pure codimension 1, it is a union of logarithmic divisors. \square

In §2.2 we described the logarithmic strata of $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$, using the cones of the tropicalization $\Sigma_{1,\alpha}(\mathbb{P}^m|H, d)$. Every divisorial logarithmic stratum is obtained as follows:

- (1) choose a combinatorial type Δ of a tropical map;
- (2) subdivide the corresponding tropical moduli space σ to align the type;
- (3) choose a ray in this subdivision.

This process contains redundancies: upon choosing a ray in the tropical moduli space, some of the edge lengths or vertex positions may get set to 0. This induces a generisation of the initial combinatorial type Δ , given by contracting the corresponding edges of the dual graph and moving the corresponding vertices from $\mathbb{R}_{>0}$ to 0. When we speak of the combinatorial type of a stratum in $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$, we will always mean this generisation. This is independent of the choice of Δ , and in fact we can and will always choose Δ to coincide with the generisation.

The logarithmic divisors contained in $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d)$ are those whose combinatorial types map the vertex of the dual graph containing x_k into the interior $\mathbb{R}_{>0} \subseteq \mathbb{R}_{\geq 0}$. Thus via the above procedure, we enumerate the components of $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H, d)$ in a combinatorial manner; the combinatorial type Δ allows us to describe the general element of such a component. The remainder of this section gives a recursive description of the components and a recursive computation of integrals over them.

4.2. Recursive description of the divisors: types I, II and III. Choose an irreducible component $\mathcal{D} \subseteq \mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d)$. By Lemma 4.2 this is a logarithmic divisor, and hence may be written as:

$$\mathcal{D} = \widetilde{\mathcal{D}} \cap \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$$

for a unique logarithmic stratum $\widetilde{\mathcal{D}} \subseteq \widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H,d)$. Since

$$\widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H,d) \rightarrow \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d)$$

is a logarithmic modification, the divisor $\widetilde{\mathcal{D}}$ is either exceptional or the strict transform of a logarithmic divisor. By the nature of the subdivision procedure these two cases correspond, respectively, to when the core is assigned zero degree or nonzero degree by the combinatorial type.

We begin with the latter. Suppose therefore that $\widetilde{\mathcal{D}}$ is the strict transform of a logarithmic divisor:

$$\mathcal{E} \subseteq \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d).$$

The birational map $\widetilde{\mathcal{D}} \rightarrow \mathcal{E}$ induces a morphism $\mathcal{D} \rightarrow \mathcal{E}$. We now show how to interpret this morphism as a desingularisation of the principal component. Since \mathcal{E} admits a recursive description in terms of relative and rubber moduli spaces, this will allow us to compute integrals over \mathcal{D} .

Lemma 4.3. Let $\mathcal{D} \subseteq \mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d)$ be an irreducible component which contributes nontrivially to the Gromov–Witten invariant, and let Δ be the corresponding combinatorial type. If Δ assigns positive degree to the core, then Δ takes one of the following forms:




FIGURE 3. I




FIGURE 4. II




FIGURE 5. III

where black vertices represent rational components, white vertices components of genus one. The degrees of vertices, expansions factors, and remaining markings are distributed arbitrarily, subject to the following constraints:

- (1) the vertices $\mathbb{C}_1, \dots, \mathbb{C}_r$ have positive degree;
- (2) the core has positive degree;
- (3) every vertex is stable;
- (4) the balancing condition is satisfied.

In [Vak00] these cases are referred to, respectively, as types A, B and C^+ .

Proof. Let $\sigma \in \Sigma_{1,\alpha}(\mathbb{P}^m|H,d)$ be the cone corresponding to $\mathcal{E} \subseteq \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d)$. By the discussion above \mathcal{D} corresponds to a ray in the subdivision of σ obtained by imposing the central alignment condition. Since the elliptic core is assigned positive degree, both radii are equal to 0 and the subdivision is trivial. We conclude that $\sigma = \mathbb{R}_{\geq 0}$. Since there must be at least one vertex mapped to higher level, the tropical target $\widetilde{\mathbb{R}}_{\geq 0}$ is obtained from $\mathbb{R}_{\geq 0}$ by subdividing at a single point $\diamond \in \mathbb{R}_{>0}$.

In order for the cone σ to be 1-dimensional, the dual graph Γ must be bipartite, with vertices over 0 and \diamond . The cases I, II and III enumerated above cover situations where there is a single vertex mapped to \diamond . If there is more than one such vertex, then the contribution to the Gromov–Witten invariant vanishes. To see this, we consider the stratum

$$\mathcal{E}^{\log} \subseteq \overline{\mathcal{M}}_{1,\alpha}^{\log}(\mathbb{P}^m|H,d)$$

to which \mathcal{E} maps under the collapsing morphism, and examine the composition $\mathcal{D} \rightarrow \mathcal{E} \rightarrow \mathcal{E}^{\log}$. If we let \mathcal{F}^{\log} denote the intersection of \mathcal{E}^{\log} with the main component of the moduli space, then we obtain a factorisation $\mathcal{D} \rightarrow \mathcal{F}^{\log} \hookrightarrow \mathcal{E}^{\log}$. Since the moduli space is generically unobstructed along \mathcal{F}^{\log} , the codimension of \mathcal{F}^{\log} is given by the dimension of the associated cone in the tropicalization Σ^{\log} . If there is more than one vertex mapped to higher level, then this cone has dimension ≥ 2 . Therefore the map $\mathcal{D} \rightarrow \mathcal{F}^{\log}$ has positive-dimensional fibres, and since all insertions are pulled back from the latter space the contribution vanishes by the projection formula. \square

Remark 4.4. The difference in dimensions between \mathcal{D} and \mathcal{F}^{\log} (or equivalently, between virtual dimensions of \mathcal{E} and \mathcal{E}^{\log}) may be interpreted as the difference in dimensions between moduli spaces of *disconnected* rubber and their images under the collapsing morphisms.

We now investigate the three types I, II and III separately, giving a recursive description of the boundary divisor in each case. For the remainder of this subsection, therefore, we fix a one-dimensional cone $\tau \in \Sigma_{1,\alpha}(\mathbb{P}^m|H,d)$, let $\mathcal{D} \subseteq \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$ be the associated logarithmic divisor, and assume that \mathcal{D} is contained in $\mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d)$ and is of type I, II or III. We let $\mathcal{E} \subseteq \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d)$ be the logarithmic stratum into which \mathcal{D} is mapped; this is indexed by a cone $\epsilon \in \Sigma^{\text{Kim}}$ corresponding to a combinatorial type Δ , and ϵ is one-dimensional since we are restricting to the type I, II, III cases.

4.2.1. *Type I.* Suppose Δ is of type I. Then \mathcal{E} admits a finite and surjective **splitting morphism** onto the fibre product:

$$\mathcal{E} \rightarrow \left(\overline{\mathcal{M}}_{1,\alpha^{(1)} \cup (m_1)}(\mathbb{P}^m|H,d_1) \times \prod_{i=2}^r \overline{\mathcal{M}}_{0,\alpha^{(i)} \cup (m_i)}(\mathbb{P}^m|H,d_i) \right) \times_{H^r} \overline{\mathcal{M}}_{0,\alpha^{(0)} \cup (-m_1, \dots, -m_r)}^{\leftrightarrow}(\mathbb{P}|H_0 + H_\infty, d_0).$$

Lemma 4.5. The divisor \mathcal{D} admits a natural splitting morphism

$$\mathcal{D} \xrightarrow{\rho} \left(\mathcal{VZ}_{1,\alpha^{(1)} \cup (m_1)}(\mathbb{P}^m|H,d_1) \times \prod_{i=2}^r \overline{\mathcal{M}}_{0,\alpha^{(i)} \cup (m_i)}(\mathbb{P}^m|H,d_i) \right) \times_{H^r} \overline{\mathcal{M}}_{0,\alpha^{(0)} \cup (-m_1, \dots, -m_r)}^{\leftrightarrow}(\mathbb{P}|H_0 + H_\infty, d_0)$$

such that the map $\mathcal{D} \rightarrow \mathcal{E}$ lifts the map on fibre products obtained by desingularising the main component of the first factor.

Proof. We first consider the statement of the lemma in the category of fine but not necessarily saturated logarithmic schemes, so that a relative stable map to an expansion admits a unique logarithmic lifting. The fibre product description preceding the lemma is an isomorphism in this category [AMW14, Lemma 4.2.2].

The map $\mathcal{D} \rightarrow \mathcal{E}$ is an isomorphism away from the exceptional centres, and the latter is contained in the locus where the core is contracted. Given an element of \mathcal{D} we can split it along the nodes q_1, \dots, q_r . It is then clear that \sqsubset_1 is aligned. We claim that \sqsubset satisfies the factorisation property if and only if \sqsubset_1 does. This immediately implies the lemma.

On \sqsubset there are two contraction radii δ_F, δ_B . Let $\delta \in \{\delta_F, \delta_B\}$. An examination of the stratification by combinatorial type of \mathcal{D} shows that $\lambda(\sqsubset') > \delta$ for any component \sqsubset' of \sqsubset_0 . This means that $\sqsubset_0, \sqsubset_2, \sqsubset_3, \dots, \sqsubset_r$ lie outside both contraction radii, and so the aligned curve \sqsubset satisfies the factorisation condition if and only if \sqsubset_1 does. \square

Lemma 4.5 provides a means to calculate integrals over the class $\lambda_{\mathcal{D}}[\mathcal{D}]$ appearing on the right-hand side of Theorem 3.3, provided that we can calculate $\lambda_{\mathcal{D}}$ and the degree of the splitting morphism. Simple closed formulae for these are given in §4.2.4 below.

4.2.2. *Type II.* Now suppose Δ is of type II. In this case, it is impossible for the core to be contracted. Hence \mathcal{E} is disjoint from the blown-up locus, and the map $\mathcal{D} \rightarrow \mathcal{E}$ is an isomorphism. We obtain the following description, entirely in terms of genus zero data:

Lemma 4.6. \mathcal{D} admits a finite and surjective splitting morphism:

$$\mathcal{D} \xrightarrow{\rho} \left(\overline{\mathcal{M}}_{0,\alpha^{(1)} \cup (m_1, m_2)}(\mathbb{P}^m | H, d_1) \times \prod_{i=3}^r \overline{\mathcal{M}}_{0,\alpha^{(i)} \cup (m_i)}(\mathbb{P}^m | H, d_i) \right) \times_{H^r} \overline{\mathcal{M}}_{0,\alpha^{(0)} \cup (-m_1, \dots, -m_r)}^{\leftrightarrow}(\mathbb{P} | H_0 + H_\infty, d_0).$$

4.2.3. *Type III.* Finally suppose that Δ is of type III. As before we have a finite and surjective morphism:

$$\mathcal{E} \rightarrow \left(\prod_{i=1}^r \overline{\mathcal{M}}_{0,\alpha^{(i)} \cup (m_i)}(\mathbb{P}^m | H, d_i) \right) \times_{H^r} \overline{\mathcal{M}}_{1,\alpha^{(0)} \cup (-m_1, \dots, -m_r)}^{\leftrightarrow}(\mathbb{P} | H_0 + H_\infty, d_0).$$

The same arguments as in §4.2.1 then apply to give:

Lemma 4.7. \mathcal{D} admits a finite splitting morphism

$$\mathcal{D} \xrightarrow{\rho} \left(\prod_{i=1}^r \overline{\mathcal{M}}_{0,\alpha^{(i)} \cup (m_i)}(\mathbb{P}^m | H, d_i) \right) \times_{H^r} \mathcal{VZ}_{1,\alpha^{(0)} \cup (-m_1, \dots, -m_r)}^{\leftrightarrow}(\mathbb{P} | H_0 + H_\infty, d_0)$$

such that $\mathcal{D} \rightarrow \mathcal{E}$ corresponds to the obvious map on fibre products.

The final factor in the fibre product is the logarithmic blowup of the moduli space of rubber maps constructed in §1.10. We will compute integrals over such spaces as part of the recursion (see §5).

4.2.4. *Splitting degree and vanishing order.* In each of the subsections 4.2.1–4.2.3 above, we obtained a finite splitting morphism ρ from \mathcal{D} to a fibre product of moduli spaces with smaller numerical data. Here we describe the degree of ρ and calculate the vanishing order $\lambda_{\mathcal{D}}$ of the section s_k constructed in §3. This gives a complete recursive description of the terms appearing on the right-hand side of Theorem 3.3 which are of type I, II or III.

Lemma 4.8. The degree of ρ is given by:

$$\frac{\prod_{i=1}^r m_i}{\text{lcm}(m_1, \dots, m_r)}.$$

Proof. This calculation is well known, see for instance [Che14a, Section 7.9], [ACGS17, Section 5.3]. \square

Lemma 4.9. The section s_k vanishes along the divisor \mathcal{D} with order given by

$$\lambda_{\mathcal{D}} = \text{lcm}(m_1, \dots, m_r).$$

Proof. Let $\tau \in \Sigma_{1,\alpha}(\mathbb{P}^m | H, d)$ be the cone corresponding to \mathcal{D} and let

$$\varphi: \Sigma_{1,\alpha}(\mathbb{P}^m | H, d) \rightarrow \Sigma^{\log} \rightarrow \mathbb{R}_{\geq 0}$$

be the piecewise-linear function constructed in §3. It follows from the tropical description that $\lambda_{\mathcal{D}}$ is equal to the index of the map of integral cones $\tau \rightarrow \mathbb{R}_{\geq 0}$ obtained by restricting φ . Observe that $\tau \subseteq \mathbb{R}_{\geq 0}^r$ with integral generator:

$$w = \left(\frac{\text{lcm}(m_1, \dots, m_r)}{m_1}, \dots, \frac{\text{lcm}(m_1, \dots, m_r)}{m_r} \right).$$

The map $\tau \rightarrow \mathbb{R}_{\geq 0}$ is given by projecting onto the i th factor and then multiplying by m_i . Note that by piecewise linear continuity of the tropical map, this is independent of i . We conclude that the index is equal to $\text{lcm}(m_1, \dots, m_r)$ as claimed. \square

In summary, the contribution of each term $\lambda_{\mathcal{D}}[\mathcal{D}]$ is given by integrating over the appropriate fibre product and then multiplying the result by $\lambda_{\mathcal{D}} \cdot \deg \rho = \prod_{i=1}^r m_i$.

4.3. Recursive description of the divisors: type \dagger . The treatment of the type I, II and III cases above is a fairly straightforward extension of the ideas of Vakil and Gathmann (albeit updated with logarithmic and tropical technology, which streamlines the presentation and simplifies the proofs). The remaining type \dagger case, however, is entirely new. In this section we provide a recursive description of these boundary divisors. This constitutes the technical heart of the paper.

4.3.1. Possible combinatorial types. A boundary divisor $\mathcal{D} \subseteq \mathcal{D}_{1,\alpha,k}(\mathbb{P}^m|H,d)$ is said to have **type \dagger** if the core is assigned degree zero by the combinatorial type. Divisors of this type occur as principal components in compactified torus bundles over strata in the space of relative maps. As a result, their combinatorial types exhibit very degenerate behaviour. Precisely, over the generic point of such a divisor, the target may expand multiple times, and the source tropical curve need not be bipartite with a single interior vertex; see the left-hand side of Figure 8 for a concrete example. We stress once again that such degenerate contributions are at the heart of the reduced theory and cannot be removed.

We begin with a series of lemmas describing the combinatorial types of rays associated to divisorial loci \mathcal{D} of type \dagger contributing nontrivially to the Gromov–Witten invariants.

Lemma 4.10. For every nonzero vertex \diamond of the tropical target, the fibre of the tropical map over \diamond contains exactly one stable vertex, which either is the core, or lies on the circle.

Lemma 4.11. All the vertices on the circle are stable (as opposed to semistable vertices introduced by subdivision).

Proof of Lemma 4.10. Stability of maps to an expansion guarantees that at least one stable vertex lies over each vertex of the target. If there is more than one stable vertex, then we can show as in the proof of Lemma 4.3 that the corresponding locus \mathcal{F} in the moduli space of unexpanded maps has high codimension, i.e. the collapsing map has fibres of positive dimension. As both insertions and the factorisation property for the collapsed map to \mathbb{P}^m are pulled back from the moduli space of maps without expansions, the contribution vanishes by the projection formula. \square

Proof of Lemma 4.11. Since the corresponding family of tropical maps is one-dimensional, for every vertex of the target there must be at least one vertex above it which lies on the circle. If the vertex on the circle is strictly semistable – so that in particular the collapsed map f_B is constant along the corresponding component – the factorisation condition for f_B is unaffected by \mathbb{G}_m -scaling the coordinate corresponding to that vertex in the moduli space of attaching data [Smy11b, §2.2], so the map $\mathcal{D} \rightarrow \mathcal{E}$ has generic fibre of positive dimension and the contribution vanishes. \square

4.3.2. Lines. We begin with a description of the boundary divisor corresponding to a comb whose teeth all have degree 1. This is a concrete but atypical case, since the teeth are generically transverse (not just dimensionally) to the hyperplane H . For simplicity, we assume that the contact order is (d) , concentrated at a marking x_1 supported on the elliptic curve E contracted to H . The combinatorial type is as in Figure 6.

Start with the moduli space of logarithmic stable maps. There is a finite splitting morphism:

$$\mathcal{E} \rightarrow \left(\overline{\mathcal{M}}_{0,(1)}(\mathbb{P}^m|H,1)^{\times m} \times_{H^m} H \right) \times \overline{\mathcal{M}}_{1,(-1,\dots,-1,d)}^{\leftrightarrow}(\mathbb{P}^1|0+\infty,d),$$

where the latter is the moduli space of genus one rubber maps. To obtain $\tilde{\mathcal{D}}$ from \mathcal{E} we replace the final factor by $\mathcal{VZ}_{1,(-1,\dots,-1,d)}^{\leftrightarrow}(\mathbb{P}^1|0+\infty,d)$ (see §1.10).

The first factor can be described very concretely as the choice of a point p of H , and d lines through it. A very important observation is that, at least generically, the vertices which lie on the




FIGURE 6. The combinatorial type of a comb with degree 1 teeth.

circle of radius δ and map to $0 \in \mathbb{R}_{\geq 0}$ are automatically aligned. Indeed, if R_i denotes the i th tail of the curve, glued to E at the node q_i , then the stable map provides us with identifications:

$$df_{i,q_i}: T_{R_i,q_i} \cong N_{H/\mathbb{P}^m,p}$$

which we may compose to obtain attaching data θ_{ij} (see §1.4). This produces a contraction $C \rightarrow \bar{C}$ to an elliptic d -fold point. The dimension of $\tilde{\mathcal{D}}$ coincides with the dimension of \mathcal{E} , which is $dm+m-1$.

Assume first that $d \leq m$. For the absolute factorisation to hold, it is necessary (but not sufficient) for the d lines $f_i(R_i)$ to span a subspace of $T_{\mathbb{P}^m,p}$ of dimension at most $d-1$. The locus of maps satisfying this condition has dimension:

$$(dm+m-1) - (m-d+1) = dm+d-2.$$

Without loss of generality, we may assume that $H = \{z_0 = 0\}$, $p = [0, \dots, 0, 1]$, and that the d lines span the subspace $\{z_{d-1} = \dots = z_{m-1} = 0\}$. The sections z_i for $i = d-1, \dots, m$ vanish identically near E , so descend automatically to the singularity. On the other hand the section z_0 descends to the singularity by the construction of \bar{C} . Finally, the sections z_i for $i = 1, \dots, d-2$ impose independent conditions to descend to \bar{C} . Imposing these conditions, we find that \mathcal{D} has dimension

$$(dm+d-2) - (d-2) = dm$$

making it into a divisor in $\mathcal{VZ}_{1,(d)}(\mathbb{P}^m|H,d)$. The case $d > m$ is similar, except that the lines $f_i(R_i)$ are automatically linearly dependent and so this condition does not need to be imposed.

We can express the factorisation condition tautologically by combining the isomorphisms θ_{ij} above with the differential of f . We see that factorisation amounts to the vanishing of the following vector bundle map:

$$T_{R_1,q_1} \xrightarrow{(1,\theta_{12},\dots,\theta_{1d})} \bigoplus_{i=1}^d T_{R_i,q_i} \xrightarrow{\Sigma \circ df = dz_1 + \dots + dz_{d-2}} N_{H \cap \mathbb{P}^{d-1} | \mathbb{P}^{d-1}}|_p.$$

For strata with higher tangency, we will see that the automatic alignment only occurs for vertices on the circle which map to $0 \in \mathbb{R}_{\geq 0}$. For internal components which lie on the circle, the alignment must be imposed separately. This means that $\tilde{\mathcal{D}}$ is a compactified torus bundle over \mathcal{E} (see §1.4). On the other hand, for higher contact order, the vector bundle map $\Sigma \circ df$ to $T_{\mathbb{P}^m,p}$ factors naturally through $T_{H,p}$. This is discussed in detail in §§4.3.5 and 4.3.6.

4.3.3. *Recursive description of \mathcal{E} .* Recall that we have maps

$$\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d) \hookrightarrow \widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H,d) \rightarrow \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d).$$

The first morphism is a closed embedding cut out by the two factorisation properties. The second morphism is a logarithmic modification. Let $\tau \in \Sigma_{1,\alpha}(\mathbb{P}^m|H,d)$ be the ray corresponding to $\mathcal{D} \subseteq \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d)$, and to the virtual divisor $\widetilde{\mathcal{D}} \subseteq \widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H,d)$.

Let $\sigma \in \Sigma^{\text{Kim}}$ be the minimal cone containing the image of τ . The dimension of this cone could be large (see again the left-hand side of Figure 8) – its dimension is equal to the number of levels of the target expansion in the type Δ . The cone σ determines a stratum $\mathcal{E} \subseteq \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d)$, and the morphism above restricts to a morphism

$$\mathcal{D} \rightarrow \mathcal{E}.$$

Splitting in the ordinary geometry. The locus \mathcal{E} has the following description. Let Λ_0 denote the collection of stable vertices of the combinatorial type which are mapped to $0 \in \mathbb{R}_{\geq 0}$, and use $\Lambda_{>0}$ for the stable vertices which are mapped to $\mathbb{R}_{>0}$. Each vertex v has associated discrete data Γ_v (genus, degree, contact order). The vertices over 0 determine moduli spaces of relative stable maps, while the higher level vertices determine maps to rubber. The stratum \mathcal{E} admits a finite splitting morphism

$$(3) \quad \mathcal{E} \xrightarrow{\rho} \left(\prod_{v \in \Lambda_0} \overline{\mathcal{M}}_{\Gamma_v}(\mathbb{P}^m|H) \times \prod_{v \in \Lambda_{>0}} \overline{\mathcal{M}}_{\Gamma_v}^{\hookrightarrow}(\mathbb{P}|H_0 + H_\infty) \right) \times_{H^{2\epsilon}} H^\epsilon$$

where ϵ is the number of edges of the stabilised dual graph, and the fibre product is taken over the appropriate evaluation maps. Splitting the diagonal of $H^{2\epsilon}$ we see that tautological integrals over \mathcal{E} are completely determined in terms of integrals over moduli spaces with “smaller” combinatorial data.

Exportation to the reduced geometry. There is a virtual divisor $\widetilde{\mathcal{D}} \subseteq \widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H,d)$ such that

$$\mathcal{D} = \widetilde{\mathcal{D}} \cap \mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H,d).$$

We wish to use the splitting of \mathcal{E} to recursively compute integrals on \mathcal{D} . Noting that \mathcal{D} is cut out of the excess-dimensional space $\widetilde{\mathcal{D}}$ by the factorisation properties, we make the reduction in two steps:

- (1) in §4.3.4 we explain how to construct $\widetilde{\mathcal{D}}$ from \mathcal{E} ;
- (2) in §4.3.5 we explain how to express the class of $\mathcal{D} \subseteq \widetilde{\mathcal{D}}$ in terms of tautological classes.

To compute integrals over \mathcal{D} we simply push down the appropriate classes from $\widetilde{\mathcal{D}}$ to \mathcal{E} and then integrate over \mathcal{E} using the ordinary splitting axiom.

Remark 4.12 (Splitting degree and vanishing order, revisited). Similarly to §4.2.4 above, we can calculate the degree of ρ and the vanishing order $\lambda_{\mathcal{D}}$ of the section s_k along \mathcal{D} . The degree of ρ is given exactly as in Lemma 4.8, where now we obtain one such factor for each bounded edge of the tropical target. The vanishing order $\lambda_{\mathcal{D}}$ is slightly more combinatorially delicate, but may be computed as the index of an explicit morphism of one-dimensional lattices, as explained in the proof of Lemma 4.9.

4.3.4. *Compactified bundles over strata: $\widetilde{\mathcal{D}}$ from \mathcal{E} .* Recall that the map $\widetilde{\mathcal{D}} \rightarrow \mathcal{E}$ is obtained by restricting the logarithmic blowup

$$\widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H,d) \rightarrow \overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d)$$

to the locus $\widetilde{\mathcal{D}}$. Toroidal geometry gives a construction of $\widetilde{\mathcal{D}}$ as a compactification of a torus bundle over a locally closed stratum in $\overline{\mathcal{M}}_{1,\alpha}(\mathbb{P}^m|H,d)$.




FIGURE 7. From \mathcal{E} to $\tilde{\mathcal{E}}$. Typical subdivision of a two-dimensional cone σ along the image of a ray τ .

Logarithmic structures on strata. Let $W \subseteq X$ be an irreducible stratum in a toroidal embedding X . The stratum W is itself toroidal, or, equivalently, when equipped with the divisorial logarithmic structure coming from intersecting with the toroidal boundary of X , the resulting structure makes W logarithmically smooth. Note that this differs from the logarithmic structure pulled back from X along the inclusion of W into X .

If X and W are merely logarithmic and not toroidal, there is an analogue of this. There exists a strict map

$$X \rightarrow \mathcal{A}_X,$$

where \mathcal{A}_X is the Artin fan of X , which is logarithmically étale over a point. The logarithmic structure of \mathcal{A}_X is divisorial in the smooth topology. The Artin fan \mathcal{A}_X possesses closed strata. Fix an irreducible stratum \mathcal{A}_W . In identical fashion to the paragraph above, this stack \mathcal{A}_W possesses a logarithmic structure by restricting the divisors on \mathcal{A}_X and taking the induced logarithmic structure. This equips a logarithmic stratum $W \hookrightarrow X$ with a logarithmic structure by pullback from the Artin fan, mirroring the construction for toroidal embeddings above.

Remark 4.13. Note that the inclusion of a stratum $W \hookrightarrow X$ is *not* a logarithmic morphism. The pullback logarithmic structure on a closed stratum includes residual information about the (virtual) normal bundle directions; the construction above effectively removes these directions. Using the equivalence of Artin fans with cone complexes [CCUW20], the above construction can be combinatorialised. Given a cone σ in a cone stack Σ , its **star** consists of all cones of Σ that contain σ ; these naturally form a new cone complex. The associated Artin fan gives rise to the logarithmic structure on W above.

Equipped with the strata logarithmic structures, there is a logarithmic morphism

$$\tilde{\mathcal{D}} \rightarrow \mathcal{E}$$

whose generic fibre is a toric variety.

Flattening the map to \mathcal{E} . The morphism $\tilde{\mathcal{D}} \rightarrow \mathcal{E}$ is not flat. In order to push integrals forward, it will be convenient to flatten the map by blowing up \mathcal{E} . This can be done in a universal manner, following [AK00, Mol16].

The combinatorics of the flattening can be described explicitly. Consider the map on tropicalizations

$$\tau = \tilde{\mathcal{D}}^{\text{trop}} \rightarrow \mathcal{E}^{\text{trop}} = \sigma.$$

The tropical curves parametrised on the left inherit a central alignment from $\widetilde{\mathcal{VZ}}_{1,\alpha}(\mathbb{P}^m|H, d)$, which orders a certain subset of the edge lengths. The polyhedral criterion for equidimensionality is that every cone of the source maps surjectively onto a cone of the target. There is a universal logarithmic modification $\widetilde{\mathcal{E}} \rightarrow \mathcal{E}$ such that $\widetilde{\mathcal{D}} \rightarrow \widetilde{\mathcal{E}}$ is flat, constructed by taking the minimal subdivision of $\mathcal{E}^{\text{trop}}$ such that cones map surjectively to cones (see Figure 7).

This subdivision may be described explicitly. For this, it is convenient to employ the formalism of extended tropicalizations [Thu07, ACP15]. The subdividing cones are obtained from cones in the star of τ under the limit $e_1 = \dots = e_r = \infty$. The cones in the star of τ are given by (in)equalities relating distances of pairs of vertices, and there are three cases to consider:

- (1) both vertices are components of the subgraph $\sqsubset \setminus \sqsubset_0$;
- (2) both vertices are components of the subgraph \sqsubset_0 ;
- (3) one vertex is a component of \sqsubset_0 and one is a component of $\sqsubset \setminus \sqsubset_0$.

As we pass to the limit, the first relations impose a central alignment condition on the genus zero moduli, the second relations impose a radial alignment condition on the genus one moduli, and the third relations disappear entirely. We are therefore led to the following moduli theoretic description of the logarithmic modification $\widetilde{\mathcal{E}}$.

Keeping in mind the fibre product description of \mathcal{E} , construct a modification $\widetilde{\mathcal{E}}$ from \mathcal{E} as follows:

- (1) Given a logarithmic map of type Δ with splitting nodes q_1, \dots, q_r , remove the genus one piece \sqsubset_0 and formally glue together the r vertices of the tropicalization which carry the splitting nodes, and declare the resulting vertex to be the root of the resulting tree. Impose the central alignment condition, where the special root takes on the rôle previously occupied by the core.
- (2) Replace the fibre product factor in \mathcal{E} associated to the vertex \sqsubset_0

$$\overline{\mathcal{M}}_{\Gamma_{\sqsubset_0}}^{\leftrightarrow}(\mathbb{P}|H_0 + H_\infty)$$

with the radially aligned rubber space:

$$\mathcal{VZ}_{\Gamma_{\sqsubset_0}}^{\leftrightarrow}(\mathbb{P}|H_0 + H_\infty).$$

(Here we are imposing fibre-direction factorisation in order to isolate the main component.)

This procedure induces a logarithmic modification $\widetilde{\mathcal{E}} \rightarrow \mathcal{E}$ which admits a modular description as a fibre product of genus zero centrally aligned maps coming from the vertices $\sqsubset_1, \dots, \sqsubset_r$, and genus one aligned rubber maps satisfying fibre-direction factorisation from the vertex \sqsubset_0 .

We have produced a factorisation

$$\widetilde{\mathcal{D}} \rightarrow \widetilde{\mathcal{E}} \rightarrow \mathcal{E},$$

where the first map is flat and the second map is a logarithmic modification. $\widetilde{\mathcal{D}}$ is a compactification of a torus bundle over $\widetilde{\mathcal{E}}$ whose strata can be read off from the tropicalization; in particular, we have the information of how (polynomials in) the classes of boundary strata push forward to $\widetilde{\mathcal{E}}$. In fact, we can say more and describe the generic fibre of $\widetilde{\mathcal{D}} \rightarrow \widetilde{\mathcal{E}}$:

Lemma 4.14. Generically over $\widetilde{\mathcal{E}}$, the morphism $\widetilde{\mathcal{D}} \rightarrow \widetilde{\mathcal{E}}$ is the projectivisation of a tautological vector bundle.

Proof. Let us assume that the curve does not degenerate, so that the dual graph is as prescribed by Δ . In particular there are vertices $\sqsubset_1, \dots, \sqsubset_r$ generically lying on the circle of radius δ . The moduli space of attaching data for the elliptic singularity is the complement of the coordinate hyperplanes in:

$$(4) \quad \mathbb{P} \left(\bigoplus_{i=1}^r T_{q_i} C_i \right).$$

Was r called
fore?




FIGURE 8. Moving vertices off the circle creates new tropical parameters.

This can also be thought of as parametrising compatible identifications $\theta_{ij}: T_{C_i, q_i} \cong T_{C_j, q_j}$ (see §1.4). A point in this space determines a map from the rational r -fold singularity to the elliptic r -fold singularity (this coincides with the fibre of the space of aligned curves over the space of nodal curves; see [RSPW19a, §3.4]). Along the boundary, some of the \square_i (though not all of them) may move off from the circle, so that the latter subdivides the edge between them and the core. Correspondingly, there is a destabilisation of the curve (semistable components are represented by crosses in our figures), and we obtain an elliptic singularity containing strictly semistable branches (*sprouting*). This was already identified by Smyth as a compactification of the moduli space of attaching data carrying a Gorenstein universal curve [Smy11b, §2.2].

Tropically, pushing some of the \square_i off the circle creates new finite-length edges with coordinates $\ell_i \in \mathbb{R}_{\geq 0}$ (see the right-hand side of Figure 8). Since not all of them can be strictly positive, the resulting fan is that of a projective space. This gives us a fibrewise description of $\tilde{\mathcal{D}} \rightarrow \tilde{\mathcal{E}}$ over the generic point.

A slight modification of the argument above is required to account for the fact that all vertices that lie on the circle and are mapped to $0 \in \mathbb{R}_{\geq 0}$ by the tropical map are already generically aligned on $\tilde{\mathcal{E}}$. Indeed, by the combinatorial description of rays in §4.3.1, there are no stable vertices inside the circle, so all the edges going from a vertex of Λ_0 to the core have the same expansion factor m (maximal among the contact orders at the core). Therefore, for all i such that $\square_i \in \Lambda_0$, the $T_{q_i} C_i^{\otimes m}$ can be identified to the target tangent line on the fibre product (3), and, after saturation, the $T_{q_i} C_i$ themselves are canonically isomorphic on $\tilde{\mathcal{E}}$. Thus, only one summand should be taken to represent all of them in the projective bundle (4). \square

4.3.5. \mathcal{D} from $\tilde{\mathcal{D}}$: *first way*. On $\tilde{\mathcal{D}}$ there is a tautological line bundle arising from the modular description of $\tilde{\mathcal{V}}\mathcal{Z}_{1,\alpha}(\mathbb{P}^m|H,d)$, namely $\mathcal{O}(\delta)$. Recall that the piecewise linear function δ may be viewed as a global section of the characteristic sheaf of the logarithmic structure on $\tilde{\mathcal{V}}\mathcal{Z}_{1,\alpha}(\mathbb{P}^m|H,d)$. The toric geometry dictionary gives rise to an associated Cartier divisor. Alternatively, for any logarithmic stack X , the natural coboundary map induced by

$$0 \rightarrow \mathcal{O}_X^\star \rightarrow M_X^{\text{gp}} \rightarrow \overline{M}_X^{\text{gp}} \rightarrow 0$$

associates a \mathcal{O}_X^\star -torsor to a global section of the characteristic sheaf.

h moving to
preliminary
on?

Lemma 4.15. The fibre of $\mathcal{O}(\delta)$ over a moduli point is naturally isomorphic to

$$T_p C' \otimes \mathbb{T}$$

where C' is any component of C which lies on the circle of radius δ , p is the node separating it from the core, and \mathbb{T} is the universal tangent line bundle on the moduli space for \mathbb{C}_0 .

Here we explain the meaning of \mathbb{T} . Isolating the subcurve $C_0 \subseteq C$ gives rise to a morphism from $\widetilde{\mathcal{D}}$ to the moduli space of radially aligned curves of genus one:

$$\widetilde{\mathcal{D}} \rightarrow \mathcal{VZ}_{1,n}^{\text{rad}}.$$

The latter space coincides with the Vakil–Zinger blow-up of $\overline{\mathcal{M}}_{1,n}$ [VZ08, §2.3]. In particular, after twisting by exceptional divisors, the tangent line bundles at two marking $\widetilde{\mathbb{T}}_i$ and $\widetilde{\mathbb{T}}_j$ are identified (via the dual Hodge bundle \mathbb{E}^\vee), extending the identification on the smooth locus provided by regarding the elliptic curve as an algebraic group. This gives rise to a single line bundle – the “universal tangent line bundle” – on $\mathcal{VZ}_{1,n}^{\text{rad}}$; we let \mathbb{T} denote its pullback to $\widetilde{\mathcal{D}}$.

Proof. Choose a component C' of the curve which lies on the circle. Then $\mathcal{O}(\delta)$ is equal to the tensor product of tangent line bundles along the path of edges connecting C' to the core. However, this tensor product of line bundles is telescoping: any vertex which appears in the middle of the path will have an incoming and outgoing node (and possibly other nodes that are irrelevant for this product), and since the vertex corresponds to a smooth rational curve these tangent spaces are naturally dual to each other [VZ08, §2.2]. Thus these factors cancel, and the result follows. \square

The content of Lemma 4.14 is that the choice of a point in the fibre of $\widetilde{\mathcal{D}} \rightarrow \mathcal{E}$ provides a natural identification between the inward-pointing tangent lines on the circle. According to Lemma 4.15, they can further be identified with the line bundle $\mathcal{O}(\delta) \otimes \mathbb{T}^\vee$ on $\widetilde{\mathcal{D}}$. After Corollary 1.3, we can therefore express the factorisation condition as the degeneracy of the following natural morphism of vector bundles on $\widetilde{\mathcal{D}}$:

$$\Sigma \text{d}f: \mathcal{O}(\delta) \otimes \mathbb{T}^\vee \rightarrow \text{ev}_q^* TH.$$

Here $\Sigma \text{d}f$ is defined by summing the images of the tangent vectors at the inward-pointing nodes on the circle, using the natural identification of each of these tangent spaces with $\mathcal{O}(\delta) \otimes \mathbb{T}^\vee$ established above. This morphism takes values in $\text{ev}_q^* TH$ rather than $\text{ev}_q^* T\mathbb{P}^m$ because the contact order of the exterior components with H is strictly larger than 1 (except for the case of lines, dealt with separately in §4.3.2). This gives a section of the vector bundle

$$V = (\text{ev}_q^* TH) \otimes \mathbb{T} \otimes \mathcal{O}(-\delta)$$

which is transverse and whose vanishing locus coincides with \mathcal{D} . Transversality follows by construction, because any stratum of the boundary on which $\Sigma \text{d}f$ vanishes has been (blown up and) included into $\mathbb{T} \otimes \mathcal{O}(-\delta)$ as a twist; in other words, the circle of radius δ always passes through at least one component which is not contracted by the map to the expansion. Now, if the vanishing locus of $\Sigma \text{d}f$ does not contain any stratum of the boundary, then it is the closure of its intersection with the open stratum, which can be seen to be transverse. Thus we obtain:

Proposition 4.16. $[\mathcal{D}] = e(V) \cap [\widetilde{\mathcal{D}}]$.

We claim that the class $e(V)$ is tautological and computable. The only part which needs justification is $\mathcal{O}(\delta)$. We claim that this can be obtained by pulling back a line bundle from \mathcal{E} and twisting by exceptional divisors.

Lemma 4.17. For $i \in \{1, \dots, r\}$ consider the line bundle

$$T_i = T_{q_i} C_i \otimes T_{q_i} C_0$$

on \mathcal{E} , where q_i is the splitting node. Then $\mathcal{O}(\delta)$ is isomorphic to the pull-back of T_i to $\tilde{\mathcal{D}}$ twisted by all exceptional divisors.

We illustrate this lemma in the following example, which brings out the interplay between logarithmic moduli and blowups.

Example 4.18. Consider the locus \mathcal{E} given by σ below, and the codimension-one locus $\tilde{\mathcal{D}}_\rho \subseteq \tilde{\mathcal{D}}_\sigma$ given by the two-dimensional (due to the alignment) cone ρ :




FIGURE 9. σ




FIGURE 10. ρ

On the cone ρ we have the following tropical continuity equations

$$e := e_1 = e_2, \quad f := f_1 = f_2 = 2f_3, \quad g := g_1 = g_2 = 2g_3$$

together with the equation imposed by the choice of alignment:

$$e_1 + f_1 = e_2 + f_3 + g_3 \quad (\iff f = g).$$

In this example we have:

$$\lambda(\sqsubset_1^0) = \lambda(\sqsubset_2^0) = e, \quad \lambda(\sqsubset_1^1) = \lambda(\sqsubset_1^2) = \lambda(\sqsubset_2^2) = e + f = \delta, \quad \lambda(\sqsubset_1^3) = \lambda(\sqsubset_1^4) = e + 2f.$$

From now on we use the same symbol to denote an edge length and the corresponding node on $\tilde{\mathcal{D}}_\rho$. On this locus we have line bundles pulled back from \mathcal{E}

$$T_1 = T_{e_1} C_1^0 \otimes T_{e_1} C_0$$

$$T_2 = T_{e_2} C_2^0 \otimes T_{e_2} C_0$$

as well as the logarithmic line bundle

$$\mathcal{O}(\delta) = T_{f_1} C_1^1 \otimes T_{e_1} C_0 = T_{f_2} C_1^2 \otimes T_{e_1} C_0 = T_{g_3} C_2^2 \otimes T_{e_2} C_0$$

where the identifications follow from Lemma 4.15. We claim that:

$$(5) \quad \mathcal{O}(\delta) = T_1 \otimes \mathcal{O}(\tilde{\mathcal{D}}_\rho) = T_2 \otimes \mathcal{O}(\tilde{\mathcal{D}}_\rho).$$

Notice that $\mathcal{O}(\widetilde{\mathcal{D}}_\rho) = \mathcal{O}(f) = \mathcal{O}(g)$. Thus we have

$$\begin{aligned} T_1 \otimes \mathcal{O}(\widetilde{\mathcal{D}}_\rho) &= \mathcal{O}(f_1) \otimes T_{e_1} C_1^0 \otimes T_{e_1} C_0 \\ &= T_{f_1} C_1^1 \otimes T_{f_1} C_1^0 \otimes T_{e_1} C_1^0 \otimes T_{e_1} C_0 \\ &= T_{f_1} C_1^1 \otimes T_{e_1} C_0 \quad (= \mathcal{O}(\delta)) \end{aligned}$$

where in the last line we have used the same telescoping trick used in the proof of Lemma 4.15, namely that $T_{f_1} C_1^0 \otimes T_{e_1} C_1^0 = \mathcal{O}$. On the other hand, for T_2 we may write $f = f/2 + g/2 = f_3 + g_3$ in order to obtain:

$$T_2 \otimes \mathcal{O}(\widetilde{\mathcal{D}}_\rho) = \mathcal{O}(g_3) \otimes \mathcal{O}(f_3) \otimes T_{e_2} C_2^0 \otimes T_{e_2} C_0 = T_{g_3} C_2^2 \otimes T_{e_2} C_0 = \mathcal{O}(\delta)$$

where we have used the same telescoping trick. Thus we have proven (5). This observation generalises, giving a precise relation between the intrinsically-defined line bundle $\mathcal{O}(\delta)$ on $\widetilde{\mathcal{D}}$ (which we use to impose the factorisation condition) and the line bundles T_i pulled back from \mathcal{E} . This allows us to express the Chern classes of $\mathcal{O}(\delta)$ (and hence V) in terms of pull-backs of tautological classes from \mathcal{E} and boundary strata on $\widetilde{\mathcal{D}}$, which is sufficient in order to compute integrals.

4.3.6. \mathcal{D} from $\widetilde{\mathcal{D}}$: second way. We analyse the degeneracy locus of Σdf , in order to have a more direct construction of \mathcal{D} from the fibre product (3). On \mathcal{E} consider the piecewise linear function δ_i ($i = 1, \dots, r$), defined as the minimal distance from a component of (i.e. generising to) the core \sqsubset_0 to a non-contracted component of the i -th tail \sqsubset_i , and the associated line bundle $\mathcal{O}(\delta_i)$. Construct

$$\mathcal{P} = \mathbb{P}_{\mathcal{E}} \left(\bigoplus_{i=1}^r \mathcal{O}(\delta_i) \right) \xrightarrow{p} \mathcal{E}.$$

When all fibre coordinates are non-zero, a point of \mathcal{P} induces compatible isomorphisms $\mathcal{O}(\delta_i) \cong \mathcal{O}(\delta_j)$ for all i, j . We think of this as saying that all the \sqsubset_i are at the same distance from the core. When some of the coordinates are 0, the corresponding tail is further from the core. This corresponds to an alignment of $\sqsubset_1, \dots, \sqsubset_r$ with respect to the radius δ , that is the distance of the closest of them from the core. Therefore, by the universal property, there is a morphism $\widetilde{\mathcal{D}} \rightarrow \mathcal{P}$ lifting $\widetilde{\mathcal{D}} \rightarrow \mathcal{E}$.

In order to express factorisation, consider the map of vector bundles on \mathcal{P} :

$$\mathcal{O}_{\mathcal{P}}(-1) \rightarrow p^* \left(\bigoplus_{i=1}^r \mathcal{O}(\delta_i) \right) \xrightarrow{\Sigma df} \text{ev}_q^* TH.$$

The composite, regarded as a section s of $W = \mathcal{O}_{\mathcal{P}}(1) \otimes \text{ev}_q^*(TH)$, vanishes precisely where f factors through the elliptic singularity determined by the point in the fibre of p . Generically on $\widetilde{\mathcal{E}}$, this is exactly what we are after (compare with Lemma 4.14). On the boundary, though, it may happen that a simultaneous degeneration in the base and the fibre makes this factorisation trivial, in the sense that the map \bar{f} is constant on all the branches of the elliptic singularity. These loci may be identified explicitly: let $\Lambda = (\Lambda_B, \Lambda_F)$ with $\Lambda_B \subseteq \{\sqsubset_1, \dots, \sqsubset_r\}$ and $\Lambda_F = \{\sqsubset_1, \dots, \sqsubset_r\} \setminus \Lambda_B$, and let Ξ_Λ be the locus in \mathcal{P} where:

- (B) $\forall i \in \Lambda_B$, the i -th component of the fibre product (3), namely the map $f_i: C_i \rightarrow (\mathbb{P}^m|H)$, degenerates so that it is constant near the attaching point to the core.
- (F) $\forall j \in \Lambda_F$, the j -th coordinate in the fibre of $p: \mathcal{P} \rightarrow \mathcal{E}$ vanishes.

It should be noted that the degeneracy loci of (B) are themselves Gathmann's comb loci [Gat02], similar to the loci \mathcal{D} that we are considering, only this time in genus 0, and therefore recursively computable.

Lemma 4.19. The Ξ_Λ are precisely the boundary loci on which s vanishes.

Compare with [VZ08, §3.2] for an analogous phenomenon in the non-relative case. Clearly, these loci may be in excess with respect to the expected codimension of $V(s)$, that is $N - 1$. These loci are logarithmic substrata; therefore, there exists a logarithmic blowup that principalises them. We claim that $\tilde{\mathcal{D}}$ is one.

Lemma 4.20. The pullback of Ξ_Λ is principal in $\tilde{\mathcal{D}}$.

Proof. We analyse the ideal of Ξ_Λ : it is generated by

- (1) equations on \mathcal{E} cutting the locus where C_i degenerates so that f_i is constant on a neighbourhood of q_i for $i \in \Lambda_B$: these correspond to the tropical functions $\delta_i - \lambda(q_i)$ where λ is the distance function to the core;
- (2) equations on the fibre of p for the linear subspace corresponding to $i \in \Lambda_F$: tropically, these correspond to the ℓ_i discussed in Lemma 4.14.

By construction, on $\tilde{\mathcal{D}}$ it is always possible to tell which of these tropical lengths is the shortest. Figure 11 represents the situation: the circle we have drawn on Ξ_Λ is degenerate - it passes through no non-contracted vertex - therefore we need to decide for a strictly larger circle in order to lift to $\tilde{\mathcal{D}}$. This shows that the ideal is principalised on $\tilde{\mathcal{D}}$. \square




FIGURE 11. On the left, a generic point of \mathcal{P} . On the right, a point of Ξ . The dotted circle is the one we draw on \mathcal{P} . The solid circle represents a radius δ enlarging the dotted one (a stratum of $\tilde{\mathcal{D}}$).

Finally, on $\tilde{\mathcal{D}}$ there are exceptional divisors $\tilde{\Xi}_\Lambda$ corresponding to Ξ_Λ in \mathcal{P} . Since the Ξ_Λ exhaust all the excess-dimensional components of the vanishing locus of $s \in H^0(\mathcal{P}, W)$, the latter induces a section $\tilde{s} \in H^0(\tilde{\mathcal{D}}, W(-\Sigma_\Lambda \tilde{\Xi}_\Lambda))$ by pullback and twisting, whose vanishing locus is dimensionally transverse to the boundary, and whose open part coincides with the generic point of \mathcal{D} , therefore $V(\tilde{s}) = \mathcal{D}$.

Proposition 4.21.

$$[\mathcal{D}] = e(W(-\Sigma_\Lambda \tilde{\Xi}_\Lambda)) \cap [\tilde{\mathcal{P}}]$$

where $\tilde{\mathcal{P}}$ denotes the blow-up of \mathcal{P} in the Ξ_Λ .

Applying [Ful98, Corollary 4.2.2], we get:

Corollary 4.22. In $A_*(\mathcal{P})$ the following formula holds:

$$[\mathcal{D}] = \sum_{\Lambda} \sum_i s_i(\Xi_\Lambda, \mathcal{P}) c_{N-1-i}(\text{ev}_q^* TH \otimes \mathcal{O}_{\mathcal{P}}(1))$$

It is possible to push the formula further down to \mathcal{E} , and applying [Ful98, §3.1] we obtain an expression for $[\mathcal{D}]$ in terms of:

- evaluation classes, such as $\text{ev}_q^* TH$;
- Chern classes of the logarithmic line bundles $\mathcal{O}(\delta_i)$;

- Segre classes of Gathmann's comb loci indexed by Λ_B .

Corollary 4.23. After pushforward to \mathcal{E} , the contribution of \mathcal{D} to the invariants can be computed in terms of tautological integrals on moduli spaces of maps with lower numerics.

4.4. Recursion for general (X, Y) . Now let (X, Y) be a general smooth very ample pair. In §1.11 we studied the moduli space $\mathcal{VZ}_{1,\alpha}(X|Y, \beta)$ of centrally aligned maps to (X, Y) and equipped it with a virtual fundamental class by diagonal pull-back from $\mathcal{VZ}_{1,\alpha}(\mathbb{P}^m|H, d)$.

This construction of the virtual fundamental class means that once we have established a recursion formula for (\mathbb{P}^m, H) , it immediately pulls back to give a recursion formula for (X, Y) . To be more precise, Theorem 3.3 holds *mutatis mutandis*, with the same values of $\lambda_{\mathcal{D}}$. The computation of the splitting orders is exactly the same, and the recursive description of the boundary strata pulls back to give an entirely analogous description on $\mathcal{VZ}_{1,\alpha}(X|Y, \beta)$.

5. QUANTUM LEFSCHETZ ALGORITHM

Consider a smooth pair (X, Y) with Y very ample, and let \mathbb{P} be the projective bundle $\mathbb{P} = \mathbb{P}_Y(N_{Y|X} \oplus \mathcal{O}_Y)$. We assume that all genus zero and reduced genus one Gromov–Witten invariants of X are known. From this starting data, we will apply our recursion formula to compute:

- (1) the genus one **reduced restricted absolute Gromov–Witten theory** of Y ;
- (2) the genus one **reduced relative Gromov–Witten theory** of the pair (X, Y) ;
- (3) the genus one **reduced rubber theory** of \mathbb{P} .

5.1. Reduced absolute, relative and rubber invariants. To be precise: by a reduced invariant of Y we mean an integral over $\mathcal{VZ}_{1,n}(Y, \beta)$ of a product of evaluation classes and pullbacks of psi classes along morphisms forgetting subsets of the marked points:

$$\int_{[[\mathcal{VZ}_{1,n}(Y, \beta)]]^{\text{virt}}} \prod_{i=1}^n \text{fgt}_{S_i}^* \psi_i^{k_i} \cdot \text{ev}_i^* \gamma_i.$$

(Here $S_i \subseteq \{1, \dots, n\}$ with $i \notin S_i$ for all i ; taking $S = \emptyset$ gives the ordinary psi classes.). The evaluation maps are viewed as mapping into X (hence the adjective “restricted”). Reduced relative invariants of (X, Y) are defined in the same way

$$\int_{[[\mathcal{VZ}_{1,\alpha}(X|Y, \beta)]]^{\text{virt}}} \prod_{i=1}^n \text{fgt}_{S_i}^* \psi_i^{k_i} \cdot \text{ev}_i^* \gamma_i$$

except now the forgetful morphism maps into a space of absolute stable maps:

$$\text{fgt}_S: \mathcal{VZ}_{1,\alpha}(X|Y, \beta) \rightarrow \mathcal{VZ}_{1,n-\#S}(X, \beta).$$

Note that even the case $S = \emptyset$ entails a nontrivial forgetful morphism. In particular, all the psi classes which we consider are **collapsed psi classes**, meaning that they are relative cotangent line classes for the corresponding collapsed stable map. Note that, unlike in the absolute case, in the relative case it may well be that the entire insertion is pulled back along a single forgetful map fgt_S . The reduced rubber invariants of \mathbb{P} are defined similarly (again using collapsed psi classes).

Remark 5.1. The systems of invariants defined above are equivalent to the classical systems of invariants (which do not use forgetful morphisms) by well-known topological recursion relations.

Remark 5.2. In the following discussion we will sometimes omit the adjective “reduced” when discussing genus one invariants, but it is always implicit; all genus one invariants we consider will be reduced.

5.2. Fictitious and true markings. The recursion procedure we outline is rather delicate. Roughly speaking, we will induct on the degree (meaning $Y \cdot \beta$), number of marked points and total tangency. We adopt Gathmann’s point of view on relative Gromov–Witten theory [Gat02], whereby only a subset of the contact points with the divisor are required to be marked; this allows for “total tangency” strictly smaller than $Y \cdot \beta$.

In order to make sense of this in the logarithmic setting (in which all contact points must be marked), we introduce the concept of **fictitious markings**. Consider a moduli space $\mathcal{VZ}_{1,\alpha}(X|Y,\beta)$ of reduced relative stable maps and a corresponding integrand γ . We let $F \subseteq \{1, \dots, n\}$ be the maximal subset of marked points such that the following two conditions are satisfied:

- (1) $\alpha_i = 1$ for all $i \in F$;
- (2) the entire integrand γ is pulled back along fgt_F .

This subset is uniquely determined, and its elements are referred to as **fictitious markings**. We note that $\alpha_i = 1$ is not sufficient for x_i to be fictitious; it must also be the case that all the insertions are pulled back along forgetting x_i . Markings which are not fictitious are referred to as **true markings**. When inducting on relative invariants we will always be interested in the number of true markings (as opposed to the total number of markings) and the **true tangency**

$$t := \sum_{i \notin F} \alpha_i \leq Y \cdot \beta = d$$

as opposed to the total tangency, which is always d . This formalises the idea that relative invariants with non-maximal tangency $t < d$ can be obtained by adding $d - t$ fictitious markings of tangency 1; see [Gat02, Lemma 1.15(i)].

5.3. Structure of the recursion. Given the genus zero Gromov–Witten theory of X , the results of [Gat02] give an effective algorithm to reconstruct the genus zero theories of Y and (X, Y) ; the genus zero rubber theory of \mathbb{P} is equal to the genus zero theory of Y [Gat03]. Thus we may assume that all genus zero invariants are known. We assume in addition that we know the genus one reduced theory of X . The structure of the recursion is then as follows:

for $d \geq 0$:
for $n \geq 0$:
for $t \geq 0$:
Step 1: Compute forgetful relative invariants of (X, Y) (degree d , $n + 1$ true
markings, true tangency t); see below.
Step 2: Compute absolute invariants of Y (degree d , n markings).
for $t \geq 0$:
Step 3: Compute relative invariants of (X, Y) (degree d , n true markings, true
tangency t).
for $n \geq 0$:
for $m \geq 0$:
Step 4: Compute rubber invariants of \mathbb{P} (degree d , n relative markings, m
non-relative markings).

Here each loop begins at 0 and proceeds through successively larger values, with the invariants at each step expressed recursively in terms of invariants computed in the preceding steps and genus zero invariants. Although the loops involving d, n and m have infinite length, in order to compute any single invariant it is only necessarily to iterate the preceding loops for a finite amount of time.

Remark 5.3 (Forgetful relative invariants). In Step 1 above, a **forgetful relative invariant** of (X, Y) is by definition a reduced relative invariant for which there exists a marking x_0 such that all of the insertions are pulled back along fgt_{x_0} . In our recursion, we first deal with the special class of forgetful relative invariants (with $n + 1$ true markings), before computing the absolute invariants (with n markings) and then returning to compute all of the relative invariants (with n true markings). The need to separately treat a particular subclass of the relative invariants is an inescapable feature of the genus one recursion.

Remark 5.4 (The base case). The base terms of the recursion all have $d = 0$ and as such are easy to compute: the relative invariants of (X, Y) are nothing but absolute invariants of X (i.e. there are no tangency conditions), and the absolute invariants of Y are given by obstruction bundle integrals over Deligne–Mumford spaces. The rubber invariants of \mathbb{P} in degree zero are given by integrals over the main component of the double ramification cycle (in the rubber case there may be markings with tangency even for $d = 0$, since d only records the degree covering the base); this can likewise be computed as an integral over Deligne–Mumford space, by correcting the formula for the ordinary double ramification cycle [Hai13, JPPZ17] by the obvious contribution from the boundary irreducible component (the observation which makes this easy to do in genus one is that the rubber moduli space is pure-dimensional, so that the virtual class is simply the sum of the fundamental classes of the irreducible components). We will now explain how to perform each of the four inductive steps outlined above.

Notation 5.5. Given tuples $\mathbf{a} = (a_1, \dots, a_k)$ and $\mathbf{b} = (b_1, \dots, b_k)$, we say that $\mathbf{a} < \mathbf{b}$ if there exists an $i \in \{1, \dots, k\}$ such that $a_j = b_j$ for $j < i$ and $a_i < b_i$.

Step 1. Suppose we are given a forgetful relative invariant to compute. That is, we have a relative space $\mathcal{VZ}_{1,\alpha}(X|Y, \beta)$ of degree d , with $n + 1$ true markings and true tangency t , and a marking x_0 such that the insertion $\gamma = \text{fgt}_{x_0}^* \delta$ for some class δ . We assume inductively that every absolute, relative and rubber invariant with $(d', n') < (d, n)$ has been computed, as well as every forgetful relative invariant with $(d', n', t') < (d, n + 1, t)$. As noted above, we may assume that $d \geq 1$ which means that there exists a true marking x_1 with $\alpha_1 \geq 1$. We will decrease the tangency at x_1 . Consider the vector of tangency orders

$$\alpha - e_1$$


obtained from α by replacing the entry α_1 by $\alpha_1 - 1$. Consider then the moduli space:

$$\mathcal{VZ}_{1,(\alpha-e_1)\cup(1)}(X|Y, \beta).$$

Denote the newly-introduced marking by y and take the integrand $\tilde{\gamma} = \text{fgt}_y^* \gamma$. Applying our recursion formula to x_1 , we obtain:

$$(6) \quad ((\alpha_1 - 1)\psi_1 + \text{ev}_1^* Y) \tilde{\gamma} \cap [\mathcal{VZ}_{1,(\alpha-e_1)\cup(1)}(X|Y, \beta)] = \tilde{\gamma} \cap [\mathcal{D}(1)].$$

We first examine the left-hand side. The class ψ_1 differs from $\text{fgt}_y^* \psi_1$ by the locus where x_1 and y are contained on a contracted rational bubble. This locus consists of stable maps of the form




with all other marked points contained on \mathbb{C}_1 . This is isomorphic to $\mathcal{VZ}_{1,\alpha}(X|Y, \beta)$ and when we restrict $\tilde{\gamma}$ to this locus we obtain precisely the class γ which we started with. Thus (6) may be written as

$$(7) \quad (\alpha_1 - 1)I + \text{fgt}_y^* ((\alpha_1 - 1)\psi_1 + \text{ev}_1^* Y) \tilde{\gamma} \cap [\mathcal{VZ}_{1,(\alpha-e_1)\cup(1)}(X|Y, \beta)] = \tilde{\gamma} \cap [\mathcal{D}(1)].$$

where I is the invariant we are trying to compute. We will now show that the other terms may be expressed in terms of invariants computed earlier in the recursion.

The second term on the left-hand side is a forgetful relative invariant with the same degree and number of true markings (since y is fictitious), and strictly smaller true tangency; hence it has been computed at an earlier step in the recursion. We now examine the right-hand side of (7). Recall that all of the genus zero data has already been computed, so we only need to focus on the genus one pieces. First consider the type I loci. The genus one piece has strictly smaller degree (and so has already been computed) except in the following situation (with some stable distribution of the remaining markings):



In this situation, C_1 contains at most $n + 1$ true markings. We consider three possible cases:

- (1) If C_1 has $n - 1$ or fewer true markings, then its contribution has already been computed earlier in the recursion.
- (2) If C_1 has exactly n true markings, then C_0 contains exactly one true marking besides x_1 (which itself may be true or fictitious). In this situation we must have $y \in C_1$, since otherwise we would have a moduli space for C_0 given by $\overline{\mathcal{M}}_{0,k}$ with $k \geq 4$, and applying the projection formula with fgt_y we would conclude that the contribution is zero. Thus we have $y \in C_1$, and so the genus one contribution is a forgetful relative invariant with n true markings, and so has already been computed.
- (3) Finally, if C_1 contains exactly $n + 1$ true markings, then the only additional markings on C_0 are fictitious, and by the same argument as in the previous paragraph there can only be one. Thus for each fictitious marked point we obtain a locus isomorphic to $\mathcal{VZ}_{1,\alpha}(X|Y,\beta)$ and $\tilde{\gamma}$ restricts to γ here (from the point of view of computing invariants, the fictitious marked points are indistinguishable, meaning that the contributions are all the same). Thus for each fictitious marked point (of which there is at least one, namely y) we get a contribution of $\alpha_1 I$ to the right-hand side of (7), where as before I is the invariant we are trying to compute.

The contributions of the type II loci only involve genus zero data and so can be computed algorithmically following [Gat02]. The contributions of the type \dagger loci are determined by genus zero data and tautological integrals on genus one Deligne–Mumford spaces, which can also be computed algorithmically using the string and dilaton equations.

It remains to consider type III loci. If the degree of the genus one piece is less than d then we have a rubber invariant of strictly smaller degree. The only other possibility is that the entire curve is mapped into the divisor. In this case we may apply the projection formula with fgt_y to identify this with an integral over $\mathcal{VZ}_{1,m}(Y,\beta)$ for some (possibly large) number m of marked points. But by assumption there is another marked point x_0 with all of the insertions pulled back along fgt_{x_0} , so a further application of the projection formula shows that this contribution vanishes. To conclude, we may rearrange (7) to obtain an expression of the form

$$\lambda I = \text{polynomial in previously computed invariants}$$

where λ is an explicit scalar which is always nonzero; we have thus determined the forgetful relative invariant I , which completes Step 1.

Step 2. Consider now the absolute space $\mathcal{VZ}_{1,n}(Y, \beta)$ with an insertion γ , and suppose inductively that we have computed all forgetful relative invariants with $(d', n') \leq (d, n+1)$, all relative invariants and absolute invariants with $(d', n') < (d, n)$, and all rubber invariants with $d' < d$.

Step 2a. Consider the following moduli space with $n+1$ markings:

$$\mathcal{VZ}_{1,(d,0,\dots,0)}(X|Y, \beta).$$

Let x_0 denote the relative marking and consider the integrand $\tilde{\gamma} = \text{fgt}_{x_0}^* \gamma$. Now recurse at x_1 to obtain:


$$(8) \quad \text{ev}_1^* Y \cdot \tilde{\gamma} \cap [\mathcal{VZ}_{1,(d,0,\dots,0)}(X|Y, \beta)] = \tilde{\gamma} \cap [\mathcal{D}(1)].$$

The left-hand side is a forgetful relative invariant of degree d and $\leq n+1$ true markings, and so has already been computed.

On the right-hand side, similar arguments to those in Step 1 show that the contributions of type II and type \dagger loci can be expressed in terms of invariants computed earlier in the recursion. Similarly, the contributions of type III loci have already been computed, except for the locus where the entire curve is mapped into the divisor. On this locus we apply the projection formula to fgt_{x_0} to identify the contribution with

$$d^2 \cdot \gamma \cap [\mathcal{VZ}_{1,n}(Y, \beta)] = d^2 \cdot I$$

where I is the invariant we are trying to compute. Finally, consider loci of type I. The genus one contributions from each locus have $(d', n') < (d, n)$ (and hence have already been computed) except in the following case



which gives a contribution of

$$d \cdot \gamma \cap [\mathcal{VZ}_{1,(d,0,\dots,0)}(X|Y, \beta)]$$

$\underbrace{\hspace{1.5cm}}_{n-1}$

to the right-hand side of (8). Here γ is the insertion we started with; the difference is that we are now considering relative maps to (X, Y) with maximal tangency at x_1 , rather than absolute maps to Y . Putting this all together, we obtain from (8)

$$(9) \quad I = (-\gamma/d) \cap [\mathcal{VZ}_{1,(d,0,\dots,0)}(X|Y, \beta)] + \text{polynomial in previously computed invariants}$$

$\underbrace{\hspace{1.5cm}}_{n-1}$

where on the right-hand side there are $n-1$ non-relative markings x_2, \dots, x_n , and a relative marking x_1 .

Step 2b. We now apply the recursion to the right-hand side of (9). Consider the space

$$\mathcal{VZ}_{1,(d-1,1,0,\dots,0)}(X|Y, \beta)$$

where x_1 now has tangency $d-1$ and we have introduced a new marking y with tangency 1. As usual we take the insertion $\tilde{\gamma} = \text{fgt}_y^* \gamma$. Recursing at x_1 we obtain:

$$(10) \quad ((d-1)\psi_1 + \text{ev}_1^* Y) \cdot (-\tilde{\gamma}/d) \cap [\mathcal{VZ}_{1,(d-1,1,0,\dots,0)}(X|Y, \beta)] = (-\tilde{\gamma}/d) \cap [\mathcal{D}(1)].$$

The difference between ψ_1 and $\text{fgt}_y^* \psi_1$ is given by the locus where x_1 and y belong to a collapsed rational bubble:

Step 3. Now suppose we are given a relative space $\mathcal{VZ}_{1,\alpha}(X|Y,\beta)$ with an insertion γ , and suppose inductively that we have computed all forgetful relative invariants with $(d', n') \leq (d, n+1)$, all absolute invariants with $(d', n') \leq (d, n)$, all relative invariants with $(d', n', t') < (d, n, t)$ and all rubber invariants with $d' < d$.


Choose a true marking x_1 with $\alpha_1 \geq 1$ and consider the moduli space

$$\mathcal{VZ}_{1,(\alpha-e_1)\cup(1)}(X|Y,\beta)$$

where y is the newly-introduced marking. As usual consider the insertion $\tilde{\gamma} = \text{fgt}_y^* \gamma$. Recursing at x_1 we obtain:

$$(11) \quad ((\alpha_1 - 1)\psi_1 + \text{ev}_1^* H) \tilde{\gamma} \cap [\mathcal{VZ}_{1,(\alpha-e_1)\cup(1)}(X|Y,\beta)] = \tilde{\gamma} \cap [\mathcal{D}(1)].$$

The left-hand side is a relative invariant with the same degree and number of true markings, but smaller true tangency; hence it has already been computed in the recursion. On the right-hand side, the type I contributions have already been computed except for those of the following form



where C_0 contains a single fictitious marking (if it had multiple fictitious markings then the contribution would vanish by projection formula) and all the other markings are on C_1 . There is one such locus for each fictitious marking, and each contributes $\alpha_1 I$ where I is the invariant we are trying to compute. Note that $\alpha_1 \neq 0$, and that this term appears at least once since y is a fictitious marking; so we obtain a nonzero multiple of I .

Using the same arguments as in Step 1, we see that the type II and \dagger contributions have already been computed. The type III contributions are determined by lower-degree rubber invariants, except for when the whole curve is mapped into Y ; however in this case we may apply the projection formula with fgt_y to identify the contribution with an absolute invariant of Y with degree d and n markings, which has also been computed earlier (in Step 2). We therefore see that (11) may be rearranged to give:

$$I = \text{polynomial in previously computed invariants.}$$

We have thus computed the reduced relative invariant of (X, Y) with degree d , n true markings and true tangency t , which completes Step 3.

Step 4. For the final step, consider a rubber space $\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}|Y_0 + Y_\infty, \beta)$ with insertion γ . Suppose inductively that we have computed all absolute and relative invariants with $d' \leq d$ and all rubber invariants with $(d', n', m') < (d, n, m)$.

We first note the following important reduction: if there exists a relative marking x_k such that all insertions are pulled back along fgt_k , then we may apply the projection formula, together with the following identity (arising from the torsion in the Jacobian of an elliptic curve)

$$(\text{fgt}_k)_* [\mathcal{VZ}_{1,\alpha}^{\leftrightarrow}(\mathbb{P}|Y_0 + Y_\infty, \beta)] = \alpha_k^2 \cdot [\mathcal{VZ}_{1,n+m-1}(Y, \beta)]$$

to identify the rubber invariant with a multiple of a reduced invariant of Y , which has the same degree and hence has been computed earlier.

We will deal with the general case by reducing to the one above. Consider the moduli space

$$\mathcal{VZ}_{1,\alpha\cup(0)}^{\leftrightarrow}(\mathbb{P}|Y_0 + Y_\infty, \beta)$$

obtained by introducing a marked point y with no tangency. Let x_1 be a positive-tangency marking (such a marking always exists since $d \geq 2$) and let $\tilde{\gamma}$ be the insertion obtained from γ by replacing ev_1 and ψ_1 by ev_y and ψ_y , and then pulling back along fgt_1 .

We will make use of a recursion formula for rubber spaces analogous to the recursion formula for relative spaces used in Steps 1–3. Following [Kat07], there is a line bundle

$$L_y^{\text{#bot}} \text{ on } \mathcal{VZ}_{1,\alpha \cup (0)}^{\leftarrow}(\mathbb{P}|Y_0 + Y_\infty, \beta)$$

together with a section $s_y^{\text{#bot}}$ whose vanishing locus consists of the locus $\mathcal{D}(y)$ of rubber maps where y is not mapped into the bottom level of the expanded target. As in §3, we can give a logarithmic interpretation of this: it corresponds to the piecewise-linear function on the tropical moduli space which associates, to every rubber tropical map, the distance between $\varphi(\mathbb{C}_y)$ and the leftmost vertex of the tropical target, where \mathbb{C}_y is the vertex of the source curve containing the flag corresponding to y . Using this tropical description, we may easily calculate the vanishing orders of $s_y^{\text{#bot}}$ along the various components of $\mathcal{D}(y)$, and show that

$$c_1(L_y^{\text{#bot}}) = \Psi_0 - \text{ev}_y^* Y$$

where Ψ_0 is a target psi class (see also [Kat05], where similar results in the non-reduced setting are obtained, using different methods). From this, we obtain a rubber recursion formula

$$(12) \quad (\Psi_0 - \text{ev}_y^* Y) \cdot \tilde{\gamma} \cap [\mathcal{VZ}_{1,\alpha \cup (0)}^{\leftarrow}(\mathbb{P}|Y_0 + Y_\infty, \beta)] = \tilde{\gamma} \cap [\mathcal{D}(y)]$$

where the fundamental class $[\mathcal{D}(y)]$ is weighted by vanishing orders on the components.

Step 4a. We will first show that the left-hand side can be expressed in terms of previously computed invariants. By construction the classes $\tilde{\gamma}$ and $\text{ev}_y^* Y$ are both pulled back along fgt_1 . It remains to examine Ψ_0 . If there exists a negative-tangency marking x_2 , then we have [Gat03, Construction 5.1.17]

$$\Psi_0 = -\alpha_2 \hat{\psi}_2 - \text{ev}_2^* Y$$

where $\hat{\psi}_2$ is a **non-collapsed** psi class. (If there are no negative-tangency markings, then the construction given in [MP06, §1.5.2] shows that $\Psi_0 = 0$.)

Now, $\hat{\psi}_2 - \psi_2$ is given by the locus where x_2 belongs to a trivial bubble. This entails a splitting of the curve into pieces, each of which contributes a rubber integral. Typically each such piece will have $(g', d', n', m') < (1, d, n, m)$ and hence has previously been computed. The one exception is when all of the genus and degree is concentrated on the top level of the expansion, with the bottom level containing only a single non-relative marking in addition to all the negative-tangency markings. But in this case the contribution is a rubber invariant where all of the insertions are pulled back along fgt_1 , and hence we may apply our fundamental reduction, using the projection formula to identify the rubber invariant with (a multiple of) an absolute invariant of Y , which has previously been computed. We conclude that, up to previously computed terms, we may replace $\hat{\psi}_2$ by ψ_2 in the left-hand side of (12).


If we now compare ψ_2 with $\text{fgt}_1^* \psi_2$ we see that the difference is given by the locus where x_1 and x_2 belong to a collapsed rational piece. The contribution of this locus consists of rubber invariants with strictly fewer relative markings, and hence is recursively known. So up to previously computed invariants, (12) becomes

$$(13) \quad \text{fgt}_1^*(-\alpha_2 \psi_2 - \text{ev}_y^* Y) \tilde{\gamma} \cap [\mathcal{VZ}_{1,\alpha \cup (0)}^{\leftarrow}(\mathbb{P}|Y_0 + Y_\infty, \beta)] = \tilde{\gamma} \cap [\mathcal{D}(y)]$$

and now the left-hand side can also be expressed in terms of previously computed invariants, via the projection formula.

Step 4b. Let us now examine the right-hand side of (13). The components of $\mathcal{D}(y)$ are indexed by splittings of the curve, and the contributions can be expressed in terms of previously computed invariants unless there is a piece of the curve carrying all of the genus and degree. We therefore restrict to considering these cases. Denote the piece of the curve carrying all of the genus and degree by $C' \subseteq C$.

- (1) If C' is mapped to top level, then either it contains x_1 , in which case we apply fgt_1 to compute the contribution, or it does not contain x_1 , in which case it has fewer than n relative markings and is known recursively.
- (2) If on the other hand C' is not mapped to top level, then generically it is mapped to the bottom level (since generically the core is not contracted on this locus, so the desingularisation process does nothing). One possible contribution is given by the following locus



with all other markings located on the genus one vertex. This contributes a nonzero multiple of the invariant I we are trying to compute.

For the other possibilities, first note that, unless every component at the top level contains a single positive-tangency marking and a single node (together with possibly some tangency-zero markings), then C' has fewer than n relative markings and hence the contribution is known recursively. On the other hand, if any non-relative marking other than y is mapped to the top level, C' has $\leq n$ relative markings and less than m non-relative markings, and hence again the contribution is known recursively. The only remaining possibilities are when x_1 is replaced by another positive-tangency marking x_k in the above picture; but then the contribution can be calculated by applying the projection formula with fgt_1 .

We therefore conclude that the only contribution to the right-hand side of (13) which cannot be expressed in terms of previously computed invariants is a nonzero multiple of the invariant we were trying to compute. We thus compute the reduced rubber invariant of \mathbb{P} with degree d , n relative markings and m non-relative markings, which completes Step 4 and the recursion.

REFERENCES

- [AC14] D. Abramovich and Q. Chen. Stable logarithmic maps to Deligne-Faltings pairs II. *Asian J. Math.*, 18(3):465–488, 2014. 7
- [ACGS17] D. Abramovich, Q. Chen, M. Gross, and B. Siebert. Decomposition of degenerate Gromov-Witten invariants. *arXiv e-prints*, page arXiv:1709.09864, Sep 2017. 20
- [ACP15] D. Abramovich, L. Caporaso, and S. Payne. The tropicalization of the moduli space of curves. *Ann. Sci. Éc. Norm. Supér. (4)*, 48(4):765–809, 2015. 25
- [AK00] D. Abramovich and K. Karu. Weak semistable reduction in characteristic 0. *Invent. Math.*, 139(2):241–273, 2000. 24
- [AMW14] D. Abramovich, S. Marcus, and J. Wise. Comparison theorems for Gromov-Witten invariants of smooth pairs and of degenerations. *Ann. Inst. Fourier (Grenoble)*, 64(4):1611–1667, 2014. 19
- [AW18] D. Abramovich and J. Wise. Birational invariance in logarithmic Gromov-Witten theory. *Compos. Math.*, 154(3):595–620, 2018. 8
- [BC20] Luca Battistella and Francesca Carocci. A smooth compactification of the space of genus two curves in projective space via logarithmic geometry and Gorenstein curves. *arXiv e-prints*, page arXiv:2008.13506, August 2020. 1
- [BCM20] Luca Battistella, Francesca Carocci, and Cristina Manolache. Reduced invariants from cuspidal maps. *Trans. Amer. Math. Soc.*, 373(9):6713–6756, 2020. 1
- [BN20] Luca Battistella and Navid Nabijou. Relative Quasimaps and Mirror Formulae. *International Mathematics Research Notices*, January 2020. rnz339. 13
- [CCUW20] Renzo Cavalieri, Melody Chan, Martin Ulirsch, and Jonathan Wise. A moduli stack of tropical curves. *Forum Math. Sigma*, Vol. 8:e23, 93 pages, 2020. 4, 16, 24
- [CH98] L. Caporaso and J. Harris. Counting plane curves of any genus. *Invent. Math.*, 131(2):345–392, 1998. 2
- [Che14a] Q. Chen. The degeneration formula for logarithmic expanded degenerations. *J. Algebraic Geom.*, 23(2):341–392, 2014. 11, 20
- [Che14b] Q. Chen. Stable logarithmic maps to Deligne-Faltings pairs I. *Ann. of Math. (2)*, 180(2):455–521, 2014. 7, 8

- [Ful98] William Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998. 30
- [Gat02] A. Gathmann. Absolute and relative Gromov-Witten invariants of very ample hypersurfaces. *Duke Math. J.*, 115(2):171–203, 2002. 2, 13, 16, 29, 32, 34
- [Gat03] A. Gathmann. Gromov-Witten invariants of hypersurfaces, 2003. Habilitation thesis. 32, 38
- [GS13] M. Gross and B. Siebert. Logarithmic Gromov-Witten invariants. *J. Amer. Math. Soc.*, 26(2):451–510, 2013. 4, 7
- [Hai13] R. Hain. Normal functions and the geometry of moduli spaces of curves. In *Handbook of moduli. Vol. I*, volume 24 of *Adv. Lect. Math. (ALM)*, pages 527–578. Int. Press, Somerville, MA, 2013. 33
- [HL10] Y. Hu and J. Li. Genus-one stable maps, local equations, and Vakil-Zinger’s desingularization. *Math. Ann.*, 348(4):929–963, 2010. 1
- [JPPZ17] F. Janda, R. Pandharipande, A. Pixton, and D. Zvonkine. Double ramification cycles on the moduli spaces of curves. *Publ. Math. Inst. Hautes Études Sci.*, 125:221–266, 2017. 33
- [JPPZ18] F. Janda, R. Pandharipande, A. Pixton, and D. Zvonkine. Double ramification cycles with target varieties. *arXiv e-prints*, page arXiv:1812.10136, December 2018. 3
- [Kat05] E. Katz. Line-Bundles on Stacks of Relative Maps. *arXiv Mathematics e-prints*, page math/0507322, Jul 2005. 38
- [Kat07] E. Katz. An algebraic formulation of symplectic field theory. *J. Symplectic Geom.*, 5(4):385–437, 2007. 38
- [Kim10] B. Kim. Logarithmic stable maps. In *New developments in algebraic geometry, integrable systems and mirror symmetry (RIMS, Kyoto, 2008)*, volume 59 of *Adv. Stud. Pure Math.*, pages 167–200. Math. Soc. Japan, Tokyo, 2010. 7, 8
- [Li01] J. Li. Stable morphisms to singular schemes and relative stable morphisms. *J. Differential Geom.*, 57(3):509–578, 2001. 8
- [Li02] J. Li. A degeneration formula of GW-invariants. *J. Differential Geom.*, 60(2):199–293, 2002. 8
- [Liu02] Qing Liu. *Algebraic geometry and arithmetic curves*, volume 6 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2002. Translated from the French by Reinie Ern , Oxford Science Publications. 6
- [LR18] Yoav Len and Dhruv Ranganathan. Enumerative geometry of elliptic curves on toric surfaces. *Israel J. Math.*, 226(1):351–385, 2018. 3
- [LZ07] J. Li and A. Zinger. Gromov-Witten invariants of a quintic threefold and a rigidity conjecture. *Pacific J. Math.*, 233(2):417–480, 2007. 1
- [LZ09] J. Li and A. Zinger. On the genus-one Gromov-Witten invariants of complete intersections. *J. Differential Geom.*, 82(3):641–690, 2009. 1
- [Mol16] S. Molcho. Universal Weak Semistable Reduction. *arXiv e-prints*, page arXiv:1601.00302, Jan 2016. 24
- [MP06] D. Maulik and R. Pandharipande. A topological view of Gromov-Witten theory. *Topology*, 45(5):887–918, 2006. 2, 38
- [MW20] Steffen Marcus and Jonathan Wise. Logarithmic compactification of the Abel-Jacobi section. *Proc. Lond. Math. Soc. (3)*, 121(5):1207–1250, 2020. 12
- [Ran19] D. Ranganathan. Logarithmic Gromov-Witten theory with expansions. *arXiv:1903.09006*, 2019. 3, 8
- [RSPW19a] Dhruv Ranganathan, Keli Santos-Parker, and Jonathan Wise. Moduli of stable maps in genus one and logarithmic geometry, I. *Geom. Topol.*, 23(7):3315–3366, 2019. 1, 3, 5, 6, 7, 9, 10, 11, 13, 16, 26
- [RSPW19b] Dhruv Ranganathan, Keli Santos-Parker, and Jonathan Wise. Moduli of stable maps in genus one and logarithmic geometry, II. *Algebra Number Theory*, 13(8):1765–1805, 2019. 1, 3, 5, 10, 11, 13, 14
- [RW19] D. Ranganathan and J. Wise. Rational curves in the logarithmic multiplicative group. *Proc. Amer. Math. Soc.*, to appear, 2019. 12
- [Smy11a] D. I. Smyth. Modular compactifications of the space of pointed elliptic curves I. *Compos. Math.*, 147(3):877–913, 2011. 1, 5, 6
- [Smy11b] D. I. Smyth. Modular compactifications of the space of pointed elliptic curves II. *Compos. Math.*, 147(6):1843–1884, 2011. 5, 21, 26
- [Spe14] David Speyer. Parameterizing tropical curves, i: Curves of genus zero and one. *Algebra & Number Theory*, 8(4):963–998, 2014. 3
- [Thu07] A. Thuillier. G om trie toro dale et g om trie analytique non archim dienne. Application au type d’homotopie de certains sch mas formels. *Manuscripta Math.*, 123(4):381–451, 2007. 25
- [Vak00] R. Vakil. The enumerative geometry of rational and elliptic curves in projective space. *J. Reine Angew. Math.*, 529:101–153, 2000. 2, 9, 17, 18
- [vdW10] Frederick van der Wyck. *Moduli of singular curves and crimping*. PhD thesis, Harvard University, 2010. 5
- [Vis12] M. Viscardi. Alternate compactifications of the moduli space of genus one maps. *Manuscripta Math.*, 139(1-2):201–236, 2012. 1

- [VZ08] R. Vakil and A. Zinger. A desingularization of the main component of the moduli space of genus-one stable maps into \mathbb{P}^N . *Geom. Topol.*, 12(1):1–95, 2008. [1](#), [27](#), [30](#)
- [Zin08] A. Zinger. Standard versus reduced genus-one gromov–witten invariants. *Geometry and Topology*, 12(2):1203–1241, 2008. [1](#)
- [Zin09a] A. Zinger. Reduced genus-one Gromov-Witten invariants. *J. Differential Geom.*, 83(2):407–460, 2009. [1](#)
- [Zin09b] A. Zinger. The reduced genus one Gromov-Witten invariants of Calabi-Yau hypersurfaces. *J. Amer. Math. Soc.*, (22):691–737, 2009. [1](#)

Luca Battistella
Mathematisches Institut, Universität Heidelberg
lbattistella@mathi.uni-heidelberg.de

Navid Nabijou
Department of Pure Mathematics and Mathematical Statistics, University of Cambridge
nn333@cam.ac.uk

Dhruv Ranganathan
Department of Pure Mathematics and Mathematical Statistics, University of Cambridge
dr508@cam.ac.uk