

US010138482B2

(12) **United States Patent**
Rigo

(10) **Patent No.:** **US 10,138,482 B2**
(45) **Date of Patent:** **Nov. 27, 2018**

(54) **COMPOSITIONS FOR MODULATING
C9ORF72 EXPRESSION**

(71) Applicant: **Ionis Pharmaceuticals, Inc.**, Carlsbad,
CA (US)

(72) Inventor: **Frank Rigo**, Carlsbad, CA (US)

(73) Assignee: **Ionis Pharmaceuticals, Inc.**, Carlsbad,
CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/404,979**

(22) Filed: **Jan. 12, 2017**

(65) **Prior Publication Data**

US 2017/0349897 A1 Dec. 7, 2017

Related U.S. Application Data

(63) Continuation of application No. 15/130,818, filed on
Apr. 15, 2016, now Pat. No. 9,605,263.

(60) Provisional application No. 62/239,400, filed on Oct.
9, 2015, provisional application No. 62/232,941, filed
on Sep. 25, 2015, provisional application No.
62/148,691, filed on Apr. 16, 2015.

(51) **Int. Cl.**

C07H 21/04 (2006.01)

C07H 21/02 (2006.01)

C12N 15/113 (2010.01)

(52) **U.S. Cl.**

CPC **C12N 15/113** (2013.01); **C12N 2310/11**
(2013.01); **C12N 2310/315** (2013.01); **C12N**
2310/322 (2013.01); **C12N 2310/3231**
(2013.01); **C12N 2310/3341** (2013.01); **C12N**
2310/341 (2013.01); **C12N 2310/346**
(2013.01); **C12N 2320/30** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

9,963,699 B2 5/2018 Bennett et al.
2004/0038274 A1 2/2004 Cook et al.
2010/0216864 A1 8/2010 Straarup et al.
2013/0035366 A1* 2/2013 Swayze C12N 15/1131
514/44 A
2016/0251655 A1* 9/2016 Freier G01N 33/6896
514/44 A

FOREIGN PATENT DOCUMENTS

WO WO 1996/014329 5/1996
WO WO 2005/113016 12/2005

WO WO 2005/121368 12/2005
WO WO 2007/089584 8/2007
WO WO 2007/131237 11/2007
WO WO 2008/076324 6/2008
WO WO 2009/007855 1/2009
WO WO 2012/005898 1/2012
WO WO 2012/012443 1/2012
WO WO 2012/087983 6/2012
WO WO 2012/135736 10/2012
WO WO 2013/036833 3/2013
WO WO-2015054676 A2* 4/2015 G01N 33/6896
WO WO 2017/079291 5/2017
WO WO 2017/180835 10/2017

OTHER PUBLICATIONS

Fernandes et al., "Oligonucleotide-Based Therapy for FTD/ALS
Caused by the C9orf72 Repeat Expansion: A Perspective" Journal
of Nucleic Acids (2013) :1-11.

Gendron et al., "Poly(GP) proteins are a useful pharmacodynamic
marker for C9ORF72-associated amyotrophic lateral sclerosis" Sci
Tran Med (2017) 9(383):1-12.

Lee et al., "Antisense Therapy in Neurology" Journal of Personal-
ized Medicine (2013) 3(3): 144-176.

Sha et al., "Treatment implications of C9ORF72" Alzheimers Res
Ther (2012) 4(6): 46.

Thomsen, "Dramatically improved RNA in 1-15 situ hybridization
signals using LNA-modified probes" RNA (2005) 11(11): 1745-
1748.

Ash et al., "Unconventional Translation of C9ORF72 GGGGCC
Expansion Generates Insoluble Ploypeptides Specific to c9FTD/
ALS" Neuron (2013) 77(4): 639-646.

Extended European Search Report for application No. 14852924.1
dated Jun. 20, 2017.

International Search Report for application No. PCT/US17/27355
dated Jul. 26, 2017.

Mahoney et al., "Frontotemporal dementia with the C9ORF72
hexanucleotide repeat expansion: clinical, neuroanatomical and
neuropathological features" Brain (2012) 135: 736-750.

GenBank: Accession No. NT_008413, Jul. 24, 2012, 5 pages.

International Search Report for Application No. PCT/US2016/
060106, dated Feb. 1, 2017, 10 pages.

Bennett et al., "Antisense oligonucleotides as a tool for gene
functionalization and target validation," Biochimica et Biophysica
Acta (1999) 1489: 19-30.

Jeong et al., "Rapid Identification of Monospecific Monoclonal
Antibodies Using a Human Proteome Microarray." Mol. Cell.
Proteomics (2012) 11(6): O111.016253-1 to O111.016253-10.

Watts et al., "Silencing disease genes in the laboratory and the
clinic" J Pathol (2012) 226(2): 365-379.

* cited by examiner

Primary Examiner — Richard A Schnizer

(74) *Attorney, Agent, or Firm* — McNeill Baur PLLC

(57) **ABSTRACT**

Disclosed herein are compositions and methods for reducing
expression of C9ORF72 mRNA and protein in an animal.
Such methods are useful to treat, prevent, ameliorate, or
slow progression of neurodegenerative diseases in an indi-
vidual in need thereof.

20 Claims, No Drawings

Specification includes a Sequence Listing.

1

**COMPOSITIONS FOR MODULATING
C9ORF72 EXPRESSION**

SEQUENCE LISTING

The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled BIOL0269USC 1SEQ_ST25.txt created Jan. 12, 2017, which is 104 Kb in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

FIELD

Provided are compositions and methods for modulating expression of C9ORF72 mRNA and protein in cells and animals. Such compositions and methods are useful to treat, prevent, ameliorate, or slow progression of neurodegenerative diseases, including amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, and olivopontocerebellar degeneration (OPCD).

BACKGROUND

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease characterized clinically by progressive paralysis leading to death from respiratory failure, typically within two to three years of symptom onset (Rowland and Shneider, *N. Engl. J. Med.*, 2001, 344, 1688-1700). ALS is the third most common neurodegenerative disease in the Western world (Hirtz et al., *Neurology*, 2007, 68, 326-337), and there are currently no effective therapies. Approximately 10% of cases are familial in nature, whereas the bulk of patients diagnosed with the disease are classified as sporadic as they appear to occur randomly throughout the population (Chio et al., *Neurology*, 2008, 70, 533-537). There is growing recognition, based on clinical, genetic, and epidemiological data, that ALS and frontotemporal dementia (FTD) represent an overlapping continuum of disease, characterized pathologically by the presence of TDP-43 positive inclusions throughout the central nervous system (Lillo and Hodges, *J. Clin. Neurosci.*, 2009, 16, 1131-1135; Neumann et al., *Science*, 2006, 314, 130-133).

To date, a number of genes have been discovered as causative for classical familial ALS, for example, SOD1, TARDBP, FUS, OPTN, and VCP (Johnson et al., *Neuron*, 2010, 68, 857-864; Kwiatkowski et al., *Science*, 2009, 323, 1205-1208; Maruyama et al., *Nature*, 2010, 465, 223-226; Rosen et al., *Nature*, 1993, 362, 59-62; Sreedharan et al., *Science*, 2008, 319, 1668-1672; Vance et al., *Brain*, 2009, 129, 868-876). Recently, linkage analysis of kindreds involving multiple cases of ALS, FTD, and ALS-FTD had suggested that there was an important locus for the disease on the short arm of chromosome 9 (Boxer et al., *J. Neurol. Neurosurg. Psychiatry*, 2011, 82, 196-203; Morita et al., *Neurology*, 2006, 66, 839-844; Pearson et al. *J. Neurol.*, 2011, 258, 647-655; Vance et al., *Brain*, 2006, 129, 868-876). The mutation in the C9ORF72 gene is the most common genetic cause of ALS and FTD. The ALS-FTD causing mutation is a large hexanucleotide (GGGGCC) repeat expansion in the first intron of the C9ORF72 gene (Renton et al., *Neuron*, 2011, 72, 257-268; DeJesus-Hernandez et al., *Neuron*, 2011, 72, 245-256). A founder haplotype, covering the C9ORF72 gene, is present in the majority of cases linked to this region (Renton et al., *Neuron*, 2011, 72, 257-268). This locus on chromosome 9p21 accounts for nearly half of familial ALS

2

and nearly one-quarter of all ALS cases in a cohort of 405 Finnish patients (Laaksovirta et al, *Lancet Neurol.*, 2010, 9, 978-985).

A founder haplotype, covering the C9ORF72 gene, is present in the majority of cases linked to this region.

There are currently no effective therapies to treat such neurodegenerative diseases. Therefore, it is an object to provide compositions and methods for the treatment of such neurodegenerative diseases.

SUMMARY

Certain embodiments provide methods, compounds, and compositions for inhibiting expression of C9ORF72 mRNA and protein in cells, tissues, and animals. Certain embodiments provide methods, compounds, and compositions for reducing C9ORF72 mRNA and protein levels in cells, tissues, and animals. Certain embodiments provide antisense compounds targeted to a C9ORF72 nucleic acid. In certain embodiments, the antisense compounds are modified oligonucleotides. In certain embodiments, the modified oligonucleotides are single-stranded.

In certain embodiments, C9ORF72 associated Repeat Associated Non-ATG Translation (RAN translation) products are reduced. In certain embodiments, the C9ORF72 associated RAN translation products are poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine). In certain embodiments, certain C9ORF72 mRNA variants are preferentially reduced. In certain embodiments, the C9ORF72 mRNA variants preferentially reduced are variants processed from a pre-mRNA containing intron 1. In certain embodiments, intron 1 contains a hexanucleotide repeat expansion. In certain embodiments, the C9ORF72 mRNA variant preferentially reduced is a C9ORF72 pathogenic associated mRNA variant. In certain embodiments, the C9ORF72 pathogenic associated mRNA variant is NM_001256054.1 (SEQ ID NO: 1). In certain embodiments, the hexanucleotide repeat expansion is associated with a C9ORF72 associated disease. In certain embodiments, the hexanucleotide repeat expansion is associated with a C9ORF72 hexanucleotide repeat expansion associated disease. In certain embodiments, the hexanucleotide repeat expansion comprises at least 30 GGGGCC repeats, more than 30 GGGGCC repeats, more than 100 GGGGCC repeats, more than 500 GGGGCC repeats, or more than 1000 GGGGCC repeats. In certain embodiments, the hexanucleotide repeat expansion is associated with nuclear foci. In certain embodiments, C9ORF72 associated RAN translation products are associated with nuclear foci. In certain embodiments, the C9ORF72 associated RAN translation products are poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine). In certain embodiments, the compositions and methods described herein are useful for reducing C9ORF72 mRNA levels, C9ORF72 protein levels, C9ORF72 RAN translation products, and nuclear foci. In certain embodiments, the compositions and methods described herein are useful for selectively reducing C9ORF72 pathogenic associated mRNA variants. Such reduction can occur in a time-dependent manner or in a dose-dependent manner.

Also provided are methods useful for preventing, treating, ameliorating, and slowing progression of diseases associated with C9ORF72. In certain embodiments, such C9ORF72 associated diseases are neurodegenerative diseases. In certain embodiments, the neurodegenerative disease is amyotrophic lateral sclerosis (ALS), frontotemporal dementia

(FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, and olivopontocerebellar degeneration (OPCD).

Such diseases can have one or more risk factors, causes, or outcomes in common. Certain risk factors and causes for development of a neurodegenerative disease, and, in particular, ALS and FTD, include genetic predisposition and older age.

In certain embodiments, methods of treatment include administering a C9ORF72 antisense compound to an individual in need thereof. In certain embodiments, the antisense compound is a single-stranded modified oligonucleotide. In certain embodiments, the single-stranded modified oligonucleotide is complementary to a C9ORF72 nucleic acid.

DETAILED DESCRIPTION

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed. Herein, the use of the singular includes the plural unless specifically stated otherwise. As used herein, the use of “or” means “and/or” unless stated otherwise. Additionally, as used herein, the use of “and” means “and/or” unless stated otherwise. Furthermore, the use of the term “including” as well as other forms, such as “includes” and “included”, is not limiting. Also, terms such as “element” or “component” encompass both elements and components comprising one unit and elements and components that comprise more than one subunit, unless specifically stated otherwise.

Compounds of the invention include variations of the disclosed compounds in which one or more hydrogen, carbon, nitrogen, oxygen, or sulfur atoms is replaced with a stable isotope of the same element.

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All documents, or portions of documents, cited in this disclosure, including, but not limited to, patents, patent applications, published patent applications, articles, books, treatises, and GENBANK Accession Numbers and associated sequence information obtainable through databases such as National Center for Biotechnology Information (NCBI) and other data referred to throughout in the disclosure herein are hereby expressly incorporated by reference for the portions of the document discussed herein, as well as in their entirety.

Definitions

Unless specific definitions are provided, the nomenclature utilized in connection with, and the procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical synthesis, and chemical analysis.

Unless otherwise indicated, the following terms have the following meanings:

“2'-O-methoxyethyl” (also 2'-MOE and 2'-OCH₂CH₂—OCH₃ and MOE) refers to an O-methoxy-ethyl modification of the 2' position of a furanose ring. A 2'-O-methoxyethyl modified sugar is a modified sugar.

“2'-MOE nucleoside” (also 2'-O-methoxyethyl nucleoside) means a nucleoside comprising a MOE modified sugar moiety.

“2'-substituted nucleoside” means a nucleoside comprising a substituent at the 2'-position of the furanose ring other than H or OH. In certain embodiments, 2'-substituted nucleosides include nucleosides with bicyclic sugar modifications.

“5-methylcytosine” means a cytosine modified with a methyl group attached to the 5' position. A 5-methylcytosine is a modified nucleobase.

“Administered concomitantly” refers to the co-administration of two pharmaceutical agents in any manner in which the pharmacological effects of both are manifest in the patient at the same time. Concomitant administration does not require that both pharmaceutical agents be administered in a single pharmaceutical composition, in the same dosage form, or by the same route of administration. The effects of both pharmaceutical agents need not manifest themselves at the same time. The effects need only be overlapping for a period of time and need not be coextensive.

“Administering” means providing a pharmaceutical agent to an animal, and includes, but is not limited to administering by a medical professional and self-administering.

“Amelioration” refers to a lessening, slowing, stopping, or reversing of at least one indicator of the severity of a condition or disease. The severity of indicators may be determined by subjective or objective measures, which are known to those skilled in the art.

“Animal” refers to a human or non-human animal, including, but not limited to, mice, rats, rabbits, dogs, cats, pigs, and non-human primates, including, but not limited to, monkeys and chimpanzees.

“Antibody” refers to a molecule characterized by reacting specifically with an antigen in some way, where the antibody and the antigen are each defined in terms of the other. Antibody may refer to a complete antibody molecule or any fragment or region thereof, such as the heavy chain, the light chain, Fab region, and Fc region.

“Antisense activity” means any detectable or measurable activity attributable to the hybridization of an antisense compound to its target nucleic acid. In certain embodiments, antisense activity is a decrease in the amount or expression of a target nucleic acid or protein encoded by such target nucleic acid.

“Antisense compound” means an oligomeric compound that is capable of undergoing hybridization to a target nucleic acid through hydrogen bonding. Examples of antisense compounds include single-stranded and double-stranded compounds, such as, antisense oligonucleotides, siRNAs, shRNAs, ssRNAs, and occupancy-based compounds.

“Antisense inhibition” means reduction of target nucleic acid levels in the presence of an antisense compound complementary to a target nucleic acid compared to target nucleic acid levels or in the absence of the antisense compound.

“Antisense mechanisms” are all those mechanisms involving hybridization of a compound with a target nucleic acid, wherein the outcome or effect of the hybridization is either target degradation or target occupancy with concomitant stalling of the cellular machinery involving, for example, transcription or splicing.

“Antisense oligonucleotide” means a single-stranded oligonucleotide having a nucleobase sequence that permits hybridization to a corresponding segment of a target nucleic acid.

“Base complementarity” refers to the capacity for the precise base pairing of nucleobases of an antisense oligonucleotide with corresponding nucleobases in a target

5

nucleic acid (i.e., hybridization), and is mediated by Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen binding between corresponding nucleobases.

“Bicyclic sugar” means a furanose ring modified by the bridging of two atoms. A bicyclic sugar is a modified sugar.

“Bicyclic nucleoside” (also BNA) means a nucleoside having a sugar moiety comprising a bridge connecting two carbon atoms of the sugar ring, thereby forming a bicyclic ring system. In certain embodiments, the bridge connects the 4'-carbon and the 2'-carbon of the sugar ring.

“C9ORF72 antisense transcript” means transcripts produced from the non-coding strand (also antisense strand and template strand) of the C9ORF72 gene. The C9ORF72 antisense transcript differs from the canonically transcribed “C9ORF72 sense transcript”, which is produced from the coding strand (also sense strand) of the C9ORF72 gene. In certain embodiments, a C9ORF72 antisense transcript is SEQ ID NO: 18.

“C9ORF72 antisense transcript specific inhibitor” refers to any agent capable of specifically inhibiting the expression of C9ORF72 antisense transcript and/or its expression products at the molecular level. As used herein, “specific” means reducing or inhibiting expression of C9ORF72 antisense transcript without reducing non-target transcript to an appreciable degree (e.g., a C9ORF72 antisense transcript specific inhibitor reduces expression of C9ORF72 antisense transcript, but does not reduce expression of C9ORF72 sense transcript to an appreciable degree). C9ORF72 specific antisense transcript inhibitors include antisense compounds, siRNAs, aptamers, antibodies, peptides, small molecules, and other agents capable of inhibiting the expression of C9ORF72 antisense transcript and/or its expression products, such as C9ORF72 antisense transcript associated RAN translation products.

“C9ORF72 associated disease” means any disease associated with any C9ORF72 nucleic acid or expression product thereof. Such diseases may include a neurodegenerative disease. Such neurodegenerative diseases may include ALS and FTD. In certain embodiments, the C9ORF72 associated disease is caused by (or is associated with) a hexanucleotide repeat expansion. In certain embodiments, the hexanucleotide repeat expansion may comprise GGGGCC, GGGGGG, GGGGGC, or GGGGCG repeated at least 30 times, more than 30 times, more than 100 times, more than 500 times, or more than 1000 times.

“C9ORF72 associated RAN translation products” means aberrant peptide or di-peptide polymers translated through RAN translation (i.e., repeat-associated, and non-ATG-dependent translation). In certain embodiments, the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).

“C9ORF72 nucleic acid” means any nucleic acid encoding C9ORF72. For example, in certain embodiments, a C9ORF72 nucleic acid includes a DNA sequence encoding C9ORF72, an RNA sequence transcribed from DNA encoding C9ORF72 including genomic DNA comprising introns and exons (i.e., pre-mRNA), and an mRNA sequence encoding C9ORF72. “C9ORF72 mRNA” means an mRNA encoding a C9ORF72 protein.

“C9ORF72 pathogenic associated mRNA variant” means the C9ORF72 mRNA variant processed from a C9ORF72 pre-mRNA variant containing the hexanucleotide repeat. A C9ORF72 pre-mRNA contains the hexanucleotide repeat when transcription of the pre-mRNA begins in the region from the start site of exon 1A to the start site of exon 1B, e.g., nucleotides 1107 to 1520 of the genomic sequence

6

(SEQ ID NO: 2, the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 27565000). In certain embodiments, the level of a C9ORF72 pathogenic associated mRNA variant is measured to determine the level of a C9ORF72 pre-mRNA containing the hexanucleotide repeat in a sample.

“C9ORF72 specific inhibitor” refers to any agent capable of specifically inhibiting the expression of C9ORF72 mRNA and/or C9ORF72 protein at the molecular level. For example, C9ORF72 specific inhibitors include nucleic acids (including antisense compounds), siRNAs, aptamers, antibodies, peptides, small molecules, and other agents capable of inhibiting the expression of C9ORF72 mRNA and/or C9ORF72 protein. Similarly, in certain embodiments, C9ORF72 specific inhibitors may affect other molecular processes in an animal.

“Cap structure” or “terminal cap moiety” means chemical modifications, which have been incorporated at either terminus of an antisense compound.

“cEt” or “constrained ethyl” means a bicyclic nucleoside having a sugar moiety comprising a bridge connecting the 4'-carbon and the 2'-carbon, wherein the bridge has the formula: 4'-CH(CH₃)—O-2'.

“Constrained ethyl nucleoside” (also cEt nucleoside) means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH(CH₃)—O-2' bridge.

“Chemically distinct region” refers to a region of an antisense compound that is in some way chemically different than another region of the same antisense compound. For example, a region having 2'-O-methoxyethyl nucleosides is chemically distinct from a region having nucleosides without 2'-O-methoxyethyl modifications.

“Chimeric antisense compound” means an antisense compound that has at least two chemically distinct regions, each position having a plurality of subunits.

“Co-administration” means administration of two or more pharmaceutical agents to an individual. The two or more pharmaceutical agents may be in a single pharmaceutical composition, or may be in separate pharmaceutical compositions. Each of the two or more pharmaceutical agents may be administered through the same or different routes of administration. Co-administration encompasses parallel or sequential administration.

“Complementarity” means the capacity for pairing between nucleobases of a first nucleic acid and a second nucleic acid.

“Comprise,” “comprises,” and “comprising” will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

“Contiguous nucleobases” means nucleobases immediately adjacent to each other.

“Designing” or “designed to” refer to the process of designing an oligomeric compound that specifically hybridizes with a selected nucleic acid molecule.

“Diluent” means an ingredient in a composition that lacks pharmacological activity, but is pharmaceutically necessary or desirable. For example, in drugs that are injected, the diluent may be a liquid, e.g. saline solution.

“Dose” means a specified quantity of a pharmaceutical agent provided in a single administration, or in a specified time period. In certain embodiments, a dose may be administered in one, two, or more boluses, tablets, or injections. For example, in certain embodiments where subcutaneous administration is desired, the desired dose requires a volume not easily accommodated by a single injection, therefore, two or more injections may be used to achieve the desired

dose. In certain embodiments, the pharmaceutical agent is administered by infusion over an extended period of time or continuously. Doses may be stated as the amount of pharmaceutical agent per hour, day, week, or month.

“Effective amount” in the context of modulating an activity or of treating or preventing a condition means the administration of that amount of pharmaceutical agent to a subject in need of such modulation, treatment, or prophylaxis, either in a single dose or as part of a series, that is effective for modulation of that effect, or for treatment or prophylaxis or improvement of that condition. The effective amount may vary among individuals depending on the health and physical condition of the individual to be treated, the taxonomic group of the individuals to be treated, the formulation of the composition, assessment of the individual’s medical condition, and other relevant factors.

“Efficacy” means the ability to produce a desired effect.

“Expression” includes all the functions by which a gene’s coded information is converted into structures present and operating in a cell. Such structures include, but are not limited to the products of transcription and translation.

“Focus” or “foci” means a nuclear or cytoplasmic body comprising a C9ORF72 transcript.

In certain embodiments, a focus comprises at least one C9ORF72 transcript. In certain embodiments, C9ORF72 foci comprise transcripts comprising any of the following hexanucleotide repeats: GGGGCC, GGGGGG, GGGGCG, and/or GGGGCG.

“Fully complementary” or “100% complementary” means each nucleobase of a first nucleic acid has a complementary nucleobase in a second nucleic acid. In certain embodiments, a first nucleic acid is an antisense compound and a target nucleic acid is a second nucleic acid.

“Gapmer” means a chimeric antisense compound in which an internal region having a plurality of nucleosides that support RNase H cleavage is positioned between external regions having one or more nucleosides, wherein the nucleosides comprising the internal region are chemically distinct from the nucleoside or nucleosides comprising the external regions. The internal region may be referred to as a “gap” and the external regions may be referred to as the “wings.”

“Hexanucleotide repeat expansion” means a series of six bases (for example, GGGGCC, GGGGGG, GGGGCG, or GGGGGC) repeated at least twice. In certain embodiments, the hexanucleotide repeat expansion may be located in intron 1 of a C9ORF72 nucleic acid. In certain embodiments, a pathogenic hexanucleotide repeat expansion includes at least 30, more than 30, more than 100, more than 500, or more than 1000 repeats of GGGGCC, GGGGGG, GGGGCG, or GGGGGC in a C9ORF72 nucleic acid and is associated with disease. In certain embodiments, the repeats are consecutive. In certain embodiments, the repeats are interrupted by 1 or more nucleobases. In certain embodiments, a wild-type hexanucleotide repeat expansion includes 23 or fewer repeats of GGGGCC, GGGGGG, GGGGCG, or GGGGGC in a C9ORF72 nucleic acid. In certain embodiments, the repeats are consecutive. In certain embodiments, the repeats are interrupted by 1 or more nucleobases.

“Hybridization” means the annealing of complementary nucleic acid molecules. In certain embodiments, complementary nucleic acid molecules include, but are not limited to, an antisense compound and a target nucleic acid. In certain embodiments, complementary nucleic acid molecules include, but are not limited to, an antisense oligonucleotide and a nucleic acid target.

“Identifying an animal having a C9ORF72 associated disease” means identifying an animal having been diagnosed with a C9ORF72 associated disease or predisposed to develop a C9ORF72 associated disease. Individuals predisposed to develop a C9ORF72 associated disease include those having one or more risk factors for developing a C9ORF72 associated disease, including, having a personal or family history or genetic predisposition of one or more C9ORF72 associated diseases. Such identification may be accomplished by any method including evaluating an individual’s medical history and standard clinical tests or assessments, such as genetic testing.

“Immediately adjacent” means there are no intervening elements between the immediately adjacent elements.

“Individual” means a human or non-human animal selected for treatment or therapy.


“Inhibiting C9ORF72” means reducing the level or expression of a C9ORF72 mRNA and/or protein. In certain embodiments, C9ORF72 mRNA and/or protein levels are inhibited in the presence of an antisense compound targeting C9ORF72, including an antisense oligonucleotide targeting C9ORF72, as compared to expression of C9ORF72 mRNA and/or protein levels in the absence of a C9ORF72 antisense compound, such as an antisense oligonucleotide.

“Inhibiting the expression or activity” refers to a reduction or blockade of the expression or activity and does not necessarily indicate a total elimination of expression or activity.

“Internucleoside linkage” refers to the chemical bond between nucleosides.


“Linked nucleosides” means adjacent nucleosides linked together by an internucleoside linkage.

“Locked nucleic acid” or “LNA” or “LNA nucleosides” means nucleic acid monomers having a bridge connecting two carbon atoms between the 4' and 2' position of the nucleoside sugar unit, thereby forming a bicyclic sugar. Examples of such bicyclic sugar include, but are not limited to (A) α -L-Methyleneoxy (4'-CH₂—O-2') LNA, (B) β -D-Methyleneoxy (4'-CH₂—O-2') LNA, (C) Ethyleneoxy (4'-(CH₂)₂—O-2') LNA, (D) Aminooxy (4'-CH₂—O—N(R)-2') LNA and (E) Oxyamino (4'-CH₂—N(R)—O-2') LNA, as depicted below.




9

-continued



(D)



(E)

As used herein, LNA compounds include, but are not limited to, compounds having at least one bridge between the 4' and the 2' position of the sugar wherein each of the bridges independently comprises 1 or from 2 to 4 linked groups independently selected from $-\text{C}(\text{R}_1)(\text{R}_2)_n-$, $-\text{C}(\text{R}_1)=\text{C}(\text{R}_2)-$, $-\text{C}(\text{R}_1)=\text{N}-$, $-\text{C}(=\text{NR}_1)-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{O}-$, $-\text{Si}(\text{R}_1)_2-$, $-\text{S}(=\text{O})_x-$ and $-\text{N}(\text{R}_1)-$; wherein: x is 0, 1, or 2; n is 1, 2, 3, or 4; each R_1 and R_2 is, independently, H, a protecting group, hydroxyl, C_1 - C_{12} alkyl, substituted C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, substituted C_2 - C_{12} alkenyl, C_2 - C_{12} alkynyl, substituted C_2 - C_{12} alkynyl, C_5 - C_{20} aryl, substituted C_5 - C_{20} aryl, a heterocycle radical, a substituted heterocycle radical, heteroaryl, substituted heteroaryl, C_5 - C_7 alicyclic radical, substituted C_5 - C_7 alicyclic radical, halogen, OJ_1 , NJ_1J_2 , SJ_1 , N_3 , COOJ_1 , acyl ($\text{C}(=\text{O})-\text{H}$), substituted acyl, CN, sulfonyl ($\text{S}(=\text{O})_2-\text{J}_1$), or sulfoxyl ($\text{S}(=\text{O})-\text{J}_1$); and each J_1 and J_2 is, independently, H, C_1 - C_{12} alkyl, substituted C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, substituted C_2 - C_{12} alkenyl, C_2 - C_{12} alkynyl, substituted C_2 - C_{12} alkynyl, C_5 - C_{20} aryl, substituted C_5 - C_{20} aryl, acyl ($\text{C}(=\text{O})-\text{H}$), substituted acyl, a heterocycle radical, a substituted heterocycle radical, C_1 - C_{12} aminoalkyl, substituted C_1 - C_{12} aminoalkyl or a protecting group.

Examples of 4'-2' bridging groups encompassed within the definition of LNA include, but are not limited to one of formulae: $-\text{C}(\text{R}_1)(\text{R}_2)_n-$, $-\text{C}(\text{R}_1)(\text{R}_2)_n-\text{O}-$, $-\text{C}(\text{R}_1\text{R}_2)-\text{N}(\text{R}_1)-\text{O}-$ or $-\text{C}(\text{R}_1\text{R}_2)-\text{O}-\text{N}(\text{R}_1)-$. Furthermore, other bridging groups encompassed within the definition of LNA are 4'- CH_2 -2', 4'-(CH_2)₂-2', 4'-(CH_2)₃-2', 4'- CH_2 -O-2', 4'-(CH_2)₂-O-2', 4'- CH_2 -O- $\text{N}(\text{R}_1)$ -2' and 4'- CH_2 - $\text{N}(\text{R}_1)$ -O-2'-bridges, wherein each R_1 and R_2 is, independently, H, a protecting group or C_1 - C_{12} alkyl.

Also included within the definition of LNA according to the invention are LNAs in which the 2'-hydroxyl group of the ribosyl sugar ring is connected to the 4' carbon atom of the sugar ring, thereby forming a methyleneoxy (4'- CH_2 -O-2') bridge to form the bicyclic sugar moiety. The bridge can also be a methylene ($-\text{CH}_2-$) group connecting the 2' oxygen atom and the 4' carbon atom, for which the term methyleneoxy (4'- CH_2 -O-2') LNA is used. Furthermore; in the case of the bicyclic sugar moiety having an ethylene bridging group in this position, the term ethyleneoxy (4'- CH_2CH_2 -O-2') LNA is used. α -L-methyleneoxy (4'- CH_2 -O-2'), an isomer of methyleneoxy (4'- CH_2 -O-2') LNA is also encompassed within the definition of LNA, as used herein.

10

"Mismatch" or "non-complementary nucleobase" refers to the case when a nucleobase of a first nucleic acid is not capable of pairing with the corresponding nucleobase of a second or target nucleic acid.

5 "Modified internucleoside linkage" refers to a substitution or any change from a naturally occurring internucleoside bond (i.e., a phosphodiester internucleoside bond).

"Modified nucleobase" means any nucleobase other than adenine, cytosine, guanine, thymidine, or uracil. An "unmodified nucleobase" means the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C), and uracil (U).

10 "Modified nucleoside" means a nucleoside having, independently, a modified sugar moiety and/or modified nucleobase.

15 "Modified nucleotide" means a nucleotide having, independently, a modified sugar moiety, modified internucleoside linkage, and/or modified nucleobase.

20 "Modified oligonucleotide" means an oligonucleotide comprising at least one modified internucleoside linkage, modified sugar, and/or modified nucleobase.

"Modified sugar" means substitution and/or any change from a natural sugar moiety.

"Monomer" means a single unit of an oligomer. Monomers include, but are not limited to, nucleosides and nucleotides, whether naturally occurring or modified.

"Motif" means the pattern of unmodified and modified nucleoside in an antisense compound.

30 "Natural sugar moiety" means a sugar moiety found in DNA (2'-H) or RNA (2'-OH).

"Naturally occurring internucleoside linkage" means a 3' to 5' phosphodiester linkage.

35 "Non-complementary nucleobase" refers to a pair of nucleobases that do not form hydrogen bonds with one another or otherwise support hybridization.

"Nucleic acid" refers to molecules composed of monomeric nucleotides. A nucleic acid includes, but is not limited to, ribonucleic acids (RNA), deoxyribonucleic acids (DNA), single-stranded nucleic acids, double-stranded nucleic acids, small interfering ribonucleic acids (siRNA), and microRNAs (miRNA).

"Nucleobase" means a heterocyclic moiety capable of pairing with a base of another nucleic acid.

45 "Nucleobase complementarity" refers to a nucleobase that is capable of base pairing with another nucleobase. For example, in DNA, adenine (A) is complementary to thymine (T). For example, in RNA, adenine (A) is complementary to uracil (U). In certain embodiments, complementary nucleobase refers to a nucleobase of an antisense compound that is capable of base pairing with a nucleobase of its target nucleic acid. For example, if a nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be complementary at that nucleobase pair.

"Nucleobase sequence" means the order of contiguous nucleobases independent of any sugar, linkage, and/or nucleobase modification.

60 "Nucleoside" means a nucleobase linked to a sugar.

65 "Nucleoside mimetic" includes those structures used to replace the sugar or the sugar and the base and not necessarily the linkage at one or more positions of an oligomeric compound such as for example nucleoside mimetics having morpholino, cyclohexenyl, cyclohexyl, tetrahydropyranyl, bicyclo, or tricyclo sugar mimetics, e.g., non furanose sugar units. Nucleotide mimetic includes those structures used to

11

replace the nucleoside and the linkage at one or more positions of an oligomeric compound such as for example peptide nucleic acids or morpholinos (morpholinos linked by —N(H)—C(=O)—O— or other non-phosphodiester linkage). Sugar surrogate overlaps with the slightly broader term nucleoside mimetic but is intended to indicate replacement of the sugar unit (furanose ring) only. The tetrahydropyran rings provided herein are illustrative of an example of a sugar surrogate wherein the furanose sugar group has been replaced with a tetrahydropyran ring system. “Mimetic” refers to groups that are substituted for a sugar, a nucleobase, and/or internucleoside linkage. Generally, a mimetic is used in place of the sugar or sugar-internucleoside linkage combination, and the nucleobase is maintained for hybridization to a selected target.

“Nucleotide” means a nucleoside having a phosphate group covalently linked to the sugar portion of the nucleoside.

“Off-target effect” refers to an unwanted or deleterious biological effect associated with modulation of RNA or protein expression of a gene other than the intended target nucleic acid.

“Oligomeric compound” or “oligomer” means a polymer of linked monomeric subunits which is capable of hybridizing to at least a region of a nucleic acid molecule.

“Oligonucleotide” means a polymer of linked nucleosides each of which can be modified or unmodified, independent one from another.

“Parenteral administration” means administration through injection (e.g., bolus injection) or infusion. Parenteral administration includes subcutaneous administration, intravenous administration, intramuscular administration, intraarterial administration, intraperitoneal administration, or intracranial administration, e.g., intrathecal or intracerebroventricular administration.

“Peptide” means a molecule formed by linking at least two amino acids by amide bonds. Without limitation, as used herein, peptide refers to polypeptides and proteins.

“Pharmaceutical agent” means a substance that provides a therapeutic benefit when administered to an individual. For example, in certain embodiments, an antisense oligonucleotide targeted to C9ORF72 is a pharmaceutical agent.

“Pharmaceutically acceptable derivative” encompasses pharmaceutically acceptable salts, conjugates, prodrugs or isomers of the compounds described herein.

“Pharmaceutically acceptable salts” means physiologically and pharmaceutically acceptable salts of antisense compounds, i.e., salts that retain the desired biological activity of the parent oligonucleotide and do not impart undesired toxicological effects thereto.

“Phosphorothioate linkage” means a linkage between nucleosides where the phosphodiester bond is modified by replacing one of the non-bridging oxygen atoms with a sulfur atom. A phosphorothioate linkage is a modified internucleoside linkage.

“Portion” means a defined number of contiguous (i.e., linked) nucleobases of a nucleic acid. In certain embodiments, a portion is a defined number of contiguous nucleobases of a target nucleic acid. In certain embodiments, a portion is a defined number of contiguous nucleobases of an antisense compound.

“Prevent” or “preventing” refers to delaying or forestalling the onset or development of a disease, disorder, or condition for a period of time from minutes to days, weeks to months, or indefinitely.

“Prodrug” means a therapeutic agent that is prepared in an inactive form that is converted to an active form within the

12

body or cells thereof by the action of endogenous enzymes or other chemicals or conditions.

“Prophylactically effective amount” refers to an amount of a pharmaceutical agent that provides a prophylactic or preventative benefit to an animal.

“Region” is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or characteristic.

“Ribonucleotide” means a nucleotide having a hydroxy at the 2' position of the sugar portion of the nucleotide. Ribonucleotides may be modified with any of a variety of substituents.

“Salts” mean a physiologically and pharmaceutically acceptable salts of antisense compounds, i.e., salts that retain the desired biological activity of the parent oligonucleotide and do not impart undesired toxicological effects thereto.

“Segments” are defined as smaller or sub-portions of regions within a target nucleic acid.

“Shortened” or “truncated” versions of antisense oligonucleotides taught herein have one, two or more nucleosides deleted.

“Side effects” means physiological responses attributable to a treatment other than desired effects. In certain embodiments, side effects include, without limitation, injection site reactions, liver function test abnormalities, renal function abnormalities, liver toxicity, renal toxicity, central nervous system abnormalities, and myopathies.

“Single-stranded oligonucleotide” means an oligonucleotide which is not hybridized to a complementary strand. A “single-stranded modified oligonucleotide” means a modified oligonucleotide which is not hybridized to a complementary strand.

“Sites,” as used herein, are defined as unique nucleobase positions within a target nucleic acid.

“Slows progression” means decrease in the development of the disease.

“Specifically hybridizable” refers to an antisense compound having a sufficient degree of complementarity between an antisense oligonucleotide and a target nucleic acid to induce a desired effect, while exhibiting minimal or no effects on non-target nucleic acids under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays and therapeutic treatments.

“Stringent hybridization conditions” or “stringent conditions” refer to conditions under which an oligomeric compound will hybridize to its target sequence, but to a minimal number of other sequences.

“Subject” means a human or non-human animal selected for treatment or therapy.

“Target” refers to a protein, the modulation of which is desired.

“Target gene” refers to a gene encoding a target.

“Targeting” or “targeted” means the process of design and selection of an antisense compound that will specifically hybridize to a target nucleic acid and induce a desired effect.

“Target nucleic acid,” “target RNA,” and “target RNA transcript” and “nucleic acid target” all mean a nucleic acid capable of being targeted by antisense compounds.

“Target region” means a portion of a target nucleic acid to which one or more antisense compounds is targeted.

“Target segment” means the sequence of nucleotides of a target nucleic acid to which an antisense compound is targeted. “5' target site” refers to the 5'-most nucleotide of a target segment. “3' target site” refers to the 3'-most nucleotide of a target segment.

13

“Therapeutically effective amount” means an amount of a pharmaceutical agent that provides a therapeutic benefit to an individual.

“Treat” or “treating” or “treatment” means administering a composition to effect an alteration or improvement of a disease or condition.

“Unmodified nucleobases” means the purine bases adenine (A) and guanine (G), and the pyrimidine bases (T), cytosine (C), and uracil (U).

“Unmodified nucleotide” means a nucleotide composed of naturally occurring nucleobases, sugar moieties, and internucleoside linkages. In certain embodiments, an unmodified nucleotide is an RNA nucleotide (i.e. β -D-ribonucleosides) or a DNA nucleotide (i.e. β -D-deoxyribonucleoside).

“Wing segment” means a plurality of nucleosides modified to impart to an oligonucleotide properties such as enhanced inhibitory activity, increased binding affinity for a target nucleic acid, or resistance to degradation by in vivo nucleases.

Certain Embodiments

Certain embodiments provide compositions and methods for reducing total C9ORF72 mRNA and protein expression.

Certain embodiments provide compositions and methods for reducing C9ORF72 pathogenic associated mRNA variants.

Certain embodiments provide methods for the treatment, prevention, amelioration, or slowing progression of diseases associated with C9ORF72 in an individual in need thereof. Also contemplated are methods for the preparation of a medicament for the treatment, prevention, or amelioration of a disease associated with C9ORF72. C9ORF72 associated diseases include neurodegenerative diseases. In certain embodiments, the neurodegenerative disease may be ALS or FTD. In certain embodiments, the neurodegenerative disease may be familial or sporadic.

The present disclosure provided the following non-limiting numbered embodiments:

Embodiment 1

A compound comprising a modified oligonucleotide consisting of 12 to 30 linked nucleosides and having a nucleobase sequence comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 consecutive nucleobases of any of the nucleobase sequences of SEQ ID NOs: 22-55.

Embodiment 2

The compound of embodiment 1, wherein the nucleobase sequence of the modified oligonucleotide is at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% complementary to SEQ ID NO: 1 or SEQ ID NO: 2.

Embodiment 3

The compound of embodiments 1 and 2, wherein the modified oligonucleotide is a single-stranded modified oligonucleotide.

14

Embodiment 4

The compound of any of embodiments 1-3, wherein the modified oligonucleotide comprises at least one modified internucleoside linkage.

Embodiment 5

The compound of embodiment 4, wherein the at least one modified internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 6

The compound of embodiments 4 and 5, wherein the modified oligonucleotide comprises at least one phosphodiester linkage.

Embodiment 7

The compound of embodiment 5, wherein each modified internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 8

The compound of any of embodiments 1-7, wherein at least one nucleoside comprises a modified nucleobase.

Embodiment 9

The compound of embodiment 8, wherein the modified nucleobase is a 5-methylcytosine.

Embodiment 10

The compound of any of embodiments 1-9, wherein at least one nucleoside of the modified oligonucleotide comprises a modified sugar.

Embodiment 11

The compound of embodiment 10, wherein each nucleoside of the modified oligonucleotide comprises a modified sugar.

Embodiment 12

The compound of embodiments 10 or 11, wherein the at least one modified sugar is a bicyclic sugar.

Embodiment 13

The compound of embodiment 12, wherein the bicyclic sugar comprises a chemical bridge between the 4' and 2' positions of the sugar, wherein the chemical bridge is selected from: 4'-CH(R)—O-2' and 4'-(CH₂)₂—O-2', wherein R is independently H, C₁-C₆ alkyl, and C₁-C₆ alkoxy.

Embodiment 14

The compound of embodiment 13, wherein the chemical bridge is 4'-CH(R)—O-2' and wherein R is methyl.

15

Embodiment 15

The compound of embodiment 13, wherein the chemical bridge is 4'-CH(R)—O-2' and wherein R is H.

Embodiment 16

The compound of embodiment 13, wherein the chemical bridge is 4'-CH(R)—O-2' and wherein R is —CH₂—O—CH₃.

Embodiment 17

The compound of embodiments 10 or 11, wherein the at least one modified sugar comprises a 2'-O-methoxyethyl group.

Embodiment 18

The compound of any of embodiments 1-10 or 12-16, wherein the modified oligonucleotide is a gapmer.

Embodiment 19

The compound of embodiment 18, wherein the gapmer is any of a 3-8-7 MOE gapmer, a 3-10-7 MOE gapmer, a 4-8-6 MOE gapmer, a 4-10-6 MOE gapmer, a 6-10-4 MOE gapmer, a 6-8-4 MOE gapmer, a 7-8-3 MOE gapmer, or a 7-10-3 MOE gapmer.

Embodiment 20

The compound of claim 5, wherein the modified oligonucleotide comprises internucleoside linkages in any of the following patterns: soossssssssssooooss, soossssssssssooooss, soooooosssssssssss, soooooosssssssssss, soooooossssssssssooss, soossssssssssooooss, soooooSssssssssoss, soossssssssssooooss, soooooosssssssssoss, soossssssssooooss, or soooooosssssssssss, wherein, s=a phosphorothioate linkage, and o=a phosphodiester linkage.

Embodiment 19

A composition comprising the compound of any preceding embodiment or salt thereof and at least one of a pharmaceutically acceptable carrier or diluent.

Embodiment 20

The composition of embodiment 19, further comprising a C9ORF72 antisense transcript specific inhibitor.

Embodiment 21

The composition of embodiment 20, wherein the C9ORF72 antisense transcript specific inhibitor is an antisense compound.

Embodiment 22

The composition of embodiment 21, wherein the antisense compound is a modified oligonucleotide.

Embodiment 23

The composition of embodiment 22, wherein the modified oligonucleotide is single-stranded.

16

Embodiment 24

The composition of embodiments 22 or 23, wherein the modified oligonucleotide has a nucleobase sequence that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% complementary to a C9ORF72 antisense transcript.

Embodiment 25

The composition of embodiment 24, wherein the C9ORF72 antisense transcript has the nucleobase sequence of SEQ ID NO: 18.

Embodiment 26

A method comprising administering to an animal the compound or composition of any preceding embodiment.

Embodiment 27

The method of embodiment 26, wherein the animal is a human.

Embodiment 28

The method of embodiments 26 and 27, wherein administering the compound prevents, treats, ameliorates, or slows progression of a C9ORF72 associated disease.

Embodiment 29

The method of embodiment 28, wherein the C9ORF72 associated disease is caused by a hexanucleotide repeat expansion.

Embodiment 30

The method of embodiment 28, wherein the C9ORF72 associated disease is amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, and olivopontocerebellar degeneration (OPCD).

Embodiment 31

The method of embodiments 26-30, wherein the administering reduces nuclear foci.

Embodiment 32

The method of embodiments 26-31, wherein the administering reduces expression of C9ORF72 associated RAN translation products.


Embodiment 33

The method of embodiment 32, wherein the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).

Embodiment 34


Use of the compound or composition of any of embodiments 1-33 for the manufacture of a medicament for treating a neurodegenerative disorder.

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:



A composition consisting of the sodium salt of a modified 5' oligonucleotide according to the following formula:

(SEQ ID NO: 33)



21


Embodiment 37

22

5

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:

(SEQ ID NO: 49)



23


Embodiment 38

24

5

A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:

(SEQ ID NO: 49)



25


Embodiment 39

26

5

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:

(SEQ ID NO: 47)



27


Embodiment 40

28

5

A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:

(SEQ ID NO: 47)



29


Embodiment 41

30

5


A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:

(SEQ ID NO: 21)



31**Embodiment 42**

A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:

**Embodiment 43**

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of reducing human C9ORF72 mRNA or protein expression in a mammal.


Embodiment 44

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of ameliorating at least one symptom of amy-

32

trophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, or olivopontocerebellar degeneration (OPCD).

(SEQ ID NO: 21)

**Embodiment 45**

The compound or composition of embodiment 44, wherein the symptom of ALS is any of motor deficit, anxiety, and denervation.

Embodiment 46

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of delaying progression of disease.

33

Embodiment 47

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of extending survival.

Embodiment 48

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of reducing C9ORF72 associated RAN translation products.

Embodiment 49

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is

34

capable of selectively reducing C9ORF72 pathogenic associated mRNA variants.


Embodiment 50

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of reducing nuclear foci.

Embodiment 51

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:

(SEQ ID NO: 33)



35

Embodiment 52

A method comprising administering to an animal the compound of embodiment 51.

Embodiment 53

The method of embodiment 52, wherein the animal is a human.

Embodiment 54

The method of embodiment 53, wherein the administering inhibits C9ORF72.

Embodiment 55

The method of embodiment 53, wherein the administering ²⁵ prevents, treats, ameliorates, or slows progression of a C9ORF72 associated disease.

Embodiment 56

The method of embodiment 55, wherein the C9ORF72 associated disease is caused by a hexanucleotide repeat expansion.

36

Embodiment 57

The method of embodiment 55, wherein the C9ORF72
5 associated disease is any of amyotrophic lateral sclerosis
(ALS), frontotemporal dementia (FTD), corticobasal degen-
eration syndrome (CBD), atypical Parkinsonian syndrome,
or olivopontocerebellar degeneration (OPCD).

Embodiment 58

The method of embodiment 53, wherein the administering
15 reduces nuclear foci.

Embodiment 59

20 The method of embodiment 53, wherein the administering
reduces expression of C9ORF72 associated RAN translation
products.




Embodiment 60

The method of embodiment 59, wherein the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).


Embodiment 61

A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:

(SEQ ID NO: 33)



37



Embodiment 64

The method of embodiment 63, wherein the administering inhibits C9ORF72.

Embodiment 65

The method of embodiment 62, wherein the administering prevents, treats, ameliorates, or slows progression of a C9ORF72 associated disease.

Embodiment 66

The method of embodiment 65, wherein the C9ORF72 associated disease is caused by a hexanucleotide repeat expansion.

Embodiment 67

The method of embodiment 65, wherein the C9ORF72 associated disease is any of amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, or olivopontocerebellar degeneration (OPCD).

Embodiment 68

The method of embodiment 63, wherein the administering reduces nuclear foci.


Embodiment 69

The method of embodiment 63, wherein the administering reduces expression of C9ORF72 associated RAN translation products.

Embodiment 70

The method of embodiment 69, wherein the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).

38



25

Antisense Compounds

Oligomeric compounds include, but are not limited to, oligonucleotides, oligonucleosides, oligonucleotide analogs, oligonucleotide mimetics, antisense compounds, antisense oligonucleotides, and siRNAs. An oligomeric compound may be "antisense" to a target nucleic acid, meaning that is capable of undergoing hybridization to a target nucleic acid through hydrogen bonding.

30

35

In certain embodiments, an antisense compound has a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of the target segment of a target nucleic acid to which it is targeted. In certain such embodiments, an antisense oligonucleotide has a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of the target segment of a target nucleic acid to which it is targeted.

40

45

In certain embodiments, an antisense compound targeted to a C9ORF72 nucleic acid is 12 to 30 subunits in length. In other words, such antisense compounds are from 12 to 30 linked subunits. In certain embodiments, the antisense compound is 8 to 80, 12 to 50, 15 to 30, 18 to 24, 19 to 22, or 20 linked subunits. In certain embodiments, the antisense compounds are 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 linked subunits in length, or a range defined by any two of the above values. In some embodiments the antisense compound is an antisense oligonucleotide, and the linked subunits are nucleosides.

50

55

60

In certain embodiments antisense oligonucleotides targeted to a C9ORF72 nucleic acid may be shortened or truncated. For example, a single subunit may be deleted from the 5' end (5' truncation), or alternatively from the 3' end (3' truncation). A shortened or truncated antisense compound targeted to a C9ORF72 nucleic acid may have two subunits deleted from the 5' end, or alternatively may have two subunits deleted from the 3' end, of the antisense compound. Alternatively, the deleted nucleosides may be dispersed throughout the antisense compound, for example,

65

in an antisense compound having one nucleoside deleted from the 5' end and one nucleoside deleted from the 3' end.

When a single additional subunit is present in a lengthened antisense compound, the additional subunit may be located at the 5' or 3' end of the antisense compound. When two or more additional subunits are present, the added subunits may be adjacent to each other, for example, in an antisense compound having two subunits added to the 5' end (5' addition), or alternatively to the 3' end (3' addition), of the antisense compound. Alternatively, the added subunits may be dispersed throughout the antisense compound, for example, in an antisense compound having one subunit added to the 5' end and one subunit added to the 3' end.

It is possible to increase or decrease the length of an antisense compound, such as an antisense oligonucleotide, and/or introduce mismatch bases without eliminating activity. For example, in Woolf et al. (Proc. Natl. Acad. Sci. USA 89:7305-7309, 1992), a series of antisense oligonucleotides 13-25 nucleobases in length were tested for their ability to induce cleavage of a target RNA in an oocyte injection model. Antisense oligonucleotides 25 nucleobases in length with 8 or 11 mismatch bases near the ends of the antisense oligonucleotides were able to direct specific cleavage of the target mRNA, albeit to a lesser extent than the antisense oligonucleotides that contained no mismatches. Similarly, target specific cleavage was achieved using 13 nucleobase antisense oligonucleotides, including those with 1 or 3 mismatches.

Gautschi et al (J. Natl. Cancer Inst. 93:463-471, March 2001) demonstrated the ability of an oligonucleotide having 100% complementarity to the bcl-2 mRNA and having 3 mismatches to the bcl-xL mRNA to reduce the expression of both bcl-2 and bcl-xL in vitro and in vivo. Furthermore, this oligonucleotide demonstrated potent anti-tumor activity in vivo.

Maher and Dolnick (Nuc. Acid. Res. 16:3341-3358, 1988) tested a series of tandem 14 nucleobase antisense oligonucleotides, and a 28 and 42 nucleobase antisense oligonucleotides comprised of the sequence of two or three of the tandem antisense oligonucleotides, respectively, for their ability to arrest translation of human DHFR in a rabbit reticulocyte assay. Each of the three 14 nucleobase antisense oligonucleotides alone was able to inhibit translation, albeit at a more modest level than the 28 or 42 nucleobase antisense oligonucleotides.

Antisense Compound Motifs

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid have chemically modified subunits arranged in patterns, or motifs, to confer to the antisense compounds properties such as enhanced inhibitory activity, increased binding affinity for a target nucleic acid, or resistance to degradation by in vivo nucleases.

Chimeric antisense compounds typically contain at least one region modified so as to confer increased resistance to nuclease degradation, increased cellular uptake, increased binding affinity for the target nucleic acid, and/or increased inhibitory activity. A second region of a chimeric antisense compound may optionally serve as a substrate for the cellular endonuclease RNase H, which cleaves the RNA strand of an RNA:DNA duplex.

Antisense compounds having a gapmer motif are considered chimeric antisense compounds. In a gapmer an internal region having a plurality of nucleotides that supports RNaseH cleavage is positioned between external regions having a plurality of nucleotides that are chemically distinct from the nucleosides of the internal region. In the case of an antisense oligonucleotide having a gapmer motif, the gap

segment generally serves as the substrate for endonuclease cleavage, while the wing segments comprise modified nucleosides. In certain embodiments, the regions of a gapmer are differentiated by the types of sugar moieties comprising each distinct region. The types of sugar moieties that are used to differentiate the regions of a gapmer may in some embodiments include β -D-ribonucleosides, β -D-deoxyribonucleosides, 2'-modified nucleosides (such 2'-modified nucleosides may include 2'-MOE, and 2'-O—CH₃, among others), and bicyclic sugar modified nucleosides (such bicyclic sugar modified nucleosides may include those having a 4'-(CH₂)_n-O-2' bridge, where n=1 or n=2 and 4'-CH₂—O—CH₂-2'). Preferably, each distinct region comprises uniform sugar moieties. The wing-gap-wing motif is frequently described as "X—Y—Z", where "X" represents the length of the 5' wing region, "Y" represents the length of the gap region, and "Z" represents the length of the 3' wing region. As used herein, a gapmer described as "X—Y—Z" has a configuration such that the gap segment is positioned immediately adjacent to each of the 5' wing segment and the 3' wing segment. Thus, no intervening nucleotides exist between the 5' wing segment and gap segment, or the gap segment and the 3' wing segment. Any of the antisense compounds described herein can have a gapmer motif. In some embodiments, X and Z are the same, in other embodiments they are different. In a preferred embodiment, Y is between 8 and 15 nucleotides. X, Y or Z can be any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30 or more nucleotides. Thus, gapmers described herein include, but are not limited to, for example 5-10-5, 5-10-4, 4-10-4, 4-10-3, 3-10-3, 2-10-2, 5-9-5, 5-9-4, 4-9-5, 5-8-5, 5-8-4, 4-8-5, 5-7-5, 4-7-5, 5-7-4, or 4-7-4.

In certain embodiments, the antisense compound has a "wingmer" motif, having a wing-gap or gap-wing configuration, i.e. an X—Y or Y—Z configuration as described above for the gapmer configuration. Thus, wingmer configurations described herein include, but are not limited to, for example 5-10, 8-4, 4-12, 12-4, 3-14, 16-2, 18-1, 10-3, 2-10, 1-10, 8-2, 2-13, 5-13, 5-8, or 6-8.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess a 5-10-5 gapmer motif.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess a 5-8-5 gapmer motif.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess sugar modifications in any of the following patterns: eekkkdddddckkeee, eek-kdddddckkeee, ekddddddekekeee, kekeddddddde-keke, and ekekdddddckkeee; wherein,

e=a 2'-O-methoxyethyl modified nucleoside

d=a 2'-deoxynucleoside, and

k=a cEt nucleoside.

In certain embodiments, an antisense compound targeted to a C9ORF72 nucleic acid has a gap-narrowed motif. In certain embodiments, a gap-narrowed antisense oligonucleotide targeted to a C9ORF72 nucleic acid has a gap segment of 9, 8, 7, or 6 2'-deoxynucleotides positioned immediately adjacent to and between wing segments of 5, 4, 3, 2, or 1 chemically modified nucleosides. In certain embodiments, the chemical modification comprises a bicyclic sugar. In certain embodiments, the bicyclic sugar comprises a 4' to 2' bridge selected from among: 4'-(CH₂)_n—O-2' bridge, wherein n is 1 or 2; and 4'-CH₂—O—CH₂-2'. In certain embodiments, the bicyclic sugar is comprises a 4'-CH(CH₃)—O-2' bridge. In certain embodiments, the chemical modification comprises a non-bicyclic 2'-modified sugar

moiety. In certain embodiments, the non-bicyclic 2'-modified sugar moiety comprises a 2'-O-methylethyl group or a 2'-O-methyl group.

Target Nucleic Acids, Target Regions and Nucleotide Sequences

Nucleotide sequences that encode C9ORF72 include, without limitation, the following: the complement of GENBANK Accession No. NM_001256054.1 (incorporated herein as SEQ ID NO: 1), the complement of GENBANK Accession No. NT_008413.18 truncated from nucleobase 27535000 to 27565000 (incorporated herein as SEQ ID NO: 2), GENBANK Accession No. BQ068108.1 (incorporated herein as SEQ ID NO: 3), GENBANK Accession No. NM_018325.3 (incorporated herein as SEQ ID NO: 4), GENBANK Accession No. DN993522.1 (incorporated herein as SEQ ID NO: 5), GENBANK Accession No. NM_145005.5 (incorporated herein as SEQ ID NO: 6), GENBANK Accession No. DB079375.1 (incorporated herein as SEQ ID NO: 7), GENBANK Accession No. BU194591.1 (incorporated herein as SEQ ID NO: 8), Sequence Identifier 4141_014_A (incorporated herein as SEQ ID NO: 9), and Sequence Identifier 4008_73_A (incorporated herein as SEQ ID NO: 10), and GENBANK Accession No. NW_001101662.1 truncated from nucleosides 8522000 to U.S. Pat. No. 8,552,000 (incorporated herein as SEQ ID NO: 11).

It is understood that the sequence set forth in each SEQ ID NO in the Examples contained herein is independent of any modification to a sugar moiety, an internucleoside linkage, or a nucleobase. As such, antisense compounds defined by a SEQ ID NO may comprise, independently, one or more modifications to a sugar moiety, an internucleoside linkage, or a nucleobase. Antisense compounds described by Isis Number (Isis No) indicate a combination of nucleobase sequence and motif.

In certain embodiments, a target region is a structurally defined region of the target nucleic acid. For example, a target region may encompass a 3' UTR, a 5' UTR, an exon, an intron, an exon/intron junction, a coding region, a translation initiation region, translation termination region, or other defined nucleic acid region. The structurally defined regions for C9ORF72 can be obtained by accession number from sequence databases such as NCBI and such information is incorporated herein by reference. In certain embodiments, a target region may encompass the sequence from a 5' target site of one target segment within the target region to a 3' target site of another target segment within the same target region.

Targeting includes determination of at least one target segment to which an antisense compound hybridizes, such that a desired effect occurs. In certain embodiments, the desired effect is a reduction in mRNA target nucleic acid levels. In certain embodiments, the desired effect is reduction of levels of protein encoded by the target nucleic acid or a phenotypic change associated with the target nucleic acid.

A target region may contain one or more target segments. Multiple target segments within a target region may be overlapping. Alternatively, they may be non-overlapping. In certain embodiments, target segments within a target region are separated by no more than about 300 nucleotides. In certain embodiments, target segments within a target region are separated by a number of nucleotides that is, is about, is no more than, is no more than about, 250, 200, 150, 100, 90, 80, 70, 60, 50, 40, 30, 20, or 10 nucleotides on the target nucleic acid, or is a range defined by any two of the preceding values. In certain embodiments, target segments

within a target region are separated by no more than, or no more than about, 5 nucleotides on the target nucleic acid. In certain embodiments, target segments are contiguous. Contemplated are target regions defined by a range having a starting nucleic acid that is any of the 5' target sites or 3' target sites listed herein.

Suitable target segments may be found within a 5' UTR, a coding region, a 3' UTR, an intron, an exon, or an exon/intron junction. Target segments containing a start codon or a stop codon are also suitable target segments. A suitable target segment may specifically exclude a certain structurally defined region such as the start codon or stop codon.

The determination of suitable target segments may include a comparison of the sequence of a target nucleic acid to other sequences throughout the genome. For example, the BLAST algorithm may be used to identify regions of similarity amongst different nucleic acids. This comparison can prevent the selection of antisense compound sequences that may hybridize in a non-specific manner to sequences other than a selected target nucleic acid (i.e., non-target or off-target sequences).

There may be variation in activity (e.g., as defined by percent reduction of target nucleic acid levels) of the antisense compounds within a target region. In certain embodiments, reductions in C9ORF72 mRNA levels are indicative of inhibition of C9ORF72 expression. Reductions in levels of a C9ORF72 protein are also indicative of inhibition of target mRNA expression. Reduction in the presence of expanded C9ORF72 RNA foci are indicative of inhibition of C9ORF72 expression. Further, phenotypic changes are indicative of inhibition of C9ORF72 expression. For example, improved motor function and respiration may be indicative of inhibition of C9ORF72 expression.

Hybridization

In some embodiments, hybridization occurs between an antisense compound disclosed herein and a C9ORF72 nucleic acid. The most common mechanism of hybridization involves hydrogen bonding (e.g., Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding) between complementary nucleobases of the nucleic acid molecules.

Hybridization can occur under varying conditions. Stringent conditions are sequence-dependent and are determined by the nature and composition of the nucleic acid molecules to be hybridized.

Methods of determining whether a sequence is specifically hybridizable to a target nucleic acid are well known in the art. In certain embodiments, the antisense compounds provided herein are specifically hybridizable with a C9ORF72 nucleic acid.

Complementarity

An antisense compound and a target nucleic acid are complementary to each other when a sufficient number of nucleobases of the antisense compound can hydrogen bond with the corresponding nucleobases of the target nucleic acid, such that a desired effect will occur (e.g., antisense inhibition of a target nucleic acid, such as a C9ORF72 nucleic acid).

Non-complementary nucleobases between an antisense compound and a C9ORF72 nucleic acid may be tolerated provided that the antisense compound remains able to specifically hybridize to a target nucleic acid. Moreover, an antisense compound may hybridize over one or more segments of a C9ORF72 nucleic acid such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure, mismatch or hairpin structure).

In certain embodiments, the antisense compounds provided herein, or a specified portion thereof, are, or are at least, 70%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% complementary to a C9ORF72 nucleic acid, a target region, target segment, or specified portion thereof. Percent complementarity of an antisense compound with a target nucleic acid can be determined using routine methods.

For example, an antisense compound in which 18 of 20 nucleobases of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleobases may be clustered or interspersed with complementary nucleobases and need not be contiguous to each other or to complementary nucleobases. As such, an antisense compound which is 18 nucleobases in length having 4 (four) noncomplementary nucleobases which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid and would thus fall within the scope of the present invention. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., J. Mol. Biol., 1990, 215, 403 410; Zhang and Madden, Genome Res., 1997, 7, 649 656). Percent homology, sequence identity or complementarity, can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., 1981, 2, 482 489).

In certain embodiments, the antisense compounds provided herein, or specified portions thereof, are fully complementary (i.e., 100% complementary) to a target nucleic acid, or specified portion thereof. For example, an antisense compound may be fully complementary to a C9ORF72 nucleic acid, or a target region, or a target segment or target sequence thereof. As used herein, "fully complementary" means each nucleobase of an antisense compound is capable of precise base pairing with the corresponding nucleobases of a target nucleic acid. For example, a 20 nucleobase antisense compound is fully complementary to a target sequence that is 400 nucleobases long, so long as there is a corresponding 20 nucleobase portion of the target nucleic acid that is fully complementary to the antisense compound. Fully complementary can also be used in reference to a specified portion of the first and/or the second nucleic acid. For example, a 20 nucleobase portion of a 30 nucleobase antisense compound can be "fully complementary" to a target sequence that is 400 nucleobases long. The 20 nucleobase portion of the 30 nucleobase oligonucleotide is fully complementary to the target sequence if the target sequence has a corresponding 20 nucleobase portion wherein each nucleobase is complementary to the 20 nucleobase portion of the antisense compound. At the same time, the entire 30 nucleobase antisense compound may or may not be fully complementary to the target sequence, depending on whether the remaining 10 nucleobases of the antisense compound are also complementary to the target sequence.

The location of a non-complementary nucleobase may be at the 5' end or 3' end of the antisense compound. Alternatively, the non-complementary nucleobase or nucleobases may be at an internal position of the antisense compound. When two or more non-complementary nucleobases are present, they may be contiguous (i.e., linked) or non-

contiguous. In one embodiment, a non-complementary nucleobase is located in the wing segment of a gapmer antisense oligonucleotide.

In certain embodiments, antisense compounds that are, or are up to 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleobases in length comprise no more than 4, no more than 3, no more than 2, or no more than 1 non-complementary nucleobase(s) relative to a target nucleic acid, such as a C9ORF72 nucleic acid, or specified portion thereof.

In certain embodiments, antisense compounds that are, or are up to 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleobases in length comprise no more than 6, no more than 5, no more than 4, no more than 3, no more than 2, or no more than 1 non-complementary nucleobase(s) relative to a target nucleic acid, such as a C9ORF72 nucleic acid, or specified portion thereof.

The antisense compounds provided herein also include those which are complementary to a portion of a target nucleic acid. As used herein, "portion" refers to a defined number of contiguous (i.e. linked) nucleobases within a region or segment of a target nucleic acid. A "portion" can also refer to a defined number of contiguous nucleobases of an antisense compound. In certain embodiments, the antisense compounds, are complementary to at least an 8 nucleobase portion of a target segment. In certain embodiments, the antisense compounds are complementary to at least a 9 nucleobase portion of a target segment. In certain embodiments, the antisense compounds are complementary to at least a 10 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least an 11 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 12 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 13 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 14 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 15 nucleobase portion of a target segment. Also contemplated are antisense compounds that are complementary to at least a 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more nucleobase portion of a target segment, or a range defined by any two of these values.

Identity

The antisense compounds provided herein may also have a defined percent identity to a particular nucleotide sequence, SEQ ID NO, or compound represented by a specific Isis number, or portion thereof. As used herein, an antisense compound is identical to the sequence disclosed herein if it has the same nucleobase pairing ability. For example, a RNA which contains uracil in place of thymidine in a disclosed DNA sequence would be considered identical to the DNA sequence since both uracil and thymidine pair with adenine. Shortened and lengthened versions of the antisense compounds described herein as well as compounds having non-identical bases relative to the antisense compounds provided herein also are contemplated. The non-identical bases may be adjacent to each other or dispersed throughout the antisense compound. Percent identity of an antisense compound is calculated according to the number of bases that have identical base pairing relative to the sequence to which it is being compared.

In certain embodiments, the antisense compounds, or portions thereof, are at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to one or

more of the antisense compounds or SEQ ID NOs, or a portion thereof, disclosed herein.

In certain embodiments, a portion of the antisense compound is compared to an equal length portion of the target nucleic acid. In certain embodiments, an 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleobase portion is compared to an equal length portion of the target nucleic acid.

In certain embodiments, a portion of the antisense oligonucleotide is compared to an equal length portion of the target nucleic acid. In certain embodiments, an 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleobase portion is compared to an equal length portion of the target nucleic acid.

Modifications

A nucleoside is a base-sugar combination. The nucleobase (also known as base) portion of the nucleoside is normally a heterocyclic base moiety. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to the 2', 3' or 5' hydroxyl moiety of the sugar. Oligonucleotides are formed through the covalent linkage of adjacent nucleosides to one another, to form a linear polymeric oligonucleotide. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside linkages of the oligonucleotide.

Modifications to antisense compounds encompass substitutions or changes to internucleoside linkages, sugar moieties, or nucleobases. Modified antisense compounds are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target, increased stability in the presence of nucleases, or increased inhibitory activity.

Chemically modified nucleosides may also be employed to increase the binding affinity of a shortened or truncated antisense oligonucleotide for its target nucleic acid. Consequently, comparable results can often be obtained with shorter antisense compounds that have such chemically modified nucleosides.

Modified Internucleoside Linkages

The naturally occurring internucleoside linkage of RNA and DNA is a 3' to 5' phosphodiester linkage. Antisense compounds having one or more modified, i.e. non-naturally occurring, internucleoside linkages are often selected over antisense compounds having naturally occurring internucleoside linkages because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for target nucleic acids, and increased stability in the presence of nucleases.

Oligonucleotides having modified internucleoside linkages include internucleoside linkages that retain a phosphorus atom as well as internucleoside linkages that do not have a phosphorus atom. Representative phosphorus containing internucleoside linkages include, but are not limited to, phosphodiester, phosphotriester, methylphosphonates, phosphoramidate, and phosphorothioates. Methods of preparation of phosphorous-containing and non-phosphorous-containing linkages are well known.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid comprise one or more modified internucleoside linkages. In certain embodiments, the modified internucleoside linkages are interspersed throughout the antisense compound. In certain embodiments, the modified internucleoside linkages are phosphorothioate linkages. In certain embodiments, each internucleoside linkage of an antisense compound is a phosphorothioate internucleoside

linkage. In certain embodiments, the antisense compounds targeted to a C9ORF72 nucleic acid comprise at least one phosphodiester linkage and at least one phosphorothioate linkage.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess internucleoside linkages in any of the following patterns: sooooooooooooo, sooooooooooooo, sooooooooooooo, and sooooooooooooo; wherein,

s=a phosphorothioate linkage, and
o=a phosphodiester linkage.

Modified Sugar Moieties

Antisense compounds of the invention can optionally contain one or more nucleosides wherein the sugar group has been modified. Such sugar modified nucleosides may impart enhanced nuclease stability, increased binding affinity, or some other beneficial biological property to the antisense compounds. In certain embodiments, nucleosides comprise chemically modified ribofuranose ring moieties. Examples of chemically modified ribofuranose rings include without limitation, addition of substituent groups (including 5' and 2' substituent groups, bridging of non-geminal ring atoms to form bicyclic nucleic acids (BNA), replacement of the ribosyl ring oxygen atom with S, N(R), or C(R₁)(R₂) (R, R₁ and R₂ are each independently H, C₁-C₁₂ alkyl or a protecting group) and combinations thereof. Examples of chemically modified sugars include 2'-F-5'-methyl substituted nucleoside (see PCT International Application WO 2008/101157 Published on Aug. 21, 2008 for other disclosed 5',2'-bis substituted nucleosides) or replacement of the ribosyl ring oxygen atom with S with further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on Jun. 16, 2005) or alternatively 5'-substitution of a BNA (see PCT International Application WO 2007/134181 Published on Nov. 22, 2007 wherein LNA is substituted with for example a 5'-methyl or a 5'-vinyl group).

Examples of nucleosides having modified sugar moieties include without limitation nucleosides comprising 5'-vinyl, 5'-methyl (R or S), 4'-S, 2'-F, 2'-OCH₃, 2'-OCH₂CH₃, 2'-OCH₂CH₂F and 2'-O(CH₂)₂₀CH₃ substituent groups. The substituent at the 2' position can also be selected from allyl, amino, azido, thio, O-allyl, O—C₁-C₁₀ alkyl, OCF₃, OCH₂F, O(CH₂)₂SCH₃, O(CH₂)₂—O—N(R_m)(R_n), O—CH₂—C(=O)—N(R_m)(R_n), and O—CH₂—C(=O)—N(R₁)—(CH₂)₂—N(R_m)(R_n), where each R₁, R_m and R_n is, independently, H or substituted or unsubstituted C₁-C₁₀ alkyl.

As used herein, "bicyclic nucleosides" refer to modified nucleosides comprising a bicyclic sugar moiety. Examples of bicyclic nucleic acids (BNAs) include without limitation nucleosides comprising a bridge between the 4' and the 2' ribosyl ring atoms. In certain embodiments, antisense compounds provided herein include one or more BNA nucleosides wherein the bridge comprises one of the formulas: 4'-(CH₂)—O-2' (LNA); 4'-(CH₂)—S-2'; 4'-(CH₂)₂—O-2' (ENA); 4'-CH(CH₃)—O-2' and 4'-CH(CH₂OCH₃)—O-2' (and analogs thereof see U.S. Pat. No. 7,399,845, issued on Jul. 15, 2008); 4'-C(CH₃)(CH₃)—O-2' (and analogs thereof see PCT/US2008/068922 published as WO/2009/006478, published Jan. 8, 2009); 4'-CH₂—N(OCH₃)-2' (and analogs thereof see PCT/US2008/064591 published as WO/2008/150729, published Dec. 11, 2008); 4'-CH₂—O—N(CH₃)-2' (see published U.S. Patent Application US2004-0171570, published Sep. 2, 2004); 4'-CH₂—N(R)—O-2', wherein R is H, C₁-C₁₂ alkyl, or a protecting group (see U.S. Pat. No. 7,427,672, issued on Sep. 23, 2008); 4'-CH₂—C(H)(CH₃)-2'

47

(see Chattopadhyaya et al., *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH₂-C(=CH₂)-2' (and analogs thereof see PCT/US2008/066154 published as WO 2008/154401, published on Dec. 8, 2008).

Further bicyclic nucleosides have been reported in published literature (see for example: Srivastava et al., *J. Am. Chem. Soc.*, 2007, 129(26) 8362-8379; Frieden et al., *Nucleic Acids Research*, 2003, 21, 6365-6372; Elayadi et al., *Curr. Opin. Inven. Drugs*, 2001, 2, 558-561; Braasch et al., *Chem. Biol.*, 2001, 8, 1-7; Orum et al., *Curr. Opin. Mol. Ther.*, 2001, 3, 239-243; Wahlestedt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2000, 97, 5633-5638; Singh et al., *Chem. Commun.*, 1998, 4, 455-456; Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630; Kumar et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222; Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039; U.S. Pat. Nos. 7,399,845; 7,053,207; 7,034,133; 6,794,499; 6,770,748; 6,670,461; 6,525,191; 6,268,490; U.S. Patent Publication Nos.: US2008-0039618; US2007-0287831; US2004-0171570; U.S. patent application Ser. Nos. 12/129,154; 61/099,844; 61/097,787; 61/086,231; 61/056,564; 61/026,998; 61/026,995; 60/989,574; International applications WO 2007/134181; WO 2005/021570; WO 2004/106356; and PCT International Applications Nos.: PCT/US2008/068922; PCT/US2008/066154; and PCT/US2008/064591). Each of the foregoing bicyclic nucleosides can be prepared having one or more stereochemical sugar configurations including for example α -L-ribofuranose and β -D-ribofuranose (see PCT international application PCT/DK98/00393, published on Mar. 25, 1999 as WO 99/14226).

As used herein, "monocyclic nucleosides" refer to nucleosides comprising modified sugar moieties that are not bicyclic sugar moieties. In certain embodiments, the sugar moiety, or sugar moiety analogue, of a nucleoside may be modified or substituted at any position.

As used herein, "4'-2' bicyclic nucleoside" or "4' to 2' bicyclic nucleoside" refers to a bicyclic nucleoside comprising a furanose ring comprising a bridge connecting two carbon atoms of the furanose ring connects the 2' carbon atom and the 4' carbon atom of the sugar ring.

In certain embodiments, bicyclic sugar moieties of BNA nucleosides include, but are not limited to, compounds having at least one bridge between the 4' and the 2' carbon atoms of the pentofuranosyl sugar moiety including without limitation, bridges comprising 1 or from 1 to 4 linked groups independently selected from —[C(R_a)(R_b)]_n—, —C(R_a)=C(R_b)—, —C(R_a)=N—, —C(=NR_a)—, —C(=O)—, —C(=S)—, —O—, —Si(R_a)₂—, —S(=O)_x—, and —N(R_a)—; wherein: x is 0, 1, or 2; n is 1, 2, 3, or 4; each R_a and R_b is, independently, H, a protecting group, hydroxyl, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, heterocycle radical, substituted heterocycle radical, heteroaryl, substituted heteroaryl, C₅-C₇ alicyclic radical, substituted C₅-C₇ alicyclic radical, halogen, OJ₁, NJ₁J₂, SJ₁, N₃, COOJ₁, acyl (C(=O)—H), substituted acyl, CN, sulfonyl (S(=O)₂-J₁), or sulfoxyl (S(=O)-J₁); and

each J₁ and J₂ is, independently, H, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, acyl (C(=O)—H), substituted acyl, a heterocycle radical, a substituted heterocycle radical, C₁-C₁₂ aminoalkyl, substituted C₁-C₁₂ aminoalkyl or a protecting group.

In certain embodiments, the bridge of a bicyclic sugar moiety is, —[C(R_a)(R_b)]_n—, —[C(R_a)(R_b)]_n—O—,


48

—C(R_aR_b)—N(R)—O— or —C(R_aR_b)—O—N(R)—. In certain embodiments, the bridge is 4'-CH₂-2', 4'-(CH₂)₂-2', 4'-(CH₂)₃-2', 4'-CH₂-O-2', 4'-(CH₂)₂-O-2', 4'-CH₂-O—N(R)-2' and 4'-CH₂-N(R)—O-2' wherein each R is, independently, H, a protecting group or C₁-C₁₂ alkyl.

In certain embodiments, bicyclic nucleosides are further defined by isomeric configuration. For example, a nucleoside comprising a 4'-(CH₂)—O-2' bridge, may be in the α -L configuration or in the β -D configuration. Previously, α -L-methyleneoxy (4'-CH₂—O-2') BNA's have been incorporated into antisense oligonucleotides that showed antisense activity (Frieden et al., *Nucleic Acids Research*, 2003, 21, 6365-6372).

In certain embodiments, bicyclic nucleosides include those having a 4' to 2' bridge wherein such bridges include without limitation, α -L-4'-(CH₂)—O-2', β -D-4'-CH₂—O-2', 4'-(CH₂)₂—O-2', 4'-CH₂—O—N(R)-2', 4'-CH₂—N(R)—O-2', 4'-CH(CH₃)—O-2', 4'-CH₂—S-2', 4'-CH₂—N(R)-2', 4'-CH₂—CH(CH₃)-2', and 4'-(CH₂)₃-2', wherein R is H, a protecting group or C₁-C₁₂ alkyl.

In certain embodiment, bicyclic nucleosides have the formula:



wherein:


Bx is a heterocyclic base moiety;

—Q_a-Q_b-Q_c— is —CH₂—N(R_c)—CH₂—, —C(=O)—N(R_c)—CH₂—, —CH₂—O—N(R_c)—, —CH₂—N(R_c)—O— or —N(R_c)—O—CH₂;

R_c is C₁-C₁₂ alkyl or an amino protecting group; and

T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium.

In certain embodiments, bicyclic nucleosides have the formula:



wherein:

Bx is a heterocyclic base moiety;


T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

Z_a is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, substituted C₁-C₆ alkyl, substituted C₂-C₆ alkenyl, substituted C₂-C₆ alkynyl, acyl, substituted acyl, substituted amide, thiol or substituted thiol.

In one embodiment, each of the substituted groups, is, independently, mono or poly substituted with substituent groups independently selected from halogen, oxo, hydroxyl, OJ_c, NJ_cJ_d, SJ_c, N₃, OC(=X)J_e, and NJ_eC(=X)NJ_cJ_d, wherein each J_c, J_d and J_e is, independently, H, C₁-C₆ alkyl, or substituted C₁-C₆ alkyl and X is O or NJ_c.

49

In certain embodiments, bicyclic nucleosides have the formula:




wherein:

Bx is a heterocyclic base moiety;

Ta and Tb are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

Zb is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, substituted C₁-C₆ alkyl, substituted C₂-C₆ alkenyl, substituted C₂-C₆ alkynyl or substituted acyl (C(=O)—).

In certain embodiments, bicyclic nucleosides have the formula:



wherein:


Bx is a heterocyclic base moiety;

Ta and Tb are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

Rd is C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl;

each qa, qb, qc and qd is, independently, H, halogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl, C₁-C₆ alkoxy, substituted C₁-C₆ alkoxy, acyl, substituted acyl, C₁-C₆ aminoalkyl or substituted C₁-C₆ aminoalkyl;

In certain embodiments, bicyclic nucleosides have the formula:



wherein:

Bx is a heterocyclic base moiety;

Ta and Tb are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

qa, qb, qc and qd are each, independently, hydrogen, halogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₁-C₁₂ alkoxy, substituted C₁-C₁₂ alkoxy,

50


OJ_j, SJ_j, SOJ_j, SO₂J_j, NJ_jJ_k, N₃, CN, C(=O)OJ_j, C(=O)NJ_jJ_k, C(=O)J_j, O—C(=O)NJ_jJ_k, N(H)C(=NH)NJ_jJ_k, N(H)C(=O)NJ_jJ_k or N(H)C(=S)NJ_jJ_k; or q_e and q_f together are =C(q_g)(q_h);

5 q_g and q_h are each, independently, H, halogen, C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl.

The synthesis and preparation of adenine, cytosine, guanine, 5-methyl-cytosine, thymine and uracil bicyclic nucleosides having a 4'-CH₂—O-2' bridge, along with their oligomerization, and nucleic acid recognition properties have been described (Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630). The synthesis of bicyclic nucleosides has also been described in WO 98/39352 and WO 99/14226.

Analogous of various bicyclic nucleosides that have 4' to 2' bridging groups such as 4'-CH₂—O-2' and 4'-CH₂—S-2', have also been prepared (Kumar et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222). Preparation of oligodeoxyribonucleotide duplexes comprising bicyclic nucleosides for use as substrates for nucleic acid polymerases has also been described (Wengel et al., WO 99/14226). Furthermore, synthesis of 2'-amino-BNA, a novel conformationally restricted high-affinity oligonucleotide analog has been described in the art (Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039). In addition, 2'-amino- and 2'-methylamino-BNA's have been prepared and the thermal stability of their duplexes with complementary RNA and DNA strands has been previously reported.

In certain embodiments, bicyclic nucleosides have the formula:



wherein:

Bx is a heterocyclic base moiety;

Ta and Tb are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

each q₁, q₂, q₃ and q₄ is, independently, H, halogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₁-C₁₂ alkoxy, substituted C₁-C₁₂ alkoxy, OJ_j, SJ_j, SOJ_j, SO₂J_j, NJ_jJ_k, N₃, CN, C(=O)OJ_j, C(=O)NJ_jJ_k, C(=O)J_j, O—C(=O)NJ_jJ_k, N(H)C(=NH)NJ_jJ_k, N(H)C(=O)NJ_jJ_k or N(H)C(=S)NJ_jJ_k; and








50 q₁ and q₂ or q₃ and q₄ together are =C(q_g)(q_h), wherein q_g and q_h are each, independently, H, halogen, C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl.

One carbocyclic bicyclic nucleoside having a 4'-(CH₂)₃-2' bridge and the alkenyl analog bridge 4'-CH=CH—CH₂-2' have been described (Frier et al., *Nucleic Acids Research*, 1997, 25(22), 4429-4443 and Albaek et al., *J. Org. Chem.*, 2006, 71, 7731-7740). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (Srivastava et al., *J. Am. Chem. Soc.* 2007, 129(26), 8362-8379).





In certain embodiments, bicyclic nucleosides include, but are not limited to, (A) α-L-methyleneoxy (4'-CH₂—O-2') BNA, (B) β-D-methyleneoxy (4'-CH₂—O-2') BNA, (C) ethyleneoxy (4'-(CH₂)₂—O-2') BNA, (D) aminoxy (4'-CH₂—O—N(R)-2') BNA, (E) oxyamino (4'-CH₂—N(R)—O-2') BNA, (F) methyl(methyleneoxy) (4'-CH(CH₃)—O-2') BNA (also referred to as constrained ethyl or cEt), (G)

51

methylene-thio (4'-CH₂—S-2') BNA, (H) methylene-amino (4'-CH₂—N(R)-2') BNA, (I) methyl carbocyclic (4'-CH₂—CH(CH₃)-2') BNA, (J) propylene carbocyclic (4'-(CH₂)₃-2') BNA, and (K) vinyl BNA as depicted below.

**52**

-continued



(A)

(B)

(C)

(D)

(E)

(F)

(G)

(H)

(I)


(J)

(K)

(L)


wherein Bx is the base moiety and R is, independently, H, a protecting group, C₁-C₆ alkyl or C₁-C₆ alkoxy.

As used herein, the term “modified tetrahydropyran nucleoside” or “modified THP nucleoside” means a nucleoside having a six-membered tetrahydropyran “sugar” substituted for the pentofuranosyl residue in normal nucleosides and can be referred to as a sugar surrogate. Modified THP nucleosides include, but are not limited to, what is referred to in the art as hexitol nucleic acid (HNA), anitol nucleic acid (ANA), manitol nucleic acid (MNA) (see Leumann, *Bioorg. Med. Chem.*, 2002, 10, 841-854) or fluoro HNA (F-HNA) having a tetrahydropyranyl ring system as illustrated below.



In certain embodiment, sugar surrogates are selected having the formula:

53



wherein:

Bx is a heterocyclic base moiety;


T₃ and T₄ are each, independently, an internucleoside linking group linking the tetrahydropyran nucleoside analog to the oligomeric compound or one of T₃ and T₄ is an internucleoside linking group linking the tetrahydropyran nucleoside analog to an oligomeric compound or oligonucleotide and the other of T₃ and T₄ is H, a hydroxyl protecting group, a linked conjugate group or a 5' or 3'-terminal group;

q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each independently, H, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl; and

one of R₁ and R₂ is hydrogen and the other is selected from halogen, substituted or unsubstituted alkoxy, NJ₁J₂, SJ₁, N₃, OC(=X)J₁, OC(=X)NJ₁J₂, NJ₃C(=X)NJ₁J₂ and CN, wherein X is O, S or NJ₁ and each J₁, J₂ and J₃ is, independently, H or C₁-C₆ alkyl.

In certain embodiments, q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each H. In certain embodiments, at least one of q₁, q₂, q₃, q₄, q₅, q₆ and q₇ is other than H. In certain embodiments, at least one of q₁, q₂, q₃, q₄, q₅, q₆ and q₇ is methyl. In certain embodiments, THP nucleosides are provided wherein one of R₁ and R₂ is F. In certain embodiments, R₁ is fluoro and R₂ is H; R₁ is methoxy and R₂ is H, and R₁ is methoxyethoxy and R₂ is H.

In certain embodiments, sugar surrogates comprise rings having more than 5 atoms and more than one heteroatom. For example nucleosides comprising morpholino sugar moieties and their use in oligomeric compounds has been reported (see for example: Braasch et al., *Biochemistry*, 2002, 41, 4503-4510; and U.S. Pat. Nos. 5,698,685; 5,166,315; 5,185,444; and 5,034,506). As used here, the term “morpholino” means a sugar surrogate having the following formula:




In certain embodiments, morpholinos may be modified, for example by adding or altering various substituent groups from the above morpholino structure. Such sugar surrogates are referred to herein as “modified morpholinos.”

Combinations of modifications are also provided without limitation, such as 2'-F-5'-methyl substituted nucleosides (see PCT International Application WO 2008/101157 published on Aug. 21, 2008 for other disclosed 5', 2'-bis substituted nucleosides) and replacement of the ribosyl ring oxygen atom with S and further substitution at the 2'-posi-

54

tion (see published U.S. Patent Application US2005-0130923, published on Jun. 16, 2005) or alternatively 5'-substitution of a bicyclic nucleic acid (see PCT International Application WO 2007/134181, published on Nov. 22, 2007 wherein a 4'-CH₂—O-2' bicyclic nucleoside is further substituted at the 5' position with a 5'-methyl or a 5'-vinyl group). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (see, e.g., Srivastava et al., *J. Am. Chem. Soc.* 2007, 129(26), 8362-8379).

In certain embodiments, antisense compounds comprise one or more modified cyclohexenyl nucleosides, which is a nucleoside having a six-membered cyclohexenyl in place of the pentofuranosyl residue in naturally occurring nucleosides. Modified cyclohexenyl nucleosides include, but are not limited to those described in the art (see for example commonly owned, published PCT Application WO 2010/036696, published on Apr. 10, 2010, Robeyns et al., *J. Am. Chem. Soc.*, 2008, 130(6), 1979-1984; Horvath et al., *Tetrahedron Letters*, 2007, 48, 3621-3623; Nauwelaerts et al., *J. Am. Chem. Soc.*, 2007, 129(30), 9340-9348; Gu et al., *Nucleosides, Nucleotides & Nucleic Acids*, 2005, 24(5-7), 993-998; Nauwelaerts et al., *Nucleic Acids Research*, 2005, 33(8), 2452-2463; Robeyns et al., *Acta Crystallographica, Section F: Structural Biology and Crystallization Communications*, 2005, F61(6), 585-586; Gu et al., *Tetrahedron*, 2004, 60(9), 2111-2123; Gu et al., *Oligonucleotides*, 2003, 13(6), 479-489; Wang et al., *J. Org. Chem.*, 2003, 68, 4499-4505; Verbeure et al., *Nucleic Acids Research*, 2001, 29(24), 4941-4947; Wang et al., *J. Org. Chem.*, 2001, 66, 8478-82; Wang et al., *Nucleosides, Nucleotides & Nucleic Acids*, 2001, 20(4-7), 785-788; Wang et al., *J. Am. Chem.*, 2000, 122, 8595-8602; Published PCT application, WO 06/047842; and Published PCT Application WO 01/049687; the text of each is incorporated by reference herein, in their entirety). Certain modified cyclohexenyl nucleosides have Formula X.



X

wherein independently for each of said at least one cyclohexenyl nucleoside analog of Formula X:

Bx is a heterocyclic base moiety;

T₃ and T₄ are each, independently, an internucleoside linking group linking the cyclohexenyl nucleoside analog to an antisense compound or one of T₃ and T₄ is an internucleoside linking group linking the tetrahydropyran nucleoside analog to an antisense compound and the other of T₃ and T₄ is H, a hydroxyl protecting group, a linked conjugate group, or a 5'- or 3'-terminal group; and

q₁, q₂, q₃, q₄, q₅, q₆, q₇, q₈ and q₉ are each, independently, H, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl, substituted C₂-C₆ alkynyl or other sugar substituent group.

Many other monocyclic, bicyclic and tricyclic ring systems are known in the art and are suitable as sugar surrogates that can be used to modify nucleosides for incorporation into

oligomeric compounds as provided herein (see for example review article: Leumann, Christian J. *Bioorg. & Med. Chem.*, 2002, 10, 841-854). Such ring systems can undergo various additional substitutions to further enhance their activity.

As used herein, "2'-modified sugar" means a furanosyl sugar modified at the 2' position. In certain embodiments, such modifications include substituents selected from: a halide, including, but not limited to substituted and unsubstituted alkoxy, substituted and unsubstituted thioalkyl, substituted and unsubstituted amino alkyl, substituted and unsubstituted alkyl, substituted and unsubstituted allyl, and substituted and unsubstituted alkynyl. In certain embodiments, 2' modifications are selected from substituents including, but not limited to: $O[(CH_2)_nO]_mCH_3$, $O(CH_2)_nNH_2$, $O(CH_2)_nCH_3$, $O(CH_2)_nF$, $O(CH_2)_nONH_2$, $OCH_2C(=O)N(H)CH_3$, and $O(CH_2)_nON[(CH_2)_nCH_3]_2$, where n and m are from 1 to about 10. Other 2'-substituent groups can also be selected from: C_1 - C_{12} alkyl, substituted alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH_3 , OCN, Cl, Br, CN, F, CF_3 , OCF_3 , $SOCH_3$, SO_2CH_3 , ONO_2 , NO_2 , N_3 , NH_2 , heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving pharmacokinetic properties, or a group for improving the pharmacodynamic properties of an antisense compound, and other substituents having similar properties. In certain embodiments, modified nucleosides comprise a 2'-MOE side chain (Baker et al., *J. Biol. Chem.*, 1997, 272, 11944-12000). Such 2'-MOE substitution have been described as having improved binding affinity compared to unmodified nucleosides and to other modified nucleosides, such as 2'-O-methyl, O-propyl, and O-amino-propyl. Oligonucleotides having the 2'-MOE substituent also have been shown to be antisense inhibitors of gene expression with promising features for in vivo use (Martin, *Helv. Chim. Acta*, 1995, 78, 486-504; Altmann et al., *Chimia*, 1996, 50, 168-176; Altmann et al., *Biochem. Soc. Trans.*, 1996, 24, 630-637; and Altmann et al., *Nucleosides Nucleotides*, 1997, 16, 917-926).

As used herein, "2'-modified" or "2'-substituted" refers to a nucleoside comprising a sugar comprising a substituent at the 2' position other than H or OH. 2'-modified nucleosides, include, but are not limited to, bicyclic nucleosides wherein the bridge connecting two carbon atoms of the sugar ring connects the 2' carbon and another carbon of the sugar ring; and nucleosides with non-bridging 2'-substituents, such as allyl, amino, azido, thio, O-allyl, $O-C_1-C_{10}$ alkyl, $-OCF_3$, $O-(CH_2)_2-O-CH_3$, $2'-O(CH_2)_2SCH_3$, $O-(CH_2)_2-O-N(R_m)(R_n)$, or $O-CH_2-C(=O)-N(R_m)(R_n)$, where each R_m and R_n is, independently, H or substituted or unsubstituted C_1-C_{10} alkyl. 2'-modified nucleosides may further comprise other modifications, for example at other positions of the sugar and/or at the nucleobase.

As used herein, "2'-F" refers to a nucleoside comprising a sugar comprising a fluoro group at the 2' position of the sugar ring.

As used herein, "2'-OMe" or "2'-OCH₃", "2'-O-methyl" or "2'-methoxy" each refers to a nucleoside comprising a sugar comprising an $-OCH_3$ group at the 2' position of the sugar ring.

As used herein, "MOE" or "2'-MOE" or "2'-OCH₂CH₂OCH₃" or "2'-O-methoxyethyl" each refers to a nucleoside comprising a sugar comprising a $-OCH_2CH_2OCH_3$ group at the 2' position of the sugar ring.

Methods for the preparations of modified sugars are well known to those skilled in the art. Some representative U.S.

patents that teach the preparation of such modified sugars include without limitation, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,670,633; 5,700,920; 5,792,847 and 6,600,032 and International Application PCT/US2005/019219, filed Jun. 2, 2005 and published as WO 2005/121371 on Dec. 22, 2005, and each of which is herein incorporated by reference in its entirety.

As used herein, "oligonucleotide" refers to a compound comprising a plurality of linked nucleosides. In certain embodiments, one or more of the plurality of nucleosides is modified. In certain embodiments, an oligonucleotide comprises one or more ribonucleosides (RNA) and/or deoxyribonucleosides (DNA).

In nucleotides having modified sugar moieties, the nucleobase moieties (natural, modified or a combination thereof) are maintained for hybridization with an appropriate nucleic acid target.

In certain embodiments, antisense compounds comprise one or more nucleosides having modified sugar moieties. In certain embodiments, the modified sugar moiety is 2'-MOE. In certain embodiments, the 2'-MOE modified nucleosides are arranged in a gapmer motif. In certain embodiments, the modified sugar moiety is a bicyclic nucleoside having a (4'-CH(CH₃)-O-2') bridging group. In certain embodiments, the (4'-CH(CH₃)-O-2') modified nucleosides are arranged throughout the wings of a gapmer motif.

Compositions and Methods for Formulating Pharmaceutical Compositions

Antisense oligonucleotides may be admixed with pharmaceutically acceptable active or inert substances for the preparation of pharmaceutical compositions or formulations. Compositions and methods for the formulation of pharmaceutical compositions are dependent upon a number of criteria, including, but not limited to, route of administration, extent of disease, or dose to be administered.

An antisense compound targeted to a C9ORF72 nucleic acid can be utilized in pharmaceutical compositions by combining the antisense compound with a suitable pharmaceutically acceptable diluent or carrier. A pharmaceutically acceptable diluent includes phosphate-buffered saline (PBS). PBS is a diluent suitable for use in compositions to be delivered parenterally. Accordingly, in one embodiment, employed in the methods described herein is a pharmaceutical composition comprising an antisense compound targeted to a C9ORF72 nucleic acid and a pharmaceutically acceptable diluent. In certain embodiments, the pharmaceutically acceptable diluent is PBS. In certain embodiments, the antisense compound is an antisense oligonucleotide.

Pharmaceutical compositions comprising antisense compounds encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other oligonucleotide which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to pharmaceutically acceptable salts of antisense compounds, prodrugs, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents. Suitable pharmaceutically acceptable salts include, but are not limited to, sodium and potassium salts.

A prodrug can include the incorporation of additional nucleosides at one or both ends of an antisense compound which are cleaved by endogenous nucleases within the body, to form the active antisense compound.

Conjugated Antisense Compounds

Antisense compounds may be covalently linked to one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the resulting antisense oligonucleotides. Typical conjugate groups include cholesterol moieties and lipid moieties. Additional conjugate groups include carbohydrates, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes.

Antisense compounds can also be modified to have one or more stabilizing groups that are generally attached to one or both termini of antisense compounds to enhance properties such as, for example, nuclease stability. Included in stabilizing groups are cap structures. These terminal modifications protect the antisense compound having terminal nucleic acid from exonuclease degradation, and can help in delivery and/or localization within a cell. The cap can be present at the 5'-terminus (5'-cap), or at the 3'-terminus (3'-cap), or can be present on both termini. Cap structures are well known in the art and include, for example, inverted deoxy abasic caps. Further 3' and 5'-stabilizing groups that can be used to cap one or both ends of an antisense compound to impart nuclease stability include those disclosed in WO 03/004602 published on Jan. 16, 2003.

Cell Culture and Antisense Compounds Treatment

The effects of antisense compounds on the level, activity or expression of C9ORF72 nucleic acids can be tested in vitro in a variety of cell types. Cell types used for such analyses are available from commercial vendors (e.g. American Type Culture Collection, Manassas, Va.; Zen-Bio, Inc., Research Triangle Park, N.C.; Clonetics Corporation, Walkersville, Md.) and are cultured according to the vendor's instructions using commercially available reagents (e.g. Invitrogen Life Technologies, Carlsbad, Calif.). Illustrative cell types include, but are not limited to, HepG2 cells, Hep3B cells, and primary hepatocytes.

In Vitro Testing of Antisense Oligonucleotides

Described herein are methods for treatment of cells with antisense oligonucleotides, which can be modified appropriately for treatment with other antisense compounds.

In general, cells are treated with antisense oligonucleotides when the cells reach approximately 60-80% confluency in culture.

One reagent commonly used to introduce antisense oligonucleotides into cultured cells includes the cationic lipid transfection reagent LIPOFECTIN (Invitrogen, Carlsbad, Calif.). Antisense oligonucleotides are mixed with LIPOFECTIN in OPTI-MEM 1 (Invitrogen, Carlsbad, Calif.) to achieve the desired final concentration of antisense oligonucleotide and a LIPOFECTIN concentration that typically ranges 2 to 12 ug/mL per 100 nM antisense oligonucleotide.

Another reagent used to introduce antisense oligonucleotides into cultured cells includes LIPOFECTAMINE (Invitrogen, Carlsbad, Calif.). Antisense oligonucleotide is mixed with LIPOFECTAMINE in OPTI-MEM 1 reduced serum medium (Invitrogen, Carlsbad, Calif.) to achieve the desired concentration of antisense oligonucleotide and a LIPOFECTAMINE concentration that typically ranges 2 to 12 ug/mL per 100 nM antisense oligonucleotide.

Another technique used to introduce antisense oligonucleotides into cultured cells includes electroporation.

Cells are treated with antisense oligonucleotides by routine methods. Cells are typically harvested 16-24 hours after antisense oligonucleotide treatment, at which time RNA or protein levels of target nucleic acids are measured by methods known in the art and described herein. In general,

when treatments are performed in multiple replicates, the data are presented as the average of the replicate treatments.

The concentration of antisense oligonucleotide used varies from cell line to cell line. Methods to determine the optimal antisense oligonucleotide concentration for a particular cell line are well known in the art. Antisense oligonucleotides are typically used at concentrations ranging from 1 nM to 300 nM when transfected with LIPOFECTAMINE. Antisense oligonucleotides are used at higher concentrations ranging from 625 to 20,000 nM when transfected using electroporation.

RNA Isolation

RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are well known in the art. RNA is prepared using methods well known in the art, for example, using the TRIZOL Reagent (Invitrogen, Carlsbad, Calif.) according to the manufacturer's recommended protocols.

Analysis of Inhibition of Target Levels or Expression

Inhibition of levels or expression of a C9ORF72 nucleic acid can be assayed in a variety of ways known in the art. For example, target nucleic acid levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or quantitative real-time PCR. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are well known in the art. Northern blot analysis is also routine in the art. Quantitative real-time PCR can be conveniently accomplished using the commercially available ABI PRISM 7600, 7700, or 7900 Sequence Detection System, available from PE-Applied Biosystems, Foster City, Calif. and used according to manufacturer's instructions.

Quantitative Real-Time PCR Analysis of Target RNA Levels

Quantitation of target RNA levels may be accomplished by quantitative real-time PCR using the ABI PRISM 7600, 7700, or 7900 Sequence Detection System (PE-Applied Biosystems, Foster City, Calif.) according to manufacturer's instructions. Methods of quantitative real-time PCR are well known in the art.

Prior to real-time PCR, the isolated RNA is subjected to a reverse transcriptase (RT) reaction, which produces complementary DNA (cDNA) that is then used as the substrate for the real-time PCR amplification. The RT and real-time PCR reactions are performed sequentially in the same sample well. RT and real-time PCR reagents are obtained from Invitrogen (Carlsbad, Calif.). RT real-time-PCR reactions are carried out by methods well known to those skilled in the art.

Gene (or RNA) target quantities obtained by real time PCR are normalized using either the expression level of a gene whose expression is constant, such as cyclophilin A, or by quantifying total RNA using RIBOGREEN (Invitrogen, Inc. Carlsbad, Calif.). Cyclophilin A expression is quantified by real time PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RIBOGREEN RNA quantification reagent (Invitrogen, Inc. Eugene, Oreg.). Methods of RNA quantification by RIBOGREEN are taught in Jones, L. J., et al, (Analytical Biochemistry, 1998, 265, 368-374). A CYTOFLUOR 4000 instrument (PE Applied Biosystems) is used to measure RIBOGREEN fluorescence.

Probes and primers are designed to hybridize to a C9ORF72 nucleic acid. Methods for designing real-time PCR probes and primers are well known in the art, and may include the use of software such as PRIMER EXPRESS Software (Applied Biosystems, Foster City, Calif.).

Analysis of Protein Levels

Antisense inhibition of C9ORF72 nucleic acids can be assessed by measuring C9ORF72 protein levels. Protein levels of C9ORF72 can be evaluated or quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), enzyme-linked immunosorbent assay (ELISA), quantitative protein assays, protein activity assays (for example, caspase activity assays), immunohistochemistry, immunocytochemistry or fluorescence-activated cell sorting (FACS). Antibodies directed to a target can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, Mich.), or can be prepared via conventional monoclonal or polyclonal antibody generation methods well known in the art. Antibodies useful for the detection of mouse, rat, monkey, and human C9ORF72 are commercially available.

In Vivo Testing of Antisense Compounds

Antisense compounds, for example, antisense oligonucleotides, are tested in animals to assess their ability to inhibit expression of C9ORF72 and produce phenotypic changes, such as, improved motor function and respiration. In certain embodiments, motor function is measured by rotarod, grip strength, pole climb, open field performance, balance beam, hindpaw footprint testing in the animal. In certain embodiments, respiration is measured by whole body plethysmograph, invasive resistance, and compliance measurements in the animal. Testing may be performed in normal animals, or in experimental disease models. For administration to animals, antisense oligonucleotides are formulated in a pharmaceutically acceptable diluent, such as phosphate-buffered saline (PBS) or artificial cerebrospinal fluid (aCSF). Administration includes parenteral routes of administration, such as intraperitoneal, intravenous, and subcutaneous, as well as central routes of administration such as intracerebroventricular or intrathecal. Calculation of antisense oligonucleotide dosage and dosing frequency is within the abilities of those skilled in the art, and depends upon factors such as route of administration and animal body weight. Following a period of treatment with antisense oligonucleotides, RNA is isolated from CNS tissue or CSF and changes in C9ORF72 nucleic acid expression are measured.

Targeting C9ORF72

Antisense oligonucleotides described herein may hybridize to a C9ORF72 nucleic acid in any stage of RNA processing. For example, described herein are antisense oligonucleotides that are complementary to a pre-mRNA or a mature mRNA. Additionally, antisense oligonucleotides described herein may hybridize to any element of a C9ORF72 nucleic acid. For example, described herein are antisense oligonucleotides that are complementary to an exon, an intron, the 5' UTR, the 3' UTR, a repeat region, a hexanucleotide repeat expansion, a splice junction, an exon: exon splice junction, an exonic splicing silencer (ESS), an exonic splicing enhancer (ESE), exon 1a, exon 1b, exon 1c, exon 1d, exon 1e, exon 2, exon 3, exon 4, exon 5, exon 6, exon 7, exon 8, exon 9, exon 10, exon 11, intron 1, intron 2, intron 3, intron 4, intron 5, intron 6, intron 7, intron 8, intron 9, or intron 10 of a C9ORF72 nucleic acid.

In certain embodiments, antisense oligonucleotides described herein hybridize to all variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein selectively hybridize to certain variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein selectively hybridize to variants of C9ORF72 containing a hexanucleotide repeat expansion. In certain embodiments, the antisense oligonucleotides

described herein selectively hybridize to pre-mRNA variants containing the hexanucleotide repeat. In certain embodiments, pre-mRNA variants of C9ORF72 containing a hexanucleotide repeat expansion include SEQ ID NO: 1-3 and 6-10. In certain embodiments, such hexanucleotide repeat expansion comprises at least 24 repeats of any of GGGGCC, GGGGGG, GGGGGC, or GGGGCG.

In certain embodiments, the antisense oligonucleotides described herein inhibit expression of all variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein inhibit expression of all variants of C9ORF72 equally. In certain embodiments, the antisense oligonucleotides described herein preferentially inhibit expression of one or more variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein preferentially inhibit expression of variants of C9ORF72 containing a hexanucleotide repeat expansion. In certain embodiments, the antisense oligonucleotides described herein selectively inhibit expression of pre-mRNA variants containing the hexanucleotide repeat. In certain embodiments, the antisense oligonucleotides described herein selectively inhibit expression of C9ORF72 pathogenic associated mRNA variants. In certain embodiments, pre-mRNA variants of C9ORF72 containing a hexanucleotide repeat expansion include SEQ ID NO: 1-3 and 6-10. In certain embodiments, such hexanucleotide repeat expansion comprises at least 24 repeats of any of GGGGCC, GGGGGG, GGGGGC, or GGGGCG. In certain embodiments, the hexanucleotide repeat expansion forms nuclear foci. In certain embodiments, antisense oligonucleotides described herein are useful for reducing nuclear foci. Nuclear foci may be reduced in terms of percent of cells with foci as well as number of foci per cell.

Selective Inhibition of Certain Pathogenic Associated Variants

In certain examples herein, primer probe set RTS3905 detects an mRNA variant (e.g. NM_001256054.1) processed from a pre-mRNA variant containing the hexanucleotide repeat. The mRNA variant processed from a pre-mRNA variant containing the hexanucleotide repeat (i.e., the "C9ORF72 pathogenic associated mRNA variant"). A pre-mRNA contains the hexanucleotide repeat when transcription of the pre-mRNA begins in the region from the start site of exon 1A to the start site of exon 1B, e.g., nucleotides 1107 to 1520 of the genomic sequence (SEQ ID NO: 2, the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 27565000). Oligonucleotides were designed in this region to selectively target the pre-mRNA variant containing the hexanucleotide repeat. RTS3905 measures an mRNA product (i.e. the C9ORF72 pathogenic associated mRNA variant) of the pre-mRNA variant containing the hexanucleotide repeat and, therefore, measures the reduction of the pre-mRNA variant containing the hexanucleotide repeat.

C9ORF72 Features

Antisense oligonucleotides described herein may hybridize to any C9ORF72 variant at any state of processing within any element of the C9ORF72 gene. For example, antisense oligonucleotides described herein may hybridize to an exon, an intron, the 5' UTR, the 3' UTR, a repeat region, a hexanucleotide repeat expansion, a splice junction, an exon: exon splice junction, an exonic splicing silencer (ESS), an exonic splicing enhancer (ESE), exon 1a, exon 1b, exon 1c, exon 1d, exon 1e, exon 2, exon 3, exon 4, exon 5, exon 6, exon 7, exon 8, exon 9, exon 10, exon 11, intron 1, intron 2, intron 3, intron 4, intron 5, intron 6, intron 7, intron 8, intron 9, or intron 10. For example, antisense oligonucleotides may

61

target any of the exons characterized below in Tables 1-5 for the various C9ORF72 variants described below. Antisense oligonucleotides described herein may also target variants not characterized below and such variants are characterized in GENBANK. Moreover, antisense oligonucleotides described herein may also target elements other than exons and such elements are characterized in GENBANK.

TABLE 1

Functional Segments for NM_001256054.1 (SEQ ID NO: 1)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1C	1	158	1137	1294
exon 2	159	646	7839	8326
exon 3	647	706	9413	9472
exon 4	707	802	12527	12622
exon 5	803	867	13354	13418
exon 6	868	940	14704	14776
exon 7	941	1057	16396	16512
exon 8	1058	1293	18207	18442
exon 9	1294	1351	24296	24353
exon 10	1352	1461	26337	26446
exon 11	1462	3339	26581	28458

TABLE 2

Functional Segments for NM_018325.3 (SEQ ID NO: 4)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1B	1	63	1510	1572
exon 2	64	551	7839	8326
exon 3	552	611	9413	9472
exon 4	612	707	12527	12622
exon 5	708	772	13354	13418
exon 6	773	845	14704	14776
exon 7	846	962	16396	16512
exon 8	963	1198	18207	18442
exon 9	1199	1256	24296	24353
exon 10	1257	1366	26337	26446
exon 11	1367	3244	26581	28458

TABLE 3

Functional Segments for NM_145005.5 (SEQ ID NO: 6)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1A	1	80	1137	1216
exon 2	81	568	7839	8326
exon 3	569	628	9413	9472
exon 4	629	724	12527	12622
exon 5B (exon 5 into intron 5)	725	1871	13354	14500

62

TABLE 4

Functional Segments for DB079375.1 (SEQ ID NO: 7)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1E	1	35	1135	1169
exon 2	36	524	7839	8326
exon 3 (EST ends before end of full exon)	525	562	9413	9450

TABLE 5

Functional Segments for BU194591.1 (SEQ ID NO: 8)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1D	1	36	1241	1279
exon 2	37	524	7839	8326
exon 3	525	584	9413	9472
exon 4	585	680	12527	12622
exon 5B (exon 5 into intron 5)	681	798	13354	13465

Certain Indications

In certain embodiments, provided herein are methods of treating an individual comprising administering one or more pharmaceutical compositions described herein. In certain embodiments, the individual has a neurodegenerative disease. In certain embodiments, the individual is at risk for developing a neurodegenerative disease, including, but not limited to, ALS or FTD. In certain embodiments, the individual has been identified as having a C9ORF72 associated disease. In certain embodiments, provided herein are methods for prophylactically reducing C9ORF72 expression in an individual. Certain embodiments include treating an individual in need thereof by administering to an individual a therapeutically effective amount of an antisense compound targeted to a C9ORF72 nucleic acid.

In one embodiment, administration of a therapeutically effective amount of an antisense compound targeted to a C9ORF72 nucleic acid is accompanied by monitoring of C9ORF72 levels in an individual, to determine an individual's response to administration of the antisense compound. An individual's response to administration of the antisense compound may be used by a physician to determine the amount and duration of therapeutic intervention.

In certain embodiments, administration of an antisense compound targeted to a C9ORF72 nucleic acid results in reduction of C9ORF72 expression by at least 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 99%, or a range defined by any two of these values. In certain embodiments, administration of an antisense compound targeted to a C9ORF72 nucleic acid results in improved motor function and respiration in an animal. In certain embodiments, administration of a C9ORF72 antisense compound improves motor function and respiration by at least 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 99%, or a range defined by any two of these values.

In certain embodiments, pharmaceutical compositions comprising an antisense compound targeted to C9ORF72 are used for the preparation of a medicament for treating a patient suffering or susceptible to a neurodegenerative disease including ALS and FTD.

Certain Combination Therapies

In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with one or more other pharmaceutical agents. In certain embodiments, such one or more other pharmaceutical agents are designed to treat the same disease, disorder, or condition as the one or more pharmaceutical compositions described herein. In certain embodiments, such one or more other pharmaceutical agents are designed to treat a different disease, disorder, or condition as the one or more pharmaceutical compositions described herein. In certain embodiments, such one or more other pharmaceutical agents are designed to treat an undesired side effect of one or more pharmaceutical compositions described herein. In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with another pharmaceutical agent to treat an undesired effect of that other pharmaceutical agent. In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with another pharmaceutical agent to produce a combinational effect. In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with another pharmaceutical agent to produce a synergistic effect.

In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are administered at the same time. In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are administered at different times. In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are prepared together in a single formulation. In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are prepared separately.

In certain embodiments, pharmaceutical agents that may be co-administered with a pharmaceutical composition described herein include Riluzole (Rilutek), Lioresal (Lioresal), and Dexpropomid.

In certain embodiments, pharmaceutical agents that may be co-administered with a C9ORF72 specific inhibitor described herein include, but are not limited to, an additional C9ORF72 inhibitor.

In certain embodiments, pharmaceutical agents that may be co-administered with a C9ORF72 specific inhibitor described herein include, but are not limited to, a C9ORF72 antisense transcript specific inhibitor. In certain embodiments, the C9ORF72 antisense transcript specific inhibitor is an antisense compound. In certain embodiments, the antisense compound is a modified oligonucleotide. In certain embodiments, the modified oligonucleotide is single-stranded.

In certain embodiments, the co-administered pharmaceutical agent is administered prior to administration of a pharmaceutical composition described herein. In certain

embodiments, the co-administered pharmaceutical agent is administered following administration of a pharmaceutical composition described herein. In certain embodiments the co-administered pharmaceutical agent is administered at the same time as a pharmaceutical composition described herein. In certain embodiments the dose of a co-administered pharmaceutical agent is the same as the dose that would be administered if the co-administered pharmaceutical agent was administered alone. In certain embodiments the dose of a co-administered pharmaceutical agent is lower than the dose that would be administered if the co-administered pharmaceutical agent was administered alone. In certain embodiments the dose of a co-administered pharmaceutical agent is greater than the dose that would be administered if the co-administered pharmaceutical agent was administered alone.

In certain embodiments, the co-administration of a second compound enhances the effect of a first compound, such that co-administration of the compounds results in an effect that is greater than the effect of administering the first compound alone. In other embodiments, the co-administration results in effects that are additive of the effects of the compounds when administered alone. In certain embodiments, the co-administration results in effects that are supra-additive of the effects of the compounds when administered alone. In certain embodiments, the first compound is an antisense compound. In certain embodiments, the second compound is an antisense compound.

Certain Comparator Compositions

In certain embodiments, compounds described herein are more tolerable than ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, and/or ISIS 577083. ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, and ISIS 577083 were selected as comparator compounds because they exhibited high levels of dose-dependent reduction of C9ORF72 mRNA in various studies described in WO2014/062691. Thus, ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, and ISIS 577083 were deemed highly efficacious and potent compounds. Additionally, ISIS 577065, ISIS 577056, and ISIS 576816 described in WO2014/062691 are structurally similar as compounds described herein. For example, ISIS 577065 has a 16 nucleobase overlap with ISIS 801287; ISIS 577056 has a 16 nucleobase overlap with ISIS 806679; ISIS 576816 has an 18 nucleobase overlap with ISIS 802473 (18-mer); and ISIS 576816 has an 18 nucleobase overlap with ISIS 802459.

In certain embodiments, ISIS 576816, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') GCCT-TACTCTAGGACCAAGA (incorporated herein as SEQ ID NO: 21), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 576816 achieved an average FOB score of 7.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS

802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 576816.

In certain embodiments, ISIS 576974, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') GGGACAC-TACAAGGTAGTAT (incorporated herein as SEQ ID NO: 56), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 576974 achieved an average FOB score of 5.67 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 576974.

In certain embodiments, ISIS 577061, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') TACAGGCT-GCGGTTGTTTCC (incorporated herein as SEQ ID NO: 57), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577061 achieved an average FOB score of 7.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577061.

In certain embodiments, ISIS 577065, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') CCCGGC-CCCTAGCGCGCGAC (incorporated herein as SEQ ID NO: 58), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577065 achieved an average FOB score of 6.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577065.

In certain embodiments, ISIS 577083, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') GGTAAC-TCAAACCTCTTGGG (incorporated herein as SEQ ID NO: 59), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577083 achieved an average FOB score of 7.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described

herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577083.

In certain embodiments, ISIS 577056, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') AATCTT-TATCAGGTCTTTTC (incorporated herein as SEQ ID NO: 60), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577056 achieved an average FOB score of 6.5 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577056.

Certain Human Therapeutics

The human C9ORF72 antisense oligonucleotides described herein are human therapeutics. Various parameters of potency, efficacy, and/or tolerability are being examined. Such parameters include in vitro inhibition of total C9ORF72 RNA expression, in vitro inhibition of C9ORF72 pathogenic associated RNA variant expression, in vitro dose response (IC50), in vivo inhibition of total or pathogenic RNA and/or protein in a transgenic animal containing a human C9ORF72 transgene in relevant tissues (e.g., brain and/or spinal cord), tolerability in mouse, tolerability in rat, and/or tolerability in a primate. Tolerability markers that may be measured include blood and serum chemistry parameters, CSF chemistry parameters, body and organ weights, general observations and/or behavioral tests, and/or biochemical markers such as GFAP and/or AIF1. Acute or long term tolerability may be measured.

Certain Compositions

1. ISIS 801287

In certain embodiments, ISIS 801287 is characterized as a 4-8-6 MOE gapmer, having a sequence of (from 5' to 3') GCCCCTAGCGCGCGACTC (incorporated herein as SEQ ID NO: 33), wherein each of nucleosides 1-4 and 13-18 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 5-12 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 13 to 14, 14 to 15, and 15 to 16 are phosphodiester linkages and the internucleoside linkages between nucleosides 1 to 2, 4 to 5, 5 to 6, 6 to 7, 7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to 12, 12 to 13, 16 to 17, and 17 to 18 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.

In certain embodiments, ISIS 801287 is described by the following chemical notation: Ges mCeo mCeo mCes mCds Tds Ads Gds mCds Gds mCds Gds mCeo Geo Aeo mCes Tes mCe; wherein,

A=an adenine,

mC=a 5'-methylcytosine

67


G=a guanine,
 T=a thymine,
 e=a 2'-O-methoxyethylribose modified sugar
 d=a 2'-deoxyribose sugar,

68

s=a phosphorothioate internucleoside linkage, and
 o=a phosphodiester internucleoside linkage.

In certain embodiments, ISIS 801287 is described by the
 following chemical structure, or a salt thereof:


(SEQ ID NO: 33)



69

Structure 1. ISIS 801287


In certain embodiments, the sodium salt of ISIS 801287 is described by the following chemical structure:



70

7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to 12, 12 to 13, 13 to 14, 14 to 15, 15 to 16, 16 to 17, 18 to 19, and 19 to 20 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.

(SEQ ID NO: 33)



55

Structure 2. The Sodium Salt of ISIS 801287

2. ISIS 806679

In certain embodiments, ISIS 806679 is characterized as a 6-10-4 MOE gapmer, having a sequence of (from 5' to 3') 60 GGTAAATCTTTATCAGGTCT (incorporated herein as SEQ ID NO: 49), wherein each of nucleosides 1-6 and 17-20 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 7-16 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 65 4 to 5, 5 to 6, 6 to 7, and 17 to 18 are phosphodiester linkages and the internucleoside linkages between nucleosides 1 to 2,

In certain embodiments, ISIS 806679 is described by the following chemical notation: Ges Geo Teo Teo AeO AeO Tds mCds Tds Tds Tds Ads Tds mCds Ads Gds Geo Tes mCes Te; wherein,


A=an adenine,
mC=a 5'-methylcytosine
G=a guanine,
T=a thymine,
e=a 2'-O-methoxyethylribose modified sugar
d=a 2'-deoxyribose sugar,
s=a phosphorothioate internucleoside linkage, and
o=a phosphodiester internucleoside linkage.

71

In certain embodiments, ISIS 806679 is described by the following chemical structure, or a salt thereof:

72

(SEQ ID NO: 49)




73

Structure 3. ISIS 806679

In certain embodiments, the sodium salt of ISIS 806679 is described by the following chemical structure:

(SEQ ID NO: 49)



Structure 4. The Sodium Salt of ISIS 806679

3. ISIS 802473

In certain embodiments, ISIS 802473 is characterized as a 4-8-6 MOE gapmer, having a sequence of (from 5' to 3') GCCTTACTCTAGGACCAA (incorporated herein as SEQ ID NO: 47), wherein each of nucleosides 1-4 and 13-18 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 5-12 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 13 to 14, 14 to 15, and 15 to 16 are phosphodiester linkages and the internucleoside linkages between nucleosides are 1 to 2, 4 to 5, 5 to 6, 6 to 7, 7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to

12, 12 to 13, 16 to 17, and 17 to 18 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.

In certain embodiments, ISIS 802473 is described by the following chemical notation: Ges mCeo mCeo Tes Tds Ads mCds Tds mCds Tds Ads Gds Geo Aeo mCeo mCes Aes Ae; wherein,

A=an adenine,

mC=a 5'-methylcytosine

G=a guanine,

T=a thymine,


e=a 2'-O-methoxyethylribose modified sugar

d=a 2'-deoxyribose sugar,

s=a phosphorothioate internucleoside linkage, and

o=a phosphodiester internucleoside linkage.

76



The image displays three chemical structures of polyphosphate nucleic acid analogs, labeled (a), (b), and (c). Each structure consists of a sugar-phosphate backbone with various nitrogenous bases attached. The structures are shown in a schematic representation, with the sugar rings and phosphate groups connected by oxygen atoms. The bases include purines (adenine, guanine) and pyrimidines (cytosine, uracil, thymine). The structures are labeled (a), (b), and (c) at the bottom.

In certain embodiments, ISIS 802459 is described by the following chemical notation: Ges mCeo mCeo Tds Tds Ads mCds Tds mCds Tds Ads Gds Gds Ae0 mCeo mCeo Ae0 Aes Ges Ae; wherein,
A=an adenine,
mC=a 5'-methylcytosine

79


G=a guanine,
 T=a thymine,
 e=a 2'-O-methoxyethylribose modified sugar
 d=a 2'-deoxyribose sugar,

80

s=a phosphorothioate internucleoside linkage, and
 o=a phosphodiester internucleoside linkage.

In certain embodiments, ISIS 802459 is described by the following chemical structure, or a salt thereof:

(SEQ ID NO: 21)




81

Structure 7 ISIS 802459

In certain embodiments, the sodium salt of ISIS 802459 is described by the following chemical structure:

(SEQ ID NO: 21)



Structure 8. The Sodium Salt of ISIS 802459

EXAMPLES

Non-Limiting Disclosure and Incorporation by Reference

While certain compounds, compositions, and methods described herein have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds described herein and are not intended to limit the same. Each of the

references recited in the present application is incorporated herein by reference in its entirety.

Example 1: Antisense Oligonucleotides Targeting Human C9ORF72

The antisense oligonucleotides in the table below were designed as MOE gapmers. The central gap segment of each gapmer contains 2'-deoxynucleosides and is flanked by wing segments on both the 5' end and on the 3' end containing nucleosides that each comprise a 2'-MOE group. The specific motif of each gapmer is listed in table below, repre-

sented by three numbers separated by hyphens. The numbers represent the number of nucleosides in the 5'-wing, the gap, and the 3'-wing, respectively. All cytosine residues throughout each oligonucleotide are 5-methylcytosines. The internucleoside linkages for the gapmers are mixed phosphorothioate and phosphodiester linkages. The internucleoside linkages for each gapmer are presented in the Linkage column, where 'o' indicates a phosphodiester linkage and 's' indicates a phosphorothioate linkage.

Each antisense oligonucleotide listed in the table below is targeted to the human C9ORF72 genomic sequence, designated herein as SEQ ID NO: 2 (the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 27565000). "Start site" indicates the 5'-most nucleoside to which the gapmer is targeted in the human genomic sequence. "Stop site" indicates the 3'-most nucleoside to which the gapmer is targeted human genomic sequence.

TABLE 6

Antisense oligonucleotides targeting human C9ORF72						
Isis No.	Start Site	Stop Site	Sequence	Linkage	Motif	SEQ ID NO
791656	1445	1464	CCGGCCCCCTAGCGCGCGACT	sooooooooooooooss	3-10-7	22
791657	1445	1464	CCGGCCCCCTAGCGCGCGACT	sooooooooooooooss	4-10-6	22
791658	1445	1464	CCGGCCCCCTAGCGCGCGACT	sooooooooooooooss	6-10-4	22
791659	1445	1464	CCGGCCCCCTAGCGCGCGACT	sooooooooooooooss	7-10-3	22
791660	1445	1463	CGGCCCCCTAGCGCGCGACT	sooooooooooooooss	5-9-5	23
791661	1446	1464	CCGGCCCCCTAGCGCGCGAC	sooooooooooooooss	5-9-5	24
791662	1445	1463	CGGCCCCCTAGCGCGCGACT	sooooooooooooooss	4-9-6	23
791663	1446	1464	CCGGCCCCCTAGCGCGCGAC	sooooooooooooooss	4-9-6	24
791664	1445	1463	CGGCCCCCTAGCGCGCGACT	sooooooooooooooss	6-9-4	23
791665	1446	1464	CCGGCCCCCTAGCGCGCGAC	sooooooooooooooss	6-9-4	24
801274	1440	1459	CCCTAGCGCGCGACTCCTGA	sooooooooooooooss	4-10-6	25
801275	1441	1460	CCCCTAGCGCGCGACTCCTG	sooooooooooooooss	4-10-6	26
801276	1442	1461	GCCCCCTAGCGCGCGACTCCT	sooooooooooooooss	4-10-6	27
801277	1443	1462	GGCCCCCTAGCGCGCGACTCC	sooooooooooooooss	4-10-6	28
801278	1444	1463	CGGCCCCCTAGCGCGCGACTC	sooooooooooooooss	4-10-6	29
801279	1440	1459	CCCTAGCGCGCGACTCCTGA	sooooooooooooooss	6-10-4	25
801280	1441	1460	CCCCTAGCGCGCGACTCCTG	sooooooooooooooss	6-10-4	26
801281	1442	1461	GCCCCCTAGCGCGCGACTCCT	sooooooooooooooss	6-10-4	27
801282	1443	1462	GGCCCCCTAGCGCGCGACTCC	sooooooooooooooss	6-10-4	28
801283	1444	1463	CGGCCCCCTAGCGCGCGACTC	sooooooooooooooss	6-10-4	29
801284	1441	1458	CCTAGCGCGCGACTCCTG	sooooooooooooooss	4-8-6	30
801285	1442	1459	CCCTAGCGCGCGACTCCT	sooooooooooooooss	4-8-6	31
801286	1443	1460	CCCCTAGCGCGCGACTCC	sooooooooooooooss	4-8-6	32
801287	1444	1461	GCCCCCTAGCGCGCGACTC	sooooooooooooooss	4-8-6	33
801288	1445	1462	GGCCCCCTAGCGCGCGACT	sooooooooooooooss	4-8-6	34
801289	1446	1463	CGGCCCCCTAGCGCGCGAC	sooooooooooooooss	4-8-6	35
801290	1441	1458	CCTAGCGCGCGACTCCTG	sooooooooooooooss	6-8-4	30
801291	1442	1459	CCCTAGCGCGCGACTCCT	sooooooooooooooss	6-8-4	31
801292	1443	1460	CCCCTAGCGCGCGACTCC	sooooooooooooooss	6-8-4	32
801293	1444	1461	GCCCCCTAGCGCGCGACTC	sooooooooooooooss	6-8-4	33
801294	1445	1462	GGCCCCCTAGCGCGCGACT	sooooooooooooooss	6-8-4	34
801295	1446	1463	CGGCCCCCTAGCGCGCGAC	sooooooooooooooss	6-8-4	35

TABLE 6-continued

Antisense oligonucleotides targeting human C9ORF72						
Isis No.	Start Site	Stop Site Sequence	Linkage	Motif	SEQ ID NO	
801296	1403	1422 AGGCTGCGGTTGTTTCCCTC	sooooooooooooooooo	4-10-6	36	
801297	1404	1423 CAGGCTGCGGTTGTTTCCCT	sooooooooooooooooo	4-10-6	37	
801298	1403	1421 GGCTGCGGTTGTTTCCCTC	sooooooooooooooooo	5-9-5	38	
801299	1404	1422 AGGCTGCGGTTGTTTCCCT	sooooooooooooooooo	5-9-5	39	
801300	1405	1423 CAGGCTGCGGTTGTTTCCC	sooooooooooooooooo	5-9-5	40	
801301	1403	1421 GGCTGCGGTTGTTTCCCTC	sooooooooooooooooo	4-9-6	38	
801302	1404	1422 AGGCTGCGGTTGTTTCCCT	sooooooooooooooooo	4-9-6	39	
801303	1405	1423 CAGGCTGCGGTTGTTTCCC	sooooooooooooooooo	4-9-6	40	
801304	1403	1421 GGCTGCGGTTGTTTCCCTC	sooooooooooooooooo	6-9-4	38	
801305	1404	1422 AGGCTGCGGTTGTTTCCCT	sooooooooooooooooo	6-9-4	39	
801306	1405	1423 CAGGCTGCGGTTGTTTCCC	sooooooooooooooooo	6-9-4	40	
801307	1403	1420 GCTGCGGTTGTTTCCCTC	sooooooooooooooooo	4-8-6	41	
801308	1404	1421 GGCTGCGGTTGTTTCCCT	sooooooooooooooooo	4-8-6	42	
801309	1405	1422 AGGCTGCGGTTGTTTCCC	sooooooooooooooooo	4-8-6	43	
801310	1406	1423 CAGGCTGCGGTTGTTTCC	sooooooooooooooooo	4-8-6	44	
801311	1403	1420 GCTGCGGTTGTTTCCCTC	sooooooooooooooooo	6-8-4	41	
801312	1404	1421 GGCTGCGGTTGTTTCCCT	sooooooooooooooooo	6-8-4	42	
801313	1405	1422 AGGCTGCGGTTGTTTCCC	sooooooooooooooooo	6-8-4	43	
801314	1406	1423 CAGGCTGCGGTTGTTTCC	sooooooooooooooooo	6-8-4	44	
801315	1403	1422 AGGCTGCGGTTGTTTCCCTC	sooooooooooooooooo	6-10-4	36	
801316	1404	1423 CAGGCTGCGGTTGTTTCCCT	sooooooooooooooooo	6-10-4	37	
802459	7990	8009 GCCTTACTCTAGGACCAAGA	sooooooooooooooooo	3-10-7	21	
802460	8012	8031 TCTGTCTTTGGAGCCCAAAT	sooooooooooooooooo	3-10-7	45	
802461	8186	8205 CTGCGATCCCCATTCCAGTT	sooooooooooooooooo	3-10-7	46	
802462	7990	8009 GCCTTACTCTAGGACCAAGA	sooooooooooooooooo	4-10-6	21	
802463	8012	8031 TCTGTCTTTGGAGCCCAAAT	sooooooooooooooooo	4-10-6	45	
802464	8186	8205 CTGCGATCCCCATTCCAGTT	sooooooooooooooooo	4-10-6	46	
802465	7990	8009 GCCTTACTCTAGGACCAAGA	sooooooooooooooooo	6-10-4	21	
802466	8012	8031 TCTGTCTTTGGAGCCCAAAT	sooooooooooooooooo	6-10-4	45	
802467	8186	8205 CTGCGATCCCCATTCCAGTT	sooooooooooooooooo	6-10-4	46	
802468	7990	8009 GCCTTACTCTAGGACCAAGA	sooooooooooooooooo	7-10-3	21	
802469	8012	8031 TCTGTCTTTGGAGCCCAAAT	sooooooooooooooooo	7-10-3	45	
802470	8186	8205 CTGCGATCCCCATTCCAGTT	sooooooooooooooooo	7-10-3	46	
802471	7992	8009 GCCTTACTCTAGGACCAA	sooooooooooooooooo	3-8-7	47	
802472	8014	8031 TCTGTCTTTGGAGCCCAA	sooooooooooooooooo	3-8-7	48	
802473	7992	8009 GCCTTACTCTAGGACCAA	sooooooooooooooooo	4-8-6	47	
802474	8014	8031 TCTGTCTTTGGAGCCCAA	sooooooooooooooooo	4-8-6	48	
802475	7992	8009 GCCTTACTCTAGGACCAA	sooooooooooooooooo	6-8-4	47	

TABLE 6-continued

Antisense oligonucleotides targeting human C9ORF72					
Isis No.	Start Site	Stop Site	Sequence	Linkage	SEQ ID NO
802476	8014	8031	TCTGTCTTTGGAGCCCAA	sooooooossssssooss	48
802477	7992	8009	GCCTTACTCTAGGACCAA	sooooooossssssooss	47
802478	8014	8031	TCTGTCTTTGGAGCCCAA	sooooooossssssooss	48
806673	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooooooooooss	49
806674	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooooooooooss	50
806675	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooooooooooss	51
806676	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooooooooooss	49
806677	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooooooooooss	50
806678	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooooooooooss	51
806679	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooossssssooss	49
806680	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooossssssooss	50
806681	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooossssssooss	51
806682	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooossssssooss	49
806683	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooossssssooss	50
806684	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooossssssooss	51
806685	1371	1388	GTTAATCTTTATCAGGTC	sooooooooooooooss	52
806686	1372	1389	GGTTAATCTTTATCAGGT	sooooooooooooooss	53
806687	1373	1390	TGGTTAATCTTTATCAGG	sooooooooooooooss	54
806688	1440	1457	CTAGCGCGCGACTCCTGA	sooooooooooooooss	55
806689	1371	1388	GTTAATCTTTATCAGGTC	sooooooossssssooss	52
806690	1372	1389	GGTTAATCTTTATCAGGT	sooooooossssssooss	53
806691	1373	1390	TGGTTAATCTTTATCAGG	sooooooossssssooss	54
806692	1440	1457	CTAGCGCGCGACTCCTGA	sooooooossssssooss	55

Example 2: Tolerability of Antisense Oligonucleotides Targeting Human C9ORF72 in Mice

Antisense oligonucleotides described above were tested in mice to assess tolerability of the oligonucleotides. Wild type C57/Bl6 mice each received a single ICV dose of 700 µg of an antisense oligonucleotide listed in the table below. Each treatment group consisted of 4 mice. At 3 hours post injection, each mouse was evaluated according to 7 different criteria. The 7 criteria are (1) the mouse was bright, alert, and responsive; (2) the mouse was standing or hunched without stimuli; (3) the mouse showed any movement without stimuli; (4) the mouse demonstrated forward movement after it was lifted; (5) the mouse demonstrated any movement after it was lifted; (6) the mouse responded to a tail pinch; (7) regular breathing. For each of the 7 different criteria, each mouse was given a sub-score of 0 if it met the criteria or 1 if it did not. After all of the 7 criteria were evaluated, the sub-scores were summed for each mouse and then averaged for each group. For example, if a mouse was

dose, and met all other criteria, it would get a summed score of 0. If another mouse was not bright, alert, and responsive 3 hours after the 700 µg ICV dose but met all other criteria, it would receive a score of 1. The results are presented as the average score for each treatment group.

TABLE 7

Acute tolerability scores	
Isis No.	Score
791656	3.25
791657	4.25
791658	3.50
791659	2.00
791660	4.00
791661	4.50
791662	5.25
791663	6.00
791664	3.00
791665	5.75
801274	1.75
801275	3.75
801276	1.25

TABLE 7-continued

Acute tolerability scores	
Isis No.	Score
801277	1.50
801278	1.75
801279	1.25
801280	6.00
801281	0.00
801282	1.25
801283	1.25
801284	3.00
801285	1.00
801286	1.00
801287	1.50
801288	2.50
801289	7.00
801290	6.00
801291	3.00
801292	2.00
801293	1.00
801294	3.25
801295	4.50
801296	6.25
801297	4.75
801298	5.50
801299	6.25
801300	5.00
801301	6.00
801302	6.50
801303	4.00
801304	5.25
801305	6.00
801306	6.00
801307	5.00
801308	6.00
801309	7.00
801310	3.50
801311	5.50
801312	2.50
801313	5.25
801314	4.50
801315	4.00
801316	2.00
802459	2.00
802460	6.75
802461	1.75
802462	5.75
802463	6.75
802464	1.75
802465	2.25
802466	4.25
802467	0.25
802468	3.25
802469	2.00
802470	0.25
802471	1.25
802472	4.00
802473	0.25
802474	5.25
802475	1.00
802476	5.50
802477	2.50
802478	6.25
806673	0.00
806674	0.25
806675	0.00
806676	0.00
806677	1.00
806678	0.00
806679	1.50
806680	1.00
806681	0.00
806682	5.75
806683	3.75
806684	2.25
806685	1.00
806686	1.00
806687	3.25
806688	3.25

TABLE 7-continued

Acute tolerability scores	
Isis No.	Score
806689	3.00
806690	1.25
806691	6.25

Example 3: Tolerability of Oligonucleotides from WO 2014/062691

Oligonucleotides described in WO 2014/062691 were tested in an acute tolerability study in mice. Groups of 3 wild type C57/Bl6 mice were treated and analyzed as described in Example 2. The tested oligonucleotides include those listed in the table below, which are 5-10-5 MOE gapmers with a full phosphorothioate backbone and each cytosine is a 5-methylcytosine. The start and stop sites on SEQ ID NO: 2 that each oligonucleotide is targeted to are shown. The results are presented as the average score for each treatment group in the table below. These results demonstrate that ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, ISIS 577083, and ISIS 577056 were poorly tolerated.

TABLE 8

Acute tolerability scores 3h after treatment with antisense oligonucleotides from WO 2014/062691					
Isis No.	Start site	Stop site	Sequence	Score	SEQ ID NO
576816	7990	8009	GCCTTACTCTAGGACCAAGA	7.00	21
576974	28251	28270	GGGACACTACAAGGTAGTAT	5.67	56
577061	1406	1425	TACAGGCTGCGGTTGTTTCC	7.00	57
577065	1446	1465	CCCGGCCCTAGCGCGCGAC	6.00	58
577083	3452	3471	GGTAACTTCAAACCTCTGGG	7.00	59

TABLE 9

Acute tolerability scores 3h after treatment with antisense oligonucleotides from WO 2014/062691					
Isis No.	Start site	Stop site	Sequence	Score	SEQ ID NO
577056	1366	1385	AATCTTTATCAGGTCTTTTC	6.5	60

Example 4: Antisense Inhibition of a Human C9ORF72 mRNA Variant in HepG2 Cells

Antisense oligonucleotides described above are tested for their effects on C9ORF72 mRNA in vitro. Cultured HepG2 cells at a density of 20,000 cells per well are electroporated with an antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Human primer probe set RTS3905 (forward primer sequence GGGTCTAGCAAGAGCAGGTG, designated herein as SEQ ID NO: 12; reverse primer sequence GTCTTGGCAACAGCTGGAGAT, designated herein as

SEQ ID NO: 13; probe sequence TGATGTCGACTCTTT-GCCCACCGC, designated herein as SEQ ID NO: 14—a TAQ-man primer probe set) are used. RTS3905 detects an mRNA variant (e.g. NM_001256054.1) processed from a pre-mRNA variant containing the hexanucleotide repeat. The mRNA variant processed from a pre-mRNA variant containing the hexanucleotide repeat is herein the “C9ORF72 pathogenic associated mRNA variant.” A pre-mRNA contains the hexanucleotide repeat when transcription of the pre-mRNA begins in the region from the start site of exon 1A to the start site of exon 1B (generally nucleotides 1107 to 1520 of the genomic sequence: SEQ ID NO: 2, the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 2756500. Therefore, oligonucleotides designed in this region selectively target the pre-mRNA variant containing the hexanucleotide repeat. RTS3905 measures an mRNA product (i.e. the C9ORF72 pathogenic associated mRNA variant) of the pre-mRNA variant containing the hexanucleotide repeat and, therefore, measures the reduction of the pre-mRNA variant containing the hexanucleotide repeat. The levels of the C9ORF72 pathogenic associated mRNA variant are normalized to the total RNA content of the cell, as measured by RIBOGREEN®, then the normalized mRNA variant levels are compared to those of cells that were not treated with antisense oligonucleotide.

Example 5: Dose-Dependent Antisense Inhibition of a Human C9ORF72 mRNA Variant

Antisense oligonucleotides described above are tested at various doses in HepG2 cells. Cells are plated at a density of 20,000 cells per well and electroporated with antisense oligonucleotide. After a treatment period of approximately 16 hours or 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Human C9ORF72 primer probe set RTS3905 is used to measure the C9ORF72 pathogenic associated mRNA variant. The levels of the C9ORF72 pathogenic associated mRNA variant are adjusted according to total RNA content, as measured by RIBOGREEN. The half maximal inhibitory concentration (IC_{50}) of each oligonucleotide is calculated based on the inhibition of mRNA variant levels observed at each individual dose of antisense oligonucleotide.

Example 6: Antisense Inhibition of C9ORF72 by Human-Rhesus Cross-Reactive Antisense Oligonucleotides in LLC-MK2 Cells

Antisense oligonucleotides described above that are fully cross-reactive with a rhesus C9ORF72 nucleic acid are tested for their effects on rhesus C9ORF72 mRNA in vitro. Cultured rhesus LLC-MK2 cells at a density of 20,000 cells per well are electroporated with antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Primer probe set RTS3750 (forward sequence TGTGACAGTTGGAAT-GCAGTGA, designated herein as SEQ ID NO: 15; reverse sequence GCCACTTAAAGCAATCTCTGTCTTG, designated herein as SEQ ID NO: 16; probe sequence TCGACTCTTTGCCACCGCCA, designated herein as SEQ ID NO: 17—a TAQ-man primer probe set) is used to measure total C9ORF72 mRNA levels. RTS3750 targets exon 2 of the mRNA transcripts and, therefore, measures total mRNA transcripts. C9ORF72 mRNA levels are

adjusted according to total RNA content, as measured by RIBOGREEN®, then the normalized mRNA variant levels are compared to those of cells that were not treated with antisense oligonucleotide.

Example 7: Dose-Dependent Antisense Inhibition of Human C9ORF72 mRNA in LLC-MK2

Antisense oligonucleotides described above are tested at various doses in LLC-MK2 cells. Cells are plated at a density of 20,000 cells per well and electroporated with antisense oligonucleotide. After a treatment period of approximately 16 hours or 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Primer probe set RTS3750 is used to measure total C9ORF72 mRNA levels. C9ORF72 mRNA levels are adjusted according to total RNA content, as measured by RIBOGREEN®, then the normalized mRNA variant levels are compared to those of cells that were not treated with antisense oligonucleotide.

Example 8: Antisense Inhibition of Human C9ORF72 mRNA in a Transgenic Mouse Model

Antisense oligonucleotides described above are tested in two BAC transgenic mouse lines, designated herein as C9B41 and C9B183, that each express a truncated human C9ORF72 gene comprising exons 1-5. The truncated human C9ORF72 genes of the C9B41 and C9B183 mouse lines comprise 110 and 450 hexanucleotide repeats, respectively. Each mouse in each treatment group receives 350 μ g of an antisense oligonucleotide, then expression levels of human C9ORF72 RNA are analyzed by RT-PCR. Dose responses are also performed.

Example 9: Antisense Inhibition of C9ORF72 mRNA in Patient Fibroblasts

Antisense oligonucleotides described above were tested for their effects on C9ORF72 mRNA in vitro. The antisense oligonucleotides listed in the table below were added to a plate before patient fibroblasts F09-152 were added at a density of 20,000 cells per well. The final concentrations of the antisense oligonucleotides after the addition of the cells are listed in the table below. After 30 seconds of shaking, the cells were electroporated then transferred to a Primaria coated culture plate. After 16 hours, RNA was isolated from the cells, and levels of total C9ORF72 mRNA (i.e., mRNA starting from exon 1A and mRNA starting from exon 1B) and levels of the pathogenic associated mRNA (see Example 4) were measured by RT-qPCR. Human primer probe sets RTS 3750 (see Example 6) and RTS3905 (see Example 4) were used to detect the total C9ORF72 mRNA and C9ORF72 pathogenic associated mRNA, respectively. The levels of the C9ORF72 total mRNA and C9ORF72 pathogenic associated mRNA were normalized to the total RNA content of the cell, as measured by RIBOGREEN®, then the normalized mRNA levels were compared to those of cells that were not treated with antisense oligonucleotide. The results are shown in the table below. An entry of “nd” means not determined. IC_{50} values listed as “nd” were not determined, because the target was not inhibited sufficiently to determine an IC_{50} . The results show that all of the oligonucleotides listed below inhibited the pathogenic associated C9ORF72 mRNA variant. Oligonucleotides that do not specifically target the repeat variant pre-mRNA inhibited both the pathogenic associated C9ORF72 mRNA and total

C9ORF72 mRNA. Oligonucleotides that specifically target the repeat variant pre-mRNA selectively inhibited the pathogenic associated C9ORF72 mRNA.

TABLE 10

C9ORF72 mRNA levels following antisense inhibition in patient fibroblasts					
Isis No.	Concentration (μM)	Total C9ORF72 mRNA		Pathogenic associated C9ORF72 mRNA	
		Level (% UTC)	IC ₅₀ (μM)	Level (% UTC)	IC ₅₀ (μM)
791658	0.12	87	nd	26	<0.1
	0.60	80		9	
	3.00	87		nd	
	15.00	105		4	
791664	0.12	85	nd	13	<0.1
	0.60	101		7	
	3.00	125		4	
	15.00	114		3	
801278	0.12	91	nd	19	<0.1
	0.60	71		7	
	3.00	85		nd	
	15.00	90		nd	
801279	0.12	98	nd	57	0.12
	0.60	77		27	
	3.00	77		3	
	15.00	86		5	
801282	0.12	100	nd	51	0.08
	0.60	84		12	
	3.00	97		6	
	15.00	190		nd	
801283	0.12	87	nd	40	<0.1
	0.60	75		8	
	3.00	94		4	
	15.00	111		4	
801285	0.12	103	nd	35	<0.1
	0.60	78		23	
	3.00	75		7	
	15.00	79		6	
801286	0.12	94	nd	33	<0.1
	0.60	85		14	
	3.00	97		5	
	15.00	88		3	
801287	0.12	85	nd	31	<0.1
	0.60	76		11	
	3.00	77		9	
	15.00	87		2	
801288	0.12	77	nd	13	<0.1
	0.60	86		11	
	3.00	123		5	
	15.00	177		3	
801292	0.12	69	nd	29	<0.1
	0.60	71		12	
	3.00	82		3	
	15.00	71		nd	
801293	0.12	77	nd	51	0.09
	0.60	70		10	
	3.00	74		3	
	15.00	81		1	
801294	0.12	75	nd	27	<0.1
	0.60	67		9	
	3.00	107		7	
	15.00	146		2	
801316	0.12	73	nd	35	<0.1
	0.60	68		7	
	3.00	65		1	
	15.00	78		1	
802459	0.12	58	0.10	75	0.18
	0.60	15		6	
	3.00	2		11	
	15.00	1		6	
802464	0.12	81	0.39	69	0.21
	0.60	31		23	
	3.00	6		7	
	15.00	2		3	

TABLE 10-continued

C9ORF72 mRNA levels following antisense inhibition in patient fibroblasts					
Isis No.	Concentration (μM)	Total C9ORF72 mRNA		Pathogenic associated C9ORF72 mRNA	
		Level (% UTC)	IC ₅₀ (μM)	Level (% UTC)	IC ₅₀ (μM)
802465	0.12	40	<0.1	35	<0.1
	0.60	7		7	
	3.00	2		nd	
	15.00	1		nd	
802468	0.12	52	0.10	40	<0.1
	0.60	15		13	
	3.00	3		7	
	15.00	2		nd	
802469	0.12	69	0.17	57	~0.5
	0.60	16		9	
	3.00	4		nd	
	15.00	1		nd	
802471	0.12	71	0.27	54	0.12
	0.60	29		20	
	3.00	6		6	
	15.00	7		nd	
802473	0.12	63	0.18	48	<0.1
	0.60	27		14	
	3.00	8		11	
	15.00	4		nd	
802477	0.12	54	0.12	32	<0.1
	0.60	16		12	
	3.00	4		11	
	15.00	3		3	
806676	0.12	66	nd	20	<0.1
	0.60	66		4	
	3.00	76		2	
	15.00	64		nd	
806679	0.12	71	nd	23	<0.1
	0.60	68		2	
	3.00	78		1	
	15.00	84		1	
806680	0.12	89	nd	41	<0.1
	0.60	76		13	
	3.00	65		7	
	15.00	84		nd	
806690	0.12	99	nd	44	<0.1
	0.60	88		17	
	3.00	85		2	
	15.00	77		1	

Example 10: Antisense Inhibition of Human C9ORF72 mRNA in a Transgenic Mouse Model

Antisense oligonucleotides described above were tested in a BAC transgenic mouse line, C9B41 (see Example 8), that expresses a human C9ORF72 gene comprising the promoter region through exon 5 and 110 hexanucleotide repeats. Each treatment group consisted of 2-3 mice. Each mouse received a single ICVB of 350 μg of an antisense oligonucleotide listed in the tables below or PBS. Two weeks later, the mice were euthanized, and expression levels of the human pathogenic associated C9ORF72 mRNA variant and/or total human C9ORF72 mRNA were analyzed by RT-qPCR as described in Example 9. Analysis of the pathogenic associated C9ORF72 variant mRNA levels was not completed for the oligonucleotides that do not specifically target the C9ORF72 repeat variant pre-mRNA. The results in the tables below show the average percent normalized human C9ORF72 mRNA levels relative to the normalized average for the PBS treated group.

TABLE 11

Human C9ORF72 mRNA levels following antisense inhibition in transgenic mice				
Spinal Cord (% PBS treated)			Cortex (% PBS treated)	
Isis No.	Pathogenic variant	Total	Pathogenic variant	Total
791658	6	69	12	47
791659	18	72	27	57
791664	13	55	15	39
801274	49	65	27	37
801276	10	47	12	29
801277	9	44	10	28
801278	8	39	10	27
801279	33	55	40	47
801281	16	48	17	32
801282	18	49	31	42
801283	15	46	17	31
801285	29	51	24	36
801286	37	61	37	47
801287	13	47	30	40
801288	18	52	39	48
801292	25	49	33	40
801293	20	50	29	38
801310	70	80	68	70
801312	43	61	46	55
801315	39	59	41	51
801316	27	56	50	58
806673	38	66	66	66
806674	78	88	98	96
806675	67	86	81	85
806676	29	64	52	58
806677	61	77	69	68
806678	83	96	95	102
806679	24	63	26	44
806680	29	62	41	56
806681	39	68	36	54
806684	44	69	62	64
806685	69	87	56	58
806686	59	75	57	68
806687	74	88	77	80
806688	34	62	42	51
806689	67	92	87	88
806690	28	61	57	68
806692	45	64	44	50

TABLE 12

Human C9ORF72 mRNA levels following antisense inhibition in transgenic mice		
Total human C9ORF72 mRNA (% PBS)		
Isis No.	Spinal Cord	Cortex
802459	28	30
802461	22	19
802464	30	28
802465	25	25
802467	13	14
802468	24	26
802469	24	20
802470	18	20
802471	36	44
802473	28	38
802475	15	15
802477	14	15

Example 11: Dose Dependent Antisense Inhibition of Human C9ORF72 mRNA in a Transgenic Mouse Model

Antisense oligonucleotides described above were tested in two BAC transgenic mouse lines, C9B41 and C9B183, that each express a truncated human C9ORF72 gene comprising exons 1-5. The truncated human C9ORF72 genes of the C9B41 and C9B183 mouse lines comprise 110 and 450 hexanucleotide repeats, respectively (see Example 8). Each

treatment group consisted of 2-4 mice. Each mouse received a single ICVB of 30 µg, 100 µg, 300 µg, or 700 µg of an antisense oligonucleotide as listed in the tables below or PBS. Two weeks later, the mice were euthanized, and expression levels of the human pathogenic associated C9ORF72 mRNA variant and/or total human C9ORF72 mRNA were analyzed by RT-qPCR as described in Example 9. The results in the tables below show the average percent normalized human C9ORF72 mRNA levels relative to the normalized average for the PBS treated group. A value of 100 or greater means the antisense oligonucleotide did not reduce mRNA or increased the amount of mRNA.

TABLE 13

Human C9ORF72 mRNA levels following dose dependnt antisense inhibition in C9B41 transgenic mice			
Isis No.	Concentration (µg)	Spinal Cord (% PBS treated) Total C9	Cortex (% PBS treated) Total C9
802459	30	84	85
	100	63	61
	300	33	26
	700	29	17
802473	30	74	90
	100	60	77
	300	37	39
	700	28	26

TABLE 14

Human C9ORF72 mRNA levels following dose dependnt antisense inhibition in C9B183 transgenic mice			
Isis No.	Concentration (µg)	Spinal Cord (% PBS treated) Pathogenic variant	Cortex (% PBS treated) Pathogenic variant
801287	30	76	10
	100	37	75
	300	21	32
	700	11	15
806679	30	72	13
	100	59	100
	300	21	61
	700	10	16
806680	30	94	118
	100	72	97
	300	34	89
	700	30	56
806690	30	52	125
	100	60	131
	300	43	119
	700	19	49

Example 12: Tolerability of Antisense Oligonucleotides Targeting Human C9Orf72in Mice

Wild type C57/Bl6 mice each received a single ICV dose of 700 µg of an antisense oligonucleotide listed in the table below, as described in Example 2. Each treatment group consisted of 4 mice. At 8 weeks post-injection, mice were evaluated according to 7 different criteria. The criteria are (1) the mouse was bright, alert, and responsive; (2) the mouse was standing or hunched without stimuli; (3) the mouse showed any movement without stimuli; (4) the mouse demonstrated forward movement after it was lifted; (5) the mouse demonstrated any movement after it was lifted; (6) the mouse responded to tail pinching; (7) regular breathing. For each of the 7 criteria, a mouse was given a subscore of 0 if it met the criteria and 1 if it did not. After all 7 criteria were evaluated, the scores were summed for

each mouse and averaged within each treatment group. The results are presented in the table below.

Animals were sacrificed at 8 weeks. The cortex and spinal cord were collected from each animal, and RT-PCR was performed. Expression levels of allograft inflammatory factor (AIF1) were determined as a measure of inflammation. Expression levels of glial fibrillary acidic protein (GFAP) were also determined as a measure of glial cell activation. Results were normalized to Gpadh and are presented relative to PBS control (1.0) in the table below. "N.D." indicates there was no data because the experiment was not performed. An asterisk indicates that the corresponding result is the average of 1-3 mice.

TABLE 15

Tolerability of antisense oligonucleotides targeting C9Orf72 in mice						
ISIS No.	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cerebellum)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
791656	3.5	1.1*	1.3*	1.0*	1.3*	22
791657	3.5	2.3*	1.0*	1.4*	0.8*	22
791658	0.0	1.9	1.2	1.4	1.3	22
791659	0.0	1.8	1.3	1.5	2.5	22
791660	1.8	2.5*	1.4*	1.9*	0.8*	23
791661	1.8	1.7*	1.3*	1.5*	1.1*	24
791662	5.3	1.1*	1.2*	1.1*	0.8*	23
791663	7.0	N.D.	N.D.	N.D.	N.D.	24
791664	0.0	1.6	1.6	1.2	0.9	23
791665	3.5	1.9*	1.5*	1.9*	0.9*	24
801274	0.0	1.2	1.3	1.3	1.1	25
801275	3.5	1.4*	1.7*	1.6*	2.1*	26
801276	0.0	13.1	2.6	3.7	1.2	27
801277	0.0	2.9	1.4	2.3	1.5	28
801278	0.0	2.0	1.3	1.8	1.1	29
801279	0.0	1.2	1.4	1.3	1.0	25
801280	5.3	1.5*	1.5*	1.4*	1.2*	26
801281	0.0	3.6	2.2	2.4	1.6	27
801282	0.0	1.2	1.2	1.2	1.0	28
801283	0.0	1.4	1.4	1.0	0.9	29
801284	1.8	1.2*	1.5*	0.9*	1.0*	30
801285	0.0	1.2	1.2	1.1	0.8	31
801286	0.0	1.1	1.2	1.0	1.1	32
801287	0.0	1.3	1.2	1.1	1.4	33
801288	0.0	1.3	1.3	1.1	0.9	34
801289	7.0	N.D.	N.D.	N.D.	N.D.	35
801290	3.5	1.2*	1.3*	1.0*	1.1*	30
801291	0.0	2.2	2.2	1.5	1.1	31
801292	0.0	1.3	1.3	1.2	0.9	32
801293	0.0	1.6	1.6	1.3	1.5	33
801294	0.0	1.2	1.2	1.0	1.1	34
801295	1.8	1.4*	1.4*	1.0*	0.9*	35
801296	7.0	N.D.	N.D.	N.D.	N.D.	36
801297	3.5	1.0*	1.0*	0.9*	0.8*	37
801307	5.3	1.2*	0.9*	1.1*	0.8*	41
801308	5.3	1.2*	1.2*	0.9*	1.1*	42
801309	7.0	N.D.	N.D.	N.D.	N.D.	43
801310	1.8	1.2*	1.2*	1.0*	1.0*	44
801311	7.0	N.D.	N.D.	N.D.	N.D.	41
801312	0.0	2.0	1.3	1.0	1.4	42
801313	3.5	1.7*	1.1*	0.9*	0.9*	43
801314	0.0	1.9	1.2	1.0	2.9	44
801315	1.8	1.4*	1.3*	1.0*	1.1*	36
801316	0.0	1.5	1.4	1.0	1.2	37
801298	3.5	1.4*	1.0*	1.1*	1.3*	38
801299	5.3	1.2*	1.2*	1.0*	1.0*	39
801300	3.5	1.2*	1.1*	0.9*	1.0*	40
801301	7.0	N.D.	N.D.	N.D.	N.D.	38
801302	5.3	1.0*	1.1*	0.8*	0.8*	39
801303	1.8	1.1*	1.0*	1.0*	1.1*	40
801304	4.8	14.3*	5.4*	2.2*	3.5*	38
801305	5.3	3.6*	1.7*	1.6*	1.6*	39
801306	0.0	4.7	1.8	1.8	1.8	40
806673	0.0	1.1	1.0	1.0	9.1	49
806674	0.0	1.0	1.0	0.8	8.2	50
806675	0.0	1.0	1.0	0.8	10.3	51
806676	0.0	1.0	1.0	0.8	1.5	49
806677	0.0	0.9	1.0	0.9	1.5	50

TABLE 15-continued

Tolerability of antisense oligonucleotides targeting C9Orf72 in mice						
ISIS No.	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cerebellum)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
806678	0.0	1.0	0.9	1.1	1.2	51
806679	0.0	1.2	1.1	1.1	1.3	49
806680	0.0	1.0	1.1	0.8	1.0	50
806681	0.0	1.0	1.0	0.9	1.4	51
806682	1.8	1.4*	1.2*	1.0*	1.2*	49
806683	0.0	1.2	1.2	0.9	1.2	50
806684	0.0	1.2	1.1	0.9	2.0	51
806685	0.0	1.0	0.9	0.8	1.5	52
806686	0.0	0.9	0.9	0.8	1.1	53
806687	0.0	1.0	1.0	0.8	1.0	54
806688	1.8	1.1*	1.0*	0.8*	1.8*	55
806689	0.0	1.1	1.0	0.8	1.3	52
806690	0.0	1.0	1.1	0.9	1.1	53
806691	5.3	N.D.	N.D.	N.D.	N.D.	54
802459	0.0	0.9	0.9	0.9	1.2	21
802460	5.3	1.1*	1.1*	1.0*	1.4*	45
802461	3.5	1.4*	1.4*	0.9*	0.9*	46
802462	3.5	0.8*	0.9*	0.9*	1.1*	21
802463	7.0	N.D.	N.D.	N.D.	N.D.	45
802464	1.8	1.4*	1.2*	1.5*	1.4*	46
802465	0.0	0.9	1.1	1.1	1.6	21
802466	1.8	1.0*	1.0*	1.0*	1.4*	45
802467	1.3	4.9	2.4	2.7	2.8	46
802468	0.0	1.4	1.3	1.2	2.2	21
802469	0.0	1.0	1.0	1.0	1.2	45
802470	1.5	3.2	3.4	2.6	3.4	46
802471	0.0	0.9	0.9	0.9	1.2	47
802472	3.5	1.6*	1.4*	0.9*	1.5*	48
802473	0.0	1.0	0.9	0.9	1.3	47
802474	1.8	1.3*	1.2*	1.0*	1.8*	48
802475	7.0	N.D.	N.D.	N.D.	N.D.	47
802476	1.8	1*	1.2*	0.8*	1.8*	48
802477	0.0	1.9	1.7	1.3	2.1	47
802478	3.5	1.4*	1.1*	1.1*	1.3*	48

Example 13: Tolerability of Antisense Oligonucleotides Targeting Human C9Orf72 in Rats

40 Sprague Dawley rats were separated into groups of 4 or 6 rats. Each rat in each group of rats was administered a single 3 mg intrathecal (IT) dose of the oligonucleotide indicated in the table below. At 3 hours and at 8 weeks following the IT dose, the movement of 7 different parts of the body was evaluated for each rat. The 7 body parts are (1) the rat's tail; (2) the rat's posterior posture; (3) the rat's hind limbs; (4) the rat's hind paws; (5) the rat's forepaws; (6) the rat's anterior posture; (7) the rat's head. For each of the 7 different body parts, each rat was given a sub-score of 0 if the body part was moving or 1 if the body part was paralyzed. After each of the 50 7 body parts were evaluated, the sub-scores were summed for each rat and then averaged for each group. For example, if a rat's tail, head, and all other evaluated body parts were moving 3 hours after the 3 mg IT dose, it would get a summed score of 0. If another rat was not moving its tail 3 hours after the 3 mg IT dose but all other evaluated body parts were moving, it would receive a score of 1. Saline treated rats generally receive a score of 0. A score of at the top end of the range would be suggestive of acute toxicity. Results are presented in the table below as the average score for each treatment group.

60 Animals were sacrificed at 8 weeks. The cortex and spinal cord were collected from each animal, and RT-PCR was performed. Expression levels of AIF1 were determined as a measure of inflammation. Expression levels of (GFAP) were also determined as a measure of glial cell activation. An asterisk indicates that the corresponding result is the average of 2-3 mice. Results were normalized to Gapdh and are presented relative to PBS control (1.0) in the table below.

TABLE 16

Tolerability of antisense oligonucleotides targeting C9Orf72 in rats							
ISIS No.	Score 3 hours after injection	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cortex)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
801287	2.5	0.0	1.9	1.2	1.3	1.3	33
801288	4.0	3.5	1.3*	1.3*	1.2*	1.3*	34
806676	2.0	1.8	1.6*	1.5*	1.3*	2.5*	49
806679	1.2	0.0	1.4	1.3	1.3	1.8	49
806680	2.0	0.3	1.4	1.4	1.3	1.3	50
806690	1.3	0.0	1.5	1.4	1.3	2.0	53
802459	2.0	0.0	1.4	1.2	1.5	1.5	21
802473	2.0	2.3	1.8*	1.6*	1.7*	1.6*	47

Example 14: Tolerability of Antisense Oligonucleotides Targeting Human C9ORF72 in Non-Human Primates

Female cynomolgus monkeys (2-6 kg) were given 3 doses of 35 mg of antisense oligonucleotide on days 1, 14, and 28 via intrathecal bolus injection (1 mL slow bolus followed by 0.25 mL flush). Each treatment group contained four monkeys. Two weeks after the final dose, animals were sacrificed and RT-PCR was performed on various CNS tissues. Expression levels of AIF1 were determined as a measure of inflammation and expression levels of GFAP were determined as a measure of glial cell activation. Results were normalized to GADPH and are presented relative to PBS control (1.0) in the table below for ISIS No. 801287, 802459, and 806679.

TABLE 17

Tolerability of antisense oligonucleotides targeting C9ORF72 in cynomolgus monkeys						
Brain Region	AIF1			GFAP		
	801287	802459	806679	801287	802459	806679
Cervical spinal cord	1.0	0.9	0.9	1.1	1.0	1.1

TABLE 17-continued

Tolerability of antisense oligonucleotides targeting C9ORF72 in cynomolgus monkeys						
Brain Region	AIF1			GFAP		
	801287	802459	806679	801287	802459	806679
Thoracic spinal cord	1.1	0.9	0.9	1.2	1.2	1.2
Temporal cortex	1.0	1.0	1.5	1.2	1.1	1.7
Motor cortex	1.0	1.0	1.2	1.2	0.7	1.3
Lumbar spinal cord	1.2	0.9	0.9	1.3	0.9	1.1
Hippocampus	1.0	1.0	1.3	1.9	1.9	1.7
Frontal cortex	1.3	0.9	1.3	1.0	0.7	1.3

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 60

<210> SEQ ID NO 1
<211> LENGTH: 3339
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 1

acgtaacctacggtgtcccgctaggaaagagaggtgcgtcaaacagcgacaagtccgcc60
cacgtaaaagatgacgcttggtgtgcagcgtccctgctgcccggttgc ttctcttttg120
ggggcggggctagcaagagcaggtgtgggtttaggagatatctccgggacatttgata180
atgtgacagttggaatgcagtgatgtcgactctttgcccacgccatctc cagctgttgc240
caagacagagattgctttaagtggcaaatc acctttatta gcagctactt ttgcttactg300
ggacaatattcttggtcctagagtaaggca catttgggctccaaagacag aacagggtact360
tctcagtgatggagaaataacttttcttgc caaccacactctaaatggag aaatccttcg420
aaatgcagagagtggtgcta tagatgtaaa gttttttgtc ttgtctgaaa agggagtgat480
tattgtttcattaatctttgatggaaactggaatggggatcgcagcacat atggactatc540

-continued

aattatactt	ccacagacag	aacttagttt	ctacctccca	cttcatagag	tgtgtgttga	600
tagattaaca	catataatcc	ggaaaggaag	aatatggatg	cataaggaaa	gacaagaaaa	660
tgtccagaag	attatcttag	aaggcacaga	gagaatggaa	gatcagggtc	agagtattat	720
tccaatgctt	actggagaag	tgattcctgt	aatggaactg	ctttcatcta	tgaaatcaca	780
cagtgttcct	gaagaaatag	atatagctga	tacagtactc	aatgatgatg	atattgggtga	840
cagctgtcat	gaaggctttc	ttctcaatgc	catcagctca	cacttgcaaa	cctgtggctg	900
ttccgttgta	gtaggtagca	gtgcagagaa	agtaaataag	atagtcagaa	cattatgcct	960
ttttctgact	ccagcagaga	gaaaatgctc	caggttatgt	gaagcagaat	catcatttaa	1020
atatgagtca	gggctctttg	tacaaggcct	gctaaaggat	tcaactggaa	gctttgtgct	1080
gcctttccgg	caagtcatgt	atgctccata	ttccaccaca	cacatagatg	tggatgtcaa	1140
tactgtgaag	cagatgccac	cctgtcatga	acatatttat	aatcagcgtg	gatacatgag	1200
atccgagctg	acagccttct	ggagagccac	ttcagaagaa	gacatggctc	aggatacgat	1260
catctacact	gacgaaagct	ttactcctga	tttgaatatt	tttcaagatg	tcttacacag	1320
agacactcta	gtgaaagcct	tcttggatca	ggtctttcag	ctgaaacctg	gcttatctct	1380
cagaagtact	ttccttgcac	agtttctact	tgctcttcac	agaaaagcct	tgacactaat	1440
aaaatatata	gaagacgata	cgcagaaggg	aaaaaagccc	tttaaactct	ttcggaacct	1500
gaagatagac	cttgatttaa	cagcagaggg	cgatcttaac	ataataatgg	ctctgggtga	1560
gaaaattaaa	ccaggcctac	actcttttat	ctttggaaga	cctttctaca	ctagtgtgca	1620
agaacgagat	gttctaata	ctttttaaat	gtgtaactta	ataagcctat	tccatcacia	1680
tcatgatcgc	tggtaaagta	gctcagtggt	gtggggaaac	gttcccctgg	atcatactcc	1740
agaattctgc	tctcagcaat	tgcagttaag	taagttacac	tacagttctc	acaagagcct	1800
gtgaggggat	gtcaggtgca	tcattacatt	gggtgtctct	tttcctagat	ttatgctttt	1860
gggatacaga	cctatgttta	caatataata	aatattattg	ctatctttta	aagatataat	1920
aataggatgt	aaacttgacc	acaactactg	tttttttgaa	atacatgatt	catgggtttac	1980
atgtgtcaag	gtgaaatctg	agttggcttt	tacagatagt	tgactttcta	tcttttgcca	2040
ttctttggtg	tgtagaatta	ctgtaatact	tctgcaatca	actgaaaact	agagccttta	2100
aatgatttca	attccacaga	aagaaagtga	gcttgaacat	aggatgagct	ttagaaagaa	2160
aattgatcaa	gcagatgttt	aattggaatt	gattattaga	tcctactttg	tggatttagt	2220
ccctgggatt	cagtctgtag	aaatgtctaa	tagttctcta	tagtccttgt	tcttggtgaa	2280
ccacagttag	ggtgttttgt	ttattttatt	gttcttgcta	ttgttgatat	tctatgtagt	2340
tgagctctgt	aaaaggaaat	tgtattttat	gttttagtaa	ttgttgccaa	ctttttaaat	2400
taattttcat	tatttttgag	ccaaattgaa	atgtgcacct	cctgtgcctt	ttttctcctt	2460
agaaaatcta	attacttgga	acaagttcag	atttcaactgg	tcagtcattt	tcatcttggt	2520
ttctttctgc	taagtcttac	catgtacctg	ctttggcaat	cattgcaact	ctgagattat	2580
aaaatgcctt	agagaatata	ctaactaata	agatcttttt	ttcagaaaca	gaaaatagtt	2640
ccttgagtac	ttccttcttg	catttctgcc	tatgtttttg	aagttgttgc	tgtttgcctg	2700
caataggcta	taaggaatag	caggagaaat	tttactgaag	tgctgttttc	ctaggtgcta	2760
ctttggcaga	gctaagttat	cttttgtttt	cttaatgcgt	ttggaccatt	ttgctggcta	2820
taaaataact	gattaatata	attctaacac	aatgttgaca	ttgtagttac	acaaacacaa	2880
ataaatatth	tatttaaaat	tctggaagta	atataaaagg	gaaaatatat	ttataagaaa	2940

-continued

```

gggataaagg taatagagcc cttctgcccc ccacccacca aatttacaca acaaaatgac 3000
atgttcgaat gtgaaaggtc ataatagctt tcccatcatg aatcagaaaag atgtggacag 3060
cttgatgttt tagacaacca ctgaactaga tgactgttgt actgtagctc agtcatttaa 3120
aaaatatata aatactacct tgtagtgtcc catactgtgt tttttacatg gtagattctt 3180
atttaagtgc taactgggta ttttctttgg ctggtttatt gtactgttat acagaatgta 3240
agttgtacag tgaaataagt tattaaagca tgtgtaaaca ttgttatata tcttttctcc 3300
taaatggaga attttgaata aaatatattht gaaatthtg 3339

```

```

<210> SEQ ID NO 2
<211> LENGTH: 30001
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

```

```

<400> SEQUENCE: 2

```

```

caaagaaaag ggggaggttt tgttaaaaaa gagaaatgtt acatagtgtc ctttgagaaa 60
attcattggc actattaagg atctgaggag ctggtgagtt tcaactgggt agtgatgggtg 120
gtagataaaa ttagagctgc agcagggtcat ttagcaact attagataaa actggtctca 180
ggtcacaacg ggcagttgca gcagctggac ttggagagaa ttacactgtg ggagcagtgt 240
catttgctct aagtgttttt ctacccctta cccccactat tttagttggg tataaaaaga 300
atgacccaat ttgtatgatc aactttcaca aagcatagaa cagtaggaaa aggggtctgtt 360
tctgcagaag gtgtagacgt tgagagccat tttgtgtatt ttttctctcc tttcttctct 420
ggatgaatgat taaaacgttc tgtgtgattt ttagtgatga aaaagattaa atgctactca 480
ctgtagtaag tgccatctca cacttgcaga tcaaaaggca cacagttaa aaaacctttg 540
tttttttaca catctgagtg gtgtaaatgc tactcatctg tagtaagtgg aatctatata 600
cctgcagacc aaaagacgca aggtttcaaa aatctttgtg ttttttacac atcaaacaga 660
atggtacgtt tttcaaaagt taaaaaaaaa caactcatcc acatattgca actagcaaaa 720
atgacattcc ccagtgtgaa aatcatgctt gagagaattc ttacatgtaa aggcaaaatt 780
gcgatgactt tgcagggggc cgtgggattc ccgcccgcag tgccggagct gtcccctacc 840
agggtttgca gtggagttht gaatgcactt aacagtgtct tacggtaaaa acaaaatttc 900
atccaccaat tatgtgttga gcgcccactg cctaccaagc acaaacaaaa ccattcaaaa 960
ccacgaaatc gtcttcactt tctccagatc cagcagctc ccctattaag gttcgcacac 1020
gctattgcgc caacgctcct ccagagcggg tcttaagata aaagaacagg acaagttgcc 1080
ccgccccatt tgcgtagcct cgtgagaaaa cgtcatcgca catagaaaac agacagacgt 1140
aacctacggt gtcccgtag gaaagagagg tgcgtcaaac agcgacaagt tccgcccacg 1200
taaaagatga cgcttggtgt gtcagccgtc cctgctgcc ggttgcttct cttttggggg 1260
cggggtctag caagagcagg tgtgggttta ggaggtgtgt gttttgttt tttccacct 1320
ctctccccac tacttgcctt cacagtactc gctgaggggtg aacaagaaaa gacctgataa 1380
agattaacca gaagaaaaa aggagggaaa caaccgcagc ctgtagcaag ctctggaact 1440
caggagtcgc gcgctagggg ccggggccgg gccggggcg tggtcggggc gggccgggg 1500
gcggggccgg ggcggggctg cggttgcggt gcctgcgcc gcggcgggcg aggcgcaggc 1560
ggtggcgagt ggggtgagtga ggaggcggca tctggcggg tggctgtttg gggttcggt 1620
gccgggaaga ggcgcgggta gaagcggggg ctctcctcag agctcgacgc atttttactt 1680
tccctctcat ttctctgacc gaagctgggt gtcgggcttt cgctctagc gactgggtga 1740

```


-continued

attgcctgca	tccgggcccc	gggcttcccc	gcggcggcgg	cggcggcggc	ggcgcaggga	1800
caagggatgg	ggatctggcc	tcttcttgc	tttcccgccc	tcagtaccgg	agctgtctcc	1860
ttcccgggga	cccgtggga	gcgctgccgc	tgcgggctcg	agaaaaggga	gcctcgggta	1920
ctgagaggcc	tcgcctgggg	gaaggccgga	gggtgggcgg	cgcgcggctt	ctgcggacca	1980
agtcgggggtt	cgctaggaac	ccgagacggg	ccctgccggc	gaggagatca	tgcgggatga	2040
gatgggggtg	tggagacgcc	tgacacaattt	cagcccaagc	ttctagagag	tggtgatgac	2100
ttgcatatga	gggcagcaat	gcaagtcggg	gtgctcccca	ttctgtggga	catgacctgg	2160
ttgcttcaca	gctccgagat	gacacagact	tgcttaaagg	aagtgactat	tgtgacttgg	2220
gcatcacttg	actgatggta	atcagttgtc	taaagaagtg	cacagattac	atgtccgtgt	2280
gctcattggg	tctatctggc	cgcgttgaac	accaccaggc	tttgtattca	gaaacaggag	2340
ggaggtcctg	cactttccca	ggaggggtgg	ccctttcaga	tgcaatcgag	attgttaggc	2400
tctgggagag	tagttgcctg	gttgtggcag	ttggtaaatt	tctattcaaa	cagttgccat	2460
gcaccagttg	ttcacaacaa	gggtacgtaa	tctgtctggc	attacttcta	cttttgtaca	2520
aaggatcaaa	aaaaaaaaag	atactgttaa	gatatgattt	ttctcagact	ttgggaaact	2580
tttaacataa	tctgtgaata	tcacagaaac	aagactatca	tataggggat	attaataacc	2640
tggagtcaga	atacttgaaa	tacgggtgtca	tttgacacgg	gcattgttgt	caccacctct	2700
gccaaggcct	gccactttag	gaaaaccctg	aatcagttgg	aaactgctac	atgctgatag	2760
tacatctgaa	acaagaacga	gagtaattac	cacattccag	attgttcact	aagccagcat	2820
ttacctgctc	caggaaaaaa	ttacaagcac	cttatgaagt	tgataaaaata	ttttgtttgg	2880
ctatgttggc	actccacaat	ttgctttcag	agaaacaaag	taaaccaagg	aggacttctg	2940
tttttcaagt	ctgcctcgg	gttctattct	acgttaatta	gatagttccc	aggaggacta	3000
ggttagccta	cctattgtct	gagaaacttg	gaactgtgag	aaatggccag	atagtgatat	3060
gaacttcacc	ttccagtctt	ccctgatgtt	gaagattgag	aaagtgttgt	gaactttctg	3120
gtactgtaaa	cagttcactg	tccttgaaagt	ggtcctgggc	agctcctgtt	gtggaaagtg	3180
gacggtttag	gatcctgctt	ctctttgggc	tgggagaaaa	taaacagcat	ggttacaagt	3240
attgagagcc	aggttggaga	agggtggctta	cacctgtaat	gccagagctt	tgggaggcgg	3300
aggcaagagg	atcacttgaa	gccaggagtt	caagctcaac	ctgggcaacg	tagaccctgt	3360
ctctacaaaa	aattaaaaac	ttagccgggc	gtggtgatgt	gcacctgtag	tcctagctac	3420
ttgggaggct	gaggcaggag	ggtcatttga	gcccagaggt	ttgaagttac	cgagagctat	3480
gatcctgcca	gtgcattcca	gcctggatga	caaaacgaga	ccctgtctct	aaaaaacaag	3540
aagtgagggc	tttatgattg	tagaattttc	actacaatag	cagtggacca	accacctttc	3600
taaataccaa	tcagggaaga	gatggttgat	tttttaacag	acgtttaaaag	aaaagcaaa	3660
acctcaaaact	tagcactcta	ctaacagttt	tagcagatgt	taattaatgt	aatcatgtct	3720
gcatgtatgg	gattatttcc	agaaagtgta	ttgggaaacc	tctcatgaac	cctgtgagca	3780
agccaccgtc	tcactcaatt	tgaatcttgg	cttccctcaa	aagactggct	aatgtttggg	3840
aactctctgg	agtagacagc	actacatgta	cgtaagatag	gtacataaac	aactattggg	3900
tttgagctga	tttttttcag	ctgcatttgc	atgtatggat	ttttctcacc	aaagacgatg	3960
acttcaagta	ttagtaaaat	aattgtacag	ctctcctgat	tatacttctc	tgtgacattt	4020
catttcccag	gctatttctt	ttggtaggat	ttaaaactaa	gcaattcagt	atgatctttg	4080

-continued

tccttcattt	tctttcttat	tctttttggt	tggttggttg	tttggttttt	tcttgaggca	4140
gagtctctct	ctgtcgccca	ggctggagtg	cagtggcgcc	atctcagctc	attgcaacct	4200
ctgccacctc	cgggttcaag	agattctcct	gcctcagcct	cccagtagc	tgggattaca	4260
ggtgtccacc	accacacccg	gctaattttt	tgtattttta	gtagaggtgg	ggtttcacca	4320
tggtggccag	gctggtcttg	agctcctgac	ctcaggtgat	ccacctgcct	cggcctacca	4380
aagagctggg	ataacaggtg	tgaccacca	tgcccgccc	atTTTTTTTT	tcttattctg	4440
ttaggagtg	gagtgtact	agcagtataa	tagttcaatt	ttcacaacgt	ggtaaaagtt	4500
tccctataat	tcaatcagat	tttgctccag	ggttcagttc	tgtttttaga	aatactttta	4560
ttttcagttt	aatgatgaaa	tattagagtt	gtaatatgct	ctttatgatt	atccaccttt	4620
ttaacctaaa	agaatgaaag	aaaaatatgt	ttgcaatata	atTTTtatgg	tgtatgttaa	4680
cttaattcat	tatgttggcc	tccagtttgc	tggtgttagt	tatgacagca	gtagtgtcat	4740
taccatttca	attcagatta	cattcctata	tttgatcatt	gtaaactgac	tgcttacatt	4800
gtattaaaaa	cagtggatat	tttaaagaag	ctgtacggct	tatatctagt	gctgtctctt	4860
aagactatta	aattgataca	acatatttaa	aagtaaatat	tacctaaatg	aatttttgaa	4920
attacaaata	cacgtgttaa	aactgtcggt	gtgttcaacc	atTTctgtac	atacttagag	4980
ttaaactggt	tgccaggctc	tgtatgcta	ctcataatat	gataaaagca	ctcatcta	5040
gctctgtaaa	tagaagtcag	tgctttccat	cagactgaac	tctcttgaca	agatgtggat	5100
gaaattcttt	aagtaaaatt	gtttactttg	tcatacatTT	acagatcaaa	tgtagctcc	5160
caaagcaatc	atatggcaaa	gataggtata	tcatagtttg	cctattagct	gctttgtatt	5220
gctattatta	taaatagact	tcacagtttt	agacttgctt	agggtgaaatt	gcaattcttt	5280
ttactttcag	tcttagataa	caagtcttca	attatagtac	aatcacacat	tgcttaggaa	5340
tgcatcatta	ggcgattttg	tcattatgca	aacatcatag	agtgtactta	cacaaaccta	5400
gatagtatag	cctttatgta	cctaggccgt	atggtatagt	ctggtgctcc	taggccacaa	5460
acctgtacaa	ctgttactgt	actgaatact	atagacagtt	gtaacacagt	ggtaaata	5520
tatctaaata	tatgcaaaca	gagaaaaggt	acagtaaaag	tatggtataa	aagataatgg	5580
tatacctgtg	taggccactt	accacgaatg	gagcttgacg	gactagaagt	tgctctgggt	5640
gagtcagtga	gtgagtgggt	aattaatgtg	aaggcctaga	acactgtaca	ccactgtaga	5700
ctataaacac	agtacgctga	agctacacca	aatttatctt	aacagtTTTT	cttcaataaa	5760
aaattataac	tttttaactt	tgtaaaactt	ttaatttttt	aactttttaa	atacttagct	5820
tgaaacacaa	atacattgta	tagctataca	aaaatatTTT	ttctttgtat	ccttattcta	5880
gaagctTTTT	tctattttct	atTTtaaat	ttttttttta	cttgttagtc	gtttttgtta	5940
aaaactaaaa	cacacacact	ttcacctagg	catagacagg	attaggatca	tcagtatcac	6000
tcccttcac	ctcactgcct	tccacctcca	catcttgctc	cactggaagg	tttttagggg	6060
caataacaca	catgtagctg	tcacctatga	taacagtgtc	ttctgttgaa	tacctcctga	6120
aggacttgcc	tgaggctgtt	ttacatttaa	cttaaaaaaa	aaaaaagtag	aaggagtgca	6180
ctctaaaata	acaataaaaag	gcatagtata	gtgaatacat	aaaccagcaa	tgtagtagtt	6240
tattatcaag	tggtgtacac	tgtaataatt	gtatgtgcta	tactttaaat	aacttgcaaa	6300
atagtactaa	gaccttatga	tggttacagt	gtcactaagg	caatagcata	ttttcaggct	6360
cattgtaatc	taatgggact	accatcatat	atgcagtcta	ccattgactg	aaacgttaca	6420
tggcacataa	ctgtatttgc	aagaatgatt	tgttttacat	taatatcaca	taggatgtac	6480

-continued

cttttttagag	tggtatgttt	atgtggatta	agatgtacaa	gttgagcaag	gggaccaaga	6540
gccctggggt	ctgtcttgga	tgtgagcggt	tatgttcttc	tctcatgtc	tgttttctca	6600
ttaaattcaa	aggcttgaac	gggccctatt	tagcccttct	gttttctacg	tgttctaaat	6660
aactaaagct	tttaattct	agccatttag	tgtagaactc	tctttgcagt	gatgaaatgc	6720
tgtattgggt	tcttggctag	catattaaat	atttttatct	ttgtcttgat	acttcaatgt	6780
cgttttaaac	atcaggatcg	ggcttcagta	ttctcataac	cagagagtcc	actgaggata	6840
caggactggt	tgcccathtt	ttgttatggc	tccagacttg	tggtatttcc	atgtcttttt	6900
tttttttttt	ttttttgacc	ttttagcggc	tttaaagtat	ttctgttggt	aggtgttgta	6960
ttacttttct	aagattactt	aacaaagcac	cacaaactga	gtggctttaa	acaacagcaa	7020
ttatttctct	cacaattcta	gaagctagaa	gtccgaaatc	aaagtgttga	caggggcatg	7080
atcttcaaga	gagaagactc	tttccttgcc	tcttcctggc	ttctgggtgg	taccagcaat	7140
cctgagtgtt	cctttcttgc	cttgtagttt	caacaatcca	gtatctgcct	tttgtcttca	7200
catggctgtc	taccatttgt	ctctgtgtct	ccaaatctct	ctccttataa	acacagcagt	7260
tattggatta	ggccccactc	taatccagta	tgaccccat	ttaacatgat	tacacttatt	7320
tctagataag	gtcacattca	cgtacaccaa	gggttaggaa	ttgaacatat	ctttttgggg	7380
gacacaattc	aaccacaag	tgtcagtctc	tagctgagcc	tttcccttcc	tgtttttctc	7440
cttttttagtt	gctatgggtt	aggggccaaa	tctccagtca	tactagaatt	gcacatggac	7500
tggatatttg	ggaatactgc	gggtctattc	tatgagcttt	agtatgtaac	atttaatatc	7560
agtgtaaaga	agcccttttt	taagttattt	ctttgaattt	ctaaatgtat	gccctgaata	7620
taagtaacaa	gttaccatgt	cttgtaaaat	gatcatatca	acaaacattt	aatgtgcacc	7680
tactgtgcta	gttgaatgtc	tttatcctga	taggagataa	caggattcca	catctttgac	7740
ttaagaggac	aaaccaaata	tgtctaaatc	atttgggggt	ttgatggata	tctttaaatt	7800
gctgaaccta	atcattgggt	tcatatgtca	ttgttttagat	atctccggag	catttggata	7860
atgtgacagt	tggaatgcag	tgatgtcgac	tctttgcccc	ccgccatctc	cagctgttgc	7920
caagacagag	attgctttta	gtggcaaatc	acctttatta	gcagctactt	ttgcttactg	7980
ggacaatatt	cttggtccta	gagtaaggca	catttgggct	ccaaagacag	aacagggtact	8040
tctcagtgat	ggagaaataa	cttttcttgc	caaccacact	ctaaatggag	aatccttcg	8100
aatgcagag	agtggtgcta	tagatgtaaa	gttttttgtc	ttgtctgaaa	aggagtgat	8160
tattgtttca	ttaatctttg	atggaaactg	gaatggggat	cgcagcacat	atggactatc	8220
aattatactt	ccacagacag	aacttagttt	ctacctccca	cttcatagag	tgtgtgttga	8280
tagattaaca	catataatcc	ggaaaggaag	aatatggatg	cataaggtaa	gtgatttttc	8340
agcttattaa	tcatgttaac	ctatctgttg	aaagcttatt	ttctggtaca	tataaatctt	8400
atttttttaa	ttatatgcag	tgaacatcaa	acaataaatg	ttatttat	tgcatttacc	8460
ctattagata	caaatacatc	tggctctgata	cctgtcatct	tcatattaac	tgtggaaggt	8520
acgaaatggg	agctccacat	tatagatgaa	aagctaaagc	ttagacaaat	aaagaaactt	8580
ttagaccctg	gattcttctt	gggagccttt	gactctaata	ccttttgttt	ccctttcatt	8640
gcacaattct	gtcttttgct	tactactatg	tgtaaagtata	acagttcaaa	gtaatagttt	8700
cataagctgt	tggctcatgta	gcctttggtc	tctttaacct	ctttgccaa	ttcccagggt	8760
cataaaatga	ggaggttgaa	tggaatgggt	cccaagagaa	ttccttttaa	tcttacagaa	8820

-continued

attattgttt	tcctaaatcc	tgtagttgaa	tatataatgc	tatttacatt	tcagtatagt	8880
tttgatgtat	ctaaagaaca	cattgaattc	tccttcctgt	gttccagttt	gataactaacc	8940
tgaaagtcca	ttaagcatta	ccagttttta	aaggcttttg	cccaatagta	aggaaaaata	9000
atatctttta	aaagaataat	tttttactat	gtttgcaggc	ttacttcctt	ttttctcaca	9060
ttatgaaact	cttaaaatca	ggagaatctt	ttaaacaaca	tcataatgtt	taatttgaaa	9120
agtgcagtc	attcttttcc	tttttgaaac	tatgcagatg	ttacattgac	tgttttctgt	9180
gaagttatct	ttttttcact	gcagaataaa	ggttgttttg	attttatttt	gtattgttta	9240
tgagaacatg	catttgttgg	gttaatttcc	taccctgcc	cccatttttt	ccctaaagta	9300
gaaagtattt	ttcttgtaaa	ctaaattact	acacaagaac	atgtctattg	aaaaataagc	9360
aagtatcaaa	atgttggtgg	ttgttttttt	aaataaattt	tctcttgctc	aggaaagaca	9420
agaaaatgtc	cagaagatta	tcttagaagg	cacagagaga	atggaagatc	aggatatatgc	9480
aaattgcata	ctgtcaaatg	tttttctcac	agcatgtatc	tgtataaggt	tgatggctac	9540
atttgtcaag	gccttggaag	catacgaata	agcctttaat	ggagctttta	tggagggtga	9600
cagaataaac	tggaggaaga	tttccatatc	ttaaacccaa	agagttaaata	cagtaaacaa	9660
aggaaaatag	taattgcatc	tacaaattaa	tatttgctcc	cttttttttt	ctgtttgccc	9720
agaataaatt	ttggataact	tgttcatagt	aaaaataaaa	aaaattgtct	ctgatatggt	9780
ctttaaggta	ctacttctcg	aacctttccc	tagaagtagc	tgtaacagaa	ggagagcata	9840
tgtacccctg	aggatctgt	ctgggggtga	ggcccaggtc	cacacaatat	ttcttctaag	9900
tcttatgttg	tatcgtaag	actcatgcaa	tttacatttt	attccataac	tatttttagta	9960
ttaaaatttg	tcagtgatat	ttcttaccct	ctcctctagg	aaaatgtgcc	atgtttatcc	10020
cttggttttg	aatgcccctc	aggaacagac	actaagagtt	tgagaagcat	ggttacaagg	10080
gtgtggcttc	ccctgcggaa	actaagtaca	gactatttca	ctgtaaagca	gagaagttct	10140
tttgaaggag	aatctccagt	gaagaaagag	ttcttcactt	ttacttccat	ttcctcttgt	10200
gggtgaccct	caatgctcct	tgtaaaactc	caatatttta	aacatggctg	ttttgccttt	10260
ctttgcttct	ttttagcatg	aatgagacag	atgatacttt	aaaaaagtaa	ttaaaaaaaa	10320
aaacttgatga	aaatacatgg	ccataataca	gaacccaata	caatgatctc	ctttaccaaa	10380
ttgttatggt	tgtacttttg	tagatagctt	tccaattcag	agacagttat	tctgtgtaaa	10440
ggtctgactt	aacaagaaaa	gatttccttt	tacccaaaga	atcccagtcc	ttatttgctg	10500
gtcaataagc	agggtcccca	ggaatggggg	aactttcagc	accctctaac	ccactagtta	10560
ttagtagact	aattaagtaa	acttatcgca	agttgaggaa	acttagaacc	aactaaaatt	10620
ctgcttttac	tgggattttg	ttttttcaaa	ccagaaacct	ttacttaagt	tgactactat	10680
taatgaattt	tggctctctt	tttaagtgtt	cttcttaaaa	atgttatctt	actgctgaga	10740
agttcaagtt	tgggaagtac	aaggaggaat	agaaacttaa	gagattttct	tttagagcct	10800
cttctgtatt	tagccctgta	ggattttttt	tttttttttt	ttttttggtg	ttgttgagct	10860
tcagtgaggc	tattcattca	cttatactga	taatgtctga	gatactgtga	atgaaatact	10920
atgtatgctt	aaacctaaga	ggaaatattt	tcccaaaatt	attcttcccg	aaaaggagga	10980
gttgcccttt	gattgagttc	ttgcaaactc	cacaacgact	ttattttgaa	caatactgtt	11040
tggggatgat	gcattagttt	gaaacaactt	cagttgtagc	tgtcatctga	taaaattgct	11100
tcacagggaa	ggaaatttaa	cacggatcta	gtcattattc	ttgttagatt	gaatgtgtga	11160
attgtaattg	taaacaggca	tgataattat	tacttttaaaa	actaaaaaca	gtgaatagtt	11220

-continued

```

agttgtggag gttactaaag gatgggtttt ttttaaataa aactttcagc attatgcaaa 11280
tgggcatatg gcttaggata aaacttccag aagtagcatc acattttaa tctcaagcaa 11340
cttaataata tggggctctg aaaaactggg taagggttact ccaaaaatgg ccctgggtct 11400
gacaaagatt ctaacttaaa gatgcttatg aagactttga gtaaaatcat ttcataaaat 11460
aagtgaggaa aaacaactag tattaaattc atcttaaata atgtatgatt taaaaaatat 11520
gtttagctaa aaatgcatag tcatttgaca atttcattta tatctcaaaa aatttactta 11580
accaagttgg tcacaaaact gatgagactg gtggtagtag tgaataaatg agggaccatc 11640
catatttgag acactttaca tttgtgatgt gttatactga attttcagtt tgattctata 11700
gactacaaat ttcaaaatta caatttcaag atgtaataag tagtaatatc ttgaaatagc 11760
tctaaagga atttttctgt tttattgatt cttaaataat atgtgctgat tttgatttgc 11820
atttgggtag attatacttt tatgagtatg gaggttaggt attgattcaa gttttcctta 11880
cctatttggg aaggatttca aagtcttttt gtgcttggtt ttcctcattt ttaaataatga 11940
aatatattga tgacctttaa caaatTTTTT ttatctcaaa ttttaaagga gatcttttct 12000
aaaagaggca tgatgactta atcattgcat gtaacagtaa acgataaacc aatgattcca 12060
tactctctaa agaataaaaag tgagctttag ggccgggcat ggtcagaaat ttgacaccaa 12120
cctggccaac atggcgaaac cccgtctcta ctaaaaatac aaaaatcagc cgggcatggt 12180
ggcggcacct atagtcccag ctacttggga ggatgagaca ggagagtcac ttgaacctgg 12240
gaggagaggt tgcagtgagc tgagatcacg ccattgcact ccagcctgag caatgaaagc 12300
aaaactocat ctcaaaaaaa aaaaaagaaa agaaagaata aaagtgagct ttggattgca 12360
tataaatcct ttagacatgt agtagacttg tttgatactg tgtttgaaca aattacgaag 12420
tattttcatc aaagaatgtt attgtttgat gttattttta ttttttattg cccagcttct 12480
ctcatattac gtgattttct tcacttcatg tcactttatt gtgcagggtc agagtattat 12540
tccaatgctt actggagaag tgattcctgt aatggaactg ctttcatcta tgaaatcaca 12600
cagtgttcc tgaagaaatag atgtaagttt aaatgagagc aattatacac tttatgagtt 12660
ttttgggggt atagtattat tatgtatatt attaatattc taattttaat agtaaggact 12720
ttgtcataca tactattcac atacagtatt agccacttta gcaaataagc acacacaaaa 12780
tcctggattt tatggcaaaa cagaggcatt tttgatcagt gatgacaaaa ttaaattcat 12840
tttgtttatt tcattacttt tataattcct aaaagtggga ggatcccagc tcttatagga 12900
gcaattaata tttaatgtag tgtcttttga aacaaaactg tgtgccaaag tagtaaccat 12960
taatggaagt ttactttag tcaaaaattt agtttcctta atcatttgtt gaggacgttt 13020
tgaatcacac actatgagtg ttaagagata cctttaggaa actattcttg ttgttttctg 13080
attttgtcat ttaggttagt ctctgattc tgacagctca gaagaggaag ttgttcttgt 13140
aaaaattggt taacctgctt gaccagcttt cacatttggt cttctgaagt ttatggtagt 13200
gcacagagat tgttttttgg ggagtcttga ttctcgaaa tgaaggcagt gtgttatatt 13260
gaatccagac ttccgaaaac ttgtatatta aaagtgttat ttcaacacta tggttacagc 13320
agactaattt ttttattttt tgatgcattt tagatagctg atacagtact caatgatgat 13380
gatattgggt acagctgtca tgaaggcttt cttctcaagt aagaattttt cttttcataa 13440
aagctggatg aagcagatac catcttatgc tcacctatga caagatttgg aagaaagaaa 13500
ataacagact gtctacttag attgttctag ggacattacg tatttgaact gttgcttaaa 13560

```


-continued

tttgtgttat	ttttcactca	ttatatattct	atatatatatt	ggtgttattc	catttgctat	13620
ttaaagaaac	cgagtttcca	tcccagacaa	gaaatcatgg	ccccttgctt	gattctgggt	13680
tcttgtttta	cttctcatta	aagctaacag	aatcctttca	tattaagttg	tactgtagat	13740
gaacttaagt	tatttaggcg	tagaacaaaa	ttattcatat	ttatactgat	ctttttccat	13800
ccagcagtgg	agtttagtac	ttaagagttt	gtgcccttaa	accagactcc	ctggattaat	13860
gctgtgtacc	cgtgggcaag	gtgcctgaat	tctctataca	cctatttcct	catctgtaaa	13920
atggcaataa	tagtaatagt	acctaatgtg	tagggttggt	ataagcattg	agtaagataa	13980
ataatataaa	gcacttagaa	cagtgcctgg	aacataaaaa	cacttaataa	tagctcatag	14040
ctaacatttc	ctatttacat	ttcttctaga	aatagccagt	atttgttgag	tgccacatg	14100
ttagttcctt	tactagttgc	ttacatgta	ttatcttata	ttctgtttta	aagtttcttc	14160
acagttacag	attttcatga	aattttactt	ttaataaaag	agaagtaaaa	gtataaagta	14220
ttcactttta	tggtcacagt	cttttccttt	aggctcatga	tggagtatca	gaggcatgag	14280
tgtgtttaac	ctaagagcct	taatggcttg	aatcagaagc	actttagtcc	tgtatctggt	14340
cagtgtcagc	ctttcataca	tcattttaaa	tcccatttga	ctttaagtaa	gtcacttaat	14400
ctctctacat	gtcaatttct	tcagctataa	aatgatggta	tttcaataaa	taaatacatt	14460
aattaaatga	tattatactg	actaattggg	ctgttttaag	gctcaataag	aaaatttctg	14520
tgaaaggctc	ctagaaaatg	taggttccta	tacaaataaa	agataacatt	gtgcttatag	14580
cttcggtggt	tatcatataa	agctattctg	agttatttga	agagctcacc	tacttttttt	14640
tgtttttagt	ttgttaaatt	gttttatagg	caatgttttt	aatctgtttt	ctttaactta	14700
cagtgccatc	agctcacact	tgcaaacctg	tggctgttcc	gttgtagtag	gtagcagtgc	14760
agagaaagta	aataaggtag	tttattttat	aatctagcaa	atgatttgac	tctttaagac	14820
tgatgatata	tcattggattg	tcattttaaat	ggtaggttgc	aattaaaatg	atctagtagt	14880
ataaggaggc	aatgtaatct	catcaaattg	ctaagacacc	ttgtggcaac	agtgagtttg	14940
aaataaactg	agtaagaatc	atztatcagt	ttattttgat	agctcggaaa	taccagtgtc	15000
agtagtgtat	aaatggtttt	gagaatatat	taaaatcaga	tatataaaaa	aaattactct	15060
tctatttccc	aatgttatct	ttaacaaatc	tgaagatagt	catgtacttt	tggtagtagt	15120
tccaaagaaa	tgttatttgt	ttattcatct	tgatttcatt	gtcttcgctt	tccttctaaa	15180
tctgtccctt	ctagggagct	attgggatta	agtggtcatt	gattattata	ctttattcag	15240
taatgtttct	gaccctttcc	ttcagtgcct	cttgagttaa	ttaaggatta	atgaacagtt	15300
acatttccaa	gcattagcta	ataaactaaa	ggattttgca	cttttcttca	ctgaccatta	15360
gtagaaaga	gttcagagat	aagtatgtgt	atctttcaat	ttcagcaaac	ctaatttttt	15420
aaaaaaagtt	ttacatagga	aatatgttgg	aaatgatact	ttacaaagat	attcataatt	15480
tttttttgta	atcagctact	ttgtatatatt	acatgagcct	taattttatat	ttctcatata	15540
accatttatg	agagcttagt	atacctgtgt	catttatattg	catctacgaa	ctagtgcctt	15600
tattccttct	gttacctcaa	acaggtggct	ttccatctgt	gatctccaaa	gccttagggt	15660
gcacagagtg	actgccgagc	tgctttatga	aggagaaaag	gctccatagt	tggagtgttt	15720
tttttttttt	ttttaaacat	ttttcccatc	ctccatcctc	ttgagggaga	atagcttacc	15780
ttttatcttg	ttttaatttg	agaaagaagt	tgccaccact	ctaggttgaa	aaccactcct	15840
ttaacataat	aactgtggat	atggtttgaa	tttcaagata	gttacatgcc	tttttatatt	15900
tcctaataga	gctgtaggtc	aaatattatt	agaatcagat	ttctaaatcc	cacccaatga	15960

-continued

cctgcttatt	ttaaatcaaa	ttcaataatt	aattctcttc	tttttgagg	atctggacat	16020
tctttgatat	ttcttacaac	gaatttcatg	tgtagacca	ctaaacagaa	gctataaaag	16080
ttgcatggtc	aaataagtct	gagaaagtct	gcagatgata	taattcacct	gaagagtcac	16140
agtatgtagc	caaagttaa	aggttttgag	atgccatata	gtaaatttac	caagcatttt	16200
ctaaatttat	ttgaccacag	aatccctatt	ttaagcaaca	actggtacat	cccatggatt	16260
ccaggtgact	aaagaatact	tatttcttag	gatatgtttt	attgataata	acaattaaaa	16320
tttcagatat	ctttcataag	caaatacgtg	gtctttttac	ttcatgtttt	aatgctaaaa	16380
tattttcttt	tatagatagt	cagaacatta	tgcttttttc	tgactccagc	agagagaaaa	16440
tgctccaggt	tatgtgaagc	agaatcatca	tttaaataatg	agtcagggct	ctttgtacaa	16500
ggcctgctaa	aggtatagtt	tctagttatc	acaagtgaac	ccacttttct	aaaatcattt	16560
ttgagactct	ttatagacaa	atcttaaata	ttagcattta	atgtatctca	tattgacatg	16620
cccagagact	gacttccttt	acacagttct	gcacatagac	tatatgtctt	atggatttat	16680
agttagtatc	atcagtgaac	caccatagaa	taccctttgt	gttccaggtg	ggteccctgtt	16740
cctacatgtc	tagcctcagg	actttttttt	ttttaacaca	tgcttaaatac	aggttgacaa	16800
tcaaaaaata	gatcatttct	ttttaactaa	atagatttga	attttattga	aaaaaaattt	16860
taaacatctt	taagaagctt	ataggattta	agcaattcct	atgtatgtgt	actaaaatat	16920
atatatttct	atatataata	tatattagaa	aaaaattgta	tttttctttt	atttgagtct	16980
actgtcaagg	agcaaaacag	agaaatgtaa	attagcaatt	atttataata	cttaaaggga	17040
agaaagtgtg	tcaccttggt	gaatctatta	ttgttatttc	aattatagtc	ccaagacgtg	17100
aagaaatagc	tttctaatg	gttatgtgat	tgtctcatag	tgactacttt	cttgaggatg	17160
tagccacggc	aaaatgaaat	aaaaaaattt	aaaaattggt	gcaaatacaa	gttatattag	17220
gcttttgtgc	atattcaata	atgtgctgct	atgaactcag	aatgatagta	tttaaataata	17280
gaaactagtt	aaaggaaacg	tagtttctat	ttgagttata	catatctgta	aattagaact	17340
tctctgttta	aaggcataat	aaagtgccta	atacttttgt	ttcctcagca	ccctctcatt	17400
taattatata	atatttagttc	tgaaagggac	ctataccaga	tgcttagagg	aaatttcaaa	17460
actatgatct	aatgaaaaaa	tatttaatat	ttctccatgc	aaatacaaat	catatagttt	17520
tccagaaaaat	acctttgaca	ttatacaaaag	atgattatca	cagcattata	atagtaaaaa	17580
aatggaaata	gcctctttct	tctgttctgt	tcatagcaca	gtgcctcata	cgcagtaggt	17640
tattattaca	tggttaactgg	ctaccccaac	tgattaggaa	agaagtaaat	ttgttttata	17700
aaaatacata	ctcattgagg	tgcatagaat	aattaagaaa	ttaaaagaca	cttgtaattt	17760
tgaatccagt	gaatacccac	tgtaaatatt	tggtatatct	ctttctagtc	tttttttccc	17820
ttttgcatgt	atattcttta	agactcccac	cccactgga	tcatctctgc	atgttctaata	17880
ctgctttttt	cacagcagat	tctaagcctc	tttgaataatc	aacacaaact	tcaacaactt	17940
catctataga	tgccaaataa	taaattcatt	tttatttact	taaccacttc	ctttggatgc	18000
ttaggtcatt	ctgatgtttt	gctattgaaa	ccaatgctat	actgaacact	tctgtcacta	18060
aaactttgca	cacactcatg	aatagcttct	taggataaat	ttttagagat	ggatttgcta	18120
aatcagagac	cattttttta	aattaaaaaa	caattattca	tatcgtttgg	catgtaagac	18180
agtaaaattt	ccttttattt	tgacaggatt	caactggaag	ctttgtgctg	cctttccggc	18240
aagtcatgta	tgctccatat	cccaccacac	acatagatgt	ggatgtcaat	actgtgaagc	18300

-continued

agatgccacc	ctgtcatgaa	catatttata	atcagcgtag	atacatgaga	tccgagctga	18360
cagccttctg	gagagccact	tcagaagaag	acatggctca	ggatacgatc	atctacactg	18420
acgaaagctt	tactcctgat	ttgtacgtaa	tgctctgcct	gctgggtactg	tagtcaagca	18480
atatgaaatt	gtgtccttta	cgaataaaaa	caaaacagaa	gttgcattha	aaaagaaaga	18540
aatattacca	gcagaattat	gottgaagaa	acatttaatc	aagcattttt	ttcttaaag	18600
ttcttctttt	tccatacaat	tgtgtttacc	ctaaaatagg	taagattaac	ccttaaagta	18660
aatatttaac	tatttgttta	ataaatatat	attgagctcc	taggcactgt	tctaggtacc	18720
gggcttaata	gtggccaacc	agacagcccc	agccccagcc	cctacattgt	gtatagtcta	18780
ttatgtaaca	gttattgaat	ggacttatta	acaaaaccaa	agaagtaatt	ctaagtcttt	18840
tttttcttga	catatgaata	taaaatacag	caaaactggt	aaaatatatt	aatggaacat	18900
ttttttactt	tgcattttat	attgttatcc	acttcttatt	tttttttaaa	aaaaaaagcc	18960
tgaacagtaa	attcaaaagg	aaaagtaatg	ataattaatt	gttgagcatg	gacccaactt	19020
gaaaaaaaaa	atgatgatga	taaatctata	atcctaaaac	cctaagtaaa	cacttaaaag	19080
atgttctgaa	atcaggaaaa	gaattatagt	atacttttgt	gtttctcttt	tatcagttga	19140
aaaaaggcac	agtagctcat	gcctgtaaga	acagagcttt	gggagtgcaa	ggcaggcgga	19200
tcacttgagg	ccaggagttc	cagaccagcc	tgggcaacat	agtgaaaccc	catctctaca	19260
aaaaataaaa	aagaattatt	ggaatgtgtt	tctgtgtgcc	tgtaatccta	gctattccga	19320
aagctgaggc	aggaggatct	tttgagccca	ggagtgtgag	gttacaggga	gttatgatgt	19380
gccagtgtac	tccagcctgg	ggaacaccga	gactctgtct	tatttaaaaa	aaaaaaaaaa	19440
aaaatgcttg	caataatgcc	tggcacatag	aaggtaacag	taagtgttaa	ctgtaataac	19500
ccagggtctaa	gtgtgtaagg	caatagaaaa	attggggcaa	ataagcctga	cctatgtatc	19560
tacagaatca	gtttgagctt	aggtaacaga	cctgtggagc	accagtaatt	acacagtaag	19620
tgtaaaccaa	aagcatagaa	taggaatatc	ttgttcaagg	gacccccagc	cttatacatc	19680
tcaagggtgca	gaaagatgac	ttaatatagg	accatttttt	tcctagtctt	ccagagtttt	19740
tattggttct	tgagaaagta	gtaggggaat	gttttagaaa	atgaattggg	ccaactgaaa	19800
ttacatgtca	gtaagttttt	atatattggg	aaattttagt	agacatgtag	aagttttcta	19860
attaatctgt	gccttgaaac	atcttctttt	ttcctaaagt	gcttagtatt	ttttccgttt	19920
tttgattggg	tacttgggag	cttttttgag	gaaatttagt	gaactgcaga	atgggtttgc	19980
aaccatttgg	tatttttggt	ttgtttttta	gaggatgtat	gtgtatttta	acatttctta	20040
atcattttta	gccagctatg	tttgttttgc	tgatttgaca	aactacagtt	agacagctat	20100
tctcattttg	ctgatcatga	caaaataata	tctgaatth	ttaaattttg	catccagctc	20160
taaattttct	aaacataaaa	ttgtccaaaa	aatagtatth	tcagccacta	gattgtgtgt	20220
taagtctatt	gtcacagagt	cattttactt	ttaagtatat	gtttttacat	gttaattatg	20280
tttgttattt	ttaattttta	cttttttaaa	taattccagt	cactgccaat	acatgaaaaa	20340
ttggctcactg	gaattttttt	tttgactttt	attttagggt	catgtgtaca	tgtgcagggtg	20400
gtttatacag	gtaaattgcg	tgtcatgagg	gtttggtgta	cagggtgattt	cattaccag	20460
gtaataagca	tagtacccaa	taggtagttt	tttgatcttc	acccttctcc	cacctcaag	20520
taggccttgg	tggtgctgtt	tccttctttg	tgtccatgta	tactcagtgt	ttagctccca	20580
cttagaagtg	agaacatgcg	gtagtgggtt	ttctgttctt	ggattagttc	acttaggata	20640
atgacctcta	gctccatctg	gtttttatgg	ctgcatagta	ttccatgggtg	tatatgtatc	20700

-continued

acat	tttctt	tatccag	tct accattg	gata ggcattt	tagg ttgattc	cct gtctttg	tta 20760	
tcat	gaatag	tgctgtg	atg aacatac	acata tgc	atgtgtc	tttatgg	tag aaaaatt	tgt 20820
attc	cttttag	gtacata	tag aataatg	ggg ttgctag	gggt gaatg	gtagt tctattt	tca 20880	
gttatt	tgag aaatctt	caa actgctt	ttc ataata	gcta aacta	attta cagtc	ccgcc 20940		
agcag	tgtat aagtgtt	ccc ttttctc	cac aaccttg	cca acatctg	tga tttttg	act 21000		
tttta	aataat	agccatt	cct agagaat	tga tttgca	attc tctatt	tagtg atatta	agca 21060	
ttttt	cata tgctttt	tag ctgtctg	tat atattc	ttct gaaaa	attt catgtc	cttt 21120		
gcccag	tttg tagtggg	tg ggttg	tttt tgctt	gttaa ttagt	tttaa gtcc	ctcca 21180		
gattct	gcat atccctt	tgt tggata	catg gtttg	cagat atttt	ctcc cattg	tgtag 21240		
gttgt	ctttt actctg	tga tagttt	cttt tgccat	gcag gagct	cgta ggtcc	attt 21300		
gtgtt	gttt ttgttg	cagt tgcttt	tggc gtctt	catca taaaat	ctgt gccag	ggcct 21360		
atgtc	cagaa tggat	tttcc taggt	tgtct tccag	ggttt ttaca	attt agatt	ttacg 21420		
tttat	gtctt taatcc	atct tgagtt	gatt tttgt	atatg gcaca	aggaa ggggt	ccagt 21480		
ttcact	ccaa ttcctat	ggc tagca	attat cccag	acca tttatt	gaat acgg	agtcct 21540		
ttcccc	attg cttgtt	tttt gtca	actttg ttga	atca gatgg	tgtga agtgt	gtggc 21600		
tttatt	ctt ggctct	tat tctcc	attgg tctat	gtg tgttt	tata acagt	accct 21660		
gctgtt	cagg ttcctat	agc ctttt	tagtat aaaat	cggt aatgt	gatgc ctcc	agcttt 21720		
gttctt	tttg cttagg	attg ctttg	gctat ttggg	ctct ttttg	gtcc atatta	attt 21780		
taaa	acagtt tttctg	gtt ttgtg	aagga tatc	attgg agttt	atagg aatag	cattg 21840		
aatct	gtaga ttgctt	tggg cagt	atggc atttt	aacaa tatta	attct tcc	atctat 21900		
gaatat	ggaa tgtttt	tcca tgtgt	ttgtg tcat	ctctt atacct	gatg tataa	agaaa 21960		
agctg	gtatt attcct	actc aatct	gttcc aaaaa	attga ggagg	aggaa ctctt	cccta 22020		
atgag	gccag catcatt	ctg atacc	aaaac ctggc	agaga caca	acagaa aaa	agaaaac 22080		
ttcagg	ccaa tatcct	tgat gaata	tagat gcaaaa	tcc tcaac	aaaat actag	caaac 22140		
caa	atccagc agcac	atcaa aaagc	tgatc tactt	tgatc aagtag	gctt tatcc	ctggg 22200		
atgca	agggt ggttca	acat acacaa	tca ataag	tgtga ttc	atcac aaac	agagct 22260		
aaaa	acaaaa accaca	agat tatct	caata ggtag	agaaa aggtt	gtcaa taaa	atttaa 22320		
catc	ctccat gttaaaa	acc ttcag	taggt caggt	gtagt gactc	acacc tgta	atccca 22380		
gcact	ttggg aggcc	aaggc gggc	atatct ctta	agccca ggagt	tcaag acgag	cctag 22440		
gcagc	atgggt gaaac	cccat ctct	acaaaa aaaaa	aaaaa aaaaa	atta gctt	gggatg 22500		
gtgac	atgca cctat	agtcc cagct	attca ggag	gttgg gtggg	aggat tgttt	gagcc 22560		
cgagg	ggcag aggtt	ggcag cgagc	tgaga tcatg	ccacc gca	ctccagc ctggg	caacg 22620		
gagt	gagacc ctgtct	caaa aaagaa	aat caca	aaaca ccta	aaaaa ctagg	cattg 22680		
aagga	acatg cctca	aaaaa ata	agaacca tctat	gacag accca	tagcc aat	atctt	ac 22740	
caaat	gggca aaagc	tggaa gtatt	ctcct tgaga	accgt aaca	agacaa ggat	gtccac 22800		
tctc	accact cctttt	cagc atag	ttcttg aagtc	ctagc cagag	caatc agg	aaagaga 22860		
aagaa	agaaa gacatt	caga tagga	agaga agaag	tcaaa ctatt	tctgt ttgc	aggcag 22920		
tata	attctg tacct	agaaa atctc	atagt ctctg	cccag aaact	cctaa atct	gttaaa 22980		
aattt	cagca aagttt	tggc attct	ctata ctcca	acacc ttcca	aagtg agag	caaaat 23040		

-continued

caagaacaca	gtccattca	caatagccgc	aaaacgaata	aaatacctag	gaatccagct	23100
aaccagggag	gtgaaagatc	tctatgagaa	ttacaaaaca	ctgctgaaag	aaatcagaga	23160
tgacacaaac	aaatggaaat	gttctttttt	aacaccttgc	tttatctaata	tcacttatga	23220
tgaagatact	cattcagtgg	aacaggtata	ataagtccac	tcgattaaat	ataagcctta	23280
ttctctttcc	agagcccaag	aaggggcact	atcagtgcc	agtcaataat	gacgaaatgc	23340
taatatTTTT	cccccttacg	gtttctttct	tctgtagtgt	ggtacactcg	tttcttaaga	23400
taaggaaact	tgaactacct	tcctgtttgc	ttctacacat	accattctc	tttttttgcc	23460
actctggtea	ggtataggat	gatccctacc	actttcagtt	aaaaactcct	cctcttacta	23520
aatgttctct	taccctctgg	cctgagtaga	acctagggaa	aatggaagag	aaaaagatga	23580
aaggagggtg	gggcctggga	aggggaataag	tagtcctgtt	tgtttggtgtg	tttgctttag	23640
cacctgctat	atcctagggtg	ctgtgttagg	cacacattat	tttaagtggc	cattatatta	23700
ctactactca	ctctggctcg	tgccaaggta	ggtagtactt	tcttgगतag	ttggttcgatg	23760
ttacttacag	atggtgggct	tggtgaggca	aaccagtggtg	ataatcatcg	gagtgtgttc	23820
tctaactctca	ctcaaatttt	tcttcacatt	ttttggtttg	ttttggtttt	tgatggtagt	23880
ggcttatTTTT	tggtgctgggt	ttgttttttg	tttttttttg	agatggcaag	aattggtagt	23940
tttatTTtatt	aattgcctaa	gggtctctac	tttttttaaa	agatgagagt	agtaaaatag	24000
attgatagat	acatacatat	ccttactggg	gactgcttat	attcttttaga	gaaaaaatta	24060
catattagcc	tgacaaacac	cagtaaaatg	taaatatatc	cttgagtaaa	taaatgaatg	24120
tatatTTTTgt	gtctccaaat	atatatatct	atattcttac	aaatgtgttt	atatgtaata	24180
tcaatttata	agaacttaaa	atgttggtctc	aagtgaggga	ttgtggaagg	tagcattata	24240
tggccatttc	aacatttgaa	cttttttctt	ttcttcattt	tcttcttttc	ttcaggaata	24300
tttttcaaga	tgtcttacac	agagacactc	tagtgaaagc	cttcctggat	caggtaaagt	24360
ttgaacttga	gattgtcaga	gtgaatgata	tgacatgttt	tcttttttaa	tatatcctac	24420
aatgcctgtt	ctatatattt	atattcccct	ggatcatgcc	ccagagtctt	gctcagcaat	24480
tgcagttaag	ttagttacac	tacagttctc	agaagagtct	gtgagggcat	gtcaagtgca	24540
tcattacatt	ggttgcctct	tgtcctagat	ttatgcttcg	ggaattcaga	cctttgttta	24600
caatataata	aatattattg	ctatctttta	aagatataat	aataagatat	aaagttgacc	24660
acaactactg	ttttttgaaa	catagaattc	ctggtttaca	tgtatcaaag	tgaaatctga	24720
cttagctttt	acagatataa	tatatacata	tatatatcct	gcaatgcttg	tactatatat	24780
gtagtacaag	tatatatata	tgtttggtgtg	tgtatatata	tatagtacga	gcatatatac	24840
atattaccag	cattgtagga	tatatatatg	tttatatat	aaaaaaaagt	tataaactta	24900
aaaccctatt	atgttatgta	gagtatatgt	tatatatgat	atgtaaaata	tataacatat	24960
actctatgat	agagtgtaat	atattttttta	tatatatttt	aacatttata	aaatgataga	25020
attaagaatt	gagtcctaata	ctgttttatt	aggtgctttt	tgtagtgtct	ggtcttttcta	25080
aagtgtctaa	atgatttttc	cttttgactt	attaatgggg	aagagcctgt	atattaacaa	25140
ttaagagtgc	agcattccat	acgtcaaaca	acaaacattt	taattcaagc	attaacctat	25200
aacaagtaag	tttttttttt	ttttttgaga	aagggagggt	gtttatttgc	ctgaaatgac	25260
tcaaaaatat	ttttgaaaca	tagtgtactt	atttaaataa	catctttatt	gtttcattct	25320
tttaaaaaat	atctacttaa	ttacacagtt	gaaggaaatc	gtagattata	tggaacttat	25380
ttcttaatat	attacagttt	gttataataa	cattctgggg	atcaggccag	gaaactgtgt	25440

-continued

catagataaa	gctttgaaat	aatgagatcc	ttatgtttac	tagaaatfff	ggattgagat	25500
ctatgaggte	tgtgacatat	tgogaagtte	aaggaaaatt	cgtaggcctg	gaatttcatg	25560
cttctcaagc	tgacataaaa	tccctcccac	tctccacctc	atcatatgca	cacattctac	25620
tcctaccacc	ccactccacc	ccctgcaaaa	gtacaggtat	atgaatgtct	caaaaccata	25680
ggctcatctt	ctaggagctt	caatgttatt	tgaagatttg	ggcagaaaaa	attaagtaat	25740
acgaaataac	ttatgtatga	gttttaaaag	tgaagtaaac	atggatgtat	tctgaagtag	25800
aatgcaaaat	ttgaatgcat	ttttaaagat	aaattagaaa	acttctaaaa	actgtcagat	25860
tgtctggggc	tggtggctta	tgectgtaat	cccagcactt	tgaggagtccg	aggtgggtgg	25920
atcacaaggt	caggagatcg	agaccatcct	gccaacatgg	tgaaaccccg	tctctactaa	25980
gtatacaaaa	attagctggg	cgtggcagcg	tgtgcctgta	atcccagcta	cctgggaggc	26040
tgaggcagga	gaatcgcttg	aaccaggag	gtgtaggttg	cagtgagtca	agatcgcgcc	26100
actgcacttt	agcctgggtga	cagagctaga	ctccgtctca	aaaaaaaaaa	aaaatatcag	26160
attgttccta	cacctagtgc	ttctatacca	cactcctggt	agggggcatc	agtggaaatg	26220
gttaaggaga	tgtttagtgt	gtattgtctg	ccaagcactg	tcaacactgt	catagaaact	26280
tctgtacgag	tagaatgtga	gcaaattatg	tgttgaaaatg	gttctctctc	ctgcaggtct	26340
ttcagctgaa	acctggctta	tctctcagaa	gtactttcct	tgcacagttt	ctacttgtcc	26400
ttcacagaaa	agccttgaca	ctaataaaaat	atatagaaga	cgatacgtga	gtaaaactcc	26460
tacacggaag	aaaaaccttt	gtacattggt	tttttgtttt	gtttcctttg	tacattttct	26520
atatcataat	ttttgcgctt	cttttttttt	tttttttttt	tttttttcca	ttatttttag	26580
gcagaagggg	aaaaagccct	ttaaatctct	tcggaacctg	aagatagacc	ttgatttaac	26640
agcagagggg	gatcttaaca	taataatggc	tctggctgag	aaaattaaac	caggcctaca	26700
ctctttttatc	tttggaagac	ctttctacac	tagtgtgcaa	gaacgagatg	ttctaataac	26760
tttttaaatg	tgtaacttaa	taagcctatt	ccatcacaat	catgatcgct	ggtaaagtag	26820
ctcagtgggtg	tggggaaaacg	ttcccttgga	tcatactcca	gaattctgct	ctcagcaatt	26880
gcagtttaagt	aagttacact	acagttctca	caagagcctg	tgaggggatg	tcaggtgcat	26940
cattacattg	gggtgtctct	ttcctagatt	tatgcttttg	ggatacagac	ctatgtttac	27000
aatataataa	atattattgc	tatcttttaa	agatataata	ataggatgta	aacttgacca	27060
caactactgt	ttttttgaaa	tacatgattc	atggttttaca	tgtgtcaagg	tgaaatctga	27120
gttggttttt	acagatagtt	gactttctat	cttttggeat	tctttgggtg	gtagaattac	27180
tgtaataactt	ctgcaatcaa	ctgaaaacta	gagcctttta	atgatttcaa	ttccacagaa	27240
agaaagtgag	cttgaacata	ggatgagctt	tagaaagaaa	attgatcaag	cagatgttta	27300
attggaattg	attattagat	cctactttgt	ggatttagtc	cctgggattc	agtctgtaga	27360
aatgtctaata	agttctctat	agtccttggt	cctgggtgaac	cacagttagg	gtgttttggt	27420
tatttttattg	ttcttgctat	tgttgatatt	ctatgtagtt	gagctctgta	aaaggaaatt	27480
gtattttatg	tttttagtaat	tgttgccaac	tttttaaat	aattttcatt	atttttgagc	27540
caaattgaaa	tgtgcacctc	ctgtgccttt	tttctcctta	gaaaatctaa	ttacttgga	27600
caagttcaga	tttactgggt	cagtcatttt	catcttggtt	tcttcttgct	aagtcttacc	27660
atgtacctgc	tttggaatc	attgcaactc	tgagattata	aaatgcctta	gagaatatac	27720
taactaataa	gatctttttt	tcagaaacag	aaaatagttc	cttgagtact	tccttcttgc	27780

-continued

atttctgcct	atgtttttga	agttgttgct	gtttgcctgc	aataggctat	aaggaatagc	27840
aggagaaatt	ttactgaagt	gctgttttcc	taggtgctac	tttggcagag	ctaagttatc	27900
ttttgttttc	ttaatgcgtt	tggaccattt	tgctggctat	aaaataactg	attaatataa	27960
ttctaacaca	atgttgacat	tgtagttaca	caaacacaaa	taaatatttt	atttaaaatt	28020
ctggaagtaa	tataaaaggg	aaaatatatt	tataagaaag	ggataaaggt	aatagagccc	28080
ttctgcccc	caccaccaa	atttacacaa	caaatgaca	tgttcgaatg	tgaaagggtca	28140
taatagcttt	cccatcatga	atcagaaaga	tgtggacagc	ttgatgtttt	agacaaccac	28200
tgaactagat	gactgttgta	ctgtagctca	gtcattttaa	aaatatataa	atactacctt	28260
gtagtgctcc	atactgtgtt	ttttacatgg	tagattctta	tttaagtgtc	aactgggttat	28320
tttctttggc	tggtttattg	tactgttata	cagaatgtaa	gttgtacagt	gaaataagtt	28380
attaaagcat	gtgtaaacad	tgttatatat	cttttctcct	aaatggagaa	ttttgaataa	28440
aatatatattg	aaattttgcc	tctttcagtt	gttcattcag	aaaaaaatac	tatgatattt	28500
gaagactgat	cagcttctgt	tcagctgaca	gtcatgctgg	atctaaactt	tttttaaaat	28560
taattttgtc	ttttcaaaga	aaaaatattt	aaagaagctt	tataatataa	tcttatgtta	28620
aaaaaacttt	ctgcttaact	ctctggattt	cattttgatt	tttcaaatta	tatattaata	28680
tttcaaattg	aaaatactat	ttagataaat	tgttttttaa	cattcttatt	attataatat	28740
taatataacc	taaactgaag	ttattcatcc	caggtagcta	atacatgtat	ccaaagtaaa	28800
aatccaagga	atctgaacac	tttcatctgc	aaagctagga	ataggtttga	cattttcact	28860
ccaagaaaaa	gttttttttt	gaaaatagaa	tagttgggat	gagaggtttc	tttaaaagaa	28920
gactaactga	tcacattact	atgattctca	aagaagaaac	caaaacttca	tataatacta	28980
taaagtaaata	ataaaatagt	tccttctata	gtatatttct	ataatgctac	agtttaaaaca	29040
gatcactctt	atataatact	attttgattt	tgatgtagaa	ttgcacaaat	tgatatttct	29100
cctatgatct	gcagggtata	gcttaaagta	acaaaaacag	tcaaccacct	ccatttaaca	29160
cacagtaaca	ctatgggact	agttttatta	cttccatttt	acaaatgagg	aaactaaagc	29220
ttaaagatgt	gtaatacacc	gcccagggtc	acacagctgg	taaagggtgga	tttcatccca	29280
gacagttaca	gtcattgcc	tgggcacagc	tcctaactta	gtaactccat	gtaactggta	29340
ctcagtgtag	ctgaattgaa	aggagagtaa	ggaagcaggt	tttacaggtc	tacttgccact	29400
attcagagcc	cgagtgtgaa	tccctgctgt	gctgcttgga	gaagttactt	aacctatgca	29460
aggttcattt	tgtaaataatt	ggaaatggag	tgataatacg	tacttcacca	gaggatttaa	29520
tgagacctta	tacgatcctt	agttcagtac	ctgactagt	cttcataaat	gctttttcat	29580
ccaatctgac	aatctccagc	ttgtaattgg	ggcatttaga	acatttaata	tgattattgg	29640
catggtaggt	taaagctgtc	atcttgctgt	tttctatttg	ttctttttgt	tttctcctta	29700
cttttggtt	tttttattct	actatgtctt	ttctattgtc	ttattaacta	tactctttga	29760
tttattttag	tggttggttt	agggttatac	ctctttctaa	tttaccagtt	tataaccagt	29820
ttatatacta	cttgacatat	agcttaagaa	acttactgtt	gttgtctttt	tgtgtttatg	29880
gtcttaacgt	ttttatttct	acaaacatta	taaactccac	actttattgt	tttttaattt	29940
tacttataca	gtcaattatc	ttttaagat	atttaaatat	aaacattcaa	aacaccccaa	30000
t						30001

-continued

<212> TYPE: DNA
<213> ORGANISM: homo sapiens
<400> SEQUENCE: 3

attccccgga tacgtaacct acggtgtccc gctaggaaag agaggtgcgt caaacagcga 60
caagttccgc ccacgtaaaa gatgacgctt ggtgtgtcag ccgtccctgc tgcccgggtg 120
cttctctttt gggggcgggg tctagcaaga gcaggtgtgg gtttaggaga tatctccgga 180
gcatttggat aatgtgacag ttggaatgca gtgatgtcga ctctttgccc accgccatct 240
ccagctgttg ccaagacaga gattgcttta agtggcaaact cacctttatt agcagctact 300
tttgcttact gggacaatat tcttggtcct agagtaaggc acatttgggc tccaaagaca 360
gaacaggtag ttctcagtga tggagaaata acttttcttg ccaaccacac tctaaatgga 420
gaaatccttc gaaatgcaga gagtgggtgct atagatgtaa agttttttgt cttgtctgaa 480
aagggagtga ttattgtttc attaatcttt gatggaaact ggaatgggga tcgcagcaca 540
tatggactat caattatact tccacagaca gaacttagtt tctacctccc acttcataga 600
gtgtgtgttg atagattaac acatataatc cggaaaggaa gaatatggat gcataaggaa 660
agacaagaaa aatgtccaga agattatctt agaaggcaca gagagaatgg aagatcaggg 720
tcagagtatt attccaatgc ttactggaga agtgattcct gtaatggaaa ctgctttcct 780
ctatgaaatt cccccgggtt cctggaggaa atagatatag gctgatacag ttacccaatg 840
atggatgaat attgggggac cgcttgggtc ttgaaaggct ttcttttctc caggaaagaa 900
atttttttcc ttttccataa aaagcttggg aatggaagac aacaattccc attctttttt 960
tgcgttcac ccctatgtga caacagaaat ttttggggaa acaacaacga aaaaatttta 1020
tcccgcgcgc a 1031

<210> SEQ ID NO 4
<211> LENGTH: 3244
<212> TYPE: DNA
<213> ORGANISM: homo sapiens
<400> SEQUENCE: 4

gggcggggct gcggttgcgg tgcttgcgcc cgcggcgggc gaggcgcagg cgggtggcgag 60
tgatatctc cggagcattt ggataatgtg acagttggaa tgcagtgatg tcgactcttt 120
gcccaccgcc atctccagct gttgccaaaga cagagattgc tttaagtggc aaatcacctt 180
tattagcagc tacttttgct tactgggaca atattcttgg tcctagagta aggcacattt 240
gggctccaaa gacagaacag gtactttctc gtgatggaga aataactttt cttgccaaac 300
acactctaaa tggagaaatc cttcgaaatg cagagagtgg tgctatagat gtaaagtttt 360
ttgtcttgtc tgaaaaggga gtgattattg tttcattaat ctttgatgga aactggaatg 420
gggatcgcag cacatatgga ctatcaatta tacttccaca gacagaactt agtttctacc 480
tcccacttca tagagtgtgt gttgatagat taacacatat aatccggaaa ggaagaatat 540
ggatgcataa ggaaagacaa gaaaatgtcc agaagattat cttagaaggc acagagagaa 600
tggaagatca gggtcagagt attattccaa tgcttactgg agaagtgatt cctgtaatgg 660
aactgctttc atctatgaaa tcacacagtg ttctgaaga aatagatata gctgatacag 720
tactcaatga tgatgatatt ggtgacagct gtcataagg ctttcttctc aatgccatca 780
gctcacactt gcaaacctgt ggctgttccg ttgtagtagg tagcagtgca gagaaagtaa 840
ataagatagt cagaacatta tgcttttttc tgactccagc agagagaaaa tgctccaggt 900

-continued

tatgtgaagc	agaatcatca	tttaaatatg	agtcagggct	ctttgtacaa	ggcctgctaa	960
aggattcaac	tggaagcttt	gtgctgcctt	tccggcaagt	catgtatgct	ccatatccca	1020
ccacacacat	agatgtggat	gtcaatactg	tgaagcagat	gccaccctgt	catgaacata	1080
tttataatca	gcgtagatac	atgagatccg	agctgacagc	cttctggaga	gccacttcag	1140
aagaagacat	ggctcaggat	acgatcatct	acactgacga	aagctttact	cctgatttga	1200
atatttttca	agatgtctta	cacagagaca	ctctagtga	agccttcctg	gatcaggtct	1260
ttcagctgaa	acctggctta	tctctcagaa	gtactttcct	tgcacagttt	ctacttgtcc	1320
ttcacagaaa	agccttgaca	ctaataaaat	atatagaaga	cgatacgag	aagggaaaaa	1380
agccctttaa	atctcttcgg	aacctgaaga	tagaccttga	tttaacagca	gagggcgatc	1440
ttaacataat	aatggctctg	gctgagaaaa	ttaaaccagg	cctacactct	tttatctttg	1500
gaagaccttt	ctacactagt	gtgcaagaac	gagatgttct	aatgactttt	taaatgtgta	1560
acttaataag	cctattccat	cacaatcatg	atcgctggta	aagtagctca	gtggtgtggg	1620
gaaacgttcc	cctggatcat	actccagaat	tctgctctca	gcaattgcag	ttaagtaagt	1680
tacactacag	ttctcacaag	agcctgtgag	gggatgtcag	gtgcatcatt	acattgggtg	1740
tctcttttcc	tagatttatg	cttttgggat	acagacctat	gtttacaata	taataaatat	1800
tattgctatc	ttttaagat	ataataatag	gatgtaaact	tgaccacaac	tactgttttt	1860
ttgaaataca	tgattcatgg	tttacetgtg	tcaagggtga	atctgagttg	gcttttacag	1920
atagttgact	ttctatcttt	tggcattctt	tgggtgtgtg	aattactgta	atacttctgc	1980
aatcaactga	aaactagagc	ctttaaatga	tttcaattcc	acagaaagaa	agtgagcttg	2040
aacataggat	gagctttaga	aagaaaattg	atcaagcaga	tgtttaattg	gaattgatta	2100
ttagatccta	ctttgtggat	ttagtccctg	ggattcagtc	tgtagaaatg	tctaatagtt	2160
ctctatagtc	cttgttcctg	gtgaaccaca	gttaggggtg	tttgtttatt	ttattgttct	2220
tgctattggt	gatattctat	gtagttgagc	tctgtaaaag	gaaattgtat	tttatgtttt	2280
agtaattggt	gccaactttt	taaattaatt	ttcattatct	ttgagccaaa	ttgaaatgtg	2340
cacctcctgt	gccttttttc	tccttagaaa	atctaattac	ttggaacaag	ttcagatttc	2400
actggtcagt	cattttcatc	ttgttttctt	cttgctaagt	cttaccatgt	acctgctttg	2460
gcaatcattg	caactctgag	attataaaat	gccttagaga	atatactaac	taataagatc	2520
tttttttcag	aaacagaaaa	tagttccttg	agtacttcct	tcttgcatct	ctgcctatgt	2580
ttttgaagtt	gttgctgttt	gcctgcaata	ggctataagg	aatagcagga	gaaattttac	2640
tgaagtgctg	ttttcctagg	tgctactttg	gcagagctaa	gttatctttt	gttttcttaa	2700
tgcgttttga	ccattttgct	ggctataaaa	taactgatta	atataattct	aacacaatgt	2760
tgacattgta	gttacacaaa	cacaaataaa	tattttatct	aaaattcttg	aagtaatata	2820
aaagggaaaa	tatatattata	agaaagggat	aaaggtaata	gagcccttct	gccccccacc	2880
caccaaatct	acacaacaaa	atgacatgtt	cgaatgtgaa	aggtcataat	agctttccca	2940
tcatgaatca	gaaagatgtg	gacagcttga	tgtttttagac	aaccactgaa	ctagatgact	3000
gttgtagctg	agctcagtca	tttaaaaaat	atataaatac	tacctttag	tgtcccatc	3060
tgtgtttttt	acatggtaga	ttcttattta	agtgttaact	ggttattttc	tttggctggg	3120
ttattgtact	gttatacaga	atgtaagttg	tacagtgaaa	taagttatta	aagcatgtgt	3180
aaacattggt	atatactttt	tctcctaaat	ggagaatctt	gaataaaata	tatttgaaat	3240
tttg						3244

-continued

```

<210> SEQ ID NO 5
<211> LENGTH: 761
<212> TYPE: DNA
<213> ORGANISM: homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (693)..(693)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (722)..(722)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 5
cacgaggctt tgatatttct tacaacgaat ttcattgtgta gaccactaa acagaagcta    60
taaaagttgc atggtcaaat aagtctgaga aagtctgcag atgatataat tcacctgaag    120
agtcacagta tgtagccaaa tgttaaaggt tttgagatgc catacagtaa atttaccag    180
cattttctaa atttatttga ccacagaatc cctattttta gcaacaactg ttacatccca    240
tggattccag gtgactaaag aatacttatt tcttaggata tgttttattg ataataacaa    300
ttaaaatttc agatatcttt cataagcaaa tcagtggctt ttttacttca tgttttaatg    360
ctaaaatatt ttcttttata gatagtcaga acattatgcc tttttctgac tccagcagag    420
agaaaatgct ccaggttatg tgaagcagaa tcatcattta aatatgagtc agggctcttt    480
gtacaaggcc tgctaaagga ttcaactgga agctttgtgc tgcctttccg gcaagtcatt    540
tatgtcccat atcccaccac acacatagat gtggatgtca atactgtgaa gcagatgcca    600
ccctgtcatg aacatattta taatcagcgt agatacatga gatccgagct gacagccttc    660
tggagagcca cttcagaaga agacatggct cangatacga tcatctacac tgacgaaagc    720
tntactcctg atttgaatat ttttcaagat gtcttacaca g                        761

```

```

<210> SEQ ID NO 6
<211> LENGTH: 1901
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 6
acgtaacctc cgggtgtcccg ctaggaaaga gaggtgcgtc aaacagcgac aagttccgcc    60
cacgtaaaag atgacgcttg atatctccgg agcatttgga taatgtgaca gttggaatgc    120
agtgatgtcg actctttgcc caccgccatc tccagctggt gccaaagacag agattgcttt    180
aagtggcaaa tcacctttat tagcagctac ttttgcttac tgggacaata ttcttggtcc    240
tagagtaagg cacatttggg ctccaaagac agaacaggta cttctcagtg atggagaaat    300
aacttttctt gccaaaccaca ctctaaatgg agaaatcctt cgaaatgcag agagtgggtgc    360
tatagatgta aagttttttg tcttgtctga aaagggagtg attattgttt cattaatctt    420
tgatggaaac tggaatgggg atcgagcac atatggacta tcaattatac ttccacagac    480
agaacttagt ttctacctcc cacttcattg agtgtgtggt gatagattaa cacatataat    540
ccggaaagga agaatatgga tgcataagga aagacaagaa aatgtccaga agattatctt    600
agaaggcaca gagagaatgg aagatcaggg tcagagtatt attccaatgc ttactggaga    660
agtgattcct gtaatggaac tgctttcatc tatgaaatca cacagtgttc ctgaagaaat    720
agatatagct gatacagtac tcaatgatga tgatattggg gacagctgtc atgaaggctt    780
tcttctcaag taagaatttt tcttttcata aaagctggat gaagcagata ccatcttatg    840

```


-continued

ctcacctatg	acaagatttg	gaagaaagaa	aataacagac	tgtctactta	gattgttcta	900
gggacattac	gtatttgaac	tggtgcttaa	atttgtgtta	tttttcactc	attatatattc	960
tatatatatt	tggtgttatt	ccatttgcta	tttaaagaaa	ccgagtttcc	atcccagaca	1020
agaaatcatg	gccccttgct	tgattctggg	ttcttgtttt	acttctcatt	aaagctaaca	1080
gaatcctttc	atattaagtt	gtactgtaga	tgaacttaag	ttatttaggc	gtagaacaaa	1140
attattcata	tttatactga	tctttttcca	tccagcagtg	gagtttagta	cttaagagtt	1200
tgtgccctta	aaccagactc	cctggattaa	tgctgtgtac	ccgtggggcaa	ggtgcctgaa	1260
ttctctatac	acctatttcc	tcatctgtaa	aatggcaata	atagtaatag	tacctaattgt	1320
gtagggttgt	tataagcatt	gagtaagata	aataatataa	agcacttaga	acagtgcctg	1380
gaacataaaa	acacttaata	atagctcata	gctaacattt	cctatttaca	tttcttctag	1440
aaatagccag	tatttggtga	gtgcctacat	gttagttcct	ttactagttg	ctttacatgt	1500
attatcttat	attctgtttt	aaagtttctt	cacagttaca	gattttcatg	aaattttact	1560
tttaataaaa	gagaagtaaa	agtataaagt	attcactttt	atgttcacag	tcttttcctt	1620
taggctcatg	atggagtatc	agaggcatga	gtgtgtttta	cctaagagcc	ttaatggctt	1680
gaatcagaag	cacttttagtc	ctgtatctgt	tcagtgtcag	cctttcatac	atcattttta	1740
atcccatttg	actttaagta	agtcacttaa	tctctctaca	tgtcaatttc	ttcagctata	1800
aaatgatggg	atttcaataa	ataaatacat	taattaaatg	atattatact	gactaattgg	1860
gctgttttta	ggcaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	a		1901
<210> SEQ ID NO 7						
<211> LENGTH: 562						
<212> TYPE: DNA						
<213> ORGANISM: homo sapiens						
<220> FEATURE:						
<221> NAME/KEY: misc_feature						
<222> LOCATION: (166)..(166)						
<223> OTHER INFORMATION: n is a, c, g, or t						
<400> SEQUENCE: 7						
agacgtaacc	tacgggtgtcc	cgctaggaaa	gagagatatc	tccggagcat	ttggataatg	60
tgacagttgg	aatgcagtga	tgtcgactct	ttgcccaccg	ccatctccag	ctgttgccaa	120
gacagagatt	gctttaagtg	gcaaatacacc	tttattagca	gctacntttt	gcttactggg	180
acaatattct	tggtcctaga	gtaaggcaca	tttgggctcc	aaagacagaa	caggtacttc	240
tcagtgatgg	agaaataact	tttcttgcca	accacactct	aaatggagaa	atccttcgaa	300
atgcagagag	tggtgctata	gatgtaaagt	tttttgtctt	gtctgaaaag	ggagtgatta	360
ttgtttcatt	aatctttgat	ggaaactgga	atggggatcg	cagcacatat	ggactatcaa	420
ttatacttcc	acagacagaa	cttagtttct	acctcccact	tcatagagtg	tgtgttgata	480
gattaacaca	tataatccgg	aaaggaagaa	tatggatgca	taaggaaaga	caagaaaatg	540
tccagaagat	tatcttagaa	gg				562
<210> SEQ ID NO 8						
<211> LENGTH: 798						
<212> TYPE: DNA						
<213> ORGANISM: homo sapiens						
<400> SEQUENCE: 8						
gggctctctt	ttgggggcgg	ggctctagcaa	gagcagatat	ctccggagca	tttggataat	60
gtgacagttg	gaatgcagtg	atgtcgactc	tttgcccacc	gccatctcca	gctgttgcca	120

-continued

```

agacagagat tgctttaagt ggcaaatcac ctttattagc agctactttt gcttactggg 180
acaatattct tggctctaga gtaaggcaca tttgggctcc aaagacagaa caggtaacttc 240
tcagtgatgg agaaataact tttcttgcca accacactct aaatggagaa atccttcgaa 300
atgcagagag tggtgctata gatgtaaagt tttttgtctt gtctgaaaag ggagtgatta 360
ttgtttcatt aatctttgat ggaaactgga atggggatcg cagcacatat ggactatcaa 420
ttatacttcc acagacagaa cttagtttct acctccact tcatagagtg tgtgttgata 480
gattaacaca tataatccgg aaaggaagaa tatggatgca taaggaaaga caagaaaatg 540
tccagaagat tatcttagaa ggcacagaga gaatggaaga tcagggtcag agtattattc 600
caatgcttac tggagaagtg attcctgtaa tgggactgct ttcactatg aaatcacaca 660
gtgttcctga agaaatagat atagctgata cagtactcca tgatgatgat atttggtgac 720
agctgtcatg aaaggctttc ttctcaagta ggaatttttt cttttcataa aagctgggat 780
gaagccagat tcccatct 798

```

```

<210> SEQ ID NO 9
<211> LENGTH: 169
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

```

```

<400> SEQUENCE: 9

```

```

aaacagcgac aagttccgcc cacgtaaaag atgatgcttg gtgtgtcagc cgtccctgct 60
gcccggttgc ttctcttttg ggggcggggg ctagcaagag cagatatctc cggagcattt 120
ggataatgtg acagttggaa tgcggtgatg tcgactcttt gccacccg 169

```

```

<210> SEQ ID NO 10
<211> LENGTH: 176
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

```

```

<400> SEQUENCE: 10

```

```

aaaacgtcat cgcacataga aaacagacag acgtaacctc cgggtgtccc ctaggaaaga 60
gaggtgcgtc aaacagcgac aagttccgcc cacgtaaaag atgacgcttg atatctccgg 120
agcatttgga taatgtgaca gttggaatgc agtgatgtcg actctttgcc caccgc 176

```

```

<210> SEQ ID NO 11
<211> LENGTH: 30001
<212> TYPE: DNA
<213> ORGANISM: macaca mulatta
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (26113)..(26155)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (28797)..(29186)
<223> OTHER INFORMATION: n is a, c, g, or t

```

```

<400> SEQUENCE: 11

```

```

aatctctaag caattttttg gggaagaaag aattgcaatt agggcatatg ttagatcag 60
atgggtcttc gtatatccaa cgacaaagaa aagggtgggag gtttcgttaa aaaagagaaa 120
tgttacatag tacttttaga gaaaattcac tggcactatt aagggtctga ggagctggta 180
agtttcaatt ggtgagtgat ggtggtagat aaaattagag ctgcagcagg tcatttcagc 240
aactatcaga taaaactggg ctcaggtcac aacgggcagt ttcagcagct agacttgaaa 300

```


-continued

gaattacact	gcgaggagcaa	tgtcatttgt	cctgcatgct	tttctacccc	ctacccccac	360
tttttttagtt	gggtataaca	agaacgaccc	aaattgtatg	atcaactttc	acaaagcata	420
gaacagtagg	aaaaggggtct	gtttctgcag	aagatgtaga	cgttgagagc	cattttatgt	480
at ttat tttct	ccctttcttc	atcgggtgaat	gattaaaatg	ttctgtatga	tttttagtga	540
tgagaaagg	taaacgccac	tcatctgtag	taagtgtaat	ctacacactt	gcagacccaa	600
aggcataagg	tttaaaaaac	ctttgttttt	ttacacatca	aacagagtgg	tataaatgct	660
actcatctgt	agtaagtga	atctatacac	ctgcagacca	acgacgcaag	gtttcaaaaa	720
tctttgtgtt	ttttacacat	caaacagaat	ggtacatttt	tcaaaagttt	aaaaaaaaaa	780
aaaatccaca	tatcacaact	agcaaaaatg	acattcccca	gtgtgaaaat	catgcttgag	840
agaattctta	catgtaaagg	caaaattgca	gtgactttac	aagggacctg	gggattcccc	900
cccacagtgt	ggagctgtcc	cctaccaggg	tttgccggcg	agttttgaat	gtacttaaca	960
gtgtctcacg	gtaaaaaaca	aacttcatcc	accaaataat	tgttgagcgc	ccactgcctg	1020
ccaagcacia	acaaaaccat	tcaaaaccac	gaaatcgtct	gcactttctc	cggatccagc	1080
agcctctgcg	attaagggtt	gcacacgcta	ttgcgccaac	gctcctccag	agcgcgctct	1140
aagataaaag	aatgggacaa	gttgcccttc	cccctttcac	gggcctcgtg	cgtaaacgtc	1200
atcgcatata	gaaaacacac	agacgtaacc	tacgggtgtc	cgctaggaaa	gagaggcgcg	1260
tcaaacagcg	acaagttccg	cccacgtaaa	agatgacgct	tggtgcgtca	gccgtccctg	1320
ctgcccgggt	ccttctctct	ggggggcggg	cctggctaga	gcagggtgtg	gtttaggagg	1380
tgtgtgtttt	tgtttttcct	accctctccc	ctctacttgc	tctcacagta	ctcgctgagg	1440
gtgaacaaga	aaagacctga	taaagattaa	ccagaagaaa	acaaagaggg	aaacaactgc	1500
agcctgtagc	gggctctgga	gcttaagaga	ggcgcgctag	gcgcccgggc	gtgggcgtgg	1560
tcggggcggg	gtcggggccag	gggcggggct	gcggttgccg	tccctgcgcc	cgcggcgggc	1620
gcggcgggcg	cagcggaggc	gcaggcggtg	gcgagtgggt	gagtgaagag	gcggcgctct	1680
ggcgggtgtc	tgtttggcgt	cgggttgccg	ggaagagacg	cgggtagcag	ccggggctct	1740
cctcagagct	cgacacattt	ttactttccc	tctcgtttct	ctgaccgaag	tcgggtgtcc	1800
ggctttcgcc	tctagcgact	ggtggaattg	cctgcatctg	ggccccgggc	ttcgcgggcg	1860
cgcagggacg	agggatggga	atctggcctc	ttcctcgctt	tcccggccgc	agtgcgctgc	1920
cccagctgtc	tccttcccgg	ggacctgctg	ggagcgctgc	cgctacagac	tcgagagaaa	1980
ggagcctcgg	gcactgagag	gcctcgcccc	ggggaaggcc	ggagggcggg	cggcgggcg	2040
cgcggcgctc	ctgcggacca	agtctgggtt	ctctgggaac	ccgagacggt	ccctgatggc	2100
gaggagatca	tgcgggggtc	tatgggggtg	tggagacgtc	tgcagaattt	tagcccaagc	2160
ttctaaggag	tgctgatgac	ttgcatatga	gggcagcaat	gccagtcggt	gtactcccta	2220
ttctgtggga	catgatgtgg	ttgcttcaca	gtcccgagat	gacacagact	tgcttaaagg	2280
aagtgaccat	tgtgacttgg	gcataccttg	actgatggta	atcagttgca	gacagaagtg	2340
cacagattac	atgtctgtgt	ccacactgga	tcagtctggc	cacgaggaac	accacaggct	2400
ttgtattgag	aaacaggagg	gaggtcctgc	actttcccag	gaggggtggc	cctttcagat	2460
gcaatcgaga	ttgttaggct	ctggtagagt	ggttgccctg	ttgtggcagt	tggcaaattc	2520
ctattcaaac	tgttgccgtg	cgtcaccagt	taacaacaag	ggtacacgat	ctgtctggca	2580
ttactttctac	tttgtacaaa	ggatcaaaaa	tactgttaga	tatgattttt	ctcagacttt	2640
gggaaacttt	taacgtaatc	tgtgaatatc	acagaagcaa	gactgtcata	tagaggatat	2700

-continued

taataacctg	gagtcagaat	acttgaaata	tggtgtcatt	tgacacgggc	tctgttatca	2760
ccacctttgc	caagcccttt	cacttgagga	aaacctcaa	tcagttggaa	actgcctcat	2820
gctgacagta	catctgaaac	aaaaacgaga	gtagttacca	cattccagat	tgttcactaa	2880
ggcagcattt	atctgctcca	ggaaaacatt	acaagcaact	tatgaagttg	ataaaatatt	2940
ttgtttggct	atgttggtac	tccaaaagtt	gctttcagag	aaacaaagta	aaccaaggag	3000
gacttctgtt	gttcacgtct	gcccttgggc	tctattctac	gttaattagg	tagttcccag	3060
gaggactaga	ttagcctacc	tattgtctga	gaaacttggg	tctgtgagaa	atggccagat	3120
agtgatacga	acttcacctc	ccagtctttc	ctgatgttta	agattgagaa	agtgttgtga	3180
actttctggg	gctgtaagca	gttcactgtc	cttaaagttg	tccctgggcag	ctcctgttgt	3240
ggaaagtggg	ccgatttagg	attctgcttg	gctttggact	gggagaaaat	aaactgcatg	3300
gttacaagta	ttgagagcca	agttggagaa	ggtggcttac	acctataatg	ccagagcctt	3360
aggaggcagg	ggcaagagga	tcactggaag	tcaggagttc	aagcccaacc	tgggcagcct	3420
agaccctgtc	tctacaaaaa	attaaaaact	tagccggggc	cggtggtgtg	cacctgtagt	3480
cctagctact	tgggaggctg	aggcaggagg	gtcttttgag	cccaggagtt	tgaagttaca	3540
gggagctatg	atcctgccag	tgcactccag	cctggatggc	aaaacgagac	cctgtctcta	3600
aaaaacaaga	agtgagggct	ttatgatcgt	agaaattttg	cttacaatag	cagtggacca	3660
accacctttc	taaataccaa	tcagggaaga	catagttgat	ttttaacaaa	catttaaaga	3720
aaaagcaaaa	cctcaaactt	agcactctac	taacagtttt	agccgatgct	aattaaggta	3780
atcatgtctg	catatatggg	attactttca	gaaagtgtat	tgggaaacct	ctcatgaacc	3840
ctgtgcaacc	ctgagcaagc	caccgtctca	ctcagtttga	atcttggctt	ccctcaaaag	3900
actctgtggc	taatgtttgg	taactctctg	gagtagccag	cactgcatgt	acataggata	3960
ggtacataaa	acaattattg	gttttgagct	gatttttttc	agctgcattt	gcgtgtatgg	4020
atttttctca	ccaaagacaa	tgacttcaag	tgtaataaaa	ataattgtac	agctctccta	4080
attataacttc	tctgtaacat	ttcattttct	agactatttc	ttttggtagg	atttaaaact	4140
aaacaattca	gtatgatctt	tgttcttcat	tttctttctt	attctttttt	ttttcgagac	4200
agagtctccc	tctgttgtgc	catctcagcc	cattgcaacc	tccgccacct	gggttcaagt	4260
gattctectg	cctcagcctc	ctgagtagct	gggattacag	gtgcccgcga	ccacacctac	4320
ctaatttttt	gtatttttag	tagaggcggg	gtttcaccat	gttggctagg	ctggctctaa	4380
actcctgacc	tcagatgatc	cacctgcctc	ggcctcccaa	agagctggga	tgataggcgt	4440
gacccaccat	gcccgcacca	ttttttttct	tattctgtta	ggagtgagag	tgtaactagc	4500
agtctataat	agttcaattt	tcacaacgtg	gtaaaaattt	tccctgtaat	tcaacgagat	4560
tttgcttcag	ggctcagttc	tgttttagga	aatactttta	ttttcagttt	gatgatgaaa	4620
tattagagtt	gtgatattgc	ctttatgatt	acctaccttt	ttaacctaaa	agaatgaaag	4680
aaaaatatgt	ttacagtata	attgtatggg	tgcgtgttaa	cttaattcat	tatgttggcc	4740
tccagtttgc	tgttgttcgt	tatgacagca	gtagtgtcat	taccatttca	attcagatta	4800
cattcctgta	tttgatcatt	gtaaactgat	tgcttaaatt	gtattaaaaa	cagtggatat	4860
tttaaacaag	ctgtactgct	tatatccagt	gctgtctcct	aagactatta	aattgatata	4920
acatatattaa	aagtaaatat	ttcctaaatg	aatttttgaa	attaaaaata	cacgtgttaa	4980
aactgtcttt	gtgttcaacc	atttctgtac	gtacttagag	ttaactgttt	tgccaggctc	5040

-continued

tgtatgccta	ctcataatgt	gataaaagca	ctcatcta	gctctataaa	tagaagtcag	5100
tgctttccat	cagactgaac	actcttggca	agatgtggat	aaaattat	aagtaaaatt	5160
gtttactttg	tcatacat	acagatcaaa	tgtagctcc	caaagcaatc	atatggcaaa	5220
gataggcata	tcataatttg	cctattagct	gctttgtatt	gctattatga	tagatttcac	5280
agtttttagat	ctgcttagat	gaaaatgtaa	ttctttttac	tgtcagtcct	agatataagt	5340
cttcaattat	agtacagtca	cacattgctt	aggaatgcat	cattaggcga	ttttgtcatt	5400
atgcaaacat	catagagtat	acttacataa	acctatatag	tacagccttt	acgtacgtag	5460
gccatatggg	atagtctatt	gctcctaggc	tacaaatctg	tacagctgtt	actgtactga	5520
atactataga	cagttgtaac	acagtgggat	ttatttatct	aaatatatcc	aaacatagaa	5580
aaggtagagt	taaagtatgg	tataaaaaat	aatgatatac	ctatataggc	cacttaccgt	5640
gaatggagct	tgaggacta	gaagttgctc	tggtgagtc	agtaagtaag	tggtgaatga	5700
atgtgaaggc	ctagaacatt	actgtacaca	ctgtagactt	tataaacaca	gtatgcttaa	5760
gctacaccaa	atttatcttt	acagtttttc	ttcaataaaa	aattaatgtg	aacctactat	5820
aactttttta	ctttgtaaac	tttttaattt	tttaactttt	aaaatactta	gcttgaaaca	5880
caaacacgca	tagctataca	aaaatat	ttctttatat	ccttattcta	gaagcttttt	5940
cctattttta	actttttttt	ttttacttgt	tagtcgtttt	tgtaaaaaac	taaaacacac	6000
acactttcac	ctaagcatag	acaggattag	gatcatcagt	ttcactccct	tcacctcac	6060
tgctttccac	ctccacatct	tgcccactg	gaacgttttt	agggggaata	acacacatgt	6120
agctgtcacc	tgctatgata	acagtgcttt	ctgttgaata	cctcctgaag	gacctgcctg	6180
aggctgtttt	acatttaact	taaaaaaaaa	aataagtaga	aggagtacac	tctaaaataa	6240
caataaaagg	tatagtctag	tgaatacata	aaccagcaac	atagtagttt	attatcaagt	6300
gttgataact	gtaataattg	tatgtgctat	actttaaatg	acttgaaaaa	ttgtactaag	6360
accttatgat	ggttacagtg	tcactaaggc	gatagcatat	tttcaggctc	attgtaatct	6420
aatgggacca	ccatcatata	tgagtcacac	cattgactga	aatgttacat	ggtacgtaac	6480
tgtatttgca	agaatgattt	gttttacatt	aatatcacat	aggatgtacc	tttttagagt	6540
gatatgttta	tgtggattaa	gatgtacaag	tgagcaagg	ggacaagagc	ccttggttct	6600
gtcttgatg	tgagctttta	tgctcttctc	atcatgtctg	tttcttatt	aaattcaaag	6660
gcttgacag	gccctattta	gcccttggtt	tctatgtgtt	ctaaataact	aaagctttta	6720
aattctagcc	atttagtgga	gaaactctct	tgcaatggta	aaatgctgta	ttggtttctt	6780
gactagcata	ttaaataat	ttatctttgt	cttgatattt	caatgtcatt	ttaaacaatca	6840
ggattgggct	ttagtattct	cataccaga	gagttcactg	aggatacagg	actgtttgcc	6900
cattttttgt	tatggctcca	gacttggtgt	atttcgatgt	cttttttttt	tttttttttt	6960
tttaaccttt	tagcagcttt	aaagtatttc	tggtgttagg	tggtgtatta	cttttctaag	7020
attactgtaa	caaagcacca	caaactgagt	ggctttaaac	aacagcaatt	tattctctca	7080
caattctaga	agctagaagt	ccgaaatgga	agtgttgatg	gggcatgatc	ctcaaaagag	7140
agaagactct	ttccttgcc	cttcctggct	tctggtggtt	accagcaatc	ctgagcgctc	7200
ctttcttgct	tcgtagtctc	agcagtcacg	tatctgcctt	ttgtcttcac	atggatgtct	7260
acctcttgct	tctgtgtctc	cagatctctc	tccttataaa	cacagaagtt	actggattag	7320
gccccactct	aatccagtat	gaccccat	taacacgatt	acacctattt	ctaaataagg	7380
tcacattcac	atataccaag	ggtaggaat	tgagcatatc	ttttgcaggg	acacaattca	7440

-continued

accacaagt	gtcagtcct	agctgagcct	ttcccttcct	ggttttctcc	tttttagttg	7500
ctgtgggtta	ggggccaaat	ctccagtcac	actagacttg	cacatggact	ggagatttgg	7560
gaatactgcy	ggctattct	atgagcttta	gtatgtaaca	tttaatatca	gtgtaaagaa	7620
gccatttttt	cagttcacta	tttctttgaa	tttcttaatg	tatgccctga	atataagtaa	7680
caagttacta	tgtctcataa	aatgatcata	tcaacaaaca	tttaatgtgc	acctactgtg	7740
ctagttgaat	gtctttatcc	tgataggaga	taacaggcct	ccgcatcttt	gacttaagag	7800
gacaaaccaa	gtatgtctga	atcatttggg	gttttgatgg	atatctttaa	attgctgaac	7860
ctaactcattg	gttttatatg	tcattgttta	gatatctcag	gagcatttgg	ataatgtgac	7920
agttggaatg	cagtgatgtc	gactctttgc	ccaccgccat	ctccagctgt	tgccaagaca	7980
gagattgctt	taagtgggtga	atcaccttta	ttagcagcta	cttttgctta	ctgggacaat	8040
attcttggtc	ctagagtaag	gcacatttgg	gtcccaaaga	cagaacaggt	acttctcagt	8100
gatggagaaa	taacttttct	tgccaaccac	actctaaatg	gagaaatcct	tcgaaatgca	8160
gagagtgggtg	ctatagatgt	aaagtttttt	gtcttgtctg	aaaagggagt	gattattggt	8220
tcattaatct	ttgatggaaa	ctggaatggg	gatcgcagca	catacggact	atcaattata	8280
cttccacaga	cagaacttag	tttctacctc	ccacttcata	gagtgtgtgt	tgatagatta	8340
acacatataa	tccggaaagg	agaatatatg	atgcataagg	taagtgattt	ttcagcttat	8400
taatcatggt	aacctatctt	ttgaaagctt	atcttctgat	acatataaat	cttattttta	8460
aatttatatgc	agtgaacatc	aaacaataga	tattatttat	tttgcattha	tcctgttaga	8520
tacaaataca	tctgggtctga	tgctgtcat	cttcatatta	actgtggaag	gtaggaaatg	8580
gtagctccac	attacagatg	aaaagctaaa	gcttaaaaca	atgcagaaac	ttttagatcc	8640
tggattcttc	ttgggagcct	ttgactctaa	taccttttgt	ttccctttca	ttgcacaatc	8700
ctgtctttcg	cttactacta	tgtgtaagta	taacagttca	aaaaaatagt	ttcataagct	8760
gttggttatg	tagcctttgg	tctctttaac	ctctttgcca	agttcccagg	ttcataaaat	8820
gaggagggtg	aaccgcatgg	ttcccaagag	aattcctttt	aattttacag	aaattattgt	8880
tttccccgaa	gtcctatagt	tcaatatata	atgatattta	catttcagta	tagttttggc	8940
atatctaaag	aacacattaa	gttctccttc	ctgtgttcca	gtttgatact	aacctggaag	9000
tccattaagc	attaccaatt	ttaaaaggct	tttgcccaat	agtaaggaaa	aataatatct	9060
tttaaaagaa	taatttttta	ctatgtttgc	aggcttactt	ccttttttct	cacattatga	9120
aactcttaaa	atcaggagaa	tcttttaaac	atcataatgt	ttaatttgaa	aagtgcaagt	9180
cattcttttc	ctttttgaaa	ctatgcagat	gttacattga	ctattttctg	tgaagttatc	9240
ttttttttcc	ctgcagaata	aagggtgttt	tgattttatt	ttgtgttggt	tataagaaca	9300
tacattcggt	gggttaattt	cctgcccctg	ccccgtttt	ttccctaaag	tagaaagtat	9360
ttttcttggtg	aactaaatta	ctacacaaga	acatgtctat	tgaaaaataa	gtatcaaaat	9420
gttggtgggtt	gtttttttta	ataaattctt	tcttgctcag	gaaagacaag	aaaatgtcca	9480
gaagattatc	ttagaaggca	cagagagaat	ggaagatcag	gtatatgcag	attgcatact	9540
gtcaaatatt	attctcatgg	catgtatctg	tgtaaagttg	atggctacat	ttgtgaaggc	9600
cttggggaca	tacagagtaa	gccttaatgg	agcttttatg	gagggtgtaca	gaataaacta	9660
gaggaagatt	tccatatctt	agacctgaag	agttaaatca	gtaaacaaag	gaaaatagta	9720
attgcatcta	caaattaata	tttgctccct	ttttttttct	gtttgaacag	aataaatttt	9780

-continued

ggataacttg	ttactagtaa	aaaattttaa	aattgtctgt	gatatgttct	ttaaggtact	9840
acttctcgaa	ctttttccta	gaagtagctg	taacaggagg	agagcatatg	taccctaag	9900
gtatctgggg	tataggccca	tgtccaaaca	atatttcttt	taagtcttgt	gttgtatctt	9960
taagactcat	gcaatttaca	ttttattcca	tgatataact	attttaatat	taaaatttgt	10020
cagtgatatt	tcttaccctc	tctcttagga	aaatgtgcc	tgtttatact	ttggctttga	10080
gtgccccga	ggaacagaca	ctagagtttg	agaagcatgg	ttacacaggc	gtggcttccc	10140
ctgcagaaat	taagtacaga	ctatttcagt	gtaaagcaga	gaagttcttt	tgaaggggaa	10200
tctccagtga	agaaaggggt	cttcactttt	acttccattt	cctcttgagg	gtgacctca	10260
ttgtctcttg	taaaactccg	atatttttaa	catggctgtt	ttgctttcct	ctggttcttt	10320
ttaacatgag	tgagacagat	gatactttaa	aaagtaattt	taaaaaaaag	tgttaaaata	10380
tatggccata	atgcagaacc	ctatgctgtg	atctccttta	ccaaattgtt	gtgtttgtac	10440
ttttgtagat	agctttccag	tccagagaca	gttattctgt	gtaaaggtct	gactcaacaa	10500
gaaaagattt	ccctttaccc	aaagaatgcc	agtctttatt	tgctgggtcaa	taagcaggggt	10560
ccccaggaaa	ggggtaactt	tcaccaccct	ctaaccact	ggttattagt	aaactaatta	10620
agtagactta	tctcaagatg	aggaaactta	aaaccaagta	aaattctgct	tttactggga	10680
ttttattttt	tgaaaccaga	aacgtttact	taagttgact	actattaatg	aattttgggtc	10740
tctcttttaa	gtactcttct	taaaaatgtt	atcctactgc	tgagaagttc	aagtttgaga	10800
agtacaagga	ggaatagaaa	cttgagagat	tttctttttc	ttttagagcc	tcttctgtat	10860
ttagccctgt	aggaattttt	tttttcccc	aagattcttc	ttcgtgaaaa	ggaggagtgt	10920
ccttttgatt	gagttcttgc	aaatctcaca	acgactttat	tttgaacaat	actgtttggg	10980
gatgatgcat	gagtctgaaa	caacttcagt	tgtagctgtc	atctgataaa	attgcttcac	11040
aggaaggaa	atttagcacg	gatctagtca	ttattcttgt	tagattgaat	gtgttaatca	11100
taattgtaaa	caggcatgat	aattattact	ttaaaaactg	aaaacagtga	atagttagtt	11160
gtggagggtta	ctaaagcatg	atttttttta	aataaaaactt	tcagcatttt	gcaaatatgc	11220
atatggttta	ggatagaact	tccagaggta	gcatcacatt	taaattctca	agcaacttag	11280
taatacgagg	ctctgaaaaa	ctgggttaaag	ttactccaga	aatggccctg	ggtctgacag	11340
acattctaac	ttaaagatgc	atatgaagac	tttgaataaa	atcatttcat	atgaagacat	11400
tgaataaaat	catttcataa	aataagttag	gaaaaacaac	tactattgaa	ttcatcttaa	11460
tgtatgattt	taaaaatatg	tttagctaaa	aattcataga	catttgacaa	tttcgtttat	11520
atctcaaaaa	gttgacttac	ccaagttgat	cacaaaactg	atgagactgg	tggtggtagt	11580
gaataaatga	gggaccacc	atatttgaga	cactttacat	ttgtgatgtg	ttatactgaa	11640
ttttcagttt	gattctataa	actaccaatt	tcaaaattac	aatttcaagg	tgtaataagt	11700
agtggtatta	tcttgaaata	ggtctaaagg	gaacttttct	gttttaaaat	attcttaaac	11760
tatatgtgct	gattttgatt	tgcatttggg	tagattatac	tcttatgaat	cggggggctg	11820
ggtattgatt	caggttttcc	ttacctattt	ggtaaggatt	tcaaagtctt	tttgtgcttg	11880
attttctctg	tttttaaata	tgaaacatat	tgatgacttt	taattaacaa	atgtttttat	11940
ctcgaataaa	ttttaaagga	gatcttttct	aaaagaggta	tgatgactta	attattgcat	12000
ataacaataa	atgagaaacc	agtgattcca	tactctctaa	agaataaaaag	tgagcttttag	12060
gcccgagcat	ggtggctcat	gcctgtaatc	ccagcacttt	ggaaggccga	ggcaggcgga	12120
tcacctgagg	tcaggaattc	gacaccagcc	tgcccaaatg	gcaaaaccct	gtctctacta	12180

-continued

caaatacaaa	aattagctgg	gcatggtggc	agcccctata	gtcccagcta	cttggaagac	12240
tgagacagga	gagtcactcg	aacccgagag	gcagaggttg	cagtaagctg	aatcacacc	12300
attgcactcc	agcctgggca	acaagagcaa	aactccgtct	caaaaaaaaa	aaaaaaaaa	12360
aaaaagaata	aaagtgagct	ttggattgcg	tataaatcct	ttagacaagt	agtagacttg	12420
tttgatactg	tgtttgaaca	aattacaaag	tattttcatc	aaagaatgtt	attgtttgct	12480
gttattttta	ttttttattg	cccagcttct	ctcatattca	ttatgtgatt	ttcttcactt	12540
catgttactt	tattgtgcag	ggtcagagta	ttattccaat	gcttactgga	gaagtgattc	12600
ctgtaatgga	actgctttca	tctatgaaat	cacacagtgt	tcctgaagaa	atagatgtaa	12660
gtttttatat	ttttaaatga	gagcaattat	accctttatc	agtttttttg	ggttatatta	12720
ttattatgta	tattattaat	attctaattt	taatactaag	cacttcgtcg	tacgtactat	12780
ccacatgcag	tattagccac	ttgaacagat	aagcacacac	aaaatcctgg	attttatggc	12840
ataacagagg	catttttgat	cagtgatgac	aaaactaaat	ttattttgtt	tatttcacta	12900
cttttataat	tcctaaaagt	gggaggatcc	cagctcttat	aggagcaatt	aatatttaat	12960
gcagtacctt	ttgaaacaaa	actgtgtgcc	aaagcagtaa	ccattaatgg	aagttgactt	13020
atagtcacaa	atttagtttc	cttaatcatt	tgttgaggat	gttttgaatc	acacactatg	13080
agtgttaaga	gatatcttta	ggacactatt	cttgttgttt	tattgtcatt	taggttagtc	13140
tcctgtctga	cagctcagaa	gaggaagttg	ttcttgtaaa	aattgtttac	acaacctgat	13200
tgaccagctt	tcacatttgt	tcttctgaaa	gctgatggta	gtgcacagat	tgttttatgg	13260
ggagtcttga	ttctcagaaa	tgaaggcagt	gtgttatatt	gaatccagac	ttcagaaaac	13320
ttgtatatta	aaagtgtttt	ttcaacacta	tgttatagcc	agactaattt	ttttattttt	13380
ttgatgcatt	ttagatagct	gatacagtac	tcaatgatga	tgatattggg	gacagttgtc	13440
atgaaggctt	tcttctcaag	taagaatttt	tcttttcata	aaacctggat	gaagcatatg	13500
ttcacctatg	acaagatttg	gaaggaagaa	aataacagac	tgtctactta	gattgttcta	13560
gggacaacat	tgcatatttg	aattgttgct	taaatgttg	ttatttttca	ttcgttatat	13620
ttctataata	tatttgatgt	tattccattt	gctatttaaa	gaaactgagt	ttccatattt	13680
cccagacaag	aatcatggc	cccttgcttg	attctgggtt	cttgttttac	ttctcattaa	13740
agctaaaaga	accctttcaa	attaagttgt	actgtagatg	aacttaagtt	atttaggcct	13800
agaaaaaaaa	aattcatatt	tatactgatc	tttttccatc	cagcagtgga	gtttagtact	13860
taagagtttg	tgcccttaaa	ccagactccc	tgggttaatg	ctgtgtacct	gtgggcaagg	13920
tcctgaatt	ctctatacac	ctatttcctc	atctgtaaaa	tggcaataat	aataatagta	13980
cctaattgat	agagttgtta	taagcattga	gtaagataaa	taatataaag	cacttagaac	14040
agtgcctgga	acataagaac	acttaataat	agtaacatt	ttctatttac	atttcttcta	14100
aggaaaagg	taacagaaat	agccaatatt	tgttcagtgc	ctacatgtta	gttcctatac	14160
taagtgcctt	acatgtatta	tcttatattc	tattttaatg	tttcttcaca	gttgcagatt	14220
atcatgaaat	tttatttttt	aaaaagaga	agtaaaagga	taaagtattc	acttttatgt	14280
ccacagtctt	ttcctttagg	ctcatgatgg	agtatcagag	gcatgaatgt	gtttaaccta	14340
agagccttaa	tggcttgaat	cagaagcact	ttagtccgtg	atctgttcag	tgtcagcctt	14400
tcaaacatca	ttttaaatcc	catttgactt	taagtaaata	acttaatctc	tctacatgtc	14460
aatttcttca	gctataaaat	gatggtat	caataaataa	atacattaat	taaatgat	14520

-continued

tttacaaact	aattgggctg	ttttaaggct	caataagaaa	atttctgtga	aaggctctcta	14580
gcaaatgtag	ggttctatac	aaataaaaaga	taacattatg	cttatatctt	cggtgtttat	14640
catgcaaagc	tcttctgagt	tttttgaaga	gctcacctac	tattttttgt	ttttagtttg	14700
ttaaattggt	ttataggcaa	tgtttttaat	ctgttttctt	taacttacag	tgccatcagc	14760
tcacacttgc	aaacctgtgg	ctgttccgtt	gtagtaggta	gcagtgcaga	gaaagtaaat	14820
aaggtagttt	attttataat	ctagcaaatg	atttgactct	ttaagactga	tgatatatca	14880
tggattgtca	tttaaattgt	aggttgcaat	taaaatgatc	taatagtata	aggaggcaat	14940
gtaatctcat	cgaattgctg	agacaacttg	tggcaacagt	gagtttgaaa	taaagtgaat	15000
aggagtcatt	tatcagttta	ttttgataac	ttgtaaatac	cagtgtcaga	tgtgtataaa	15060
tggttttgag	aatatattaa	aatcaggtat	ttaaaaaac	actattcttc	tatttcccaa	15120
tgtaatcttt	aacaaatctg	aaggtagtca	tgtactttcg	gtactagttc	tgaagaaatg	15180
ttatttgttt	attcatcttg	atttcattgt	cttggctttc	cttctaaatc	tatcccttct	15240
tgggagctat	tgggattaag	tggtcattga	tgattatact	ttattcagta	atgtttctga	15300
ccctttcctt	cagtgtact	tgagttaata	aaggattaat	gaacagttac	atttccaagc	15360
attagctaata	aaactaaagg	attttgcact	tttcttcact	gaccattagt	taaaaacagt	15420
tcagagataa	gtacatgtat	ctttcaattc	tagcaaacct	aattttttta	aagaagtttt	15480
acataggaaa	tatgttggaa	atgattatct	actttacaaa	gatattcata	atttattttt	15540
tctgtaacta	gctactttgt	atatttacat	gagccttaat	ttatcaaaat	tatatttctc	15600
atataaccat	ttatgagagc	ttagtattcc	tctgtcatta	tattgcgtct	acggactagt	15660
gatcttacta	cttctgttac	ctcgaacaag	tggcttcccg	tctgtgacct	ccaaagccgt	15720
aggttccaca	gagtgtactgc	tgagctgctt	tatgaaggga	gaaaggctcc	atagttgggt	15780
ttttggtttt	gcttttgttt	ttgttttta	catttttctt	atcctccatc	ctcttgaggc	15840
agagtagctt	accttttatc	ttgtttta	ttgagaaaga	agttgccact	gctctagatt	15900
gaaaaccact	gctttaacat	aataactctg	aatatgggtt	gaatttcaag	atagtgcacat	15960
gcctttttat	ttttactaat	agagctgtag	gtcgaatatt	attagatttc	taaacccac	16020
ccaatgacct	ccttatttta	aatcaaattt	aataattaat	tatcttctta	ttggaggatc	16080
tggacattct	ttgatgtttc	ttacaatgaa	tttcacatgt	agaccacta	aacagaagtt	16140
ataaaggttc	catggtcaaa	taagtctgag	aaagtctgca	tattatataa	ttcacctaaa	16200
gagtcacagt	atgtacccaa	atgttaaagg	ttttgagatg	ccatacagta	aatttaccaa	16260
gcattttcta	aatttatattg	accacagaat	ccctatttta	agcaacaact	gttatatccc	16320
ataggttcca	ggtgactaaa	gaataactat	tgcttaggat	atgttttatt	gataataaca	16380
attaaaatgt	cagatatctt	tcataagcag	atcagtggtc	tttttaaaac	tttgatattt	16440
aatgctaaaa	tcttttcttt	tgtagatagt	cagaacatta	tgcttttttc	tgactgcagc	16500
agagagaaaa	tgctccaggt	tatgtgaagc	agaatcatca	tttaaataatg	agtcagggtc	16560
ctttgtacag	ggcctgctaa	aggtatagtt	tctacttatc	acaagggaaa	ccaattttct	16620
aaaatcattt	ttgagactct	ttgtagacaa	atattaaata	ttagcattta	atgtatctca	16680
tattgacatg	cccagtgact	gacttccttt	gcacagttct	gcgcatagac	tatatgtctt	16740
atggatttat	agttagtatc	atcagtgtaa	caccatagaa	taccctttgt	tttccagggtg	16800
ggtccttgta	cctacatgtc	tagcatcagg	tgttggtttt	tttttttttt	tttaaaacat	16860
atgcttaaat	caggttgac	atctaaaata	agatcatttc	tttttaacta	aatagatttg	16920

-continued

aattttattg	aaaaaaattt	taaaacatct	ttaagaagca	tataggattt	aagcagttac	16980
tatgtatgtg	tactaaaata	tatatatatt	cctaaatata	tattcctata	tataatatat	17040
gtatttctat	atataatata	tattagaaaa	aacttagagt	tttctttcat	ttgagtctac	17100
tgttcaagga	gcaaaacaga	gaaatgtaaa	ttagcaatta	tttacaataa	ttaaagggaa	17160
gaaagttggt	caccttggtg	gatctattat	tggtgtttta	attatagtcc	caagacgtga	17220
agaaatagct	ttcctaattg	ttatgtgatt	gtctcatagt	gactactttc	ttgaggatgt	17280
agccacagca	aatgaaatt	taaaaaattt	aaaaattggt	gcaaataaaa	gttatattag	17340
gcttttggtg	aatttcaata	atgtgctgct	atgaactcag	aatgatagta	tttaaataata	17400
gaaactagtt	aaaggaaaca	cagtttctat	ttgagttata	caaactctga	aattagaact	17460
tctcctgtta	aggcattata	aagtgcctaa	tacttttggt	tcctcagcac	cctctcattt	17520
aattatataa	ttttagctct	gaaagggacc	tataccagat	gtgtagagga	aatttcaaaa	17580
ctatgatcta	atgaaaaaat	atttaatagt	tctccatgca	aatacaaatt	atatagtttt	17640
ctggaaaata	cctttgacat	tatacaaaga	tgattatcac	agcattataa	tagtgaaaaa	17700
atggaaatag	cctctttctt	ctgttctggt	cacagcatat	ggcacagtac	ctcatatgca	17760
gtaggttatt	atgacctggt	aactggctcc	cccaactgat	taggaaagaa	gtaaatttgt	17820
tatttataaa	aatacgtggt	cattgagatg	catagaataa	ttaagaaatt	aaaagacact	17880
tgtaatttca	aatccagtga	atacccactg	ttaatatattg	gcatactctt	ttctagtctt	17940
tttttccctt	ttgcatgtat	tttctttaag	actcccaccc	ccactggatc	atctctgcat	18000
attcetaatct	gcttttttca	cagcagattc	taagcctttt	tgcatatcaa	cacaaacttc	18060
aacaacttca	tcttttagatg	ctaaataatg	aattcatttt	tatttactta	accactttct	18120
ttggatgctc	aggttattct	gatgttttgc	cattaaaacc	aatgctatac	tgaacacttc	18180
tgtcactaaa	acttgaacac	actcatgaat	aatttcttag	gataaatttt	tagagatgga	18240
tttgctaaat	caaagaccat	tttttaaaaa	ttgaaaaaca	attatatcgt	ttggcatgta	18300
agacagtaca	ttttcctttt	attttgacag	gattcaactg	gaagctttgt	gctgcctttc	18360
cggcaagtca	tgtatgctcc	atatcccacc	acacacatag	atgtggatgt	caatactgtg	18420
aagcagatgc	caccctgtca	tgaacatatt	tataatcagc	gtagatacat	gagatccgag	18480
ctgacagcct	tctggagagc	cacttcagaa	gaagacatgg	ctcaggatac	gatcatctac	18540
actgacgaaa	gctttactcc	tgatttgtac	gtaatgctct	gcgtgctggt	actgtagtca	18600
agcaatatga	aactgtgtct	tttatgaata	aaaacaaaac	agaagttgca	ttcaaaaaga	18660
aagaaatatt	actagcagaa	ttatgcttga	agaacatttt	aatcaagcat	ttttttctta	18720
aatgttcttc	tttttccata	cgattgtggt	taccctaaaa	taagtaagat	taacccttaa	18780
agtgaatatt	taactatttg	tttaataaat	atatattgag	ctcctaagca	ctgttctagg	18840
tactgggctt	aatagtggct	aaccacacag	ctccagcccc	tacattgcat	atagtctatt	18900
gtataagtta	ctgaatggac	ttactaacia	aaccagagaa	gtaattctaa	gtcttttttt	18960
tcttgacata	tgaatataaa	atacaaaaa	actggtaaaa	tatattaata	gagcattctt	19020
ttactttgca	ttttatattg	ttactcactt	cgtattttaag	aaaaacagtc	tgatcaggaa	19080
attcaaaagg	aaaagtaatg	ataattaatt	gagcatagac	ccaacttgaa	aagaaaaaaa	19140
aggatgatga	taaatctata	atcctaaaa	cctaagtaaa	cacttaaatg	atgttctgaa	19200
atcaggaaaa	gaattatagt	atatttttgt	ggttctcttt	tattagttga	aaaaaggcac	19260

-continued

agtagctcat	gcctataaga	acagagcttt	gggattccaa	ggcaggcaga	tcaattgagg	19320
ccaggagttc	cagaccagcc	tgggcaacat	agtgaacc	catctctaca	aaaataaaaa	19380
agaattagtt	gaatgtgttt	ctgtgtgcct	ataatcctag	ctattcagaa	agctgaggca	19440
ggaggatctc	ttgagcccag	gagtttgagg	ttacatggag	ttatgatgtg	ccagtgtact	19500
ccagcctgcg	ggacaatgag	actctgtctt	gttaaaaaaa	aaagtgcctg	gaataatggt	19560
tggcatatag	aaggtaacaa	cagtaaagt	taactgtaat	aaccaggt	taagtgtgta	19620
aggtgataga	aaaattgggg	caaacaacc	tgacctgtgt	ctctacagaa	taagtttgag	19680
ttgaggcaac	agacatgtgg	agcaccagta	attacacact	aaatgttaac	caaagcggt	19740
gaatagtaac	atcttattca	agggacccc	agccttatat	atctcaaggt	gcagaaagat	19800
gacttaatat	aggaccatt	ttttccgagt	tctccagagt	ttttattggt	tcttgagaaa	19860
gtagtggggg	aattgtttta	gaaaatgaat	tggcacaact	gaaattccat	gtcagtaagt	19920
ttttacatat	tggtaaattt	tgatagacat	gtagaagttt	tctaattaat	ctgcgccttg	19980
aaacattttc	ctttttccta	aagtgcctag	tattttttcc	cttttttgat	tggttgcttg	20040
ggagcttttt	tgaggaaatt	tagtgaactg	cagaatgagt	ttgcaaccat	ttagtatttt	20100
tgttttgtgt	tttagaggag	gtatgtgtat	tttaacattt	cttaatcatt	tttagccagc	20160
tatgtttgtt	ttgctgattg	acaaactata	attaaacagc	tattctcatt	ttgctgatca	20220
tgacaaagta	atatcctgaa	tttttaaatt	ttgcatccag	ctctaaattt	tctaaatttt	20280
ctaaacataa	aattgttcaa	aaaatagtat	ttttagccac	tagattgtgt	gttggttaagt	20340
ctgttgtcac	agactcattt	tacttttcag	tgtgtgtttt	tacatgttaa	ttatgtttgt	20400
catttttaat	tttaactttt	taaaataatt	ccagtcactg	ccaaaacatg	aaaaattggg	20460
cactggaaat	ttttttttta	acttttat	taggttcactg	tgtacatgtg	caggtttggt	20520
atacaggtaa	attgcgtgtc	gtgaggggtt	ggtgtaccca	ggtaataagg	gtagtaccca	20580
ataggtagtt	ttttgatcct	tacccttctc	ccacccttct	ttcaccctcg	agtaggcctt	20640
ggtgttgctg	tttccttctt	tgtgtccatg	tgtactcaat	ggtagctcc	tacttagaag	20700
tgagaacatg	cgggtatttg	ttttctgttc	ctggattagt	tcaactcagga	taatggcctc	20760
tagctccatc	tgttttttat	ggctgcatag	tattccatgg	tgtatatgta	tcatgttttc	20820
tttatccagt	ctaccattga	tagacattta	ggttgattct	ctgtctttac	tatcatgaat	20880
agcgtgtga	tgaacatata	cacatgcatg	tgtccttatg	gtggaacaat	ttgtattcct	20940
ttaagtatat	acagaataat	ggggttgcta	gggtgaatgg	tagttctatt	gtaagttatt	21000
tgtgaaatct	tcaaactgct	tttcacaata	gctaaactaa	tttacagtcc	caccagcagt	21060
gtataagtgt	tcccttttct	ccacaacctt	gccaaacatg	tattttttta	cttttcaata	21120
ataggcattc	ctagagaatt	gatttgcaat	tctctaata	ttagtatat	tgagcatttt	21180
ttcgtatgct	ttttagctgt	gtgtatatat	tcttttgaaa	aatgttaatg	tcctttgccc	21240
agtttgtaat	ggggttggtt	gtttttgctt	gttaattaaa	gttccttcca	gattctggat	21300
atccctttgt	cagatgcgtg	gtttgcagat	atttttctcc	ccttggtgtag	gttgtctttt	21360
tactctgttg	atagtttctt	ttgccgggca	ggagctcatt	aggtctcatt	tgtgtttgtt	21420
tttgttgcag	ttgcttttg	cgtcttcac	ataaaatctg	tgccagggcc	tatgtccaga	21480
atggtatttc	ctagtttgct	ttccagggtt	tttacaattt	tagattttac	gtttatgtct	21540
ttaatccgtc	ttgagttgat	taaggaaggg	gtccagtttc	actctaattc	ctatggctaa	21600
caattatccc	agcaccattt	attgaatacg	gagtcctttc	cccattgctt	gtttttgtca	21660

-continued

at t t t t g t t g a	a g a t c t g a t g	g t t g t a g g t g	c t a t g t g g c t	t t a t t t c t t g	g c t c t c t a t t	21720
c t c c a c t g g t	c t g t c t g t t t	t t a t a c c a g t	a c c c t g c t g t	t a a g g t t c c t	a t a g c c t t t t	21780
a g t a t a a g g t	c g g c t a a t g t	g a t g c c t c c a	g c t t t g t t c t	t t t t g c t t a g	g a t t g c t t t g	21840
g c t a t t t g g g	c t c c t t t t t g	g t t c c a t a t t	a a t t t t a a a a	t a g t t t t t t c	t a g t t t t g t g	21900
a a g a a t g t c a	t t g a t a g t t t	a g a g g a a t a g	c g t t g a a t c t	g t a g a t t g c t	t t g g g c a a a t	21960
g g c c a t t t t a	a c a a t a t t g a	t t c t t c c t a t	c t a t g a a c a t	g g a a t g t t t t	t c c a t g t g t t	22020
t g t g t c a t c t	c t t t a t a c c t	g a t g t a t a a a	g a a a a a c c a g	t a t t a t t g c t	a c t c a a t c t g	22080
t t c c a a a a a a	t t g a g g a g g a	g g a a c t c t t c	c c t a a t g a g a	c c g g c t t c c t	t c t g a t a c c a	22140
a a a c c t g g c a	g a g a t a c a a c	a g a a a a a g a	a a a c t t c a g g	c c a a t a t c c t	t g a t g a a t a t	22200
a g a t g c a a a a	a t c c t c a a c a	a a a t a c t a g c	a a c c a a a t c c	a g c a g t a c g t	c a a a a a g c t a	22260
a t c t a c t t t t a	a g t a g g c t t t	a t c c c t g g g a	t g c a a g g t t g	g t t c a a c a t a	c a c a a a t c a a	22320
t a a g t g t g a t	t c a t c a c a t a	a a c a g a g c t a	a a a c a a a a a	c c a c a a g a t t	a t c t c a a t a g	22380
g t g c a g a a a a	g g c t t t c a a t	a c a a t t t a a c	a t c c t t c a t g	t t a a a a c c t	t c a g t a g g t c	22440
a g g t g c a g t g	a c t c a c a c c t	g t a a t c c c a g	c a c t t t g g g a	g g c c a a g g c g	g a c g t a t a t c	22500
t t a a g c c c a g	g a g t t c a a g a	c c a g c c t a g g	c a g c a t g g t g	a a a c c c c a t c	t c t a c a g g a a	22560
a a a a a a a a a a	a a a a a a a a a a	a a a a a a a a a a	a g c t t a a t a t	g g c g g c a t g c	a c c t a t a g t c	22620
c c a g c t a c t c	a g g a g g t t g a	g g t g g g a g g a	t t g c t t g a g c	c c a g g a g g c a	g a g g t t g c a g	22680
c g a g c t g a g a	t c g t g c c a c t	g c a c t c c a a c	c t g g g c a a t a	a a g t g a g a c c	c t g t c t c a a a	22740
a a g a a a a c a	a a a a t a a t c c	t a a a c c a a c t	a g g c a t t g a a	g g a a t a t g c c	t c a a a a a a a t	22800
a a g a a c c a t c	t a t g a c a g a c	c c a c a g c c a a	t a t c t t a c c a	a a t g g g c a a a	a g c t g g a a g t	22860
a t t c t c c t t g	a g a a c c g t a a	c a a g a c a a g g	a t a c a c a c t c	t c a t c c c t c c	t t t t c a g c a t	22920
a g t t c t g g a a	g t c c t c g c c a	g a g c a g t c a g	g a a a g a g a a a	g a a a g a a a a g	g c a t t c a g a t	22980
a g g a a g a g a a	g a a g t c a a a c	t a t t t c t g t t	t g c a g g c a g t	a t a a t t c t a t	a c c t a g a a a a	23040
t g c c a t a g t t	t c t g c c c a g a	a g c t c c t a c a	t c t g t t a a a a	a t t t c a g c a a	a g t t t t a g c a	23100
t t g t c t g t a t	t c c a a c a g c t	t c c a g g g t g a	g a g t g a a a t c	a g g a a c a c a g	t c c c g t t c a c	23160
a a t a g c c g c a	a a a a g a a t a a	a a t a c c t t g g	a a t c c a g c t a	a c c a g g g a g g	t g a a a c a t c t	23220
c t a c g a g a a t	t a c a a a a c g c	t g c t g a a a g a	a a t c a g a g a t	g a c a c a a a c a	a a t g g a a a t g	23280
t t g t t t t t a a	c a c c t t g c t t	t a t c t a a t t c	a c t t a t a a c t	a a g a t a t t c a	t t c a g t g g a a	23340
c a g g t a t a a t	a a g a c c a c t c	g a c t t a a a t a	t a a g c c t t a t	t c t c t t t t c c a	g a g c c c a a g a	23400
a g g g g c a c t a	t c a g t g c c c a	g t c a a t a a t g	a t a a a a t g c t	g a t a t t t t t c	c c c t t t a c t g	23460
t t t c t t t c t t	c t g t a g t g t g	g t a c a c t c a t	t t c t t a a g a t	t a g a a a a c t t	g a c c t a c c t t	23520
c c t g t t t g c t	t c t a c a c a c c	c c c a t t c t c t	t t t t t t g c c a	c t c c g g t c a g	g t a t a g g a t g	23580
a t c c c t a c c a	c t t t t a g t t a	a a a c c t c c t t	c c c t t a t t a a	a t g t t c t c t t	a c c a c t c t g g	23640
c c t g a g t a g a	a c c t a g g g a a	a a t g g a a g a g	a a a a g a t g a a	a g g g a g g t g g	g g g c t g g g a a	23700
g g g a a t a g t c	t t g t t t g t g t	g t t t g c t t t a	g c a c c t a c t a	t a t c c t a g g t	g c t g t g t t a g	23760
g c a c a c a t t a	t t t t a a g t g g	c c a t t a t a t t	g c t a c a t c t c	a c t c t g g t c a	t t g c c a a g g t	23820
a g g t a g t a c t	t t c t t g g a t a	g t t g g t t c a t	g t t a c t t a t a	g g t g g t g g a c	t t g t t g a g g c	23880
a a c c c c a a t g	g a t a a t c a t c	t g a g t g t g t t	c t c t a a t c t c	a g a t t t t t c t	t c a t a t t t t t	23940
t g g t t t g t t t	t g g t t t t t g a	t g g t g g t g g t	t g t g t g c t t a	t t t t t g t t g c	t g g c t t g t t t	24000

-continued

ttttgttttg	tttttgatat	ggcaagaatt	ggtagtttta	tttattaatt	gcctaagggt	24060
ctctactttt	tttaaaagat	gagagtacta	aaatagattg	ataggtacat	acataccttt	24120
atggggggact	gcttatattc	cttagagaaa	aaaattactt	attagcctga	caaacaccag	24180
taaaatgtaa	atatatccgt	gagtaaataa	atgaatgtat	gctttgtatc	tccaaatata	24240
tacatctata	ttcttacaaa	tatgttttta	tgtaatacca	atttataaga	acttaaaatg	24300
ttggctcaag	tgagggatgg	tggaaagtag	cattatatag	ccatttcaac	atttgaactt	24360
ttttcttcat	tttcttcttt	tcttcaggaa	tatttttcaa	gatgtcttac	acagagacac	24420
tctagtgaag	gccttcctgg	atcaggtaaa	tgttgaactt	gagattgtca	gagtgaatga	24480
tatgacatgt	tttctttttt	aatatatctt	acaatgcctg	ttctctctct	ctatatatat	24540
atatttatat	atttccttgg	atcatgcccc	agagttctgc	tgagcaattg	cagttaagtt	24600
agttacacta	cagttctcac	aagagtctgt	gaggggatgt	caggtgcac	attacattgg	24660
atgcctcttg	tcctagattt	atgtttcggg	aattcagacc	tatgtttaca	atataataaa	24720
tattgttgct	gccttttaca	gataaaataa	taagatataa	acttgaccac	aactactggt	24780
ttttgaaaca	tagagttcat	ggtttacatg	tatcaaagtg	aaatctgagt	tagcttttac	24840
agatataata	tatacatata	tatatcctac	aatgcttgta	ctatatatgt	agtacaagta	24900
tatatatgtg	tgtgtgtgtg	tgtatatata	ttatggcact	gtagtatata	tatgtttata	24960
tgttaaaaaa	tatataaata	tatgtttacat	atttaacata	aacatatata	catatatggt	25020
aaatatataa	catatactct	atatatgaca	aatagagtat	aatatatatt	tttatttttt	25080
atatatatat	aaaacatgat	agaattaaga	attaagtcct	aatctgtttt	attaggtgct	25140
ttttgtagtg	ttcagtcttt	ctaaagtgtc	taaatgattt	ttccttttga	cttattaatg	25200
gggaagagcc	tctatattaa	caattaaggc	tgcagcattg	attacttcaa	acaacaaaca	25260
ttttaattca	agcattaacc	tataactcaa	gtaagttttt	tttttttttt	ttttgagaaa	25320
gggaggttgt	ttatttgcct	gaattgagtc	aaaaatattt	ttgaaacatc	atgtactcat	25380
ttaaatagata	acatctttat	tgtttcattc	ttttaaaaaa	tatctactta	attacacagt	25440
tgaaggaaat	tgtagattat	atggaactta	tttcttaata	tattacagtt	ttgttataat	25500
aacattctgg	ggatcaggcc	aggaaactgt	gtcatagata	aagctttgaa	ataatgagat	25560
ccttatgttt	actagaaatt	ttggattgag	atctatgtgg	tctgtgacat	attgcaaagt	25620
tcaaggaaaa	ttcgtaggca	tggaatttct	caaactgaaa	atccctccca	ctgtccacct	25680
catcacatgc	acacattcta	ctcttaccca	cccactccac	cccttgcaaa	agtacagata	25740
tatgaatgtc	tcaaaaccat	gggctcatct	tctagaagct	tcaatgttat	ttgaagattt	25800
gggcagagga	agttaagaaa	tatgaaatag	cttacatatg	agttttaata	gtgaaacaaa	25860
catggatgta	ttctgaagta	gaatgcaaaa	tttgagtgca	tttttttttt	tttgagactg	25920
agtctggctc	tgtcgcccag	gctggagtgc	agtggccgga	tctcagctca	ctgcaagctc	25980
cacctcccgg	gtttacgcca	ttctcctgcc	tcagcctccc	gagtagctgg	gaccacaggc	26040
ggccgccact	tcgcccggct	agtttgtttg	tatttttttag	tagagatggg	gtttcaccgt	26100
gtagccagg	gannnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnccatt	26160
gagtgcattt	ttaaagataa	atcagaaaac	ttcgaaaaac	tatcagattg	gccggacatg	26220
gtggcttatg	cctgtaatcc	tagcactttg	ggaggctgag	gtgggtggat	cacgagggtca	26280
ggagatcgag	accatcctgc	caacatggtg	aaaccccatc	tctactaagt	atacaaaaat	26340
tagctgggcg	tgacagcacg	tgcttgtaat	cccagctact	tgggaggctg	aggcaggaga	26400

-continued

atcgcttgaa	cccgggaggt	ggagggttgca	gtgagtcaag	atcacaccac	tgcacttcag	26460
cttggtgaca	gagctagact	ccatatcaaa	aaaaaaaaaa	aaaaaaagaa	gtcagattgt	26520
tcctacaccc	agtgccttcta	taccacactc	ctactagggg	gcatcagtgg	aatgggttaa	26580
ggagatgttt	agtgtgtatt	gtctgccaa	gactgttaac	actgtcctag	aaacattgct	26640
gtacaagtag	aatgtgagca	aattatgtat	tgaaatgggt	cctctccctg	cagggtctttc	26700
agctgaaacc	tggcttatct	ctcaggagta	ctttccttgc	acagttttta	cttgtccttc	26760
acagaaaagc	cttgacacta	ataaaatata	tagaagatga	tacgtgagta	caactcctac	26820
atggaggaaa	aaccttttgt	acgttggttt	ttgttttatt	tcctttgtac	attttctgta	26880
tcataatttt	tgtttttttt	tttttttttt	ttttctccat	tactttcagg	cagaagggaa	26940
aaaagccctt	taaatctctt	cggaacctga	agatagacct	tgatttaaca	gcagagggcg	27000
atcttaacat	aataatggct	ctggctgaga	aaattaaacc	aggcctacac	tcttttatct	27060
ttggaagacc	tttctacact	agtgtacaag	aacgagatgt	tctaatagact	ttttaaatgt	27120
gtaacttaat	aagcctattc	catcacaatc	gtgatcgctg	ctaaagtagc	tcgggtggtg	27180
ggggaaacat	tcccctggat	catactccag	agctctgctc	ggcagttgca	gttaagttag	27240
ttacactaca	gttctcacia	gagtcctgtg	ggggatgtca	ggtgcatcat	tacattggat	27300
gtctcttttc	ctagatttat	gcttttgagg	tacagacctc	tgtttacaat	ataataggta	27360
ttattgctgt	cttttaaata	tataataata	ggatataaac	ttgaccacia	ctgctgtttt	27420
tttgaaatat	atgattcatg	gtttacatgt	attaagggtg	aatccgagtt	cgtttttaca	27480
gatattagtt	gactttctat	cttttgcat	tctttgggtg	gtggaattac	tgtaatactt	27540
ctgcaatcaa	ctgaaaatta	gagcctttta	atgatttcag	ttccacagaa	agaaagttag	27600
cttcaacata	ggataagctt	tagaaagaga	attgatcaag	cagatgttta	attggaattg	27660
attattagat	cctgctttgt	ggatttagcc	ctcgggatcc	agtctgtaga	aatgtctgat	27720
agttctctat	agtccttgct	catgggtgaac	cacagttagg	atgttttggt	tgttttattg	27780
ttgttgctat	tgttgatgtt	ctatatagtt	gagctctata	aaaggaaatt	gtattttatg	27840
ttttagtagt	tgttgccaac	tttttaaatt	aattttcatt	atttttgagc	caaattgaaa	27900
tgtgcacctc	ctgtgccttt	tttttccttg	gaaaatcgaa	ttacttgga	gaagttcaga	27960
tttcactggg	cagtcgtttt	catcttggtt	tcttcttgca	gagtcctacc	atgtacctgc	28020
tttggaatac	attgtaactc	tgagattata	aaatgcatta	gagaatatat	taactaataa	28080
gatctttttt	ttcaggaaca	gaaaatagtt	ccttgagtac	ttccttctta	catttctgcc	28140
catgtttttg	aagttgttgc	catttgccctg	caataggcta	taagggaatg	caggagaaat	28200
tttactgaag	tgctattttt	ctaggtgcta	ctttggcaga	gctaagtggg	ctgtttcttt	28260
tgtttcctta	atgcgtttgg	accattttgc	tggctgtaaa	ataactgatt	aatataattc	28320
taacacaata	ttgacattgt	agtgtacaca	aacacaaaata	ttttatttaa	aactggaagt	28380
aacataaaaag	ggaaaatata	tttataagaa	aggaataaag	gtaatagagc	tcttctgtcc	28440
cccagccacc	aaatttacac	aacaaaatca	tatgttctaa	tgtgaaaggt	cataatagct	28500
ttcccatcat	taatcagaaa	gatgtggcag	cttgattttt	tagacaaccc	ctgaactaga	28560
tgactgttgt	actgtagctc	agtcatttaa	aaaatatata	aatactatct	cgtagtgtcc	28620
catactatgt	tttttacatg	atagattctt	atttaagtgc	taactgggta	ttttctttgg	28680
ctgggtttatt	gtactgttat	atagaatgta	agttgtacag	tgaaataagt	tattaaagca	28740

-continued

tgtgtaaaca ttgttatata tcttttctcc tagatggaga attttgaata aaatatnnnn	28800
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	28860
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	28920
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	28980
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	29040
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	29100
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn	29160
nnnnnnnnnn nnnnnnnnnn nnnnnnagaa gactaattga tcatatcact atgattctca	29220
aagaagaacc aaaacttcat ataatactac aaatatgaga tagttacttc tgtagtatat	29280
ttctgtaatg ctacaggta aacaggtcac tcttatataa cactattttg attttgatgt	29340
agaattgcac aaattgatat ttcttctatg atctgtaggg tatagcttaa agtagcaaaa	29400
acagtccacc acctccagtt aacacacagt aacactatgg gactagtatt attatttcca	29460
ttttacaaag gaggaaacta aagcttaaag atgtgtaata tacagcccaa ggtcacacag	29520
ctggtaaagg tagatttcat ccagacagt tacagtcatt gccgtgggca cagctcctaa	29580
cttattaact ccatgtaact ggtactcagt ttagttgaat tgaaaggaga gtagggaagc	29640
aggtctgttt gcactattca gagcccaagt gtgaatccct gctgtgctgc ttggagaagt	29700
tacttaacct atgcaagggt cattttttaa atatttgaaa cggaatgata atacatactt	29760
caccagtggg tttaatgaga ccttataaga tcgttagtte agtacctgac cagtgttca	29820
taaatgcttt ttcattcaat ctgacaatct ctagcttgta attggggcat ttagaacatt	29880
taatatgatt attggcatgg taggttaaag ttgtcatctt gctgttttct ctttgttctt	29940
ttttctcctt tcttttgat ttttttttaa ttttactgtg tcttctctgt tgtcttatta	30000
a	30001
<210> SEQ ID NO 12	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 12	
gggtctagca agagcaggtg	20
<210> SEQ ID NO 13	
<211> LENGTH: 21	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 13	
gtcttgga cagctggaga t	21
<210> SEQ ID NO 14	
<211> LENGTH: 24	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Probe	
<400> SEQUENCE: 14	
tgatgacgac tctttgccca ccgc	24

<210> SEQ ID NO 15 <211> LENGTH: 22 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Primer <400> SEQUENCE: 15 tgtgacagtt ggaatgcagt ga		22
<210> SEQ ID NO 16 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Primer <400> SEQUENCE: 16 gccacttaaa gcaatctctg tcttg		25
<210> SEQ ID NO 17 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Probe <400> SEQUENCE: 17 tcgactcttt gccaccgcc a		21
<210> SEQ ID NO 18 <211> LENGTH: 783 <212> TYPE: DNA <213> ORGANISM: Homo sapiens <400> SEQUENCE: 18 gcctctcagt acccgaggct cccttttctc gagccgcag cggcagcgct cccagcgggt		60
ccccgggaag gagacagctc ggggtactgag ggcgggaaag caaggaagag gccagatccc		120
catcccttgt ccctgcgccg ccgccgccgc cgccgcgcc gggaagcccg gggcccgat		180
gcaggcaatt ccaccagtcg ctagaggcga aagcccgaca cccagcttcg gtcagagaaa		240
tgagagggaa agtaaaaatg cgtcgagctc tgaggagagc ccccgcttct acccgcgctt		300
cttcccggca gccgaacccc aaacagccac ccgccaggat gccgcctcct cactcaccca		360
ctcgccaccg cctgcgcctc cgccgcgcgc ggcgcaggca ccgcaaccgc agccccgccc		420
cgggcccgcc cccgggcccg ccccgaccac gccccggccc cggccccggc cccggccccg		480
gcccctagcg cgcgactcct gagttccaga gcttgetaca ggctgcggtt gtttccttcc		540
ttgttttctt ctggttaatc tttatcaggt cttttcttgt tcaccctcag cgagtactgt		600
gagagcaagt agtggggaga gaggggtggga aaaacaaaaa cacacacctc ctaaaccac		660
acctgctctt gctagacccc gccccaaaaa gagaagcaac cgggcagcag ggacggctga		720
cacaccaagc gtcacttttt acgtgggcgg aacttgtcgc tgtttgacgc acctctcttt		780
cct		783
<210> SEQ ID NO 19 <400> SEQUENCE: 19		
000		

-continued

<210> SEQ ID NO 20	
<400> SEQUENCE: 20	
000	
<210> SEQ ID NO 21	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 21	
gccttactct aggaccaaga	20
<210> SEQ ID NO 22	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 22	
ccggccccta gcgcgcgact	20
<210> SEQ ID NO 23	
<211> LENGTH: 19	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 23	
cggccccctag cgcgcgact	19
<210> SEQ ID NO 24	
<211> LENGTH: 19	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 24	
ccggccccta gcgcgcgac	19
<210> SEQ ID NO 25	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 25	
ccctagcgcg cgactcctga	20
<210> SEQ ID NO 26	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 26	
cccctagcgc gcgactcctg	20

-continued

<210> SEQ ID NO 27 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 27		
gcccctagcg cgcgactcct		20
<210> SEQ ID NO 28 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 28		
ggcccctagc gcgcgactcc		20
<210> SEQ ID NO 29 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 29		
cgggcccctag cgcgcgactc		20
<210> SEQ ID NO 30 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 30		
cctagcgcgcg gactcctg		18
<210> SEQ ID NO 31 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 31		
ccctagcgcg cgactcct		18
<210> SEQ ID NO 32 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 32		
cccctagcgcg gcgactcc		18
<210> SEQ ID NO 33 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		

-continued

<400> SEQUENCE: 33		
gcccctagcg cgcgactc		18
<210> SEQ ID NO 34		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 34		
ggcccctagc gcgcgact		18
<210> SEQ ID NO 35		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 35		
cggcccctag cgcgcgac		18
<210> SEQ ID NO 36		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 36		
aggctgcggt tgtttccctc		20
<210> SEQ ID NO 37		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 37		
caggctgcgg ttgtttccct		20
<210> SEQ ID NO 38		
<211> LENGTH: 19		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 38		
ggctgcggtt gtttcctc		19
<210> SEQ ID NO 39		
<211> LENGTH: 19		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 39		
aggctgcggt tgtttccct		19
<210> SEQ ID NO 40		
<211> LENGTH: 19		

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 40

caggctgcgg ttgtttccc 19

<210> SEQ ID NO 41
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 41

gctgcggttg tttccctc 18

<210> SEQ ID NO 42
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 42

ggctgcggtt gtttcct 18

<210> SEQ ID NO 43
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 43

aggctgcggt tgtttccc 18

<210> SEQ ID NO 44
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 44

caggctgcgg ttgtttcc 18

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 45

tctgtctttg gagcccaaat 20

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 46

-continued

ctgcgatccc cattccagtt	20
<210> SEQ ID NO 47	
<211> LENGTH: 18	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 47	
gccttactct aggaccaa	18
<210> SEQ ID NO 48	
<211> LENGTH: 18	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 48	
tctgtctttg gagcccaa	18
<210> SEQ ID NO 49	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 49	
ggttaatctt tatcaggtct	20
<210> SEQ ID NO 50	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 50	
tggttaatct ttatcaggtc	20
<210> SEQ ID NO 51	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 51	
ctggttaatc tttatcaggt	20
<210> SEQ ID NO 52	
<211> LENGTH: 18	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 52	
gttaatcttt atcaggtc	18
<210> SEQ ID NO 53	
<211> LENGTH: 18	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 53

ggttaatcctt tatcaggt 18

<210> SEQ ID NO 54
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 54

tggttaatct ttatcagg 18

<210> SEQ ID NO 55
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 55

ctagcgcgcg actcctga 18

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 56

gggacactac aaggtagtat 20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 57

tacaggctgc ggttgtttcc 20

<210> SEQ ID NO 58
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 58

cccggccct agcgcgcgac 20

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 59

ggtaacttca aactcttggg 20

-continued

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 60


aatctttatc aggtcttttc

20

What is claimed is:

1. A modified oligonucleotide according to the following formula:

(SEQ ID NO: 49)



or a salt thereof.

181

2. The modified oligonucleotide of claim 1, which is a sodium salt of the formula.

3. A pharmaceutical composition comprising the modified oligonucleotide of claim 1 and a pharmaceutically acceptable diluent or carrier.

4. The pharmaceutical composition of claim 3, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

5. The pharmaceutical composition of claim 3, wherein the pharmaceutical composition consists essentially of the modified oligonucleotide and phosphate-buffered saline (PBS).

182


6. A pharmaceutical composition comprising the modified oligonucleotide of claim 2 and a pharmaceutically acceptable diluent or carrier.

7. The pharmaceutical composition of claim 6, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

8. The pharmaceutical composition of claim 6, wherein the pharmaceutical composition consists essentially of the modified oligonucleotide and phosphate-buffered saline (PBS).

9. A modified oligonucleotide according to the following formula:

(SEQ ID NO: 49)



183

10. A pharmaceutical composition comprising the modified oligonucleotide of claim 9 and a pharmaceutically acceptable diluent or carrier.

11. The pharmaceutical composition of claim 10, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

12. The pharmaceutical composition of claim 10, wherein the pharmaceutical composition consists essentially of the modified oligonucleotide and phosphate-buffered saline (PBS).

13. A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide is a gapmer consisting of a 5' wing segment, a central gap segment, and a 3' wing segment, wherein:

the 5' wing segment consists of six 2'-O-methoxyethyl nucleosides,

the central gap segment consists of β -D-deoxyribo-nucleosides, and

the 3' wing segment consists of four 2'-O-methoxyethyl nucleosides;

wherein the modified oligonucleotide has the nucleobase sequence 5'-GGTTAATCTTTATCAGGTCT-3' (SEQ ID NO: 49), wherein each cytosine is a 5-methylcytosine; and wherein the internucleoside linkages of the modified oligo-

184

nucleotide are, from 5' to 3', soooooosssssssoss, wherein each s is a phosphorothioate linkage and each o is a phosphodiester linkage.

14. The compound of claim 13, comprising the modified oligonucleotide covalently linked to a conjugate group.

15. A pharmaceutical composition comprising the compound of claim 13 and a pharmaceutically acceptable diluent or carrier.

16. The pharmaceutical composition of claim 15, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

17. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition consists essentially of the compound and phosphate-buffered saline (PBS).

18. A pharmaceutical composition comprising the compound of claim 14 and a pharmaceutically acceptable diluent or carrier.

19. The pharmaceutical composition of claim 18, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

20. The pharmaceutical composition of claim 18, wherein the pharmaceutical composition consists essentially of the compound and phosphate-buffered saline (PBS).

* * * * *