



US009605263B2

(12) **United States Patent**
Rigo

(10) **Patent No.:** **US 9,605,263 B2**
(45) **Date of Patent:** **Mar. 28, 2017**

(54) **COMPOSITIONS FOR MODULATING
C9ORF72 EXPRESSION**

(71) Applicant: **Ionis Pharmaceuticals, Inc.**, Carlsbad,
CA (US)

(72) Inventor: **Frank Rigo**, Carlsbad, CA (US)

(73) Assignee: **Ionis Pharmaceuticals, Inc.**, Carlsbad,
CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/130,818**

(22) Filed: **Apr. 15, 2016**

(65) **Prior Publication Data**

US 2016/0304871 A1 Oct. 20, 2016

Related U.S. Application Data

(60) Provisional application No. 62/239,400, filed on Oct.
9, 2015, provisional application No. 62/232,941, filed
on Sep. 25, 2015, provisional application No.
62/148,691, filed on Apr. 16, 2015.

(51) **Int. Cl.**
C07H 21/04 (2006.01)
C07H 21/02 (2006.01)
C12N 15/113 (2010.01)

(52) **U.S. Cl.**
CPC **C12N 15/113** (2013.01); **C12N 2310/11**
(2013.01); **C12N 2310/315** (2013.01); **C12N**
2310/322 (2013.01); **C12N 2310/3341**
(2013.01); **C12N 2310/341** (2013.01); **C12N**
2320/30 (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,801,154 A 9/1998 Baracchini et al.
5,998,148 A 12/1999 Bennett et al.
6,268,490 B1 7/2001 Imanishi et al.
6,525,191 B1 2/2003 Ramasamy
6,582,908 B2 6/2003 Fodor et al.
6,670,461 B1 12/2003 Nielsen et al.
6,770,748 B2 8/2004 Imanishi et al.
6,794,499 B2 9/2004 Wengel et al.
7,034,133 B2 4/2006 Wengel et al.
7,053,207 B2 5/2006 Wengel et al.
7,399,845 B2 7/2008 Seth et al.
7,427,672 B2 9/2008 Imanishi et al.
7,547,684 B2 6/2009 Seth et al.
7,696,345 B2 4/2010 Allerson et al.
8,501,805 B2 8/2013 Seth et al.
8,530,640 B2 9/2013 Seth et al.
8,546,556 B2 10/2013 Seth et al.
9,012,421 B2 4/2015 Migawa et al.
2001/0053519 A1 12/2001 Fodor et al.
2003/0228597 A1 12/2003 Cowsert et al.
2004/0171570 A1 9/2004 Allerson et al.

2004/0181048 A1 9/2004 Wang
2005/0130923 A1 6/2005 Bhat et al.
2007/0031844 A1 2/2007 Khvorova et al.
2008/0039618 A1 2/2008 Allerson et al.
2009/0012281 A1 1/2009 Swayze et al.
2012/0149757 A1 6/2012 Krainer et al.
2012/0214865 A1 8/2012 Bennett et al.
2014/0255936 A1 9/2014 Rademakers
2014/0303238 A1 10/2014 Linsley et al.
2015/0148404 A1 5/2015 de Visser et al.
2016/0108396 A1* 4/2016 Jensen C12N 15/111
514/44 A

FOREIGN PATENT DOCUMENTS

EP 1752536 2/2007
WO WO 2016/024205 2/1916
WO WO 2016/060919 4/1916
WO WO 98/39352 9/1998
WO WO 99/14226 3/1999
WO WO 03/004602 1/2003
WO WO 2004/106356 12/2004
WO WO 2005/021570 3/2005
WO WO 2005/040180 5/2005
WO WO 2007/134181 11/2007
WO WO 2008/101157 8/2008
WO WO 2008/150729 12/2008
WO WO 2008/154401 12/2008
WO WO 2009/006478 1/2009
WO WO 2011/135396 11/2011
WO WO 2012/092367 7/2012
WO WO 2013/030588 3/2013
WO WO 2013/086207 6/2013
WO WO 2014/062691 4/2014
WO WO2015054676 * 4/2015

OTHER PUBLICATIONS

Riboldi et al (Mol Neurobiol (2014) 50:721-732).
Wojciechowska et al (Human Molecular Genetics, 2011, 1-11,
doi:10.1093/hmg/ddr299).
Donnelly et al ("RNA Toxicity from the ALS/FTD C9ORF72
Expansion Is Mitigated by Antisense Intervention", Neuron 80,
415-428, Oct. 16, 2013, plus supplementaty materials).
Lagier-Tourenne et al ("Targeted degradation of sense and antisense
C9orf72 RNA foci as therapy for ALS and frontotemporal degen-
eration", Proc. Nat. Acad. Sci. USA E4530-E4539, 2013).
Jiang et al ("Gain of Toxicity from ALS/FTD-Linked Repeat
Expansions in C9ORF72 Is Alleviated by Antisense Oligonucle-
otides Targeting GGGGCC-Containing RNAs", Neuron 90, 535-
550, May 4, 2016).
Lindquist et al ("Corticobasal and ataxia syndromes widen the
spectrum of C9ORF72 hexanucleotide expansion disease", Clin
Genet 2013: 83: 279-283).

(Continued)

Primary Examiner — Richard Schnizer
(74) *Attorney, Agent, or Firm* — Ionis Pharmaceuticals,
Inc. Patent Dept.

(57) **ABSTRACT**

Disclosed herein are compositions and methods for reducing
expression of C9ORF72 mRNA and protein in an animal.
Such methods are useful to treat, prevent, ameliorate, or
slow progression of neurodegenerative diseases in an indi-
vidual in need thereof.

20 Claims, No Drawings

(56)

References Cited

OTHER PUBLICATIONS

Sareen et al ("Targeting RNA Foci in iPSC-Derived Motor Neurons from ALS Patients with a C9ORF72 Repeat Expansion", *Science Translational Medicine* 5(208): 208ra149, 15 pages).*

"The ALS Association and the Packard Center Partner to Develop Animal Model Systems for Most Common Cause of Familial ALS", <http://www.alsa.org/news/archive/new-animal-model-systems.html> (printed Oct. 23, 2015).

Albaek et al., "Analogues of a Locked Nucleic Acid with Three-Carbon 2',4'-Linkages: Synthesis by Ring-Closing Metathesis and Influence of Nucleic Acid Duplex Stability" *J. Org. Chem.* (2006) 71:7731-7740.

Altmann et al., "Second Generation Antisense Oligonucleotides-Inhibition of PKC- α and c-raf Kinase Expression by Chimeric Oligonucleotides Incorporating 6"-Substituted Carbocyclic Nucleosides and 2"-O-Ethylene Glycol Substituted Ribonucleosides" *Nucleosides & Nucleotides*. (1997) 16:917-926.

Altmann et al., "Second Generation of Antisense Oligonucleotides: From Nuclease Resistance to Biological Efficacy in Animals" *Chimia*. (1996) 50(4):168-176.

Altmann et al., "Second-generation antisense oligonucleotides: structure-activity relationships and the design of improved signal-transduction inhibitors" *Biochem. Soc. Trans.* (1996) 24:630-637.

Altschul et al., "Basic Local Alignment Search Tool" *J. Mol. Biol.* (1990) 215:403-410.

Al-Sarraj et al., "p62 positive, TDP-43 negative, neuronal cytoplasmic and intranuclear inclusions in the cerebellum and hippocampus define the pathology of C9orf72-linked FTL and MND/ALS" *Acta Neuropathol* (2011) 122:691-702.

Baker et al., "2'-O-(2-Methoxy)ethyl-modified Anti-intercellular Adhesion Molecule 1 (ICAM-1) Oligonucleotides Selectively Increase the ICAM-1 mRNA Level and Inhibit Formation of the ICAM-1 Translation Initiation Complex in Human Umbilical Vein Endothelial Cells" *J. Biol. Chem.* (1997) 272:11994-12000.

Baloh, R.H., "Generation of Non-Integrating iPSC Cells and Motor Neurons from C9orf72 Repeat Expansion ALS Patients" 65th AAN Annual Meeting, San Diego, CA, Mar. 16-23, 2013.

Baloh, R.H., "Targeting RNA foci shows a therapeutic effect in iPSC-derived motor neurons from C9orf72 repeat patients" ALSMND meeting, Milan, Dec. 6, 2013.

Baloh, R.H., "Induced Pluripotent stem cell models from C9orf72 patients." Oral presentation, California ALS PAC10 Research Summit, Los Angeles, CA, Nov. 11, 2012.

Baughn et al., "Antisense Oligonucleotide as a Potential Therapy for Amyotrophic Lateral Sclerosis with C9orf72 Expansion" Poster Presentation, Keystone Symposia, New Frontiers in Neurodegenerative Disease Research, Santa Fe, NM, Feb. 3-8, 2013.

Baughn et al, "Sense and Anti-Sense RNA Foci in c9ALS/FTD: More Light in a House of Mirrors" *Annals of Neurology* (Dec. 2013) 74(17): pS60.

Bieniek et al., "Tau pathology in frontotemporal lobar degeneration with C9ORF72 hexanucleotide repeat expansion" *Acta Neuropathol* (2013) 125(2):289-302.

Boxer et al. "Clinical, neuroimaging and neuropathological features of a new chromosome 9p-linked FTD-ALS family" *J. Neurol. Neurosurg. Psychiatry* (2011) 82:196-203.

Braasch et al., "Locked nucleic acid (LNA): fine-tuning the recognition of DNA and RNA" *Chem. Biol.* (2001) 8:1-7.

Branch et al., "A good antisense molecule is hard to find," *TIBS* (1998) 23:45-50.

Brettschneider et al., "Microglial activation correlates with disease progression and upper motor neuron clinical symptoms in Amyotrophic Lateral Sclerosis", *PLOS ONE* (2012) 7:e39216.

Chin "On the Preparation and Utilization of Isolated and Purified Oligonucleotides" Document purportedly located on a CD-ROM and contributed to the public collection of the Katherine R. Everett Law Library of the University of North Carolina on Mar. 14, 2002.

Chio et al., "Prevalence of SOD1 mutations in the Italian ALS population" *Neurology* (2008) 70:533-537.

Crooke et al., "Basic Principles of Antisense Therapeutics" *Antisense Research and Application* (1998) Chapter 1:1-50.

Dejesus-Hernandez et al., "Expanded GGGGCC Hexanucleotide Repeat in Noncoding Region of C9ORF72 Causes Chromosome 9p-Linked FTD and ALS" *Neuron* (2011) 72:245-256.

Donnelly et al., "Development of a C9ORF72 ALS antisense therapy and a therapeutic biomarker" Abstracts of the Society for Neuroscience, Washington, DC, US, Oct. 17, 2012, Retrieved from the Internet Aug. 15, 2016: <http://www.abstractsonline.com/Plan/ViewAbstract.aspx?sKey=c4cccf5-5e4c-4d1e-9569-9a1b1eb21d80&4cKey=c5c69155-5d2b-467c-8d1f-87299c514c7f&mKey=%7b70007181-01C9-4DE9-A0A2-EEBFA14CD9F1%7d>.

Donnelly et al., "Development of C9ORF72 ALS Biomarkers and Therapeutics" American Neurological Association 2012 Annual Meeting, Poster Presentation, Boston, MA Oct. 10, 2012.

Donnelly et al., "Development of C9orf72 ALS Biomarkers and Therapeutics" *Annals of Neurology* (Oct. 10, 2012) 72(16):S67-S68.

Donnelly et al., "Limited availability of ZBP1 restricts axonal mRNA localization and nerve regeneration capacity" *EMBO J.* (2011) 30:4665-4677.

Donnelly et al., "Transcriptome analysis of C9orf72 ALS patient derived CNS iPS cells and autopsy tissue reveals a unique expression and splicing profile." Abstracts of the Society for Neuroscience, Washington, DC, US, Oct. 16, 2012, Retrieved from the Internet Aug. 19, 2016: <http://www.abstractsonline.com/Plan/ViewAbstract.aspx?sKey=99bd542e-9dff-4338-9756-dfbef1839aa6&cKey=63d1b086-9f01-43d4-ab3f-d258faa86d9e&mKey=%7b70007181-01C9-4DE9-A0A2-EEBFA14CD9F1%7d>.

Donnelly et al., "Transcriptome analysis of C9orf72 ALS patient derived CNS iPS cells and autopsy tissue reveals a unique expression and splicing profile." Oral Presentation, Neuroscience 2012, Washington, DC, US, Oct. 17, 2012.

Elayadi et al., "Application of PNA and LNA oligomers to chemotherapy" *Curr. Opinion Inven. Drugs* (2001) 2:558-561.

European Search Report for application No. 13847957.1 dated Jul. 13, 2016.

European Search Report for application No. 13846313.8 dated May 11, 2016.

European Search Report for application No. 13847099.2 dated May 25, 2016.

Freier et al., "The ups and downs of nucleic acid duplex stability: structure-stability studies on chemically-modified DNA:RNA duplexes" *Nucleic Acids Research* (1997) 25(22):4429-4443.

Frieden et al., "Expanding the design horizon of antisense oligonucleotides with alpha-L-LNA" *Nucleic Acids Research* (2003) 31(21):6365-6372.

Ganesalingam et al., "Combination of neurofilament heavy chain and complement C3 as CSF biomarkers for ALS" *Journal of Neurochemistry* (2011) 117: 528-537.

Gautschi et al., "Activity of a novel bcl-2/bcl-xLbispecific antisense oligonucleotide against tumors of diverse histologic origins" *J. Natl. Cancer Inst.* (2001) 93:463-471.

GenBank: JU333328.1 TSA: Macaca mulatta Mamu_527777 mRNA sequence. Mar. 26, 2012 (Retrieved from the internet Sep. 12, 2016: <http://www.ncbi.nlm.nih.gov/nuccore/380810415?sat=18&satkey=24474174>).

Hirtz et al., "How common are the "common" neurologic disorders?" *Neurology* (2007) 68:326-337.

Ince et al., "Molecular pathology and genetic advances in amyotrophic lateral sclerosis: an emerging molecular pathway and the significance of glial pathology," *Acta Neuro.* (2011) 122:657-671.

International Search Report for application No. PCT/US2013/065073 dated Apr. 22, 2014.

International Search Report for application No. PCT/US2013/065067 dated Jan. 24, 2014.

International Search Report for application No. PCT/US2013/065131 dated Feb. 14, 2014.

International Search Report for application on. PCT/US2014/060194 dated Apr. 14, 2015.

(56)

References Cited

OTHER PUBLICATIONS

Jiang et al., "Antisense oligonucleotide therapy for ALS/FTD caused by a gain of toxicity from C9orf72 hexanucleotide expansions." Poster Presentation, 10th Brain Research Conference, RNA Metabolism in Neurological Disease, Oct. 16, 2015.

Jiang et al., "Gain of Toxicity from ALS/FTG-Linked Repeat Expansions in C9ORF72 Is Alleviated by Antisense Oligonucleotides Targeting GGGCC-Containing RNAs." *Neuron* (2016) 90:535-550.

Johnson et al., "Exome sequencing reveals VCP mutations as a cause of familial ALS" *Neuron* (2010) 68:857-864.

Jones et al., "RNA quantitation by fluorescence-based solution assay: RiboGreen reagent characterization" *Analytical Biochemistry* (1998) 265(2):368-374.

Klein et al., "Gain of RNA function in pathological cases: Focus on myotonic dystrophy" *Biochimie* (2011) 93(11):2006-2012.

Koshkin et al., "LNA (locked nucleic acids): Synthesis of the adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil bicyclonucleoside monomers, oligomerisation, and unprecedented nucleic acid recognition" *Tetrahedron* (1998) 54:3607-3630.

Kumar et al., "The first analogues of LNA (locked nucleic acids): phosphorothioate-LNA and 2'-thio-LNA" *Bioorg Med Chem Lett* (1998) 8:2219-2222.

Kwiatkowski et al., "Mutations in the FUS/TLS gene on chromosome 16 cause familial amyotrophic lateral sclerosis" *Science* (2009) 323:1205-1208.

Laaksovirta et al., "Chromosome 9p21 in amyotrophic lateral sclerosis in Finland: a genome-wide association study" *Lancet Neural* (2010) 9:978-985.

Lagier-Tourenne, et al., "Sense and Antisense RNA Foci in C9-ALS/FTD: More Light in a House of Mirrors." Poster Presentation, American Neurological Association 2013 Annual Meeting; Oct. 14, 2013.

Lagier-Tourenne, C., "Targeted degradation of sense and antisense C9orf72 nuclear foci as therapy for ALS and FTD" Oral Presentation, 24th International Symposium on ALS/MND, Milan, Dec. 6, 2013.

Lagier-Tourenne, C., "Identifying mechanisms and therapy for ALS/FTD from C9orf72 expansion", Oral Presentation, ALSA and AFTD Symposium, Society for Neuroscience Annual Meeting, New Orleans; Oct. 15, 2012.

Lagier-Tourenne, C. "Therapy Development for ALS/MND and Frontotemporal Dementia with C9orf72 Expansion: Antisense Oligonucleotide Mediated Reduction in Nuclear RNA Foci." *ALS FD* (Nov. 4, 2013) 14(sup2): p. 17.

Leumann et al., "DNA Analogues: From Supramolecular Principles to Biological Properties" *Bioorganic & Medicinal Chemistry* (2002) 10:841-854.

Lillo et al., "Frontotemporal dementia and motor neurone disease: overlapping clinic-pathological disorders" *J. Clin. Neurosci.* (2009) 16:1131-1135.

Madson, "Antisense Against C9ORF72", <http://alsn.mda.org/article/antisense-against-c9orf72> (printed Oct. 28, 2015).

Maher et al., "Comparative hybrid arrest by tandem antisense oligodeoxyribonucleotides or oligodeoxyribonucleoside methylphosphonates in a cell-free system" *Nucl. Acid. Res.* (1998) 16(8):3341-3358.

Margolis et al., "DM2 intronic expansions: evidence for CCUG accumulation without flanking sequence or effects on ZNF9 mRNA processing or protein expression" *Hum. Mol. Genet.* (2006) 15:1808-1815.

Martin, "New acces to 2'-O-alkylated ribonucleosides and properties of 2'-O-alkylated oligoribonucleotides" *Helv. Chim. Acta.* (1995) 78:486-504.

Maruyama et al., "Mutations of optineurin in amyotrophic lateral sclerosis" *Nature* (2010) 465:223-226.

Morita et al., "A locus on chromosome 9p confers susceptibility to ALS and frontotemporal dementia" *Neurology* (2006) 66:839-844.

Mulders et al., "Triplet-repeat oligonucleotide-mediated reversal of RNA toxicity in myotonic dystrophy" *PNAS* (2009) 106(33):13915-13920.

Nelson et al., "The unstable repeats—three evolving faces of neurological disease." *Neuron* (2013) 77(5):825-43.

Neumann et al., "Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis" *Science* (2006) 314:130-133.

New England Biolabs 1998/99 Catalog (cover page and pp. 121 and 284).

O'Rourke et al., "C9orf72 BAC Transgenic Mice Display Typical Pathologic Features of ALS/FTD." *Neuron* (2015) 88(5):892-901.

Orum et al., "Locked nucleic acids: A promising molecular family for gene-function analysis and antisense drug development" *Curr. Opinion Mol. Ther.* (2001) 3:239-243.

Ostrow et al., "The C9orf72 ALS mutation causes both increased expression and aberrant splicing of the endothelin-B receptor, and its ligand endothelin-1 is increased in CNS tissue from ALS patients and mutant mice," Abstracts of the Society for Neuroscience (Oct. 17, 2012) 42: p. 1.

Pearson et al., "Familial frontotemporal dementia with amyotrophic lateral sclerosis and a shared haplotype on chromosome 9p" *J. Neurol.* (2011) 258:647-655.

Rabin et al., "Sporadic ALS has compartment-specific aberrant exon splicing and altered cell-matrix adhesion biology" *Hum Mol Genet.* (2010) 19(2):313-328.

Ravits, J., "Expanding Neurodegenerations: C9orf72-ALS/FTD" Oral Presentation, ANA Meeting, New Orleans, LA, (Oct. 15, 2013).

Ravits, J., "Regional Spread in ALS: Mechanisms and Pathogenesis." Oral Presentation, 2nd Annual Neuromuscular Colloquium, UC Irvine, Newport Beach, CA, May 4, 2012.

Renton et al., "A Hexanucleotide Repeat Expansion in C9ORF72 Is the Cause of Chromosome 9p21-Linked ALS-FTD" *Neuron* (2011) 72:257-268.

Reynolds et al., "Rational siRNA design for RNA interference" *Nature Biotechnology* (2004) 22(3) 326-330.

Rigo, F., "ASO therapy for ALS and FTD caused by a gain of toxicity from hexanucleotide expansion in the C9orf72 gene." Oral Presentation, OTS Annual Meeting, Leiden, The Netherlands; Oct. 11-14, 2015.

Rosen et al., "Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis" *Nature* (1993) 362:59-62.

Rowland et al., "Amyotrophic lateral sclerosis" *N. Engl. J. Med.* (2001) 344(22):1688-1700.

Sareen, et al., "Targeting RNA foci shows a therapeutic effect in iPSC-derived motor neurons from C9orf72 repeat patients." *ALS FD* (Nov. 4, 2013) 14(sup2): pp. 16-17.

Sanghvi et al., "Heterocyclic Base Modifications in Nucleic Acids and Their Applications in Antisense Oligonucleotides" *Antisense Research and Applications* (1993) pp. 273-288.

Singh et al., "LNA (locked nucleic acids): synthesis and high-affinity nucleic acid recognition" *Chem. Commun.* (1998) 4:455-456.

Singh et al., "Synthesis of 2'-amino-LNA: A novel conformationally restricted high-affinity oligonucleotide analogue with a handle" *J. Org. Chem.* (1998) 63: 10035-10039.

Simon-Sanchez et al., "The clinical and pathological phenotype of C9ORF72 hexanucleotide repeat expansions", *Brain: Journal of Neurology* (2012) 135:723-735.

Smith et al., "Comparison of biosequences" *Adv. Appl. Math.* (1981) 2(4):482-489.

Sreedharan et al., "TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis" *Science* (2008) 319:1668-1672.

Srivastava et al., "Five- and Six-Membered Conformationally Locked 2',4'-Carbocyclic ribo-Thymidines: Synthesis, Structure, and Biochemical Studies" *J. Am. Chem. Soc.* (2007) 129(26):8362-8379.

Todd et al., "RNA mediated neurodegeneration in repeat expansion disorders," *Annals of Neurology* (2009) 67(3):291-300.

(56)

References Cited

OTHER PUBLICATIONS

Vance et al., "Familial amyotrophic lateral sclerosis with frontotemporal dementia is linked to a locus on chromosome 9p13.2-21.3" *Brain* (2006) 129:868-876.

Wahlestedt et al., "Potent and nontoxic antisense oligonucleotide containing locked nucleic acids" *Proc. Natl. Acad. Sci. USA* (2000) 97: 5633-5638.

Woolf et al., "Specificity of antisense oligonucleotides in vivo" *PNAS* (1992) 89: 7305-7309.

Zhang et al., "PowerBLAST: A New Network BLAST Application for Interactive or Automated Sequence Analysis and Annotation" *Genome Res.* (1997) 7:649-656.

Zhang et al., "The C9orf72 repeat expansion disrupts nucleocytoplasmic transport." *Nature* (2015) 525(7567):56-61.

Zhou et al., "Fine Tuning of Electrostatics around the Internucleotidic Phosphate through Incorporation of Modified 2',4'-Carbocyclic-LNAs and -ENAs Leads to Significant Modulation of Antisense Properties" *J. Org. Chem.* (2009) 74:118-134.

* cited by examiner

1

**COMPOSITIONS FOR MODULATING
C9ORF72 EXPRESSION**

SEQUENCE LISTING

The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled BIOL0269USSEQ_ST25.txt created Apr. 15, 2016, which is 104 Kb in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

FIELD

Provided are compositions and methods for modulating expression of C9ORF72 mRNA and protein in cells and animals. Such compositions and methods are useful to treat, prevent, ameliorate, or slow progression of neurodegenerative diseases, including amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, and olivopontocerebellar degeneration (OPCD).

BACKGROUND

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative disease characterized clinically by progressive paralysis leading to death from respiratory failure, typically within two to three years of symptom onset (Rowland and Shneider, *N. Engl. J. Med.*, 2001, 344, 1688-1700). ALS is the third most common neurodegenerative disease in the Western world (Hirtz et al., *Neurology*, 2007, 68, 326-337), and there are currently no effective therapies. Approximately 10% of cases are familial in nature, whereas the bulk of patients diagnosed with the disease are classified as sporadic as they appear to occur randomly throughout the population (Chio et al., *Neurology*, 2008, 70, 533-537). There is growing recognition, based on clinical, genetic, and epidemiological data, that ALS and frontotemporal dementia (FTD) represent an overlapping continuum of disease, characterized pathologically by the presence of TDP-43 positive inclusions throughout the central nervous system (Lillo and Hodges, *J. Clin. Neurosci.*, 2009, 16, 1131-1135; Neumann et al., *Science*, 2006, 314, 130-133).

To date, a number of genes have been discovered as causative for classical familial ALS, for example, SOD1, TARDBP, FUS, OPTN, and VCP (Johnson et al., *Neuron*, 2010, 68, 857-864; Kwiatkowski et al., *Science*, 2009, 323, 1205-1208; Maruyama et al., *Nature*, 2010, 465, 223-226; Rosen et al., *Nature*, 1993, 362, 59-62; Sreedharan et al., *Science*, 2008, 319, 1668-1672; Vance et al., *Brain*, 2009, 129, 868-876). Recently, linkage analysis of kindreds involving multiple cases of ALS, FTD, and ALS-FTD had suggested that there was an important locus for the disease on the short arm of chromosome 9 (Boxer et al., *J. Neurol. Neurosurg. Psychiatry*, 2011, 82, 196-203; Morita et al., *Neurology*, 2006, 66, 839-844; Pearson et al. *J. Neurol.*, 2011, 258, 647-655; Vance et al., *Brain*, 2006, 129, 868-876). The mutation in the C9ORF72 gene is the most common genetic cause of ALS and FTD. The ALS-FTD causing mutation is a large hexanucleotide (GGGGCC) repeat expansion in the first intron of the C9ORF72 gene (Renton et al., *Neuron*, 2011, 72, 257-268; DeJesus-Hernandez et al., *Neuron*, 2011, 72, 245-256). A founder haplotype, covering the C9ORF72 gene, is present in the majority of cases linked to this region (Renton et al., *Neuron*, 2011, 72, 257-268). This locus on chromosome 9p21 accounts for nearly half of familial ALS

2

and nearly one-quarter of all ALS cases in a cohort of 405 Finnish patients (Laaksovirta et al, *Lancet Neurol.*, 2010, 9, 978-985).

A founder haplotype, covering the C9ORF72 gene, is present in the majority of cases linked to this region.

There are currently no effective therapies to treat such neurodegenerative diseases. Therefore, it is an object to provide compositions and methods for the treatment of such neurodegenerative diseases.

SUMMARY

Certain embodiments provide methods, compounds, and compositions for inhibiting expression of C9ORF72 mRNA and protein in cells, tissues, and animals. Certain embodiments provide methods, compounds, and compositions for reducing C9ORF72 mRNA and protein levels in cells, tissues, and animals. Certain embodiments provide antisense compounds targeted to a C9ORF72 nucleic acid. In certain embodiments, the antisense compounds are modified oligonucleotides. In certain embodiments, the modified oligonucleotides are single-stranded.

In certain embodiments, C9ORF72 associated Repeat Associated Non-ATG Translation (RAN translation) products are reduced. In certain embodiments, the C9ORF72 associated RAN translation products are poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine). In certain embodiments, certain C9ORF72 mRNA variants are preferentially reduced. In certain embodiments, the C9ORF72 mRNA variants preferentially reduced are variants processed from a pre-mRNA containing intron 1. In certain embodiments, intron 1 contains a hexanucleotide repeat expansion. In certain embodiments, the C9ORF72 mRNA variant preferentially reduced is a C9ORF72 pathogenic associated mRNA variant. In certain embodiments, the C9ORF72 pathogenic associated mRNA variant is NM_001256054.1 (SEQ ID NO: 1). In certain embodiments, the hexanucleotide repeat expansion is associated with a C9ORF72 associated disease. In certain embodiments, the hexanucleotide repeat expansion is associated with a C9ORF72 hexanucleotide repeat expansion associated disease. In certain embodiments, the hexanucleotide repeat expansion comprises at least 30 GGGGCC repeats, more than 30 GGGGCC repeats, more than 100 GGGGCC repeats, more than 500 GGGGCC repeats, or more than 1000 GGGGCC repeats. In certain embodiments, the hexanucleotide repeat expansion is associated with nuclear foci. In certain embodiments, C9ORF72 associated RAN translation products are associated with nuclear foci. In certain embodiments, the C9ORF72 associated RAN translation products are poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine). In certain embodiments, the compositions and methods described herein are useful for reducing C9ORF72 mRNA levels, C9ORF72 protein levels, C9ORF72 RAN translation products, and nuclear foci. In certain embodiments, the compositions and methods described herein are useful for selectively reducing C9ORF72 pathogenic associated mRNA variants. Such reduction can occur in a time-dependent manner or in a dose-dependent manner.

Also provided are methods useful for preventing, treating, ameliorating, and slowing progression of diseases associated with C9ORF72. In certain embodiments, such C9ORF72 associated diseases are neurodegenerative diseases. In certain embodiments, the neurodegenerative disease is amyotrophic lateral sclerosis (ALS), frontotemporal dementia

(FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, and olivopontocerebellar degeneration (OPCD).

Such diseases can have one or more risk factors, causes, or outcomes in common. Certain risk factors and causes for development of a neurodegenerative disease, and, in particular, ALS and FTD, include genetic predisposition and older age.

In certain embodiments, methods of treatment include administering a C9ORF72 antisense compound to an individual in need thereof. In certain embodiments, the antisense compound is a single-stranded modified oligonucleotide. In certain embodiments, the single-stranded modified oligonucleotide is complementary to a C9ORF72 nucleic acid.

DETAILED DESCRIPTION

It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed. Herein, the use of the singular includes the plural unless specifically stated otherwise. As used herein, the use of “or” means “and/or” unless stated otherwise. Additionally, as used herein, the use of “and” means “and/or” unless stated otherwise. Furthermore, the use of the term “including” as well as other forms, such as “includes” and “included”, is not limiting. Also, terms such as “element” or “component” encompass both elements and components comprising one unit and elements and components that comprise more than one subunit, unless specifically stated otherwise.

Compounds of the invention include variations of the disclosed compounds in which one or more hydrogen, carbon, nitrogen, oxygen, or sulfur atoms is replaced with a stable isotope of the same element.

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. All documents, or portions of documents, cited in this disclosure, including, but not limited to, patents, patent applications, published patent applications, articles, books, treatises, and GENBANK Accession Numbers and associated sequence information obtainable through databases such as National Center for Biotechnology Information (NCBI) and other data referred to throughout in the disclosure herein are hereby expressly incorporated by reference for the portions of the document discussed herein, as well as in their entirety.

DEFINITIONS

Unless specific definitions are provided, the nomenclature utilized in connection with, and the procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art.

Standard techniques may be used for chemical synthesis, and chemical analysis.

Unless otherwise indicated, the following terms have the following meanings:

“2'-O-methoxyethyl” (also 2'-MOE and 2'-OCH₂CH₂—OCH₃ and MOE) refers to an O-methoxyethyl modification of the 2' position of a furanose ring. A 2'-O-methoxyethyl modified sugar is a modified sugar.

“2'-MOE nucleoside” (also 2'-O-methoxyethyl nucleoside) means a nucleoside comprising a MOE modified sugar moiety.

“2'-substituted nucleoside” means a nucleoside comprising a substituent at the 2'-position of the furanose ring other than H or OH. In certain embodiments, 2'-substituted nucleosides include nucleosides with bicyclic sugar modifications.

“5-methylcytosine” means a cytosine modified with a methyl group attached to the 5' position. A 5-methylcytosine is a modified nucleobase.

“Administered concomitantly” refers to the co-administration of two pharmaceutical agents in any manner in which the pharmacological effects of both are manifest in the patient at the same time. Concomitant administration does not require that both pharmaceutical agents be administered in a single pharmaceutical composition, in the same dosage form, or by the same route of administration. The effects of both pharmaceutical agents need not manifest themselves at the same time. The effects need only be overlapping for a period of time and need not be coextensive.

“Administering” means providing a pharmaceutical agent to an animal, and includes, but is not limited to administering by a medical professional and self-administering.

“Amelioration” refers to a lessening, slowing, stopping, or reversing of at least one indicator of the severity of a condition or disease. The severity of indicators may be determined by subjective or objective measures, which are known to those skilled in the art.

“Animal” refers to a human or non-human animal, including, but not limited to, mice, rats, rabbits, dogs, cats, pigs, and non-human primates, including, but not limited to, monkeys and chimpanzees.

“Antibody” refers to a molecule characterized by reacting specifically with an antigen in some way, where the antibody and the antigen are each defined in terms of the other. Antibody may refer to a complete antibody molecule or any fragment or region thereof, such as the heavy chain, the light chain, Fab region, and Fc region.

“Antisense activity” means any detectable or measurable activity attributable to the hybridization of an antisense compound to its target nucleic acid. In certain embodiments, antisense activity is a decrease in the amount or expression of a target nucleic acid or protein encoded by such target nucleic acid.

“Antisense compound” means an oligomeric compound that is capable of undergoing hybridization to a target nucleic acid through hydrogen bonding. Examples of antisense compounds include single-stranded and double-stranded compounds, such as, antisense oligonucleotides, siRNAs, shRNAs, ssRNAs, and occupancy-based compounds.

“Antisense inhibition” means reduction of target nucleic acid levels in the presence of an antisense compound complementary to a target nucleic acid compared to target nucleic acid levels or in the absence of the antisense compound.

“Antisense mechanisms” are all those mechanisms involving hybridization of a compound with a target nucleic acid, wherein the outcome or effect of the hybridization is either target degradation or target occupancy with concomitant stalling of the cellular machinery involving, for example, transcription or splicing.

“Antisense oligonucleotide” means a single-stranded oligonucleotide having a nucleobase sequence that permits hybridization to a corresponding segment of a target nucleic acid.

“Base complementarity” refers to the capacity for the precise base pairing of nucleobases of an antisense oligonucleotide with corresponding nucleobases in a target

5

nucleic acid (i.e., hybridization), and is mediated by Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen binding between corresponding nucleobases.

“Bicyclic sugar” means a furanose ring modified by the bridging of two atoms. A bicyclic sugar is a modified sugar.

“Bicyclic nucleoside” (also BNA) means a nucleoside having a sugar moiety comprising a bridge connecting two carbon atoms of the sugar ring, thereby forming a bicyclic ring system. In certain embodiments, the bridge connects the 4'-carbon and the 2'-carbon of the sugar ring.

“C9ORF72 antisense transcript” means transcripts produced from the non-coding strand (also antisense strand and template strand) of the C9ORF72 gene. The C9ORF72 antisense transcript differs from the canonically transcribed “C9ORF72 sense transcript”, which is produced from the coding strand (also sense strand) of the C9ORF72 gene. In certain embodiments, a C9ORF72 antisense transcript is SEQ ID NO: 18.

“C9ORF72 antisense transcript specific inhibitor” refers to any agent capable of specifically inhibiting the expression of C9ORF72 antisense transcript and/or its expression products at the molecular level. As used herein, “specific” means reducing or inhibiting expression of C9ORF72 antisense transcript without reducing non-target transcript to an appreciable degree (e.g., a C9ORF72 antisense transcript specific inhibitor reduces expression of C9ORF72 antisense transcript, but does not reduce expression of C9ORF72 sense transcript to an appreciable degree). C9ORF72 specific antisense transcript inhibitors include antisense compounds, siRNAs, aptamers, antibodies, peptides, small molecules, and other agents capable of inhibiting the expression of C9ORF72 antisense transcript and/or its expression products, such as C9ORF72 antisense transcript associated RAN translation products.

“C9ORF72 associated disease” means any disease associated with any C9ORF72 nucleic acid or expression product thereof. Such diseases may include a neurodegenerative disease. Such neurodegenerative diseases may include ALS and FTD. In certain embodiments, the C9ORF72 associated disease is caused by (or is associated with) a hexanucleotide repeat expansion. In certain embodiments, the hexanucleotide repeat expansion may comprise GGGGCC, GGGGGG, GGGGGC, or GGGGCG repeated at least 30 times, more than 30 times, more than 100 times, more than 500 times, or more than 1000 times.

“C9ORF72 associated RAN translation products” means aberrant peptide or di-peptide polymers translated through RAN translation (i.e., repeat-associated, and non-ATG-dependent translation). In certain embodiments, the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).

“C9ORF72 nucleic acid” means any nucleic acid encoding C9ORF72. For example, in certain embodiments, a C9ORF72 nucleic acid includes a DNA sequence encoding C9ORF72, an RNA sequence transcribed from DNA encoding C9ORF72 including genomic DNA comprising introns and exons (i.e., pre-mRNA), and an mRNA sequence encoding C9ORF72. “C9ORF72 mRNA” means an mRNA encoding a C9ORF72 protein.

“C9ORF72 pathogenic associated mRNA variant” means the C9ORF72 mRNA variant processed from a C9ORF72 pre-mRNA variant containing the hexanucleotide repeat. A C9ORF72 pre-mRNA contains the hexanucleotide repeat when transcription of the pre-mRNA begins in the region from the start site of exon 1A to the start site of exon 1B, e.g., nucleotides 1107 to 1520 of the genomic sequence

6

(SEQ ID NO: 2, the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 27565000). In certain embodiments, the level of a C9ORF72 pathogenic associated mRNA variant is measured to determine the level of a C9ORF72 pre-mRNA containing the hexanucleotide repeat in a sample.

“C9ORF72 specific inhibitor” refers to any agent capable of specifically inhibiting the expression of C9ORF72 mRNA and/or C9ORF72 protein at the molecular level. For example, C9ORF72 specific inhibitors include nucleic acids (including antisense compounds), siRNAs, aptamers, antibodies, peptides, small molecules, and other agents capable of inhibiting the expression of C9ORF72 mRNA and/or C9ORF72 protein. Similarly, in certain embodiments, C9ORF72 specific inhibitors may affect other molecular processes in an animal.

“Cap structure” or “terminal cap moiety” means chemical modifications, which have been incorporated at either terminus of an antisense compound.

“cEt” or “constrained ethyl” means a bicyclic nucleoside having a sugar moiety comprising a bridge connecting the 4'-carbon and the 2'-carbon, wherein the bridge has the formula: 4'-CH(CH₃)—O-2'.

“Constrained ethyl nucleoside” (also cEt nucleoside) means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH(CH₃)—O-2' bridge.

“Chemically distinct region” refers to a region of an antisense compound that is in some way chemically different than another region of the same antisense compound. For example, a region having 2'-O-methoxyethyl nucleosides is chemically distinct from a region having nucleosides without 2'-O-methoxyethyl modifications.

“Chimeric antisense compound” means an antisense compound that has at least two chemically distinct regions, each position having a plurality of subunits.

“Co-administration” means administration of two or more pharmaceutical agents to an individual. The two or more pharmaceutical agents may be in a single pharmaceutical composition, or may be in separate pharmaceutical compositions. Each of the two or more pharmaceutical agents may be administered through the same or different routes of administration. Co-administration encompasses parallel or sequential administration.

“Complementarity” means the capacity for pairing between nucleobases of a first nucleic acid and a second nucleic acid.

“Comprise,” “comprises,” and “comprising” will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

“Contiguous nucleobases” means nucleobases immediately adjacent to each other.

“Designing” or “designed to” refer to the process of designing an oligomeric compound that specifically hybridizes with a selected nucleic acid molecule.

“Diluent” means an ingredient in a composition that lacks pharmacological activity, but is pharmaceutically necessary or desirable. For example, in drugs that are injected, the diluent may be a liquid, e.g. saline solution.

“Dose” means a specified quantity of a pharmaceutical agent provided in a single administration, or in a specified time period. In certain embodiments, a dose may be administered in one, two, or more boluses, tablets, or injections. For example, in certain embodiments where subcutaneous administration is desired, the desired dose requires a volume not easily accommodated by a single injection, therefore, two or more injections may be used to achieve the desired

dose. In certain embodiments, the pharmaceutical agent is administered by infusion over an extended period of time or continuously. Doses may be stated as the amount of pharmaceutical agent per hour, day, week, or month.

“Effective amount” in the context of modulating an activity or of treating or preventing a condition means the administration of that amount of pharmaceutical agent to a subject in need of such modulation, treatment, or prophylaxis, either in a single dose or as part of a series, that is effective for modulation of that effect, or for treatment or prophylaxis or improvement of that condition. The effective amount may vary among individuals depending on the health and physical condition of the individual to be treated, the taxonomic group of the individuals to be treated, the formulation of the composition, assessment of the individual’s medical condition, and other relevant factors.

“Efficacy” means the ability to produce a desired effect.

“Expression” includes all the functions by which a gene’s coded information is converted into structures present and operating in a cell. Such structures include, but are not limited to the products of transcription and translation.

“Focus” or “foci” means a nuclear or cytoplasmic body comprising a C9ORF72 transcript. In certain embodiments, a focus comprises at least one C9ORF72 transcript. In certain embodiments, C9ORF72 foci comprise transcripts comprising any of the following hexanucleotide repeats: GGGGCC, GGGGGG, GGGGCG, and/or GGGGCG.

“Fully complementary” or “100% complementary” means each nucleobase of a first nucleic acid has a complementary nucleobase in a second nucleic acid. In certain embodiments, a first nucleic acid is an antisense compound and a target nucleic acid is a second nucleic acid.

“Gapmer” means a chimeric antisense compound in which an internal region having a plurality of nucleosides that support RNase H cleavage is positioned between external regions having one or more nucleosides, wherein the nucleosides comprising the internal region are chemically distinct from the nucleoside or nucleosides comprising the external regions. The internal region may be referred to as a “gap” and the external regions may be referred to as the “wings.”

“Hexanucleotide repeat expansion” means a series of six bases (for example, GGGGCC, GGGGGG, GGGGCG, or GGGGGC) repeated at least twice. In certain embodiments, the hexanucleotide repeat expansion may be located in intron 1 of a C9ORF72 nucleic acid. In certain embodiments, a pathogenic hexanucleotide repeat expansion includes at least 30, more than 30, more than 100, more than 500, or more than 1000 repeats of GGGGCC, GGGGGG, GGGGCG, or GGGGGC in a C9ORF72 nucleic acid and is associated with disease. In certain embodiments, the repeats are consecutive. In certain embodiments, the repeats are interrupted by 1 or more nucleobases. In certain embodiments, a wild-type hexanucleotide repeat expansion includes 23 or fewer repeats of GGGGCC, GGGGGG, GGGGCG, or GGGGGC in a C9ORF72 nucleic acid. In certain embodiments, the repeats are consecutive. In certain embodiments, the repeats are interrupted by 1 or more nucleobases.

“Hybridization” means the annealing of complementary nucleic acid molecules. In certain embodiments, complementary nucleic acid molecules include, but are not limited to, an antisense compound and a target nucleic acid. In certain embodiments, complementary nucleic acid molecules include, but are not limited to, an antisense oligonucleotide and a nucleic acid target.

“Identifying an animal having a C9ORF72 associated disease” means identifying an animal having been diagnosed

with a C9ORF72 associated disease or predisposed to develop a C9ORF72 associated disease. Individuals predisposed to develop a C9ORF72 associated disease include those having one or more risk factors for developing a C9ORF72 associated disease, including, having a personal or family history or genetic predisposition of one or more C9ORF72 associated diseases. Such identification may be accomplished by any method including evaluating an individual’s medical history and standard clinical tests or assessments, such as genetic testing.

“Immediately adjacent” means there are no intervening elements between the immediately adjacent elements.

“Individual” means a human or non-human animal selected for treatment or therapy.


“Inhibiting C9ORF72” means reducing the level or expression of a C9ORF72 mRNA and/or protein. In certain embodiments, C9ORF72 mRNA and/or protein levels are inhibited in the presence of an antisense compound targeting C9ORF72, including an antisense oligonucleotide targeting C9ORF72, as compared to expression of C9ORF72 mRNA and/or protein levels in the absence of a C9ORF72 antisense compound, such as an antisense oligonucleotide.

“Inhibiting the expression or activity” refers to a reduction or blockade of the expression or activity and does not necessarily indicate a total elimination of expression or activity.

“Internucleoside linkage” refers to the chemical bond between nucleosides.



“Linked nucleosides” means adjacent nucleosides linked together by an internucleoside linkage.

“Locked nucleic acid” or “LNA” or “LNA nucleosides” means nucleic acid monomers having a bridge connecting two carbon atoms between the 4' and 2'-position of the nucleoside sugar unit, thereby forming a bicyclic sugar. Examples of such bicyclic sugar include, but are not limited to A) α -L-Methyleneoxy (4'-CH₂—O-2') LNA, (B) β -D-Methyleneoxy (4'-CH₂—O-2') LNA, (C) Ethyleneoxy (4'-(CH₂)₂—O-2') LNA, (D) Aminooxy (4'-CH₂—O—N(R)-2') LNA and (E) Oxyamino (4'-CH₂—N(R)—O-2') LNA, as depicted below.



9

-continued



As used herein, LNA compounds include, but are not limited to, compounds having at least one bridge between the 4' and the 2' position of the sugar wherein each of the bridges independently comprises 1 or from 2 to 4 linked groups independently selected from $-\text{C}(\text{R}_1)(\text{R}_2)_n-$, $-\text{C}(\text{R}_1)=\text{C}(\text{R}_2)-$, $-\text{C}(\text{R}_1)=\text{N}-$, $-\text{C}(=\text{NR}_1)-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{O}-$, $-\text{Si}(\text{R}_1)_2-$, $-\text{S}(=\text{O})-$, and $-\text{N}(\text{R}_1)-$; wherein: x is 0, 1, or 2; n is 1, 2, 3, or 4; each R_1 and R_2 is, independently, H, a protecting group, hydroxyl, C_1 - C_{12} alkyl, substituted C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, substituted C_2 - C_{12} alkenyl, C_2 - C_{12} alkynyl, substituted C_2 - C_{12} alkynyl, C_5 - C_{20} aryl, substituted C_5 - C_{20} aryl, a heterocycle radical, a substituted heterocycle radical, heteroaryl, substituted heteroaryl, C_5 - C_7 alicyclic radical, substituted C_5 - C_7 alicyclic radical, halogen, OJ_1 , NJ_1J_2 , SJ_1 , N_3 , COOJ_1 , $\text{acyl}(\text{C}(=\text{O})-\text{H})$, substituted acyl, CN, sulfonyl ($\text{S}(=\text{O})_2-\text{J}_1$), or sulfoxyl ($\text{S}(=\text{O})-\text{J}_1$); and each J_1 and J_2 is, independently, H, C_1 - C_{12} alkyl, substituted C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, substituted C_2 - C_{12} alkenyl, C_2 - C_{12} alkynyl, substituted C_2 - C_{12} alkynyl, C_5 - C_{20} aryl, substituted C_5 - C_{20} aryl, acyl ($\text{C}(=\text{O})-\text{H}$), substituted acyl, a heterocycle radical, a substituted heterocycle radical, C_1 - C_{12} aminoalkyl, substituted C_1 - C_{12} aminoalkyl or a protecting group.

Examples of 4'-2' bridging groups encompassed within the definition of LNA include, but are not limited to one of formulae: $-\text{C}(\text{R}_1)(\text{R}_2)_n-$, $-\text{C}(\text{R}_1)(\text{R}_2)_n-\text{O}-$, $-\text{C}(\text{R}_1\text{R}_2)-\text{N}(\text{R}_1)-\text{O}-$ or $-\text{C}(\text{R}_1\text{R}_2)-\text{O}-\text{N}(\text{R}_1)-$. Furthermore, other bridging groups encompassed within the definition of LNA are 4'- CH_2 -2', 4'-(CH_2)₂-2', 4'-(CH_2)₃-2', 4'- CH_2 -O-2', 4'-(CH_2)₂-O-2', 4'- CH_2 -O-N(R_1)-2' and 4'- CH_2 -N(R_1)-O-2'- bridges, wherein each R_1 and R_2 is, independently, H, a protecting group or C_1 - C_{12} alkyl.

Also included within the definition of LNA according to the invention are LNAs in which the 2'-hydroxyl group of the ribosyl sugar ring is connected to the 4' carbon atom of the sugar ring, thereby forming a methyleneoxy (4'- CH_2 -O-2') bridge to form the bicyclic sugar moiety. The bridge can also be a methylene ($-\text{CH}_2-$) group connecting the 2' oxygen atom and the 4' carbon atom, for which the term methyleneoxy (4'- CH_2 -O-2') LNA is used. Furthermore; in the case of the bicyclic sugar moiety having an ethylene bridging group in this position, the term ethyleneoxy (4'- CH_2CH_2 -O-2') LNA is used. α -L-methyleneoxy (4'- CH_2 -O-2'), an isomer of methyleneoxy (4'- CH_2 -O-2') LNA is also encompassed within the definition of LNA, as used herein.

"Mismatch" or "non-complementary nucleobase" refers to the case when a nucleobase of a first nucleic acid is not

10

capable of pairing with the corresponding nucleobase of a second or target nucleic acid.

"Modified internucleoside linkage" refers to a substitution or any change from a naturally occurring internucleoside bond (i.e., a phosphodiester internucleoside bond).

"Modified nucleobase" means any nucleobase other than adenine, cytosine, guanine, thymidine, or uracil. An "unmodified nucleobase" means the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C), and uracil (U).

"Modified nucleoside" means a nucleoside having, independently, a modified sugar moiety and/or modified nucleobase.

"Modified nucleotide" means a nucleotide having, independently, a modified sugar moiety, modified internucleoside linkage, and/or modified nucleobase.

"Modified oligonucleotide" means an oligonucleotide comprising at least one modified internucleoside linkage, modified sugar, and/or modified nucleobase.

"Modified sugar" means substitution and/or any change from a natural sugar moiety.

"Monomer" means a single unit of an oligomer. Monomers include, but are not limited to, nucleosides and nucleotides, whether naturally occurring or modified.

"Motif" means the pattern of unmodified and modified nucleoside in an antisense compound.

"Natural sugar moiety" means a sugar moiety found in DNA (2'-H) or RNA (2'-OH).

"Naturally occurring internucleoside linkage" means a 3' to 5' phosphodiester linkage.

"Non-complementary nucleobase" refers to a pair of nucleobases that do not form hydrogen bonds with one another or otherwise support hybridization.

"Nucleic acid" refers to molecules composed of monomeric nucleotides. A nucleic acid includes, but is not limited to, ribonucleic acids (RNA), deoxyribonucleic acids (DNA), single-stranded nucleic acids, double-stranded nucleic acids, small interfering ribonucleic acids (siRNA), and microRNAs (miRNA).

"Nucleobase" means a heterocyclic moiety capable of pairing with a base of another nucleic acid.

"Nucleobase complementarity" refers to a nucleobase that is capable of base pairing with another nucleobase. For example, in DNA, adenine (A) is complementary to thymine (T). For example, in RNA, adenine (A) is complementary to uracil (U). In certain embodiments, complementary nucleobase refers to a nucleobase of an antisense compound that is capable of base pairing with a nucleobase of its target nucleic acid. For example, if a nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be complementary at that nucleobase pair.

"Nucleobase sequence" means the order of contiguous nucleobases independent of any sugar, linkage, and/or nucleobase modification.

"Nucleoside" means a nucleobase linked to a sugar.

"Nucleoside mimetic" includes those structures used to replace the sugar or the sugar and the base and not necessarily the linkage at one or more positions of an oligomeric compound such as for example nucleoside mimetics having morpholino, cyclohexenyl, cyclohexyl, tetrahydropyranyl, bicyclo, or tricyclo sugar mimetics, e.g., non furanose sugar units. Nucleotide mimetic includes those structures used to replace the nucleoside and the linkage at one or more positions of an oligomeric compound such as for example

peptide nucleic acids or morpholinos (morpholinos linked by —N(H)—C(=O)—O— or other non-phosphodiester linkage). Sugar surrogate overlaps with the slightly broader term nucleoside mimetic but is intended to indicate replacement of the sugar unit (furanose ring) only. The tetrahydropyranyl rings provided herein are illustrative of an example of a sugar surrogate wherein the furanose sugar group has been replaced with a tetrahydropyranyl ring system. “Mimetic” refers to groups that are substituted for a sugar, a nucleobase, and/or internucleoside linkage. Generally, a mimetic is used in place of the sugar or sugar-internucleoside linkage combination, and the nucleobase is maintained for hybridization to a selected target.

“Nucleotide” means a nucleoside having a phosphate group covalently linked to the sugar portion of the nucleoside.

“Off-target effect” refers to an unwanted or deleterious biological effect associated with modulation of RNA or protein expression of a gene other than the intended target nucleic acid.

“Oligomeric compound” or “oligomer” means a polymer of linked monomeric subunits which is capable of hybridizing to at least a region of a nucleic acid molecule.

“Oligonucleotide” means a polymer of linked nucleosides each of which can be modified or unmodified, independent one from another.

“Parenteral administration” means administration through injection (e.g., bolus injection) or infusion. Parenteral administration includes subcutaneous administration, intravenous administration, intramuscular administration, intraarterial administration, intraperitoneal administration, or intracranial administration, e.g., intrathecal or intracerebroventricular administration.

“Peptide” means a molecule formed by linking at least two amino acids by amide bonds. Without limitation, as used herein, peptide refers to polypeptides and proteins.

“Pharmaceutical agent” means a substance that provides a therapeutic benefit when administered to an individual. For example, in certain embodiments, an antisense oligonucleotide targeted to C9ORF72 is a pharmaceutical agent.

“Pharmaceutically acceptable derivative” encompasses pharmaceutically acceptable salts, conjugates, prodrugs or isomers of the compounds described herein.

“Pharmaceutically acceptable salts” means physiologically and pharmaceutically acceptable salts of antisense compounds, i.e., salts that retain the desired biological activity of the parent oligonucleotide and do not impart undesired toxicological effects thereto.

“Phosphorothioate linkage” means a linkage between nucleosides where the phosphodiester bond is modified by replacing one of the non-bridging oxygen atoms with a sulfur atom. A phosphorothioate linkage is a modified internucleoside linkage.

“Portion” means a defined number of contiguous (i.e., linked) nucleobases of a nucleic acid. In certain embodiments, a portion is a defined number of contiguous nucleobases of a target nucleic acid. In certain embodiments, a portion is a defined number of contiguous nucleobases of an antisense compound.

“Prevent” or “preventing” refers to delaying or forestalling the onset or development of a disease, disorder, or condition for a period of time from minutes to days, weeks to months, or indefinitely.

“Prodrug” means a therapeutic agent that is prepared in an inactive form that is converted to an active form within the body or cells thereof by the action of endogenous enzymes or other chemicals or conditions.

“Prophylactically effective amount” refers to an amount of a pharmaceutical agent that provides a prophylactic or preventative benefit to an animal.

“Region” is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or characteristic.

“Ribonucleotide” means a nucleotide having a hydroxy at the 2' position of the sugar portion of the nucleotide. Ribonucleotides may be modified with any of a variety of substituents.

“Salts” mean a physiologically and pharmaceutically acceptable salts of antisense compounds, i.e., salts that retain the desired biological activity of the parent oligonucleotide and do not impart undesired toxicological effects thereto.

“Segments” are defined as smaller or sub-portions of regions within a target nucleic acid.

“Shortened” or “truncated” versions of antisense oligonucleotides taught herein have one, two or more nucleosides deleted.

“Side effects” means physiological responses attributable to a treatment other than desired effects. In certain embodiments, side effects include, without limitation, injection site reactions, liver function test abnormalities, renal function abnormalities, liver toxicity, renal toxicity, central nervous system abnormalities, and myopathies.

“Single-stranded oligonucleotide” means an oligonucleotide which is not hybridized to a complementary strand. A “single-stranded modified oligonucleotide” means a modified oligonucleotide which is not hybridized to a complementary strand.

“Sites,” as used herein, are defined as unique nucleobase positions within a target nucleic acid.

“Slows progression” means decrease in the development of the disease.

“Specifically hybridizable” refers to an antisense compound having a sufficient degree of complementarity between an antisense oligonucleotide and a target nucleic acid to induce a desired effect, while exhibiting minimal or no effects on non-target nucleic acids under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays and therapeutic treatments.

“Stringent hybridization conditions” or “stringent conditions” refer to conditions under which an oligomeric compound will hybridize to its target sequence, but to a minimal number of other sequences.

“Subject” means a human or non-human animal selected for treatment or therapy.

“Target” refers to a protein, the modulation of which is desired.

“Target gene” refers to a gene encoding a target.

“Targeting” or “targeted” means the process of design and selection of an antisense compound that will specifically hybridize to a target nucleic acid and induce a desired effect.

“Target nucleic acid,” “target RNA,” and “target RNA transcript” and “nucleic acid target” all mean a nucleic acid capable of being targeted by antisense compounds.

“Target region” means a portion of a target nucleic acid to which one or more antisense compounds is targeted.

“Target segment” means the sequence of nucleotides of a target nucleic acid to which an antisense compound is targeted. “5' target site” refers to the 5'-most nucleotide of a target segment. “3' target site” refers to the 3'-most nucleotide of a target segment.

“Therapeutically effective amount” means an amount of a pharmaceutical agent that provides a therapeutic benefit to an individual.

13

“Treat” or “treating” or “treatment” means administering a composition to effect an alteration or improvement of a disease or condition.

“Unmodified nucleobases” means the purine bases adenine (A) and guanine (G), and the pyrimidine bases (T), cytosine (C), and uracil (U).

“Unmodified nucleotide” means a nucleotide composed of naturally occurring nucleobases, sugar moieties, and internucleoside linkages. In certain embodiments, an unmodified nucleotide is an RNA nucleotide (i.e. β -D-ribonucleosides) or a DNA nucleotide (i.e. β -D-deoxyribonucleoside).

“Wing segment” means a plurality of nucleosides modified to impart to an oligonucleotide properties such as enhanced inhibitory activity, increased binding affinity for a target nucleic acid, or resistance to degradation by *in vivo* nucleases.

Certain Embodiments

Certain embodiments provide compositions and methods for reducing total C9ORF72 mRNA and protein expression.

Certain embodiments provide compositions and methods for reducing C9ORF72 pathogenic associated mRNA variants.

Certain embodiments provide methods for the treatment, prevention, amelioration, or slowing progression of diseases associated with C9ORF72 in an individual in need thereof. Also contemplated are methods for the preparation of a medicament for the treatment, prevention, or amelioration of a disease associated with C9ORF72. C9ORF72 associated diseases include neurodegenerative diseases. In certain embodiments, the neurodegenerative disease may be ALS or FTD. In certain embodiments, the neurodegenerative disease may be familial or sporadic.

The present disclosure provided the following non-limiting numbered embodiments:

Embodiment 1

A compound comprising a modified oligonucleotide consisting of 12 to 30 linked nucleosides and having a nucleobase sequence comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, or at least 20 consecutive nucleobases of any of the nucleobase sequences of SEQ ID NOs: 22-55.

Embodiment 2

The compound of embodiment 1, wherein the nucleobase sequence of the modified oligonucleotide is at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% complementary to SEQ ID NO: 1 or SEQ ID NO: 2.

Embodiment 3

The compound of embodiments 1 and 2, wherein the modified oligonucleotide is a single-stranded modified oligonucleotide.

14

Embodiment 4

The compound of any of embodiments 1-3, wherein the modified oligonucleotide comprises at least one modified internucleoside linkage.

Embodiment 5

The compound of embodiment 4, wherein the at least one modified internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 6

The compound of embodiments 4 and 5, wherein the modified oligonucleotide comprises at least one phosphodiester linkage.

Embodiment 7

The compound of embodiment 5, wherein each modified internucleoside linkage is a phosphorothioate internucleoside linkage.

Embodiment 8

The compound of any of embodiments 1-7, wherein at least one nucleoside comprises a modified nucleobase.

Embodiment 9

The compound of embodiment 8, wherein the modified nucleobase is a 5-methylcytosine.

Embodiment 10

The compound of any of embodiments 1-9, wherein at least one nucleoside of the modified oligonucleotide comprises a modified sugar.

Embodiment 11

The compound of embodiment 10, wherein each nucleoside of the modified oligonucleotide comprises a modified sugar.

Embodiment 12

The compound of embodiments 10 or 11, wherein the at least one modified sugar is a bicyclic sugar.

Embodiment 13

The compound of embodiment 12, wherein the bicyclic sugar comprises a chemical bridge between the 4' and 2' positions of the sugar, wherein the chemical bridge is selected from: 4'-CH(R)—O-2' and 4'-(CH₂)₂—O-2', wherein R is independently H, C₁-C₆ alkyl, and C₁-C₆ alkoxy.

Embodiment 14

The compound of embodiment 13, wherein the chemical bridge is 4'-CH(R)—O-2' and wherein R is methyl.

15

Embodiment 15

The compound of embodiment 13, wherein the chemical bridge is 4'-CH(R)—O-2' and wherein R is H.

Embodiment 16

The compound of embodiment 13, wherein the chemical bridge is 4'-CH(R)—O-2' and wherein R is —CH₂—O—CH₃.

Embodiment 17

The compound of embodiments 10 or 11, wherein the at least one modified sugar comprises a 2'-O-methoxyethyl group.

Embodiment 18

The compound of any of embodiments 1-10 or 12-16, wherein the modified oligonucleotide is a gapmer.

Embodiment 19

The compound of embodiment 18, wherein the gapmer is any of a 3-8-7 MOE gapmer, a 3-10-7 MOE gapmer, a 4-8-6 MOE gapmer, a 4-10-6 MOE gapmer, a 6-10-4 MOE gapmer, a 6-8-4 MOE gapmer, a 7-8-3 MOE gapmer, or a 7-10-3 MOE gapmer. ²⁵

Embodiment 20

The compound of claim 5, wherein the modified oligonucleotide comprises internucleoside linkages in any of the following patterns: sooooooooooooooss, sooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, sooooooooooooooooooss, or sooooooooooooooooooss, wherein,
s=a phosphorothioate linkage, and
o=a phosphodiester linkage.

Embodiment 19

A composition comprising the compound of any preceding embodiment or salt thereof and at least one of a pharmaceutically acceptable carrier or diluent.

Embodiment 20

The composition of embodiment 19, further comprising a C9ORF72 antisense transcript specific inhibitor.

Embodiment 21

The composition of embodiment 20, wherein the C9ORF72 antisense transcript specific inhibitor is an anti-sense compound.

Embodiment 22

The composition of embodiment 21, wherein the anti-sense compound is a modified oligonucleotide.

Embodiment 23

The composition of embodiment 22, wherein the modified oligonucleotide is single-stranded.

16

Embodiment 24

The composition of embodiments 22 or 23, wherein the modified oligonucleotide has a nucleobase sequence that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% complementary to a C9ORF72 antisense transcript.

Embodiment 25

The composition of embodiment 24, wherein the C9ORF72 antisense transcript has the nucleobase sequence of SEQ ID NO: 18.

Embodiment 26

A method comprising administering to an animal the compound or composition of any preceding embodiment.

Embodiment 27

The method of embodiment 26, wherein the animal is a human.

Embodiment 28

The method of embodiments 26 and 27, wherein administering the compound prevents, treats, ameliorates, or slows progression of a C9ORF72 associated disease.

Embodiment 29

The method of embodiment 28, wherein the C9ORF72 associated disease is caused by a hexanucleotide repeat expansion.

Embodiment 30

40 The method of embodiment 28, wherein the C9ORF72 associated disease is amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, and olivopontocerebellar degeneration (OPCD).

Embodiment 31

The method of embodiments 26-30, wherein the administering reduces nuclear foci.

Embodiment 32

The method of embodiments 26-31, wherein the administering reduces expression of C9ORF72 associated RAN translation products.

Embodiment 33

The method of embodiment 32, wherein the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).


Embodiment 34

65 Use of the compound or composition of any of embodiments 1-33 for the manufacture of a medicament for treating a neurodegenerative disorder.

17

Embodiment 35

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:



18


19

Embodiment 36

A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:

20

(SEQ ID NO: 33)




21
Embodiment 37

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:


22

(SEQ ID NO: 49)



A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:


(SEQ ID NO: 49)



25


Embodiment 39

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:




26

(SEQ ID NO: 47)



A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:


(SEQ ID NO: 47)



29

Embodiment 41

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:



The figure displays the chemical structures of the three RNA subunits of the 30S ribosomal subunit: 16S, 23S, and 16.5S. Each subunit is shown as a complex of ribose sugars, phosphate groups, and nitrogenous bases (adenine, guanine, cytosine, uracil, and thymine). The 16S subunit is the smallest, followed by 16.5S, and then 23S, which is the largest. The structures are arranged in a grid, with the 16S subunit at the top, 16.5S in the middle, and 23S at the bottom. The 16S subunit is shown in a single column, while the 16.5S and 23S subunits are shown in two columns each. The structures are connected by lines, indicating their relative positions and interactions within the 30S subunit. The 16S subunit is shown in a single column, while the 16.5S and 23S subunits are shown in two columns each. The structures are connected by lines, indicating their relative positions and interactions within the 30S subunit.

33

Embodiment 43

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of reducing human C9ORF72 mRNA or protein expression in a mammal.

Embodiment 44

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of ameliorating at least one symptom of amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, or olivopontocerebellar degeneration (OPCD).

Embodiment 45

The compound or composition of embodiment 44, wherein the symptom of ALS is any of motor deficit, anxiety, and denervation.

Embodiment 46

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of delaying progression of disease.

34

Embodiment 47

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of extending survival.

Embodiment 48

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of reducing C9ORF72 associated RAN translation products.

Embodiment 49

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of selectively reducing C9ORF72 pathogenic associated mRNA variants.


Embodiment 50

The compound or composition of any of embodiments 1-25 or 35-42, wherein the compound or composition is capable of reducing nuclear foci.


Embodiment 51

A compound consisting of a modified oligonucleotide according to the following formula, or a salt thereof:

(SEQ ID NO: 33)




35



36

-continued



Embodiment 52

A method comprising administering to an animal the compound of embodiment 51.

(ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, or olivopontocerebellar degeneration (OPCD).

Embodiment 53

The method of embodiment 52, wherein the animal is a human.

Embodiment 58

The method of embodiment 53, wherein the administering reduces nuclear foci.

Embodiment 54

The method of embodiment 53, wherein the administering inhibits C9ORF72.

Embodiment 59

The method of embodiment 53, wherein the administering reduces expression of C9ORF72 associated RAN translation products.

Embodiment 55

The method of embodiment 53, wherein the administering prevents, treats, ameliorates, or slows progression of a C9ORF72 associated disease.

Embodiment 60

The method of embodiment 59, wherein the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).

Embodiment 56

The method of embodiment 55, wherein the C9ORF72 associated disease is caused by a hexanucleotide repeat expansion.

Embodiment 61

A composition consisting of the sodium salt of a modified oligonucleotide according to the following formula:


Embodiment 57

The method of embodiment 55, wherein the C9ORF72 associated disease is any of amyotrophic lateral sclerosis

37

38

(SEQ ID NO: 33)



Embodiment 62

Embodiment 64

A method comprising administering to an animal the composition of embodiment 60.

The method of embodiment 63, wherein the administering inhibits C9ORF72.

Embodiment 63

Embodiment 65

The method of embodiment 62, wherein the animal is a human.

The method of embodiment 62, wherein the administering prevents, treats, ameliorates, or slows progression of a C9ORF72 associated disease.

39

Embodiment 66

The method of embodiment 65, wherein the C9ORF72 associated disease is caused by a hexanucleotide repeat expansion.

Embodiment 67

The method of embodiment 65, wherein the C9ORF72 associated disease is any of amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), corticobasal degeneration syndrome (CBD), atypical Parkinsonian syndrome, or olivopontocerebellar degeneration (OPCD).

Embodiment 68

The method of embodiment 63, wherein the administering reduces nuclear foci.

Embodiment 69

The method of embodiment 63, wherein the administering reduces expression of C9ORF72 associated RAN translation products.

Embodiment 70

The method of embodiment 69, wherein the C9ORF72 associated RAN translation products are any of poly-(glycine-proline), poly-(glycine-alanine), and poly-(glycine-arginine).

Antisense Compounds

Oligomeric compounds include, but are not limited to, oligonucleotides, oligonucleosides, oligonucleotide analogs, oligonucleotide mimetics, antisense compounds, antisense oligonucleotides, and siRNAs. An oligomeric compound may be "antisense" to a target nucleic acid, meaning that it is capable of undergoing hybridization to a target nucleic acid through hydrogen bonding.

In certain embodiments, an antisense compound has a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of the target segment of a target nucleic acid to which it is targeted. In certain such embodiments, an antisense oligonucleotide has a nucleobase sequence that, when written in the 5' to 3' direction, comprises the reverse complement of the target segment of a target nucleic acid to which it is targeted.

In certain embodiments, an antisense compound targeted to a C9ORF72 nucleic acid is 12 to 30 subunits in length. In other words, such antisense compounds are from 12 to 30 linked subunits. In certain embodiments, the antisense compound is 8 to 80, 12 to 50, 15 to 30, 18 to 24, 19 to 22, or 20 linked subunits. In certain embodiments, the antisense compounds are 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 linked subunits in length, or a range defined by any two of the above values. In some embodiments the antisense compound is an antisense oligonucleotide, and the linked subunits are nucleosides.

In certain embodiments antisense oligonucleotides targeted to a C9ORF72 nucleic acid may be shortened or truncated. For example, a single subunit may be deleted from the 5' end (5' truncation), or alternatively from the 3' end (3' truncation). A shortened or truncated antisense

40

compound targeted to a C9ORF72 nucleic acid may have two subunits deleted from the 5' end, or alternatively may have two subunits deleted from the 3' end, of the antisense compound. Alternatively, the deleted nucleosides may be dispersed throughout the antisense compound, for example, in an antisense compound having one nucleoside deleted from the 5' end and one nucleoside deleted from the 3' end.

When a single additional subunit is present in a lengthened antisense compound, the additional subunit may be located at the 5' or 3' end of the antisense compound. When two or more additional subunits are present, the added subunits may be adjacent to each other, for example, in an antisense compound having two subunits added to the 5' end (5' addition), or alternatively to the 3' end (3' addition), of the antisense compound. Alternatively, the added subunits may be dispersed throughout the antisense compound, for example, in an antisense compound having one subunit added to the 5' end and one subunit added to the 3' end.

It is possible to increase or decrease the length of an antisense compound, such as an antisense oligonucleotide, and/or introduce mismatch bases without eliminating activity. For example, in Woolf et al. (Proc. Natl. Acad. Sci. USA 89:7305-7309, 1992), a series of antisense oligonucleotides 13-25 nucleobases in length were tested for their ability to induce cleavage of a target RNA in an oocyte injection model. Antisense oligonucleotides 25 nucleobases in length with 8 or 11 mismatch bases near the ends of the antisense oligonucleotides were able to direct specific cleavage of the target mRNA, albeit to a lesser extent than the antisense oligonucleotides that contained no mismatches. Similarly, target specific cleavage was achieved using 13 nucleobase antisense oligonucleotides, including those with 1 or 3 mismatches.

Gautschi et al (J. Natl. Cancer Inst. 93:463-471, March 2001) demonstrated the ability of an oligonucleotide having 100% complementarity to the bcl-2 mRNA and having 3 mismatches to the bcl-xL mRNA to reduce the expression of both bcl-2 and bcl-xL in vitro and in vivo. Furthermore, this oligonucleotide demonstrated potent anti-tumor activity in vivo.

Maher and Dolnick (Nuc. Acid. Res. 16:3341-3358, 1988) tested a series of tandem 14 nucleobase antisense oligonucleotides, and a 28 and 42 nucleobase antisense oligonucleotides comprised of the sequence of two or three of the tandem antisense oligonucleotides, respectively, for their ability to arrest translation of human DHFR in a rabbit reticulocyte assay. Each of the three 14 nucleobase antisense oligonucleotides alone was able to inhibit translation, albeit at a more modest level than the 28 or 42 nucleobase antisense oligonucleotides.

Antisense Compound Motifs

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid have chemically modified subunits arranged in patterns, or motifs, to confer to the antisense compounds properties such as enhanced inhibitory activity, increased binding affinity for a target nucleic acid, or resistance to degradation by in vivo nucleases.

Chimeric antisense compounds typically contain at least one region modified so as to confer increased resistance to nuclease degradation, increased cellular uptake, increased binding affinity for the target nucleic acid, and/or increased inhibitory activity. A second region of a chimeric antisense compound may optionally serve as a substrate for the cellular endonuclease RNase H, which cleaves the RNA strand of an RNA:DNA duplex.

Antisense compounds having a gapmer motif are considered chimeric antisense compounds. In a gapmer an internal

region having a plurality of nucleotides that supports RNaseH cleavage is positioned between external regions having a plurality of nucleotides that are chemically distinct from the nucleosides of the internal region. In the case of an antisense oligonucleotide having a gapmer motif, the gap segment generally serves as the substrate for endonuclease cleavage, while the wing segments comprise modified nucleosides. In certain embodiments, the regions of a gapmer are differentiated by the types of sugar moieties comprising each distinct region. The types of sugar moieties that are used to differentiate the regions of a gapmer may in some embodiments include β -D-ribonucleosides, β -D-deoxyribonucleosides, 2'-modified nucleosides (such as 2'-modified nucleosides may include 2'-MOE, and 2'-O—CH₃, among others), and bicyclic sugar modified nucleosides (such as bicyclic sugar modified nucleosides may include those having a 4'-(CH₂)_n-O-2' bridge, where n=1 or n=2 and 4'-CH₂—O—CH₂-2'). Preferably, each distinct region comprises uniform sugar moieties. The wing-gap-wing motif is frequently described as "X-Y-Z", where "X" represents the length of the 5' wing region, "Y" represents the length of the gap region, and "Z" represents the length of the 3' wing region. As used herein, a gapmer described as "X-Y-Z" has a configuration such that the gap segment is positioned immediately adjacent to each of the 5' wing segment and the 3' wing segment. Thus, no intervening nucleotides exist between the 5' wing segment and gap segment, or the gap segment and the 3' wing segment. Any of the antisense compounds described herein can have a gapmer motif. In some embodiments, X and Z are the same, in other embodiments they are different. In a preferred embodiment, Y is between 8 and 15 nucleotides. X, Y or Z can be any of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30 or more nucleotides. Thus, gapmers described herein include, but are not limited to, for example 5-10-5, 5-10-4, 4-10-4, 4-10-3, 3-10-3, 2-10-2, 5-9-5, 5-9-4, 4-9-5, 5-8-5, 5-8-4, 4-8-5, 5-7-5, 4-7-5, 5-7-4, or 4-7-4.

In certain embodiments, the antisense compound has a "wingmer" motif, having a wing-gap or gap-wing configuration, i.e. an X-Y or Y-Z configuration as described above for the gapmer configuration. Thus, wingmer configurations described herein include, but are not limited to, for example 5-10, 8-4, 4-12, 12-4, 3-14, 16-2, 18-1, 10-3, 2-10, 1-10, 8-2, 2-13, 5-13, 5-8, or 6-8.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess a 5-10-5 gapmer motif.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess a 5-8-5 gapmer motif.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess sugar modifications in any of the following patterns: eekkkddddddekeee, eekkkddddddekeee, ekddddddekeee, kekedddddddekeee, and eekkkddddddekeee; wherein,

e=a 2'-O-methoxyethyl modified nucleoside

d=a 2'-deoxynucleoside, and

k=a cEt nucleoside.

In certain embodiments, an antisense compound targeted to a C9ORF72 nucleic acid has a gap-narrowed motif. In certain embodiments, a gap-narrowed antisense oligonucleotide targeted to a C9ORF72 nucleic acid has a gap segment of 9, 8, 7, or 6 2'-deoxynucleotides positioned immediately adjacent to and between wing segments of 5, 4, 3, 2, or 1 chemically modified nucleosides. In certain embodiments, the chemical modification comprises a bicyclic sugar. In certain embodiments, the bicyclic sugar comprises a 4' to 2' bridge selected from among: 4'-(CH₂)_n-O-2' bridge, wherein n is 1 or 2; and 4'-CH₂—O—CH₂-2'. In certain

embodiments, the bicyclic sugar is comprises a 4'-CH(CH₃)—O-2' bridge. In certain embodiments, the chemical modification comprises a non-bicyclic 2'-modified sugar moiety. In certain embodiments, the non-bicyclic 2'-modified sugar moiety comprises a 2'-O-methylethyl group or a 2'-O-methyl group.

Target Nucleic Acids, Target Regions and Nucleotide Sequences

Nucleotide sequences that encode C9ORF72 include, without limitation, the following: the complement of GENBANK Accession No. NM_001256054.1 (incorporated herein as SEQ ID NO: 1), the complement of GENBANK Accession No. NT_008413.18 truncated from nucleobase 27535000 to 27565000 (incorporated herein as SEQ ID NO: 2), GENBANK Accession No. BQ068108.1 (incorporated herein as SEQ ID NO: 3), GENBANK Accession No. NM_018325.3 (incorporated herein as SEQ ID NO: 4), GENBANK Accession No. DN993522.1 (incorporated herein as SEQ ID NO: 5), GENBANK Accession No. NM_145005.5 (incorporated herein as SEQ ID NO: 6), GENBANK Accession No. DB079375.1 (incorporated herein as SEQ ID NO: 7), GENBANK Accession No. BU194591.1 (incorporated herein as SEQ ID NO: 8), Sequence Identifier 4141_014_A (incorporated herein as SEQ ID NO: 9), and Sequence Identifier 4008_73_A (incorporated herein as SEQ ID NO: 10), and GENBANK Accession No. NW_001101662.1 truncated from nucleobases 8522000 to U.S. Pat. No. 8,552,000 (incorporated herein as SEQ ID NO: 11).

It is understood that the sequence set forth in each SEQ ID NO in the Examples contained herein is independent of any modification to a sugar moiety, an internucleoside linkage, or a nucleobase. As such, antisense compounds defined by a SEQ ID NO may comprise, independently, one or more modifications to a sugar moiety, an internucleoside linkage, or a nucleobase. Antisense compounds described by Isis Number (Isis No) indicate a combination of nucleobase sequence and motif.

In certain embodiments, a target region is a structurally defined region of the target nucleic acid. For example, a target region may encompass a 3' UTR, a 5' UTR, an exon, an intron, an exon/intron junction, a coding region, a translation initiation region, translation termination region, or other defined nucleic acid region. The structurally defined regions for C9ORF72 can be obtained by accession number from sequence databases such as NCBI and such information is incorporated herein by reference. In certain embodiments, a target region may encompass the sequence from a 5' target site of one target segment within the target region to a 3' target site of another target segment within the same target region.

Targeting includes determination of at least one target segment to which an antisense compound hybridizes, such that a desired effect occurs. In certain embodiments, the desired effect is a reduction in mRNA target nucleic acid levels. In certain embodiments, the desired effect is reduction of levels of protein encoded by the target nucleic acid or a phenotypic change associated with the target nucleic acid.

A target region may contain one or more target segments. Multiple target segments within a target region may be overlapping. Alternatively, they may be non-overlapping. In certain embodiments, target segments within a target region are separated by no more than about 300 nucleotides. In certain embodiments, target segments within a target region are separated by a number of nucleotides that is, is about, is no more than, is no more than about, 250, 200, 150, 100, 90,

80, 70, 60, 50, 40, 30, 20, or 10 nucleotides on the target nucleic acid, or is a range defined by any two of the preceding values. In certain embodiments, target segments within a target region are separated by no more than, or no more than about, 5 nucleotides on the target nucleic acid. In certain embodiments, target segments are contiguous. Contemplated are target regions defined by a range having a starting nucleic acid that is any of the 5' target sites or 3' target sites listed herein.

Suitable target segments may be found within a 5' UTR, a coding region, a 3' UTR, an intron, an exon, or an exon/intron junction. Target segments containing a start codon or a stop codon are also suitable target segments. A suitable target segment may specifically exclude a certain structurally defined region such as the start codon or stop codon.

The determination of suitable target segments may include a comparison of the sequence of a target nucleic acid to other sequences throughout the genome. For example, the BLAST algorithm may be used to identify regions of similarity amongst different nucleic acids. This comparison can prevent the selection of antisense compound sequences that may hybridize in a non-specific manner to sequences other than a selected target nucleic acid (i.e., non-target or off-target sequences).

There may be variation in activity (e.g., as defined by percent reduction of target nucleic acid levels) of the antisense compounds within a target region. In certain embodiments, reductions in C9ORF72 mRNA levels are indicative of inhibition of C9ORF72 expression. Reductions in levels of a C9ORF72 protein are also indicative of inhibition of target mRNA expression. Reduction in the presence of expanded C9ORF72 RNA foci are indicative of inhibition of C9ORF72 expression. Further, phenotypic changes are indicative of inhibition of C9ORF72 expression. For example, improved motor function and respiration may be indicative of inhibition of C9ORF72 expression.

Hybridization

In some embodiments, hybridization occurs between an antisense compound disclosed herein and a C9ORF72 nucleic acid. The most common mechanism of hybridization involves hydrogen bonding (e.g., Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding) between complementary nucleobases of the nucleic acid molecules.

Hybridization can occur under varying conditions. Stringent conditions are sequence-dependent and are determined by the nature and composition of the nucleic acid molecules to be hybridized.

Methods of determining whether a sequence is specifically hybridizable to a target nucleic acid are well known in the art. In certain embodiments, the antisense compounds provided herein are specifically hybridizable with a C9ORF72 nucleic acid.

Complementarity

An antisense compound and a target nucleic acid are complementary to each other when a sufficient number of nucleobases of the antisense compound can hydrogen bond with the corresponding nucleobases of the target nucleic acid, such that a desired effect will occur (e.g., antisense inhibition of a target nucleic acid, such as a C9ORF72 nucleic acid).

Non-complementary nucleobases between an antisense compound and a C9ORF72 nucleic acid may be tolerated provided that the antisense compound remains able to specifically hybridize to a target nucleic acid. Moreover, an antisense compound may hybridize over one or more segments of a C9ORF72 nucleic acid such that intervening or

adjacent segments are not involved in the hybridization event (e.g., a loop structure, mismatch or hairpin structure).

In certain embodiments, the antisense compounds provided herein, or a specified portion thereof, are, or are at least, 70%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% complementary to a C9ORF72 nucleic acid, a target region, target segment, or specified portion thereof. Percent complementarity of an antisense compound with a target nucleic acid can be determined using routine methods.

For example, an antisense compound in which 18 of 20 nucleobases of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleobases may be clustered or interspersed with complementary nucleobases and need not be contiguous to each other or to complementary nucleobases. As such, an antisense compound which is 18 nucleobases in length having 4 (four) noncomplementary nucleobases which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid and would thus fall within the scope of the present invention. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., *J. Mol. Biol.*, 1990, 215, 403 410; Zhang and Madden, *Genome Res.*, 1997, 7, 649 656). Percent homology, sequence identity or complementarity, can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (*Adv. Appl. Math.*, 1981, 2, 482 489).

In certain embodiments, the antisense compounds provided herein, or specified portions thereof, are fully complementary (i.e., 100% complementary) to a target nucleic acid, or specified portion thereof. For example, an antisense compound may be fully complementary to a C9ORF72 nucleic acid, or a target region, or a target segment or target sequence thereof. As used herein, "fully complementary" means each nucleobase of an antisense compound is capable of precise base pairing with the corresponding nucleobases of a target nucleic acid. For example, a 20 nucleobase antisense compound is fully complementary to a target sequence that is 400 nucleobases long, so long as there is a corresponding 20 nucleobase portion of the target nucleic acid that is fully complementary to the antisense compound. Fully complementary can also be used in reference to a specified portion of the first and/or the second nucleic acid. For example, a 20 nucleobase portion of a 30 nucleobase antisense compound can be "fully complementary" to a target sequence that is 400 nucleobases long. The 20 nucleobase portion of the 30 nucleobase oligonucleotide is fully complementary to the target sequence if the target sequence has a corresponding 20 nucleobase portion wherein each nucleobase is complementary to the 20 nucleobase portion of the antisense compound. At the same time, the entire 30 nucleobase antisense compound may or may not be fully complementary to the target sequence, depending on whether the remaining 10 nucleobases of the antisense compound are also complementary to the target sequence.

The location of a non-complementary nucleobase may be at the 5' end or 3' end of the antisense compound. Alternatively, the non-complementary nucleobase or nucleobases may be at an internal position of the antisense compound.

When two or more non-complementary nucleobases are present, they may be contiguous (i.e., linked) or non-contiguous. In one embodiment, a non-complementary nucleobase is located in the wing segment of a gapmer antisense oligonucleotide.

In certain embodiments, antisense compounds that are, or are up to 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleobases in length comprise no more than 4, no more than 3, no more than 2, or no more than 1 non-complementary nucleobase(s) relative to a target nucleic acid, such as a C9ORF72 nucleic acid, or specified portion thereof.

In certain embodiments, antisense compounds that are, or are up to 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleobases in length comprise no more than 6, no more than 5, no more than 4, no more than 3, no more than 2, or no more than 1 non-complementary nucleobase(s) relative to a target nucleic acid, such as a C9ORF72 nucleic acid, or specified portion thereof.

The antisense compounds provided herein also include those which are complementary to a portion of a target nucleic acid. As used herein, "portion" refers to a defined number of contiguous (i.e. linked) nucleobases within a region or segment of a target nucleic acid. A "portion" can also refer to a defined number of contiguous nucleobases of an antisense compound. In certain embodiments, the antisense compounds, are complementary to at least an 8 nucleobase portion of a target segment. In certain embodiments, the antisense compounds are complementary to at least a 9 nucleobase portion of a target segment. In certain embodiments, the antisense compounds are complementary to at least a 10 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least an 11 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 12 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 13 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 14 nucleobase portion of a target segment. In certain embodiments, the antisense compounds, are complementary to at least a 15 nucleobase portion of a target segment. Also contemplated are antisense compounds that are complementary to at least a 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more nucleobase portion of a target segment, or a range defined by any two of these values.

Identity

The antisense compounds provided herein may also have a defined percent identity to a particular nucleotide sequence, SEQ ID NO, or compound represented by a specific Isis number, or portion thereof. As used herein, an antisense compound is identical to the sequence disclosed herein if it has the same nucleobase pairing ability. For example, a RNA which contains uracil in place of thymidine in a disclosed DNA sequence would be considered identical to the DNA sequence since both uracil and thymidine pair with adenine. Shortened and lengthened versions of the antisense compounds described herein as well as compounds having non-identical bases relative to the antisense compounds provided herein also are contemplated. The non-identical bases may be adjacent to each other or dispersed throughout the antisense compound. Percent identity of an antisense compound is calculated according to the number of bases that have identical base pairing relative to the sequence to which it is being compared.

In certain embodiments, the antisense compounds, or portions thereof, are at least 70%, 75%, 80%, 85%, 90%,

95%, 96%, 97%, 98%, 99% or 100% identical to one or more of the antisense compounds or SEQ ID NOs, or a portion thereof, disclosed herein.

In certain embodiments, a portion of the antisense compound is compared to an equal length portion of the target nucleic acid. In certain embodiments, an 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleobase portion is compared to an equal length portion of the target nucleic acid.

In certain embodiments, a portion of the antisense oligonucleotide is compared to an equal length portion of the target nucleic acid. In certain embodiments, an 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleobase portion is compared to an equal length portion of the target nucleic acid.

Modifications

A nucleoside is a base-sugar combination. The nucleobase (also known as base) portion of the nucleoside is normally a heterocyclic base moiety. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to the 2', 3' or 5' hydroxyl moiety of the sugar. Oligonucleotides are formed through the covalent linkage of adjacent nucleosides to one another, to form a linear polymeric oligonucleotide. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside linkages of the oligonucleotide.

Modifications to antisense compounds encompass substitutions or changes to internucleoside linkages, sugar moieties, or nucleobases. Modified antisense compounds are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target, increased stability in the presence of nucleases, or increased inhibitory activity.

Chemically modified nucleosides may also be employed to increase the binding affinity of a shortened or truncated antisense oligonucleotide for its target nucleic acid. Consequently, comparable results can often be obtained with shorter antisense compounds that have such chemically modified nucleosides.

Modified Internucleoside Linkages

The naturally occurring internucleoside linkage of RNA and DNA is a 3' to 5' phosphodiester linkage. Antisense compounds having one or more modified, i.e. non-naturally occurring, internucleoside linkages are often selected over antisense compounds having naturally occurring internucleoside linkages because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for target nucleic acids, and increased stability in the presence of nucleases.

Oligonucleotides having modified internucleoside linkages include internucleoside linkages that retain a phosphorus atom as well as internucleoside linkages that do not have a phosphorus atom. Representative phosphorus containing internucleoside linkages include, but are not limited to, phosphodiester, phosphotriester, methylphosphonates, phosphoramidate, and phosphorothioates. Methods of preparation of phosphorous-containing and non-phosphorous-containing linkages are well known.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid comprise one or more modified internucleoside linkages. In certain embodiments, the modified internucleoside linkages are interspersed throughout the antisense compound. In certain embodiments, the modified internucleoside linkages are phosphorothioate linkages. In certain embodiments, each internucleoside linkage of an

antisense compound is a phosphorothioate internucleoside linkage. In certain embodiments, the antisense compounds targeted to a C9ORF72 nucleic acid comprise at least one phosphodiester linkage and at least one phosphorothioate linkage.

In certain embodiments, antisense compounds targeted to a C9ORF72 nucleic acid possess internucleoside linkages in any of the following patterns: sooooooooooooo, sooooooooooooo, sooooooooooooo, and sooooooooooooo; wherein,

s=a phosphorothioate linkage, and

o=a phosphodiester linkage.

Modified Sugar Moieties

Antisense compounds of the invention can optionally contain one or more nucleosides wherein the sugar group has been modified. Such sugar modified nucleosides may impart enhanced nuclease stability, increased binding affinity, or some other beneficial biological property to the antisense compounds. In certain embodiments, nucleosides comprise chemically modified ribofuranose ring moieties. Examples of chemically modified ribofuranose rings include without limitation, addition of substituent groups (including 5' and 2' substituent groups, bridging of non-geminal ring atoms to form bicyclic nucleic acids (BNA), replacement of the ribosyl ring oxygen atom with S, N(R), or C(R₁)(R₂) (R, R₁ and R₂ are each independently H, C₁-C₁₂ alkyl or a protecting group) and combinations thereof. Examples of chemically modified sugars include 2'-F-5'-methyl substituted nucleoside (see PCT International Application WO 2008/101157 Published on Aug. 21, 2008 for other disclosed 5',2'-bis substituted nucleosides) or replacement of the ribosyl ring oxygen atom with S with further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on Jun. 16, 2005) or alternatively 5'-substitution of a BNA (see PCT International Application WO 2007/134181 Published on Nov. 22, 2007 wherein LNA is substituted with for example a 5'-methyl or a 5'-vinyl group).

Examples of nucleosides having modified sugar moieties include without limitation nucleosides comprising 5'-vinyl, 5'-methyl (R or S), 4'-S, 2'-F, 2'-OCH₃, 2'-OCH₂CH₃, 2'-OCH₂CH₂F and 2'-O(CH₂)₂₀CH₃ substituent groups. The substituent at the 2' position can also be selected from allyl, amino, azido, thio, O-allyl, O—C₁-C₁₀ alkyl, OCF₃, OCH₂F, O(CH₂)₂SCH₃, O(CH₂)₂—O—N((R_m)(R_n), O—CH₂—C(=O)—N((R_m)(R_n), and O—CH₂—C(=O)—N(R₁)—(CH₂)₂—N((R_m)(R_n), where each R₁, (R_m) and R_n is, independently, H or substituted or unsubstituted C₁-C₁₀ alkyl.

As used herein, “bicyclic nucleosides” refer to modified nucleosides comprising a bicyclic sugar moiety. Examples of bicyclic nucleic acids (BNAs) include without limitation nucleosides comprising a bridge between the 4' and the 2' ribosyl ring atoms. In certain embodiments, antisense compounds provided herein include one or more BNA nucleosides wherein the bridge comprises one of the formulas: 4'-(CH₂)—O-2' (LNA); 4'-(CH₂)—S-2'; 4'-(CH₂)₂—O-2' (ENA); 4'-CH(CH₃)—O-2' and 4'-CH(CH₂OCH₃)—O-2' (and analogs thereof see U.S. Pat. No. 7,399,845, issued on Jul. 15, 2008); 4'-C(CH₃)(CH₃)—O-2' (and analogs thereof see PCT/US2008/068922 published as WO/2009/006478, published Jan. 8, 2009); 4'-CH₂—N(OCH₃)-2' (and analogs thereof see PCT/US2008/064591 published as WO/2008/150729, published Dec. 11, 2008); 4'-CH₂—O—N(CH₃)-2' (see published U.S. Patent Application US2004-0171570, published Sep. 2, 2004); 4'-CH₂—N(R)—O-2', wherein R is H, C₁-C₁₀ alkyl, or a protecting group (see U.S. Pat. No.

7,427,672, issued on Sep. 23, 2008); 4'-CH₂—C(H)(CH₃)-2' (see Chattopadhyaya et al., *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH₂—C(=CH₂)-2' (and analogs thereof see PCT/US2008/066154 published as WO 2008/154401, published on Dec. 8, 2008).

Further bicyclic nucleosides have been reported in published literature (see for example: Srivastava et al., *J. Am. Chem. Soc.*, 2007, 129(26) 8362-8379; Frieden et al., *Nucleic Acids Research*, 2003, 21, 6365-6372; Elayadi et al., *Curr. Opinion Inven. Drugs*, 2001, 2, 558-561; Braasch et al., *Chem. Biol.*, 2001, 8, 1-7; Orum et al., *Curr. Opinion Mol. Ther.*, 2001, 3, 239-243; Wahlestedt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2000, 97, 5633-5638; Singh et al., *Chem. Commun.*, 1998, 4, 455-456; Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630; Kumar et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222; Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039; U.S. Pat. Nos. 7,399,845; 7,053,207; 7,034,133; 6,794,499; 6,770,748; 6,670,461; 6,525,191; 6,268,490; U.S. Patent Publication Nos.: US2008-0039618; US2007-0287831; US2004-0171570; U.S. patent application Ser. No. 12/129,154; 61/099,844; 61/097,787; 61/086,231; 61/056,564; 61/026,998; 61/026,995; 60/989,574; International applications WO 2007/134181; WO 2005/021570; WO 2004/106356; and PCT International Applications Nos.: PCT/US2008/068922; PCT/US2008/066154; and PCT/US2008/064591). Each of the foregoing bicyclic nucleosides can be prepared having one or more stereochemical sugar configurations including for example α-L-ribofuranose and β-D-ribofuranose (see PCT international application PCT/DK98/00393, published on Mar. 25, 1999 as WO 99/14226).

As used herein, “monocyclic nucleosides” refer to nucleosides comprising modified sugar moieties that are not bicyclic sugar moieties. In certain embodiments, the sugar moiety, or sugar moiety analogue, of a nucleoside may be modified or substituted at any position.

As used herein, “4'-2' bicyclic nucleoside” or “4' to 2' bicyclic nucleoside” refers to a bicyclic nucleoside comprising a furanose ring comprising a bridge connecting two carbon atoms of the furanose ring connects the 2' carbon atom and the 4' carbon atom of the sugar ring.

In certain embodiments, bicyclic sugar moieties of BNA nucleosides include, but are not limited to, compounds having at least one bridge between the 4' and the 2' carbon atoms of the pentofuranosyl sugar moiety including without limitation, bridges comprising 1 or from 1 to 4 linked groups independently selected from —[C(R_a)(R_b)]_n—, —C(R_a)=C(R_b)—, —C(R_a)=N—, —C(=NR_a)—, —C(=O)—, —C(=S)—, —O—, —Si(R_a)₂—, —S(=O)_x—, and —N(R_a)—; wherein: x is 0, 1, or 2; n is 1, 2, 3, or 4; each R_a and R_b is, independently, H, a protecting group, hydroxyl, C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, heterocycle radical, substituted heterocycle radical, heteroaryl, substituted heteroaryl, C₅-C₇ alicyclic radical, substituted C₅-C₇ alicyclic radical, halogen, OJ₁, NJ₁J₂, SJ₁, N₃, COOJ₁, acyl (C(=O)—H), substituted acyl, CN, sulfonyl (S(=O)₂-J₁), or sulfoxyl (S(=O)-J₁); and

each J₁ and J₂ is, independently, H, C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, acyl (C(=O)—H), substituted acyl, a heterocycle radical, a substituted heterocycle radical, C₁-C₁₀ aminoalkyl, substituted C₁-C₁₀ aminoalkyl or a protecting group.


49

In certain embodiments, the bridge of a bicyclic sugar moiety is, $—[C(R_a)(R_b)]_n—$, $—[C(R_a)(R_b)]_n—O—$, $—C(R_aR_b)—N(R)—O—$ or $—C(R_aR_b)—O—N(R)—$. In certain embodiments, the bridge is 4'-CH₂-2', 4'-(CH₂)₂-2', 4'-(CH₂)₃-2', 4'-CH₂-O-2', 4'-(CH₂)₂-O-2', 4'-CH₂-O-N(R)-2' and 4'-CH₂-N(R)-O-2' wherein each R is, independently, H, a protecting group or C₁-C₁₀ alkyl.

In certain embodiments, bicyclic nucleosides are further defined by isomeric configuration. For example, a nucleoside comprising a 4'-(CH₂)—O-2' bridge, may be in the α-L configuration or in the β-D configuration. Previously, α-L-methyleneoxy (4'-CH₂—O-2') BNA's have been incorporated into antisense oligonucleotides that showed antisense activity (Frieden et al., *Nucleic Acids Research*, 2003, 21, 6365-6372).

In certain embodiments, bicyclic nucleosides include those having a 4' to 2' bridge wherein such bridges include without limitation, α-L-4'-(CH₂)—O-2', β-D-4'-CH₂—O-2', 4'-(CH₂)₂—O-2', 4'-CH₂—O—N(R)-2', 4'-CH₂—N(R)—O-2', 4'-CH(CH₃)—O-2', 4'-CH₂—S-2', 4'-CH₂—N(R)-2', 4'-CH₂—CH(CH₃)-2', and 4'-(CH₂)₃-2', wherein R is H, a protecting group or C₁-C₁₂ alkyl.

In certain embodiment, bicyclic nucleosides have the formula:



wherein:


Bx is a heterocyclic base moiety;

-Q_a-Q_b-Q_c- is $—CH_2—N(R_c)—CH_2—$, $—(=O)—N(R_c)—CH_2—$, $—CH_2—O—N(R_c)—$, $—CH_2—N(R_c)—O—$ or $—N(R_c)—O—CH_2—$;

R_c is C₁-C₁₀ alkyl or an amino protecting group; and

T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium.

In certain embodiments, bicyclic nucleosides have the formula:



wherein:

Bx is a heterocyclic base moiety;


T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

Z_a is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, substituted C₁-C₆ alkyl, substituted C₂-C₆ alkenyl, substituted C₂-C₆ alkynyl, acyl, substituted acyl, substituted amide, thiol or substituted thiol.

50

In one embodiment, each of the substituted groups, is, independently, mono or poly substituted with substituent groups independently selected from halogen, oxo, hydroxyl, OJ_c, NJ_cJ_d, SJ_c, N₃, OC(=X)J_c, and NJ_eC(=X)NJ_cJ_d, wherein each J_c, J_d and J_e is, independently, H, C₁-C₆ alkyl, or substituted C₁-C₆ alkyl and X is O or NJ_c.

In certain embodiments, bicyclic nucleosides have the formula:




wherein:

Bx is a heterocyclic base moiety;

T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

Z_b is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, substituted C₁-C₆ alkyl, substituted C₂-C₆ alkenyl, substituted C₂-C₆ alkynyl or substituted acyl (C(=O)—).

In certain embodiments, bicyclic nucleosides have the formula:



wherein:

Bx is a heterocyclic base moiety;


T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

R_d is C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl;

each q_a, q_b, q_c and q_d is, independently, H, halogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl, C₁-C₆ alkoxyl, substituted C₁-C₆ alkoxyl, acyl, substituted acyl, C₁-C₆ aminoalkyl or substituted C₁-C₆ aminoalkyl;

In certain embodiments, bicyclic nucleosides have the formula:

51



wherein:

Bx is a heterocyclic base moiety;

T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

q_a, q_b, q_e and q_f are each, independently, hydrogen, halogen, C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₁-C₁₀ alkoxy, substituted C₁-C₁₀ alkoxy, OJ_j, SJ_j, SOJ_j, SO₂J_j, NJ_jJ_k, N₃, CN, C(=O)OJ_j, C(=O)NJ_jJ_k, C(=O)J_j, O—C(=O)NJ_jJ_k, N(H)C(=NH)NJ_jJ_k, N(H)C(=O)NJ_jJ_k or N(H)C(=S)NJ_jJ_k;


or q_e and q_f together are =C(q_g)(q_h);

q_g and q_h are each, independently, H, halogen, C₁-C₁₀ alkyl or substituted C₁-C₁₀ alkyl.

The synthesis and preparation of adenine, cytosine, guanine, 5-methyl-cytosine, thymine and uracil bicyclic nucleosides having a 4'-CH₂—O-2' bridge, along with their oligomerization, and nucleic acid recognition properties have been described (Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630). The synthesis of bicyclic nucleosides has also been described in WO 98/39352 and WO99/14226.

Analogues of various bicyclic nucleosides that have 4' to 2' bridging groups such as 4'-CH₂—O-2' and 4'-CH₂—S-2', have also been prepared (Kumar et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222). Preparation of oligodeoxyribonucleotide duplexes comprising bicyclic nucleosides for use as substrates for nucleic acid polymerases has also been described (Wengel et al., WO 99/14226). Furthermore, synthesis of 2'-amino-BNA, a novel conformationally restricted high-affinity oligonucleotide analog has been described in the art (Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039). In addition, 2'-amino- and 2'-methylamino-BNA's have been prepared and the thermal stability of their duplexes with complementary RNA and DNA strands has been previously reported.

In certain embodiments, bicyclic nucleosides have the formula:



wherein:

Bx is a heterocyclic base moiety;

T_a and T_b are each, independently H, a hydroxyl protecting group, a conjugate group, a reactive phosphorus group, a phosphorus moiety or a covalent attachment to a support medium;

each q_i, q_j, q_k and q_l is, independently, H, halogen, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted


52

C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₁-C₁₂ alkoxy, substituted C₁-C₁₂ alkoxy, OJ_j, SJ_j, SOJ_j, SO₂J_j, NJ_jJ_k, N₃, CN, C(=O)OJ_j, C(=O)NJ_jJ_k, C(=O)J_j, O—C(=O)NJ_jJ_k, N(H)C(=NH)NJ_jJ_k, N(H)C(=O)NJ_jJ_k or N(H)C(=S)NJ_jJ_k; and

q_i and q_j or q_i and q_k together are =C(q_g)(q_h), wherein q_g and q_h are each, independently, H, halogen, C₁-C₁₂ alkyl or substituted C₁-C₁₂ alkyl.


One carbocyclic bicyclic nucleoside having a 4'-(CH₂)₃-2' bridge and the alkenyl analog bridge 4'-CH=CH—CH₂-2' have been described (Frier et al., *Nucleic Acids Research*, 1997, 25(22), 4429-4443 and Albaek et al., *J. Org. Chem.*, 2006, 71, 7731-7740). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (Srivastava et al., *J. Am. Chem. Soc.* 2007, 129(26), 8362-8379).

In certain embodiments, bicyclic nucleosides include, but are not limited to, (A) α-L-methyleneoxy (4'-CH₂—O-2') BNA, (B) β-D-methyleneoxy (4'-CH₂—O-2') BNA, (C) ethyleneoxy (4'-(CH₂)₂—O-2') BNA, (D) aminooxy (4'-CH₂—O—N(R)-2') BNA, (E) oxyamino (4'-CH₂—N(R)—O-2') BNA, (F) methyl(methyleneoxy) (4'-CH(CH₃)—O-2') BNA (also referred to as constrained ethyl or cEt), (G) methylene-thio (4'-CH₂—S-2') BNA, (H) methylene-amino (4'-CH₂—N(R)-2') BNA, (I) methyl carbocyclic (4'-CH₂—CH(CH₃)-2') BNA, (J) propylene carbocyclic (4'-(CH₂)₃-2') BNA, and (K) vinyl BNA as depicted below.



53

-continued



wherein Bx is the base moiety and R is, independently, H, a protecting group, C₁-C₆ alkyl or C₁-C₆ alkoxy.

As used herein, the term “modified tetrahydropyran nucleoside” or “modified THP nucleoside” means a nucleoside having a six-membered tetrahydropyran “sugar” substituted for the pentofuranosyl residue in normal nucleosides and can be referred to as a sugar surrogate. Modified THP nucleosides include, but are not limited to, what is referred to in the art as hexitol nucleic acid (HNA), anitol nucleic

54

acid (ANA), manitol nucleic acid (MNA) (see Leumann, *Bioorg. Med. Chem.*, 2002, 10, 841-854) or fluoro HNA (F-HNA) having a tetrahydropyran ring system as illustrated below.

(E)

5

(F)

10

15

(G)

20

(H)

25

30

(I)

35

(J)

45


(K)

50


55

60

65



In certain embodiment, sugar surrogates are selected having the formula:



wherein:

Bx is a heterocyclic base moiety;

T₃ and T₄ are each, independently, an internucleoside linking group linking the tetrahydropyran nucleoside analog to the oligomeric compound or one of T₃ and T₄ is an internucleoside linking group linking the tetrahydropyran nucleoside analog to an oligomeric compound or oligonucleotide and the other of T₃ and T₄ is H, a hydroxyl protecting group, a linked conjugate group or a 5' or 3'-terminal group;


q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each independently, H, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl or substituted C₂-C₆ alkynyl; and

one of R₁ and R₂ is hydrogen and the other is selected from halogen, substituted or unsubstituted alkoxy, NJ₁J₂, SJ₁, N₃, OC(=X)J₁, OC(=X)NJ₁J₂, NJ₃C(=X)NJ₁J₂ and CN, wherein X is O, S or NJ₁ and each J₁, J₂ and J₃ is, independently, H or C₁-C₆ alkyl.

In certain embodiments, q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each H. In certain embodiments, at least one of q₁, q₂, q₃, q₄, q₅, q₆ and q₇ is other than H. In certain embodiments, at least one of q₁, q₂, q₃, q₄, q₅, q₆ and q₇ is methyl. In certain embodiments, THP nucleosides are provided wherein one of R₁ and R₂ is F. In certain embodiments, R₁ is fluoro and R₂ is H; R₁ is methoxy and R₂ is H, and R₁ is methoxyethoxy and R₂ is H.

In certain embodiments, sugar surrogates comprise rings having more than 5 atoms and more than one heteroatom. For example nucleosides comprising morpholino sugar moieties and their use in oligomeric compounds has been reported (see for example: Braasch et al., *Biochemistry*, 2002, 41, 4503-4510; and U.S. Pat. Nos. 5,698,685; 5,166,315; 5,185,444; and 5,034,506). As used here, the term “morpholino” means a sugar surrogate having the following formula:


55



In certain embodiments, morpholinos may be modified, for example by adding or altering various substituent groups from the above morpholino structure. Such sugar surrogates are referred to herein as “modified morpholinos.”

Combinations of modifications are also provided without limitation, such as 2'-F-5'-methyl substituted nucleosides (see PCT International Application WO 2008/101157 published on Aug. 21, 2008 for other disclosed 5',2'-bis substituted nucleosides) and replacement of the ribosyl ring oxygen atom with S and further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on Jun. 16, 2005) or alternatively 5'-substitution of a bicyclic nucleic acid (see PCT International Application WO 2007/134181, published on Nov. 22, 2007 wherein a 4'-CH₂—O-2' bicyclic nucleoside is further substituted at the 5' position with a 5'-methyl or a 5'-vinyl group). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (see, e.g., Srivastava et al., *J. Am. Chem. Soc.* 2007, 129 (26), 8362-8379).

In certain embodiments, antisense compounds comprise one or more modified cyclohexenyl nucleosides, which is a nucleoside having a six-membered cyclohexenyl in place of the pentofuranosyl residue in naturally occurring nucleosides. Modified cyclohexenyl nucleosides include, but are not limited to those described in the art (see for example commonly owned, published PCT Application WO 2010/036696, published on Apr. 10, 2010, Robeyns et al., *J. Am. Chem. Soc.*, 2008, 130(6), 1979-1984; Horváth et al., *Tetrahedron Letters*, 2007, 48, 3621-3623; Nauwelaerts et al., *J. Am. Chem. Soc.*, 2007, 129(30), 9340-9348; Gu et al., *Nucleosides, Nucleotides & Nucleic Acids*, 2005, 24(5-7), 993-998; Nauwelaerts et al., *Nucleic Acids Research*, 2005, 33(8), 2452-2463; Robeyns et al., *Acta Crystallographica, Section F: Structural Biology and Crystallization Communications*, 2005, F61(6), 585-586; Gu et al., *Tetrahedron*, 2004, 60(9), 2111-2123; Gu et al., *Oligonucleotides*, 2003, 13(6), 479-489; Wang et al., *J. Org. Chem.*, 2003, 68, 4499-4505; Verbeure et al., *Nucleic Acids Research*, 2001, 29(24), 4941-4947; Wang et al., *J. Org. Chem.*, 2001, 66, 8478-82; Wang et al., *Nucleosides, Nucleotides & Nucleic Acids*, 2001, 20(4-7), 785-788; Wang et al., *J. Am. Chem.*, 2000, 122, 8595-8602; Published PCT application, WO 06/047842; and Published PCT Application WO 01/049687; the text of each is incorporated by reference herein, in their entirety). Certain modified cyclohexenyl nucleosides have Formula X.



56

wherein independently for each of said at least one cyclohexenyl nucleoside analog of Formula X:

Bx is a heterocyclic base moiety;

T₃ and T₄ are each, independently, an internucleoside linking group linking the cyclohexenyl nucleoside analog to an antisense compound or one of T₃ and T₄ is an internucleoside linking group linking the tetrahydropyran nucleoside analog to an antisense compound and the other of T₃ and T₄ is H, a hydroxyl protecting group, a linked conjugate group, or a 5'- or 3'-terminal group; and

q₁, q₂, q₃, q₄, q₅, q₆, q₇, q₈ and q₉ are each, independently, H, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl, substituted C₂-C₆ alkynyl or other sugar substituent group.

Many other monocyclic, bicyclic and tricyclic ring systems are known in the art and are suitable as sugar surrogates that can be used to modify nucleosides for incorporation into oligomeric compounds as provided herein (see for example review article: Leumann, Christian J. *Bioorg. & Med. Chem.*, 2002, 10, 841-854). Such ring systems can undergo various additional substitutions to further enhance their activity.

As used herein, “2'-modified sugar” means a furanosyl sugar modified at the 2' position. In certain embodiments, such modifications include substituents selected from: a halide, including, but not limited to substituted and unsubstituted alkoxy, substituted and unsubstituted thioalkyl, substituted and unsubstituted amino alkyl, substituted and unsubstituted alkyl, substituted and unsubstituted allyl, and substituted and unsubstituted alkynyl. In certain embodiments, 2' modifications are selected from substituents including, but not limited to: O[(CH₂)_nO]_mCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nF, O(CH₂)_nONH₂, OCH₂C(=O)N(H)CH₃, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. Other 2'-substituent groups can also be selected from: C₁-C₁₂ alkyl, substituted alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, F, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving pharmacokinetic properties, or a group for improving the pharmacodynamic properties of an antisense compound, and other substituents having similar properties. In certain embodiments, modified nucleosides comprise a 2'-MOE side chain (Baker et al., *J. Biol. Chem.*, 1997, 272, 11944-12000). Such 2'-MOE substitution have been described as having improved binding affinity compared to unmodified nucleosides and to other modified nucleosides, such as 2'-O-methyl, O-propyl, and O-aminopropyl. Oligonucleotides having the 2'-MOE substituent also have been shown to be antisense inhibitors of gene expression with promising features for in vivo use (Martin, *Helv. Chim. Acta*, 1995, 78, 486-504; Altmann et al., *Chimia*, 1996, 50, 168-176; Altmann et al., *Biochem. Soc. Trans.*, 1996, 24, 630-637; and Altmann et al., *Nucleosides Nucleotides*, 1997, 16, 917-926).

As used herein, “2'-modified” or “2'-substituted” refers to a nucleoside comprising a sugar comprising a substituent at the 2' position other than H or OH. 2'-modified nucleosides, include, but are not limited to, bicyclic nucleosides wherein the bridge connecting two carbon atoms of the sugar ring connects the 2' carbon and another carbon of the sugar ring; and nucleosides with non-bridging 2' substituents, such as allyl, amino, azido, thio, O-allyl, O—C₁-C₁₀ alkyl, —OCF₃, O—(CH₂)₂—O—CH₃, 2'-O(CH₂)₂SCH₃, O—(CH₂)₂—O—N((R_m)(R_n)), or O—CH₂—C(=O)—N((R_m)(R_n)), where

each (R_m) and R_n is, independently, H or substituted or unsubstituted C_1 - C_{10} alkyl. 2'-modified nucleosides may further comprise other modifications, for example at other positions of the sugar and/or at the nucleobase.

As used herein, "2'-F" refers to a nucleoside comprising a sugar comprising a fluoro group at the 2' position of the sugar ring.

As used herein, "2'-OMe" or "2'-OCH₃", "2'-O-methyl" or "2'-methoxy" each refers to a nucleoside comprising a sugar comprising an —OCH₃ group at the 2' position of the sugar ring.

As used herein, "MOE" or "2'-MOE" or "2'-OCH₂CH₂OCH₃" or "2'-O-methoxyethyl" each refers to a nucleoside comprising a sugar comprising a —OCH₂CH₂OCH₃ group at the 2' position of the sugar ring.

Methods for the preparations of modified sugars are well known to those skilled in the art. Some representative U.S. patents that teach the preparation of such modified sugars include without limitation, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,670,633; 5,700,920; 5,792,847 and 6,600,032 and International Application PCT/US2005/019219, filed Jun. 2, 2005 and published as WO 2005/121371 on Dec. 22, 2005, and each of which is herein incorporated by reference in its entirety.

As used herein, "oligonucleotide" refers to a compound comprising a plurality of linked nucleosides. In certain embodiments, one or more of the plurality of nucleosides is modified. In certain embodiments, an oligonucleotide comprises one or more ribonucleosides (RNA) and/or deoxyribonucleosides (DNA).

In nucleotides having modified sugar moieties, the nucleobase moieties (natural, modified or a combination thereof) are maintained for hybridization with an appropriate nucleic acid target.

In certain embodiments, antisense compounds comprise one or more nucleosides having modified sugar moieties. In certain embodiments, the modified sugar moiety is 2'-MOE. In certain embodiments, the 2'-MOE modified nucleosides are arranged in a gapmer motif. In certain embodiments, the modified sugar moiety is a bicyclic nucleoside having a (4'-CH(CH₃)—O-2') bridging group. In certain embodiments, the (4'-CH(CH₃)—O-2') modified nucleosides are arranged throughout the wings of a gapmer motif.

Compositions and Methods for Formulating Pharmaceutical Compositions

Antisense oligonucleotides may be admixed with pharmaceutically acceptable active or inert substances for the preparation of pharmaceutical compositions or formulations. Compositions and methods for the formulation of pharmaceutical compositions are dependent upon a number of criteria, including, but not limited to, route of administration, extent of disease, or dose to be administered.

An antisense compound targeted to a C9ORF72 nucleic acid can be utilized in pharmaceutical compositions by combining the antisense compound with a suitable pharmaceutically acceptable diluent or carrier. A pharmaceutically acceptable diluent includes phosphate-buffered saline (PBS). PBS is a diluent suitable for use in compositions to be delivered parenterally. Accordingly, in one embodiment, employed in the methods described herein is a pharmaceutical composition comprising an antisense compound targeted to a C9ORF72 nucleic acid and a pharmaceutically acceptable diluent. In certain embodiments, the pharmaceu-

tically acceptable diluent is PBS. In certain embodiments, the antisense compound is an antisense oligonucleotide.

Pharmaceutical compositions comprising antisense compounds encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other oligonucleotide which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to pharmaceutically acceptable salts of antisense compounds, prodrugs, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents. Suitable pharmaceutically acceptable salts include, but are not limited to, sodium and potassium salts.

A prodrug can include the incorporation of additional nucleosides at one or both ends of an antisense compound which are cleaved by endogenous nucleases within the body, to form the active antisense compound.

Conjugated Antisense Compounds

Antisense compounds may be covalently linked to one or more moieties or conjugates which enhance the activity, cellular distribution or cellular uptake of the resulting antisense oligonucleotides. Typical conjugate groups include cholesterol moieties and lipid moieties. Additional conjugate groups include carbohydrates, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes.

Antisense compounds can also be modified to have one or more stabilizing groups that are generally attached to one or both termini of antisense compounds to enhance properties such as, for example, nuclease stability. Included in stabilizing groups are cap structures. These terminal modifications protect the antisense compound having terminal nucleic acid from exonuclease degradation, and can help in delivery and/or localization within a cell. The cap can be present at the 5'-terminus (5'-cap), or at the 3'-terminus (3'-cap), or can be present on both termini. Cap structures are well known in the art and include, for example, inverted deoxy abasic caps. Further 3' and 5'-stabilizing groups that can be used to cap one or both ends of an antisense compound to impart nuclease stability include those disclosed in WO 03/004602 published on Jan. 16, 2003.

Cell Culture and Antisense Compounds Treatment

The effects of antisense compounds on the level, activity or expression of C9ORF72 nucleic acids can be tested in vitro in a variety of cell types. Cell types used for such analyses are available from commercial vendors (e.g. American Type Culture Collection, Manassas, Va.; Zen-Bio, Inc., Research Triangle Park, N.C.; Clonetics Corporation, Walkersville, Md.) and are cultured according to the vendor's instructions using commercially available reagents (e.g. Invitrogen Life Technologies, Carlsbad, Calif.). Illustrative cell types include, but are not limited to, HepG2 cells, Hep3B cells, and primary hepatocytes.

In Vitro Testing of Antisense Oligonucleotides

Described herein are methods for treatment of cells with antisense oligonucleotides, which can be modified appropriately for treatment with other antisense compounds.

In general, cells are treated with antisense oligonucleotides when the cells reach approximately 60-80% confluency in culture.

One reagent commonly used to introduce antisense oligonucleotides into cultured cells includes the cationic lipid transfection reagent LIPOFECTIN (Invitrogen, Carlsbad, Calif.). Antisense oligonucleotides are mixed with LIPOFECTIN in OPTI-MEM 1 (Invitrogen, Carlsbad, Calif.) to achieve the desired final concentration of antisense oligo-

nucleotide and a LIPOFECTIN concentration that typically ranges 2 to 12 ug/mL per 100 nM antisense oligonucleotide.

Another reagent used to introduce antisense oligonucleotides into cultured cells includes LIPOFECTAMINE (Invitrogen, Carlsbad, Calif.). Antisense oligonucleotide is mixed with LIPOFECTAMINE in OPTI-MEM 1 reduced serum medium (Invitrogen, Carlsbad, Calif.) to achieve the desired concentration of antisense oligonucleotide and a LIPOFECTAMINE concentration that typically ranges 2 to 12 ug/mL per 100 nM antisense oligonucleotide.

Another technique used to introduce antisense oligonucleotides into cultured cells includes electroporation.

Cells are treated with antisense oligonucleotides by routine methods. Cells are typically harvested 16-24 hours after antisense oligonucleotide treatment, at which time RNA or protein levels of target nucleic acids are measured by methods known in the art and described herein. In general, when treatments are performed in multiple replicates, the data are presented as the average of the replicate treatments.

The concentration of antisense oligonucleotide used varies from cell line to cell line. Methods to determine the optimal antisense oligonucleotide concentration for a particular cell line are well known in the art. Antisense oligonucleotides are typically used at concentrations ranging from 1 nM to 300 nM when transfected with LIPOFECTAMINE. Antisense oligonucleotides are used at higher concentrations ranging from 625 to 20,000 nM when transfected using electroporation.

RNA Isolation

RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are well known in the art. RNA is prepared using methods well known in the art, for example, using the TRIZOL Reagent (Invitrogen, Carlsbad, Calif.) according to the manufacturer's recommended protocols.

Analysis of Inhibition of Target Levels or Expression

Inhibition of levels or expression of a C9ORF72 nucleic acid can be assayed in a variety of ways known in the art. For example, target nucleic acid levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or quantitative real-time PCR. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. Methods of RNA isolation are well known in the art. Northern blot analysis is also routine in the art. Quantitative real-time PCR can be conveniently accomplished using the commercially available ABI PRISM 7600, 7700, or 7900 Sequence Detection System, available from PE-Applied Biosystems, Foster City, Calif. and used according to manufacturer's instructions.

Quantitative Real-Time PCR Analysis of Target RNA Levels

Quantitation of target RNA levels may be accomplished by quantitative real-time PCR using the ABI PRISM 7600, 7700, or 7900 Sequence Detection System (PE-Applied Biosystems, Foster City, Calif.) according to manufacturer's instructions. Methods of quantitative real-time PCR are well known in the art.

Prior to real-time PCR, the isolated RNA is subjected to a reverse transcriptase (RT) reaction, which produces complementary DNA (cDNA) that is then used as the substrate for the real-time PCR amplification. The RT and real-time PCR reactions are performed sequentially in the same sample well. RT and real-time PCR reagents are obtained from Invitrogen (Carlsbad, Calif.). RT real-time-PCR reactions are carried out by methods well known to those skilled in the art.

Gene (or RNA) target quantities obtained by real time PCR are normalized using either the expression level of a

gene whose expression is constant, such as cyclophilin A, or by quantifying total RNA using RIBOGREEN (Invitrogen, Inc. Carlsbad, Calif.). Cyclophilin A expression is quantified by real time PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RIBOGREEN RNA quantification reagent (Invitrogen, Inc. Eugene, Oreg.). Methods of RNA quantification by RIBOGREEN are taught in Jones, L. J., et al, (Analytical Biochemistry, 1998, 265, 368-374). A CYTOFLUOR 4000 instrument (PE Applied Biosystems) is used to measure RIBOGREEN fluorescence.

Probes and primers are designed to hybridize to a C9ORF72 nucleic acid. Methods for designing real-time PCR probes and primers are well known in the art, and may include the use of software such as PRIMER EXPRESS Software (Applied Biosystems, Foster City, Calif.).

Analysis of Protein Levels

Antisense inhibition of C9ORF72 nucleic acids can be assessed by measuring C9ORF72 protein levels. Protein levels of C9ORF72 can be evaluated or quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), enzyme-linked immunosorbent assay (ELISA), quantitative protein assays, protein activity assays (for example, caspase activity assays), immunohistochemistry, immunocytochemistry or fluorescence-activated cell sorting (FACS). Antibodies directed to a target can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies (Aerie Corporation, Birmingham, Mich.), or can be prepared via conventional monoclonal or polyclonal antibody generation methods well known in the art. Antibodies useful for the detection of mouse, rat, monkey, and human C9ORF72 are commercially available.

In Vivo Testing of Antisense Compounds

Antisense compounds, for example, antisense oligonucleotides, are tested in animals to assess their ability to inhibit expression of C9ORF72 and produce phenotypic changes, such as, improved motor function and respiration. In certain embodiments, motor function is measured by rotarod, grip strength, pole climb, open field performance, balance beam, hindpaw footprint testing in the animal. In certain embodiments, respiration is measured by whole body plethysmograph, invasive resistance, and compliance measurements in the animal. Testing may be performed in normal animals, or in experimental disease models. For administration to animals, antisense oligonucleotides are formulated in a pharmaceutically acceptable diluent, such as phosphate-buffered saline (PBS) or artificial cerebrospinal fluid (aCSF). Administration includes parenteral routes of administration, such as intraperitoneal, intravenous, and subcutaneous, as well as central routes of administration such as intracerebroventricular or intrathecal. Calculation of antisense oligonucleotide dosage and dosing frequency is within the abilities of those skilled in the art, and depends upon factors such as route of administration and animal body weight. Following a period of treatment with antisense oligonucleotides, RNA is isolated from CNS tissue or CSF and changes in C9ORF72 nucleic acid expression are measured.

Targeting C9ORF72

Antisense oligonucleotides described herein may hybridize to a C9ORF72 nucleic acid in any stage of RNA processing. For example, described herein are antisense oligonucleotides that are complementary to a pre-mRNA or a mature mRNA. Additionally, antisense oligonucleotides described herein may hybridize to any element of a C9ORF72 nucleic acid. For example, described herein are antisense oligonucleotides that are complementary to an

61

exon, an intron, the 5' UTR, the 3' UTR, a repeat region, a hexanucleotide repeat expansion, a splice junction, an exon: exon splice junction, an exonic splicing silencer (ESS), an exonic splicing enhancer (ESE), exon 1a, exon 1b, exon 1c, exon 1d, exon 1e, exon 2, exon 3, exon 4, exon 5, exon 6, exon 7, exon 8, exon 9, exon 10, exon11, intron 1, intron 2, intron 3, intron 4, intron 5, intron 6, intron 7, intron 8, intron 9, or intron 10 of a C9ORF72 nucleic acid.

In certain embodiments, antisense oligonucleotides described herein hybridize to all variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein selectively hybridize to certain variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein selectively hybridize to variants of C9ORF72 containing a hexanucleotide repeat expansion. In certain embodiments, the antisense oligonucleotides described herein selectively hybridize to pre-mRNA variants containing the hexanucleotide repeat. In certain embodiments, pre-mRNA variants of C9ORF72 containing a hexanucleotide repeat expansion include SEQ ID NO: 1-3 and 6-10. In certain embodiments, such hexanucleotide repeat expansion comprises at least 24 repeats of any of GGGGCC, GGGGGG, GGGGGC, or GGGGCG.

In certain embodiments, the antisense oligonucleotides described herein inhibit expression of all variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein inhibit expression of all variants of C9ORF72 equally. In certain embodiments, the antisense oligonucleotides described herein preferentially inhibit expression of one or more variants of C9ORF72. In certain embodiments, the antisense oligonucleotides described herein preferentially inhibit expression of variants of C9ORF72 containing a hexanucleotide repeat expansion. In certain embodiments, the antisense oligonucleotides described herein selectively inhibit expression of pre-mRNA variants containing the hexanucleotide repeat. In certain embodiments, the antisense oligonucleotides described herein selectively inhibit expression of C9ORF72 pathogenic associated mRNA variants. In certain embodiments, pre-mRNA variants of C9ORF72 containing a hexanucleotide repeat expansion include SEQ ID NO: 1-3 and 6-10. In certain embodiments, such hexanucleotide repeat expansion comprises at least 24 repeats of any of GGGGCC, GGGGGG, GGGGGC, or GGGGCG. In certain embodiments, the hexanucleotide repeat expansion forms nuclear foci. In certain embodiments, antisense oligonucleotides described herein are useful for reducing nuclear foci. Nuclear foci may be reduced in terms of percent of cells with foci as well as number of foci per cell.

Selective Inhibition of Certain Pathogenic Associated Variants

In certain examples herein, primer probe set RTS3905 detects an mRNA variant (e.g. NM_001256054.1) processed from a pre-mRNA variant containing the hexanucleotide repeat. The mRNA variant processed from a pre-mRNA variant containing the hexanucleotide repeat (i.e., the "C9ORF72 pathogenic associated mRNA variant"). A pre-mRNA contains the hexanucleotide repeat when transcription of the pre-mRNA begins in the region from the start site of exon 1A to the start site of exon 1B, e.g., nucleotides 1107 to 1520 of the genomic sequence (SEQ ID NO: 2, the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 27565000). Oligonucleotides were designed in this region to selectively target the pre-mRNA variant containing the hexanucleotide repeat. RTS3905 measures an mRNA product (i.e. the C9ORF72 pathogenic associated mRNA variant) of the pre-mRNA

62

variant containing the hexanucleotide repeat and, therefore, measures the reduction of the pre-mRNA variant containing the hexanucleotide repeat.

C9ORF72 Features

Antisense oligonucleotides described herein may hybridize to any C9ORF72 variant at any state of processing within any element of the C9ORF72 gene. For example, antisense oligonucleotides described herein may hybridize to an exon, an intron, the 5' UTR, the 3' UTR, a repeat region, a hexanucleotide repeat expansion, a splice junction, an exon: exon splice junction, an exonic splicing silencer (ESS), an exonic splicing enhancer (ESE), exon 1a, exon 1b, exon 1c, exon 1d, exon 1e, exon 2, exon 3, exon 4, exon 5, exon 6, exon 7, exon 8, exon 9, exon 10, exon 11, intron 1, intron 2, intron 3, intron 4, intron 5, intron 6, intron 7, intron 8, intron 9, or intron 10. For example, antisense oligonucleotides may target any of the exons characterized below in Tables 1-5 for the various C9ORF72 variants described below. Antisense oligonucleotides described herein may also target variants not characterized below and such variants are characterized in GENBANK. Moreover, antisense oligonucleotides described herein may also target elements other than exons and such elements are characterized in GENBANK.

TABLE 1

Functional Segments for NM_001256054.1 (SEQ ID NO: 1)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1C	1	158	1137	1294
exon 2	159	646	7839	8326
exon 3	647	706	9413	9472
exon 4	707	802	12527	12622
exon 5	803	867	13354	13418
exon 6	868	940	14704	14776
exon 7	941	1057	16396	16512
exon 8	1058	1293	18207	18442
exon 9	1294	1351	24296	24353
exon 10	1352	1461	26337	26446
exon 11	1462	3339	26581	28458

TABLE 2

Functional Segments for NM_018325.3 (SEQ ID NO: 4)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1B	1	63	1510	1572
exon 2	64	551	7839	8326
exon 3	552	611	9413	9472
exon 4	612	707	12527	12622
exon 5	708	772	13354	13418
exon 6	773	845	14704	14776
exon 7	846	962	16396	16512
exon 8	963	1198	18207	18442
exon 9	1199	1256	24296	24353
exon 10	1257	1366	26337	26446
exon 11	1367	3244	26581	28458

TABLE 3

Functional Segments for NM_145005.5 (SEQ ID NO: 6)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1A	1	80	1137	1216
exon 2	81	568	7839	8326
exon 3	569	628	9413	9472
exon 4	629	724	12527	12622
exon 5B (exon 5 into intron 5)	725	1871	13354	14500

TABLE 4

Functional Segments for DB079375.1 (SEQ ID NO: 7)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1E	1	35	1135	1169
exon 2	36	524	7839	8326
exon 3 (EST ends before end of full exon)	525	562	9413	9450

TABLE 5

Functional Segments for BU194591.1 (SEQ ID NO: 8)				
Exon Number	mRNA start site	mRNA stop site	Start site in reference to SEQ ID NO: 2	Stop site in reference to SEQ ID NO: 2
exon 1D	1	36	1241	1279
exon 2	37	524	7839	8326
exon 3	525	584	9413	9472
exon 4	585	680	12527	12622
exon 5B (exon 5 into intron 5)	681	798	13354	13465

Certain Indications

In certain embodiments, provided herein are methods of treating an individual comprising administering one or more pharmaceutical compositions described herein. In certain embodiments, the individual has a neurodegenerative disease. In certain embodiments, the individual is at risk for developing a neurodegenerative disease, including, but not limited to, ALS or FTD. In certain embodiments, the individual has been identified as having a C9ORF72 associated disease. In certain embodiments, provided herein are methods for prophylactically reducing C9ORF72 expression in an individual. Certain embodiments include treating an individual in need thereof by administering to an individual a therapeutically effective amount of an antisense compound targeted to a C9ORF72 nucleic acid.

In one embodiment, administration of a therapeutically effective amount of an antisense compound targeted to a C9ORF72 nucleic acid is accompanied by monitoring of C9ORF72 levels in an individual, to determine an individual's response to administration of the antisense compound. An individual's response to administration of the antisense compound may be used by a physician to determine the amount and duration of therapeutic intervention.

In certain embodiments, administration of an antisense compound targeted to a C9ORF72 nucleic acid results in reduction of C9ORF72 expression by at least 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 99%, or a range defined by any two of these values. In certain embodiments, administration of an antisense compound targeted to a C9ORF72 nucleic acid results in improved motor function and respiration in an animal. In certain embodiments, administration of a C9ORF72 antisense compound improves motor function and respiration by at least 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 99%, or a range defined by any two of these values.

In certain embodiments, pharmaceutical compositions comprising an antisense compound targeted to C9ORF72 are used for the preparation of a medicament for treating a patient suffering or susceptible to a neurodegenerative disease including ALS and FTD.

Certain Combination Therapies

In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with one or more other pharmaceutical agents. In certain embodiments, such one or more other pharmaceutical agents are designed to treat the same disease, disorder, or condition as the one or more pharmaceutical compositions described herein. In certain embodiments, such one or more other pharmaceutical agents are designed to treat a different disease, disorder, or condition as the one or more pharmaceutical compositions described herein. In certain embodiments, such one or more other pharmaceutical agents are designed to treat an undesired side effect of one or more pharmaceutical compositions described herein. In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with another pharmaceutical agent to treat an undesired effect of that other pharmaceutical agent. In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with another pharmaceutical agent to produce a combinational effect. In certain embodiments, one or more pharmaceutical compositions described herein are co-administered with another pharmaceutical agent to produce a synergistic effect.

In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are administered at the same time. In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are administered at different times. In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are prepared together in a single formulation. In certain embodiments, one or more pharmaceutical compositions described herein and one or more other pharmaceutical agents are prepared separately.

In certain embodiments, pharmaceutical agents that may be co-administered with a pharmaceutical composition described herein include Riluzole (Rilutek), Lioresal (Lioresal), and Dexamipexole.

In certain embodiments, pharmaceutical agents that may be co-administered with a C9ORF72 specific inhibitor described herein include, but are not limited to, an additional C9ORF72 inhibitor.

In certain embodiments, pharmaceutical agents that may be co-administered with a C9ORF72 specific inhibitor described herein include, but are not limited to, a C9ORF72 antisense transcript specific inhibitor. In certain embodiments, the C9ORF72 antisense transcript specific inhibitor is an antisense compound. In certain embodiments, the anti-

sense compound is a modified oligonucleotide. In certain embodiments, the modified oligonucleotide is single-stranded.

In certain embodiments, the co-administered pharmaceutical agent is administered prior to administration of a pharmaceutical composition described herein. In certain embodiments, the co-administered pharmaceutical agent is administered following administration of a pharmaceutical composition described herein. In certain embodiments the co-administered pharmaceutical agent is administered at the same time as a pharmaceutical composition described herein. In certain embodiments the dose of a co-administered pharmaceutical agent is the same as the dose that would be administered if the co-administered pharmaceutical agent was administered alone. In certain embodiments the dose of a co-administered pharmaceutical agent is lower than the dose that would be administered if the co-administered pharmaceutical agent was administered alone. In certain embodiments the dose of a co-administered pharmaceutical agent is greater than the dose that would be administered if the co-administered pharmaceutical agent was administered alone.

In certain embodiments, the co-administration of a second compound enhances the effect of a first compound, such that co-administration of the compounds results in an effect that is greater than the effect of administering the first compound alone. In other embodiments, the co-administration results in effects that are additive of the effects of the compounds when administered alone. In certain embodiments, the co-administration results in effects that are supra-additive of the effects of the compounds when administered alone. In certain embodiments, the first compound is an antisense compound. In certain embodiments, the second compound is an antisense compound.

Certain Comparator Compositions

In certain embodiments, compounds described herein are more tolerable than ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, and/or ISIS 577083. ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, and ISIS 577083 were selected as comparator compounds because they exhibited high levels of dose-dependent reduction of C9ORF72 mRNA in various studies described in WO2014/062691. Thus, ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, and ISIS 577083 were deemed highly efficacious and potent compounds. Additionally, ISIS 577065, ISIS 577056, and ISIS 576816 described in WO2014/062691 are structurally similar as compounds described herein. For example, ISIS 577065 has a 16 nucleobase overlap with ISIS 801287; ISIS 577056 has a 16 nucleobase overlap with ISIS 806679; ISIS 576816 has an 18 nucleobase overlap with ISIS 802473 (18-mer); and ISIS 576816 has an 18 nucleobase overlap with ISIS 802459.

In certain embodiments, ISIS 576816, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') GCCT-TACTCTAGGACCAAGA (incorporated herein as SEQ ID NO: 21), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 576816 achieved an average FOB score of 7.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 576816.

In certain embodiments, ISIS 576974, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') GGGACAC-TACAAGGTAGTAT (incorporated herein as SEQ ID NO:

56), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 576974 achieved an average FOB score of 5.67 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 576974.

In certain embodiments, ISIS 577061, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') TACAGGCT-GCGGTTGTTTCC (incorporated herein as SEQ ID NO: 57), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577061 achieved an average FOB score of 7.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577061.

In certain embodiments, ISIS 577065, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') CCCGGC-CCCTAGCGCGCGAC (incorporated herein as SEQ ID NO: 58), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577065 achieved an average FOB score of 6.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577065.

In certain embodiments, ISIS 577083, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') GGTAAC-TCAAACCTCTTGGG (incorporated herein as SEQ ID NO: 59), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577083 achieved an average FOB score of 7.00 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577083.

In certain embodiments, ISIS 577056, a 5-10-5 MOE gapmer, having a sequence of (from 5' to 3') AATCTT-TATCAGGTCTTTTC (incorporated herein as SEQ ID NO: 60), wherein each internucleoside linkage is a phosphorothioate linkage, each cytosine is a 5-methylcytosine, and each of nucleosides 1-5 and 16-20 comprise a 2'-O-methoxyethyl group, which was previously described in WO2014/062691, incorporated herein by reference, is a comparator compound. ISIS 577056 achieved an average

FOB score of 6.5 in a study of acute tolerability in mice (see Example 3 hereinbelow). Certain compounds described herein achieved a lower FOB score in a similar study of acute tolerability in mice (see Example 2 hereinbelow), including ISIS 801287, ISIS 806679, ISIS 802473, and ISIS 802459. Therefore, certain compounds described herein are more tolerable than comparator compound ISIS 577056.

Certain Human Therapeutics

The human C9ORF72 antisense oligonucleotides described herein are human therapeutics. Various parameters of potency, efficacy, and/or tolerability are being examined. Such parameters include in vitro inhibition of total C9ORF72 RNA expression, in vitro inhibition of C9ORF72 pathogenic associated RNA variant expression, in vitro dose response (IC50), in vivo inhibition of total or pathogenic RNA and/or protein in a transgenic animal containing a human C9ORF72 transgene in relevant tissues (e.g., brain and/or spinal cord), tolerability in mouse, tolerability in rat, and/or tolerability in a primate. Tolerability markers that may be measured include blood and serum chemistry parameters, CSF chemistry parameters, body and organ weights, general observations and/or behavioral tests, and/or biochemical markers such as GFAP and/or AIF1. Acute or long term tolerability may be measured.

Certain Compositions

1. ISIS 801287

In certain embodiments, ISIS 801287 is characterized as a 4-8-6 MOE gapmer, having a sequence of (from 5' to 3') GCCCCTAGCGCGCGACTC (incorporated herein as SEQ ID NO: 33), wherein each of nucleosides 1-4 and 13-18 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 5-12 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 13 to 14, 14 to 15, and 15 to 16 are phosphodiester linkages and the internucleoside linkages between nucleosides 1 to 2, 4 to 5, 5 to 6, 6 to 7, 7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to 12, 12 to 13, 16 to 17, and 17 to 18 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.


In certain embodiments, ISIS 801287 is described by the following chemical notation: Ges mCeo mCeo mCes mCds Tds Ads Gds mCds Gds mCds Gds mCeo Geo AeomCes Tes mCe;

wherein,


- A=an adenine,
- mC=a 5'-methylcytosine
- G=a guanine,
- T=a thymine,
- e=a 2'-O-methoxyethylribose modified sugar
- d=a 2'-deoxyribose sugar,
- s=a phosphorothioate internucleoside linkage, and
- o=a phosphodiester internucleoside linkage.

In certain embodiments, ISIS 801287 is described by the following chemical structure, or a salt thereof:

(SEQ ID NO: 33)




69



70


-continued




Structure 1. ISIS 801287

In certain embodiments, the sodium salt of ISIS 801287 is described by the following chemical structure:


(SEQ ID NO: 33)



71



72



Structure 2. The Sodium Salt of ISIS 801287
2. ISIS 806679


In certain embodiments, ISIS 806679 is characterized as a 6-10-4 MOE gapmer, having a sequence of (from 5' to 3') GGTAAATCTTTATCAGGTCT (incorporated herein as SEQ ID NO: 49), wherein each of nucleosides 1-6 and 17-20 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 7-16 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 4 to 5, 5 to 6, 6 to 7, and 17 to 18 are phosphodiester linkages and the internucleoside linkages between nucleosides 1 to 2, 7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to 12, 12 to 13, 13 to 14, 14 to 15, 15 to 16, 16 to 17, 18 to 19, and 19 to 20 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.


In certain embodiments, ISIS 806679 is described by the following chemical notation: Ges Geo Teo Teo AeO AeO Tds mCds Tds Tds Tds Ads Tds mCds Ads Gds Geo Tes mCes Te; wherein,

A=an adenine,
mC=a 5'-methylcytosine
G=a guanine,
T=a thymine,
e=a 2'-O-methoxyethylribose modified sugar
d=a 2'-deoxyribose sugar,
s=a phosphorothioate internucleoside linkage, and
o=a phosphodiester internucleoside linkage.

In certain embodiments, ISIS 806679 is described by the following chemical structure, or a salt thereof:


(SEQ ID NO: 49)






In certain embodiments, the sodium salt of ISIS 806679 is described by the following chemical structure:


(SEQ ID NO: 49)




75



-continued



76



Structure 4. The Sodium Salt of ISIS 806679
3. ISIS 802473

In certain embodiments, ISIS 802473 is characterized as a 4-8-6 MOE gapmer, having a sequence of (from 5' to 3') GCCTTACTCTAGGACCAA (incorporated herein as SEQ ID NO: 47), wherein each of nucleosides 1-4 and 13-18 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 5-12 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 13 to 14, 14 to 15, and 15 to 16 are phosphodiester linkages and the internucleoside linkages between nucleosides are 1 to 2, 4 to 5, 5 to 6, 6 to 7, 7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to 12, 12 to 13, 16 to 17, and 17 to 18 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.

In certain embodiments, ISIS 802473 is described by the following chemical notation: Ges mCeo mCeo Tes Tds Ads mCds Tds mCds Tds Ads Gds Geo Aeo mCeo mCes Aes Ae; wherein,

A=an adenine,

mC=a 5'-methylcytosine

G=a guanine,

T=a thymine,

e=a 2'-O-methoxyethylribose modified sugar

d=a 2'-deoxyribose sugar,

s=a phosphorothioate internucleoside linkage, and


o=a phosphodiester internucleoside linkage.

In certain embodiments, ISIS 802473 is described by the following chemical structure, or a salt thereof:

77

78


(SEQ ID NO: 47)



79


Structure 5. ISIS 802473

In certain embodiments, the sodium salt of ISIS 802473 is described by the following chemical structure:



80

(SEQ ID NO: 47)



81

Structure 6. The Sodium Salt of ISIS 802473
4. ISIS 802459

In certain embodiments, ISIS 802459 is characterized as a 3-10-7 MOE gapmer, having a sequence of (from 5' to 3') GCCTTACTCTAGGACCAAGA (incorporated herein as SEQ ID NO: 21), wherein each of nucleosides 1-3 and 14-20 are 2'-O-methoxyethylribose modified nucleosides, and each of nucleosides 4-13 are 2'-deoxynucleosides, wherein the internucleoside linkages between nucleosides 2 to 3, 3 to 4, 14 to 15, 15 to 16, 16 to 17, and 17 to 18 are phosphodiester linkages and the internucleoside linkages between nucleosides are 1 to 2, 4 to 5, 5 to 6, 6 to 7, 7 to 8, 8 to 9, 9 to 10, 10 to 11, 11 to 12, 12 to 13, 13 to 14, 18 to 19, and 19 to 20 are phosphorothioate linkages, and wherein each cytosine is a 5'-methylcytosine.

82

In certain embodiments, ISIS 802459 is described by the following chemical notation: Ges mCeo mCeo Tds Tds Ads mCds Tds mCds Tds Ads Gds Gds AeO mCeo mCeo AeO Aes Ges Ae; wherein,

A=an adenine,

mC=a 5'-methylcytosine

G=a guanine,

T=a thymine,

e=a 2'-O-methoxyethylribose modified sugar


d=a 2'-deoxyribose sugar,

s=a phosphorothioate internucleoside linkage, and


o=a phosphodiester internucleoside linkage.

In certain embodiments, ISIS 802459 is described by the following chemical structure, or a salt thereof:


(SEQ ID NO: 21)




83



-continued




84




Structure 7. ISIS 802459

In certain embodiments, the sodium salt of ISIS 802459 is described by the following chemical structure:

(SEQ ID NO: 21)





Structure 8. The Sodium Salt of ISIS 802459

EXAMPLES

Non-Limiting Disclosure and Incorporation by Reference

While certain compounds, compositions, and methods described herein have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds described herein and are not intended to limit the same. Each of the references recited in the present application is incorporated herein by reference in its entirety.

Example 1

Antisense Oligonucleotides Targeting Human C9ORF72

The antisense oligonucleotides in the table below were designed as MOE gapmers. The central gap segment of each

gapmer contains 2'-deoxynucleosides and is flanked by wing segments on both the 5' end and on the 3' end containing nucleosides that each comprise a 2'-MOE group. The specific motif of each gapmer is listed in table below, represented by three numbers separated by hyphens. The numbers represent the number of nucleosides in the 5'-wing, the gap, and the 3'-wing, respectively. All cytosine residues throughout each oligonucleotide are 5-methylcytosines. The internucleoside linkages for the gapmers are mixed phosphorothioate and phosphodiester linkages. The internucleoside linkages for each gapmer are presented in the Linkage column, where 'o' indicates a phosphodiester linkage and 's' indicates a phosphorothioate linkage.

Each antisense oligonucleotide listed in the table below is targeted to the human C9ORF72 genomic sequence, designated herein as SEQ ID NO: 2 (the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 27565000). "Start site" indicates the 5'-most nucleoside to which the gapmer is targeted in the human genomic sequence. "Stop site" indicates the 3'-most nucleoside to which the gapmer is targeted human genomic sequence.

TABLE 6

Antisense oligonucleotides targeting human C9ORF72							SEQ ID NO
Isis No.	Start Site	Stop Site	Sequence	Linkage	Motif		
791656	1445	1464	CCGGCCCCTAGCGCGCGACT	soossssssssssooooss	3-10-7		22
791657	1445	1464	CCGGCCCCTAGCGCGCGACT	sooossssssssssooooss	4-10-6		22
791658	1445	1464	CCGGCCCCTAGCGCGCGACT	soooooossssssssooss	6-10-4		22
791659	1445	1464	CCGGCCCCTAGCGCGCGACT	soooooossssssssssss	7-10-3		22
791660	1445	1463	CGGCCCTAGCGCGCGACT	soooossssssssooss	5-9-5		23
791661	1446	1464	CCGGCCCCTAGCGCGCGAC	soooossssssssooss	5-9-5		24
791662	1445	1463	CGGCCCTAGCGCGCGACT	sooossssssssooooss	4-9-6		23

Antisense oligonucleotides targeting human C9ORF72						
Isis No.	Start Site	Stop Site	Sequence	Linkage	Motif	SEQ ID NO
791663	1446	1464	CCGGCCCCCTAGCGCGCGAC	sooooooooooooooooooss	4-9-6	24
791664	1445	1463	CGGCCCCCTAGCGCGCGACT	sooooooooooooooooooss	6-9-4	23
791665	1446	1464	CCGGCCCCCTAGCGCGCGAC	sooooooooooooooooooss	6-9-4	24
801274	1440	1459	CCCTAGCGCGCGACTCCTGA	sooooooooooooooooooss	4-10-6	25
801275	1441	1460	CCCCTAGCGCGCGACTCCTG	sooooooooooooooooooss	4-10-6	26
801276	1442	1461	GCCCCCTAGCGCGCGACTCCT	sooooooooooooooooooss	4-10-6	27
801277	1443	1462	GGCCCCCTAGCGCGCGACTCC	sooooooooooooooooooss	4-10-6	28
801278	1444	1463	CGGCCCCCTAGCGCGCGACTC	sooooooooooooooooooss	4-10-6	29
801279	1440	1459	CCCTAGCGCGCGACTCCTGA	sooooooooooooooooooss	6-10-4	25
801280	1441	1460	CCCCTAGCGCGCGACTCCTG	sooooooooooooooooooss	6-10-4	26
801281	1442	1461	GCCCCCTAGCGCGCGACTCCT	sooooooooooooooooooss	6-10-4	27
801282	1443	1462	GGCCCCCTAGCGCGCGACTCC	sooooooooooooooooooss	6-10-4	28
801283	1444	1463	CGGCCCCCTAGCGCGCGACTC	sooooooooooooooooooss	6-10-4	29
801284	1441	1458	CCTAGCGCGCGACTCCTG	sooooooooooooooooooss	4-8-6	30
801285	1442	1459	CCCTAGCGCGCGACTCCT	sooooooooooooooooooss	4-8-6	31
801286	1443	1460	CCCCTAGCGCGCGACTCC	sooooooooooooooooooss	4-8-6	32
801287	1444	1461	GCCCCCTAGCGCGCGACTC	sooooooooooooooooooss	4-8-6	33
801288	1445	1462	GGCCCCCTAGCGCGCGACT	sooooooooooooooooooss	4-8-6	34
801289	1446	1463	CGGCCCCCTAGCGCGCGAC	sooooooooooooooooooss	4-8-6	35
801290	1441	1458	CCTAGCGCGCGACTCCTG	sooooooooooooooooooss	6-8-4	30
801291	1442	1459	CCCTAGCGCGCGACTCCT	sooooooooooooooooooss	6-8-4	31
801292	1443	1460	CCCCTAGCGCGCGACTCC	sooooooooooooooooooss	6-8-4	32
801293	1444	1461	GCCCCCTAGCGCGCGACTC	sooooooooooooooooooss	6-8-4	33
801294	1445	1462	GGCCCCCTAGCGCGCGACT	sooooooooooooooooooss	6-8-4	34
801295	1446	1463	CGGCCCCCTAGCGCGCGAC	sooooooooooooooooooss	6-8-4	35
801296	1403	1422	AGGCTGCGGTTGTTTCCCTC	sooooooooooooooooooss	4-10-6	36
801297	1404	1423	CAGGCTGCGGTTGTTTCCCT	sooooooooooooooooooss	4-10-6	37
801298	1403	1421	GGCTGCGGTTGTTTCCCTC	sooooooooooooooooooss	5-9-5	38
801299	1404	1422	AGGCTGCGGTTGTTTCCCT	sooooooooooooooooooss	5-9-5	39
801300	1405	1423	CAGGCTGCGGTTGTTTCCC	sooooooooooooooooooss	5-9-5	40
801301	1403	1421	GGCTGCGGTTGTTTCCCTC	sooooooooooooooooooss	4-9-6	38
801302	1404	1422	AGGCTGCGGTTGTTTCCCT	sooooooooooooooooooss	4-9-6	39
801303	1405	1423	CAGGCTGCGGTTGTTTCCC	sooooooooooooooooooss	4-9-6	40
801304	1403	1421	GGCTGCGGTTGTTTCCCTC	sooooooooooooooooooss	6-9-4	38
801305	1404	1422	AGGCTGCGGTTGTTTCCCT	sooooooooooooooooooss	6-9-4	39
801306	1405	1423	CAGGCTGCGGTTGTTTCCC	sooooooooooooooooooss	6-9-4	40
801307	1403	1420	GCTGCGGTTGTTTCCCTC	sooooooooooooooooooss	4-8-6	41

TABLE 6-continued

Antisense oligonucleotides targeting human C9ORF72						
Isis No.	Start Site	Stop Site	Sequence	Linkage	Motif	SEQ ID NO
801308	1404	1421	GGCTGCGGTTGTTTCCCT	sooooooooooooo	4-8-6	42
801309	1405	1422	AGGCTGCGGTTGTTTCCC	sooooooooooooo	4-8-6	43
801310	1406	1423	CAGGCTGCGGTTGTTTCC	sooooooooooooo	4-8-6	44
801311	1403	1420	GCTGCGGTTGTTTCCCTC	sooooooooooooo	6-8-4	41
801312	1404	1421	GGCTGCGGTTGTTTCCCT	sooooooooooooo	6-8-4	42
801313	1405	1422	AGGCTGCGGTTGTTTCCC	sooooooooooooo	6-8-4	43
801314	1406	1423	CAGGCTGCGGTTGTTTCC	sooooooooooooo	6-8-4	44
801315	1403	1422	AGGCTGCGGTTGTTTCCCTC	sooooooooooooo	6-10-4	36
801316	1404	1423	CAGGCTGCGGTTGTTTCCCT	sooooooooooooo	6-10-4	37
802459	7990	8009	GCCTTACTCTAGGACCAAGA	sooooooooooooo	3-10-7	21
802460	8012	8031	TCTGTCTTTGGAGCCCAAAT	sooooooooooooo	3-10-7	45
802461	8186	8205	CTGCGATCCCCATTCCAGTT	sooooooooooooo	3-10-7	46
802462	7990	8009	GCCTTACTCTAGGACCAAGA	sooooooooooooo	4-10-6	21
802463	8012	8031	TCTGTCTTTGGAGCCCAAAT	sooooooooooooo	4-10-6	45
802464	8186	8205	CTGCGATCCCCATTCCAGTT	sooooooooooooo	4-10-6	46
802465	7990	8009	GCCTTACTCTAGGACCAAGA	sooooooooooooo	6-10-4	21
802466	8012	8031	TCTGTCTTTGGAGCCCAAAT	sooooooooooooo	6-10-4	45
802467	8186	8205	CTGCGATCCCCATTCCAGTT	sooooooooooooo	6-10-4	46
802468	7990	8009	GCCTTACTCTAGGACCAAGA	sooooooooooooo	7-10-3	21
802469	8012	8031	TCTGTCTTTGGAGCCCAAAT	sooooooooooooo	7-10-3	45
802470	8186	8205	CTGCGATCCCCATTCCAGTT	sooooooooooooo	7-10-3	46
802471	7992	8009	GCCTTACTCTAGGACCAA	sooooooooooooo	3-8-7	47
802472	8014	8031	TCTGTCTTTGGAGCCCAA	sooooooooooooo	3-8-7	48
802473	7992	8009	GCCTTACTCTAGGACCAA	sooooooooooooo	4-8-6	47
802474	8014	8031	TCTGTCTTTGGAGCCCAA	sooooooooooooo	4-8-6	48
802475	7992	8009	GCCTTACTCTAGGACCAA	sooooooooooooo	6-8-4	47
802476	8014	8031	TCTGTCTTTGGAGCCCAA	sooooooooooooo	6-8-4	48
802477	7992	8009	GCCTTACTCTAGGACCAA	sooooooooooooo	7-8-3	47
802478	8014	8031	TCTGTCTTTGGAGCCCAA	sooooooooooooo	7-8-3	48
806673	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooooooooo	3-10-7	49
806674	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooooooooo	3-10-7	50
806675	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooooooooo	3-10-7	51
806676	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooooooooo	4-10-6	49
806677	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooooooooo	4-10-6	50
806678	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooooooooo	4-10-6	51
806679	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooooooooo	6-10-4	49
806680	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooooooooo	6-10-4	50

TABLE 6-continued

Antisense oligonucleotides targeting human C9ORF72						
Isis No.	Start Site	Stop Site	Sequence	Linkage	Motif	SEQ ID NO
806681	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooossssssssssooss	6-10-4	51
806682	1370	1389	GGTTAATCTTTATCAGGTCT	sooooooosssssssssssss	7-10-3	49
806683	1371	1390	TGGTTAATCTTTATCAGGTC	sooooooosssssssssssss	7-10-3	50
806684	1372	1391	CTGGTTAATCTTTATCAGGT	sooooooosssssssssssss	7-10-3	51
806685	1371	1388	GTTAATCTTTATCAGGTC	sooooooooooooooss	4-8-6	52
806686	1372	1389	GGTTAATCTTTATCAGGT	sooooooooooooooss	4-8-6	53
806687	1373	1390	TGGTTAATCTTTATCAGG	sooooooooooooooss	4-8-6	54
806688	1440	1457	CTAGCGCGCGACTCCTGA	sooooooooooooooss	4-8-6	55
806689	1371	1388	GTTAATCTTTATCAGGTC	sooooooossssssssssooss	6-8-4	52
806690	1372	1389	GGTTAATCTTTATCAGGT	sooooooossssssssssooss	6-8-4	53
806691	1373	1390	TGGTTAATCTTTATCAGG	sooooooossssssssssooss	6-8-4	54
806692	1440	1457	CTAGCGCGCGACTCCTGA	sooooooossssssssssooss	6-8-4	55

Example 2

Tolerability of Antisense Oligonucleotides Targeting Human C9ORF72 in Mice

Antisense oligonucleotides described above were tested in mice to assess tolerability of the oligonucleotides. Wild type C57/B16 mice each received a single ICV dose of 700 µg of an antisense oligonucleotide listed in the table below. Each treatment group consisted of 4 mice. At 3 hours post injection, each mouse was evaluated according to 7 different criteria. The 7 criteria are (1) the mouse was bright, alert, and responsive; (2) the mouse was standing or hunched without stimuli; (3) the mouse showed any movement without stimuli; (4) the mouse demonstrated forward movement after it was lifted; (5) the mouse demonstrated any movement after it was lifted; (6) the mouse responded to a tail pinch; (7) regular breathing. For each of the 7 different criteria, each mouse was given a sub-score of 0 if it met the criteria or 1 if it did not. After all of the 7 criteria were evaluated, the sub-scores were summed for each mouse and then averaged for each group. For example, if a mouse was bright, alert, and responsive 3 hours after the 700 ng ICV dose, and met all other criteria, it would get a summed score of 0. If another mouse was not bright, alert, and responsive 3 hours after the 700 µg ICV dose but met all other criteria, it would receive a score of 1. The results are presented as the average score for each treatment group.

TABLE 7

Acute tolerability scores	
Isis No.	Score
791656	3.25
791657	4.25
791658	3.50
791659	2.00

TABLE 7-continued

Acute tolerability scores	
Isis No.	Score
791660	4.00
791661	4.50
791662	5.25
791663	6.00
791664	3.00
791665	5.75
801274	1.75
801275	3.75
801276	1.25
801277	1.50
801278	1.75
801279	1.25
801280	6.00
801281	0.00
801282	1.25
801283	1.25
801284	3.00
801285	1.00
801286	1.00
801287	1.50
801288	2.50
801289	7.00
801290	6.00
801291	3.00
801292	2.00
801293	1.00
801294	3.25
801295	4.50
801296	6.25
801297	4.75
801298	5.50
801299	6.25
801300	5.00
801301	6.00
801302	6.50
801303	4.00
801304	5.25
801305	6.00
801306	6.00
801307	5.00

TABLE 7-continued

Acute tolerability scores	
Isis No.	Score
801308	6.00
801309	7.00
801310	3.50
801311	5.50
801312	2.50
801313	5.25
801314	4.50
801315	4.00
801316	2.00
802459	2.00
802460	6.75
802461	1.75
802462	5.75
802463	6.75
802464	1.75
802465	2.25
802466	4.25
802467	0.25
802468	3.25
802469	2.00
802470	0.25
802471	1.25
802472	4.00
802473	0.25
802474	5.25
802475	1.00
802476	5.50
802477	2.50
802478	6.25
806673	0.00
806674	0.25
806675	0.00
806676	0.00
806677	1.00
806678	0.00
806679	1.50
806680	1.00
806681	0.00
806682	5.75
806683	3.75
806684	2.25
806685	1.00
806686	1.00
806687	3.25
806688	3.25
806689	3.00
806690	1.25
806691	6.25

Example 3

Tolerability of Oligonucleotides from WO 2014/062691

Oligonucleotides described in WO 2014/062691 were tested in an acute tolerability study in mice. Groups of 3 wild type C57/B16 mice were treated and analyzed as described in Example 2. The tested oligonucleotides include those listed in the table below, which are 5-10-5 MOE gapmers with a full phosphorothioate backbone and each cytosine is a 5-methylcytosine. The start and stop sites on SEQ ID NO: 2 that each oligonucleotide is targeted to are shown. The results are presented as the average score for each treatment group in the table below. These results demonstrate that ISIS 576816, ISIS 576974, ISIS 577061, ISIS 577065, ISIS 577083, and ISIS 577056 were poorly tolerated.

TABLE 8

Acute tolerability scores 3h after treatment with antisense oligonucleotides from WO 2014/062691					
Isis No.	Start site	Stop site	Sequence	Score	SEQ ID NO
576816	7990	8009	GCCTTACTCTAGGACCAAGA	7.00	21
576974	28251	28270	GGGACACTACAAGGTAGTAT	5.67	56
577061	1406	1425	TACAGGCTGCGGTTGTTTCC	7.00	57
577065	1446	1465	CCCGGCCCTAGCGCGCGAC	6.00	58
577083	3452	3471	GGTAACTTCAAACCTTGGG	7.00	59

TABLE 9

Acute tolerability scores 3h after treatment with antisense oligonucleotides from WO 2014/062691					
Isis No.	Start site	Stop site	Sequence	Score	SEQ ID NO
577056	1366	1385	AATCTTTATCAGGTCTTTTC	6.5	60

Example 4

Antisense Inhibition of a Human C9ORF72 mRNA Variant in HepG2 Cells

Antisense oligonucleotides described above are tested for their effects on C9ORF72 mRNA in vitro. Cultured HepG2 cells at a density of 20,000 cells per well are electroporated with an antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Human primer probe set RTS3905 (forward primer sequence GGGTCTAGCAAGAGCAGGTG, designated herein as SEQ ID NO: 12; reverse primer sequence GTCTTGGCAACAGCTGGAGAT, designated herein as SEQ ID NO: 13; probe sequence TGATGTCGACTCTTTGCCCACCGC, designated herein as SEQ ID NO: 14—a TAQ-man primer probe set) are used. RTS3905 detects an mRNA variant (e.g. NM_001256054.1) processed from a pre-mRNA variant containing the hexanucleotide repeat. The mRNA variant processed from a pre-mRNA variant containing the hexanucleotide repeat is herein the “C9ORF72 pathogenic associated mRNA variant.” A pre-mRNA contains the hexanucleotide repeat when transcription of the pre-mRNA begins in the region from the start site of exon 1A to the start site of exon 1B (generally nucleotides 1107 to 1520 of the genomic sequence: SEQ ID NO: 2, the complement of GENBANK Accession No. NT_008413.18 truncated from nucleosides 27535000 to 2756500. Therefore, oligonucleotides designed in this region selectively target the pre-mRNA variant containing the hexanucleotide repeat. RTS3905 measures an mRNA product (i.e. the C9ORF72 pathogenic associated mRNA variant) of the pre-mRNA variant containing the hexanucleotide repeat and, therefore, measures the reduction of the pre-mRNA variant containing the hexanucleotide repeat. The levels of the C9ORF72 pathogenic associated mRNA variant are

95

normalized to the total RNA content of the cell, as measured by RIBOGREEN®, then the normalized mRNA variant levels are compared to those of cells that were not treated with antisense oligonucleotide.

Example 5

Dose-Dependent Antisense Inhibition of a Human C9ORF72 mRNA Variant

Antisense oligonucleotides described above are tested at various doses in HepG2 cells. Cells are plated at a density of 20,000 cells per well and electroporated with antisense oligonucleotide. After a treatment period of approximately 16 hours or 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Human C9ORF72 primer probe set RTS3905 is used to measure the C9ORF72 pathogenic associated mRNA variant. The levels of the C9ORF72 pathogenic associated mRNA variant are adjusted according to total RNA content, as measured by RIBOGREEN®. The half maximal inhibitory concentration (IC₅₀) of each oligonucleotide is calculated based on the inhibition of mRNA variant levels observed at each individual dose of antisense oligonucleotide.

Example 6

Antisense Inhibition of C9ORF72 by Human-Rhesus Cross-Reactive Antisense Oligonucleotides in LLC-MK2 Cells

Antisense oligonucleotides described above that are fully cross-reactive with a rhesus C9ORF72 nucleic acid are tested for their effects on rhesus C9ORF72 mRNA in vitro. Cultured rhesus LLC-MK2 cells at a density of 20,000 cells per well are electroporated with antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Primer probe set RTS3750 (forward sequence TGTGACAGTTGGAATGCAGTGA, designated herein as SEQ ID NO: 15; reverse sequence GCCACTTAAAGCAATCTCTGTCTTG, designated herein as SEQ ID NO: 16; probe sequence TCGACTCTTTGCCACCGCCA, designated herein as SEQ ID NO: 17—a TAQ-man primer probe set) is used to measure total C9ORF72 mRNA levels. RTS3750 targets exon 2 of the mRNA transcripts and, therefore, measures total mRNA transcripts. C9ORF72 mRNA levels are adjusted according to total RNA content, as measured by RIBOGREEN®, then the normalized mRNA variant levels are compared to those of cells that were not treated with antisense oligonucleotide.

Example 7

Dose-Dependent Antisense Inhibition of Human C9ORF72 mRNA in LLC-MK2

Antisense oligonucleotides described above are tested at various doses in LLC-MK2 cells. Cells are plated at a density of 20,000 cells per well and electroporated with antisense oligonucleotide. After a treatment period of approximately 16 hours or 24 hours, RNA is isolated from the cells and C9ORF72 mRNA levels are measured by quantitative real-time PCR. Primer probe set RTS3750 is used to measure total C9ORF72 mRNA levels. C9ORF72

96

mRNA levels are adjusted according to total RNA content, as measured by RIBOGREEN®, then the normalized mRNA variant levels are compared to those of cells that were not treated with antisense oligonucleotide.

Example 8

Antisense Inhibition of Human C9ORF72 mRNA in a Transgenic Mouse Model

Antisense oligonucleotides described above are tested in two BAC transgenic mouse lines, designated herein as C9B41 and C9B183, that each express a truncated human C9ORF72 gene comprising exons 1-5. The truncated human C9ORF72 genes of the C9B41 and C9B183 mouse lines comprise 110 and 450 hexanucleotide repeats, respectively. Each mouse in each treatment group receives 350 µg of an antisense oligonucleotide, then expression levels of human C9ORF72 RNA are analyzed by RT-PCR. Dose responses are also performed.

Example 9

Antisense Inhibition of C9ORF72 mRNA in Patient Fibroblasts

Antisense oligonucleotides described above were tested for their effects on C9ORF72 mRNA in vitro. The antisense oligonucleotides listed in the table below were added to a plate before patient fibroblasts F09-152 were added at a density of 20,000 cells per well. The final concentrations of the antisense oligonucleotides after the addition of the cells are listed in the table below. After 30 seconds of shaking, the cells were electroporated then transferred to a Primaria coated culture plate. After 16 hours, RNA was isolated from the cells, and levels of total C9ORF72 mRNA (i.e., mRNA starting from exon 1A and mRNA starting from exon 1B) and levels of the pathogenic associated mRNA (see Example 4) were measured by RT-qPCR. Human primer probe sets RTS 3750 (see Example 6) and RTS3905 (see Example 4) were used to detect the total C9ORF72 mRNA and C9ORF72 pathogenic associated mRNA, respectively. The levels of the C9ORF72 total mRNA and C9ORF72 pathogenic associated mRNA were normalized to the total RNA content of the cell, as measured by RIBOGREEN®, then the normalized mRNA levels were compared to those of cells that were not treated with antisense oligonucleotide. The results are shown in the table below. An entry of “nd” means not determined. IC₅₀ values listed as “nd” were not determined, because the target was not inhibited sufficiently to determine an IC₅₀. The results show that all of the oligonucleotides listed below inhibited the pathogenic associated C9ORF72 mRNA variant. Oligonucleotides that do not specifically target the repeat variant pre-mRNA inhibited both the pathogenic associated C9ORF72 mRNA and total C9ORF72 mRNA. Oligonucleotides that specifically target the repeat variant pre-mRNA selectively inhibited the pathogenic associated C9ORF72 mRNA.

97

TABLE 10

C9ORF72 mRNA levels following antisense inhibition in patient fibroblasts					
Isis No.	Concen- tration (μM)	Total C9ORF72 mRNA		Pathogenic associated C9ORF72 mRNA	
		Level (% UTC)	IC ₅₀ (μM)	Level (% UTC)	IC ₅₀ (μM)
791658	0.12	87	nd	26	<0.1
	0.60	80		9	
	3.00	87		nd	
	15.00	105		4	
791664	0.12	85	nd	13	<0.1
	0.60	101		7	
	3.00	125		4	
	15.00	114		3	
801278	0.12	91	nd	19	<0.1
	0.60	71		7	
	3.00	85		nd	
	15.00	90		nd	
801279	0.12	98	nd	57	0.12
	0.60	77		27	
	3.00	77		3	
	15.00	86		5	
801282	0.12	100	nd	51	0.08
	0.60	84		12	
	3.00	97		6	
	15.00	190		nd	
801283	0.12	87	nd	40	<0.1
	0.60	75		8	
	3.00	94		4	
	15.00	111		4	
801285	0.12	103	nd	35	<0.1
	0.60	78		23	
	3.00	75		7	
	15.00	79		6	
801286	0.12	94	nd	33	<0.1
	0.60	85		14	
	3.00	97		5	
	15.00	88		3	
801287	0.12	85	nd	31	<0.1
	0.60	76		11	
	3.00	77		9	
	15.00	87		2	
801288	0.12	77	nd	13	<0.1
	0.60	86		11	
	3.00	123		5	
	15.00	177		3	
801292	0.12	69	nd	29	<0.1
	0.60	71		12	
	3.00	82		3	
	15.00	71		nd	
801293	0.12	77	nd	51	0.09
	0.60	70		10	
	3.00	74		3	
	15.00	81		1	
801294	0.12	75	nd	27	<0.1
	0.60	67		9	
	3.00	107		7	
	15.00	146		2	
801316	0.12	73	nd	35	<0.1
	0.60	68		7	
	3.00	65		1	
	15.00	78		1	
802459	0.12	58	0.10	75	0.18
	0.60	15		6	
	3.00	2		11	
	15.00	1		6	

98

TABLE 10-continued

C9ORF72 mRNA levels following antisense inhibition in patient fibroblasts					
Isis No.	Concen- tration (μM)	Total C9ORF72 mRNA		Pathogenic associated C9ORF72 mRNA	
		Level (% UTC)	IC ₅₀ (μM)	Level (% UTC)	IC ₅₀ (μM)
802464	0.12	81	0.39	69	0.21
	0.60	31		23	
	3.00	6		7	
	15.00	2		3	
802465	0.12	40	<0.1	35	<0.1
	0.60	7		7	
	3.00	2		nd	
	15.00	1		nd	
802468	0.12	52	0.10	40	<0.1
	0.60	15		13	
	3.00	3		7	
	15.00	2		nd	
802469	0.12	69	0.17	57	~0.5
	0.60	16		9	
	3.00	4		nd	
	15.00	1		nd	
802471	0.12	71	0.27	54	0.12
	0.60	29		20	
	3.00	6		6	
	15.00	7		nd	
802473	0.12	63	0.18	48	<0.1
	0.60	27		14	
	3.00	8		11	
	15.00	4		nd	
802477	0.12	54	0.12	32	<0.1
	0.60	16		12	
	3.00	4		11	
	15.00	3		3	
806676	0.12	66	nd	20	<0.1
	0.60	66		4	
	3.00	76		2	
	15.00	64		nd	
806679	0.12	71	nd	23	<0.1
	0.60	68		2	
	3.00	78		1	
	15.00	84		1	
806680	0.12	89	nd	41	<0.1
	0.60	76		13	
	3.00	65		7	
	15.00	84		nd	
806690	0.12	99	nd	44	<0.1
	0.60	88		17	
	3.00	85		2	
	15.00	77		1	

Example 10

Antisense Inhibition of Human C9ORF72 mRNA in a Transgenic Mouse Model

Antisense oligonucleotides described above were tested in a BAC transgenic mouse line, C9B41 (see Example 8), that expresses a human C9ORF72 gene comprising the promoter region through exon 5 and 110 hexanucleotide repeats. Each treatment group consisted of 2-3 mice. Each mouse received a single ICVB of 350 μg of an antisense oligonucleotide listed in the tables below or PBS. Two weeks later, the mice were euthanized, and expression levels of the human pathogenic associated C9ORF72 mRNA variant and/or total human C9ORF72 mRNA were analyzed by RT-qPCR as described in Example 9. Analysis of the pathogenic associated C9ORF72 variant mRNA levels was not completed for the oligonucleotides that do not specifically

target the C9ORF72 repeat variant pre-mRNA. The results in the tables below show the average percent normalized human C9ORF72 mRNA levels relative to the normalized average for the PBS treated group.

TABLE 11

Human C9ORF72 mRNA levels following antisense inhibition in transgenic mice				
Isis No.	Spinal Cord (% PBS treated)		Cortex (% PBS treated)	
	Pathogenic variant	Total	Pathogenic variant	Total
791658	6	69	12	47
791659	18	72	27	57
791664	13	55	15	39
801274	49	65	27	37
801276	10	47	12	29
801277	9	44	10	28
801278	8	39	10	27
801279	33	55	40	47
801281	16	48	17	32
801282	18	49	31	42
801283	15	46	17	31
801285	29	51	24	36
801286	37	61	37	47
801287	13	47	30	40
801288	18	52	39	48
801292	25	49	33	40
801293	20	50	29	38
801310	70	80	68	70
801312	43	61	46	55
801315	39	59	41	51
801316	27	56	50	58
806673	38	66	66	66
806674	78	88	98	96
806675	67	86	81	85
806676	29	64	52	58
806677	61	77	69	68
806678	83	96	95	102
806679	24	63	26	44
806680	29	62	41	56
806681	39	68	36	54
806684	44	69	62	64
806685	69	87	56	58
806686	59	75	57	68
806687	74	88	77	80
806688	34	62	42	51
806689	67	92	87	88
806690	28	61	57	68
806692	45	64	44	50

TABLE 12

Human C9ORF72 mRNA levels following antisense inhibition in transgenic mice		
Isis No.	Total human C9ORF72 mRNA (% PBS)	
	Spinal Cord	Cortex
802459	28	30
802461	22	19
802464	30	28
802465	25	25
802467	13	14

TABLE 12-continued

Human C9ORF72 mRNA levels following antisense inhibition in transgenic mice		
Isis No.	Total human C9ORF72 mRNA (% PBS)	
	Spinal Cord	Cortex
802468	24	26
802469	24	20
802470	18	20
802471	36	44
802473	28	38
802475	15	15
802477	14	15

Example 11

Dose Dependent Antisense Inhibition of Human C9ORF72 mRNA in a Transgenic Mouse Model

Antisense oligonucleotides described above were tested in two BAC transgenic mouse lines, C9B41 and C9B183, that each express a truncated human C9ORF72 gene comprising exons 1-5. The truncated human C9ORF72 genes of the C9B41 and C9B183 mouse lines comprise 110 and 450 hexanucleotide repeats, respectively (see Example 8). Each treatment group consisted of 2-4 mice. Each mouse received a single ICVB of 30 µg, 100 µg, 300 µg, or 700 µg of an antisense oligonucleotide as listed in the tables below or PBS. Two weeks later, the mice were euthanized, and expression levels of the human pathogenic associated C9ORF72 mRNA variant and/or total human C9ORF72 mRNA were analyzed by RT-qPCR as described in Example 9. The results in the tables below show the average percent normalized human C9ORF72 mRNA levels relative to the normalized average for the PBS treated group. A value of 100 or greater means the antisense oligonucleotide did not reduce mRNA or increased the amount of mRNA.

TABLE 13

Human C9ORF72 mRNA levels following dose dependent antisense inhibition in C9B41 transgenic mice			
Isis No.	Concen- tration (µg)	Spinal Cord (% PBS treated) Total C9	Cortex (% PBS treated) Total C9
802459	30	84	85
	100	63	61
	300	33	26
	700	29	17
802473	30	74	90
	100	60	77
	300	37	39
	700	28	26

TABLE 14

Human C9ORF72 mRNA levels following dose dependent antisense inhibition in C9B183 transgenic mice			
Isis No.	Concen- tration (µg)	Spinal Cord (% PBS treated) Pathogenic variant	Cortex (% PBS treated) Pathogenic variant
801287	30	76	10
	100	37	75
	300	21	32
	700	11	15
806679	30	72	13
	100	59	100
	300	21	61
	700	10	16
806680	30	94	118
	100	72	97
	300	34	89
	700	30	56
806690	30	52	125
	100	60	131
	300	43	119
	700	19	49

Example 12

Tolerability of Antisense Oligonucleotides Targeting Human C9Orf72 in Mice

Wild type C57/B16 mice each received a single ICV dose of 700 µg of an antisense oligonucleotide listed in the table below, as described in Example 2. Each treatment group consisted of 4 mice. At 8 weeks post-injection, mice were evaluated according to 7 different criteria. The criteria are (1) the mouse was bright, alert, and responsive; (2) the mouse was standing or hunched without stimuli; (3) the mouse showed any movement without stimuli; (4) the mouse demonstrated forward movement after it was lifted; (5) the mouse demonstrated any movement after it was lifted; (6) the mouse responded to tail pinching; (7) regular breathing. For each of the 7 criteria, a mouse was given a subscore of 0 if it met the criteria and 1 if it did not. After all 7 criteria were evaluated, the scores were summed for each mouse and averaged within each treatment group. The results are presented in the table below.

Animals were sacrificed at 8 weeks. The cortex and spinal cord were collected from each animal, and RT-PCR was performed. Expression levels of allograft inflammatory factor (AIF1) were determined as a measure of inflammation. Expression levels of glial fibrillary acidic protein (GFAP) were also determined as a measure of glial cell activation. Results were normalized to Gpadh and are presented relative to PBS control (1.0) in the table below. “N.D.” indicates there was no data because the experiment was not performed. An asterisk indicates that the corresponding result is the average of 1-3 mice.

TABLE 15

Tolerability of antisense oligonucleotides targeting C9Orf72 in mice						
ISIS No.	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cerebellum)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
791656	3.5	1.1*	1.3*	1.0*	1.3*	22
791657	3.5	2.3*	1.0*	1.4*	0.8*	22
791658	0.0	1.9	1.2	1.4	1.3	22
791659	0.0	1.8	1.3	1.5	2.5	22

TABLE 15-continued

Tolerability of antisense oligonucleotides targeting C9Orf72 in mice						
ISIS No.	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cerebellum)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
791660	1.8	2.5*	1.4*	1.9*	0.8*	23
791661	1.8	1.7*	1.3*	1.5*	1.1*	24
791662	5.3	1.1*	1.2*	1.1*	0.8*	23
791663	7.0	N.D.	N.D.	N.D.	N.D.	24
791664	0.0	1.6	1.6	1.2	0.9	23
791665	3.5	1.9*	1.5*	1.9*	0.9*	24
801274	0.0	1.2	1.3	1.3	1.1	25
801275	3.5	1.4*	1.7*	1.6*	2.1*	26
801276	0.0	13.1	2.6	3.7	1.2	27
801277	0.0	2.9	1.4	2.3	1.5	28
801278	0.0	2.0	1.3	1.8	1.1	29
801279	0.0	1.2	1.4	1.3	1.0	25
801280	5.3	1.5*	1.5*	1.4*	1.2*	26
801281	0.0	3.6	2.2	2.4	1.6	27
801282	0.0	1.2	1.2	1.2	1.0	28
801283	0.0	1.4	1.4	1.0	0.9	29
801284	1.8	1.2*	1.5*	0.9*	1.0*	30
801285	0.0	1.2	1.2	1.1	0.8	31
801286	0.0	1.1	1.2	1.0	1.1	32
801287	0.0	1.3	1.2	1.1	1.4	33
801288	0.0	1.3	1.3	1.1	0.9	34
801289	7.0	N.D.	N.D.	N.D.	N.D.	35
801290	3.5	1.2*	1.3*	1.0*	1.1*	30
801291	0.0	2.2	2.2	1.5	1.1	31
801292	0.0	1.3	1.3	1.2	0.9	32
801293	0.0	1.6	1.6	1.3	1.5	33
801294	0.0	1.2	1.2	1.0	1.1	34
801295	1.8	1.4*	1.4*	1.0*	0.9*	35
801296	7.0	N.D.	N.D.	N.D.	N.D.	36
801297	3.5	1.0*	1.0*	0.9*	0.8*	37
801307	5.3	1.2*	0.9*	1.1*	0.8*	41
801308	5.3	1.2*	1.2*	0.9*	1.1*	42
801309	7.0	N.D.	N.D.	N.D.	N.D.	43
801310	1.8	1.2*	1.2*	1.0*	1.0*	44
801311	7.0	N.D.	N.D.	N.D.	N.D.	41
801312	0.0	2.0	1.3	1.0	1.4	42
801313	3.5	1.7*	1.1*	0.9*	0.9*	43
801314	0.0	1.9	1.2	1.0	2.9	44
801315	1.8	1.4*	1.3*	1.0*	1.1*	36
801316	0.0	1.5	1.4	1.0	1.2	37
801298	3.5	1.4*	1.0*	1.1*	1.3*	38
801299	5.3	1.2*	1.2*	1.0*	1.0*	39
801300	3.5	1.2*	1.1*	0.9*	1.0*	40
801301	7.0	N.D.	N.D.	N.D.	N.D.	38
801302	5.3	1.0*	1.1*	0.8*	0.8*	39
801303	1.8	1.1*	1.0*	1.0*	1.1*	40
801304	4.8	14.3*	5.4*	2.2*	3.5*	38
801305	5.3	3.6*	1.7*	1.6*	1.6*	39
801306	0.0	4.7	1.8	1.8	1.8	40
806673	0.0	1.1	1.0	1.0	9.1	49
806674	0.0	1.0	1.0	0.8	8.2	50
806675	0.0	1.0	1.0	0.8	10.3	51
806676	0.0	1.0	1.0	0.8	1.5	49
806677	0.0	0.9	1.0	0.9	1.5	50
806678	0.0	1.0	0.9	1.1	1.2	51
806679	0.0	1.2	1.1	1.1	1.3	49
806680	0.0	1.0	1.1	0.8	1.0	50
806681	0.0	1.0	1.0	0.9	1.4	51
806682	1.8	1.4*	1.2*	1.0*	1.2*	49
806683	0.0	1.2	1.2	0.9	1.2	50
806684	0.0	1.2	1.1	0.9	2.0	51
806685	0.0	1.0	0.9	0.8	1.5	52
806686	0.0	0.9	0.9	0.8	1.1	53
806687	0.0	1.0	1.0	0.8	1.0	54
806688	1.8	1.1*	1.0*	0.8*	1.8*	55
806689	0.0	1.1	1.0	0.8	1.3	52
806690	0.0	1.0	1.1	0.9	1.1	53
806691	5.3	N.D.	N.D.	N.D.	N.D.	54
802459	0.0	0.9	0.9	0.9	1.2	21
802460	5.3	1.1*	1.1*	1.0*	1.4*	45
802461	3.5	1.4*	1.4*	0.9*	0.9*	46
802462	3.5	0.8*	0.9*	0.9*	1.1*	21
802463	7.0	N.D.	N.D.	N.D.	N.D.	45
802464	1.8	1.4*	1.2*	1.5*	1.4*	46

TABLE 15-continued

Tolerability of antisense oligonucleotides targeting C9Orf72 in mice						
ISIS No.	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cerebellum)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
802465	0.0	0.9	1.1	1.1	1.6	21
802466	1.8	1.0*	1.0*	1.0*	1.4*	45
802467	1.3	4.9	2.4	2.7	2.8	46
802468	0.0	1.4	1.3	1.2	2.2	21
802469	0.0	1.0	1.0	1.0	1.2	45
802470	1.5	3.2	3.4	2.6	3.4	46

parts were moving, it would receive a score of 1. Saline treated rats generally receive a score of 0. A score of at the top end of the range would be suggestive of acute toxicity. Results are presented in the table below as the average score for each treatment group.

Animals were sacrificed at 8 weeks. The cortex and spinal cord were collected from each animal, and RT-PCR was performed. Expression levels of AIF1 were determined as a measure of inflammation. Expression levels of (GFAP) were also determined as a measure of glial cell activation. An asterisk indicates that the corresponding result is the average of 2-3 mice. Results were normalized to Gapdh and are presented relative to PBS control (1.0) in the table below.

TABLE 16

Tolerability of antisense oligonucleotides targeting C9Orf72 in rats							
ISIS No.	Score 3 hours after injection	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cortex)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
801287	2.5	0.0	1.9	1.2	1.3	1.3	33
801288	4.0	3.5	1.3*	1.3*	1.2*	1.3*	34
806676	2.0	1.8	1.6*	1.5*	1.3*	2.5*	49
806679	1.2	0.0	1.4	1.3	1.3	1.8	49
806680	2.0	0.3	1.4	1.4	1.3	1.3	50
806690	1.3	0.0	1.5	1.4	1.3	2.0	53
802459	2.0	0.0	1.4	1.2	1.5	1.5	21
802473	2.0	2.3	1.8*	1.6*	1.7*	1.6*	47

TABLE 15-continued

Tolerability of antisense oligonucleotides targeting C9Orf72 in mice						
ISIS No.	Score 8 weeks after injection	AIF1 (spinal cord)	AIF1 (cerebellum)	GFAP (spinal cord)	GFAP (cortex)	SEQ ID NO.
802471	0.0	0.9	0.9	0.9	1.2	47
802472	3.5	1.6*	1.4*	0.9*	1.5*	48
802473	0.0	1.0	0.9	0.9	1.3	47
802474	1.8	1.3*	1.2*	1.0*	1.8*	48
802475	7.0	N.D.	N.D.	N.D.	N.D.	47
802476	1.8	1*	1.2*	0.8*	1.8*	48
802477	0.0	1.9	1.7	1.3	2.1	47
802478	3.5	1.4*	1.1*	1.1*	1.3*	48

Example 14

Tolerability of Antisense Oligonucleotides Targeting Human C9ORF72 in Non-Human Primates

Female cynomolgus monkeys (2-6 kg) were given 3 doses of 35 mg of antisense oligonucleotide on days 1, 14, and 28 via intrathecal bolus injection (1 mL slow bolus followed by 0.25 mL flush). Each treatment group contained four monkeys. Two weeks after the final dose, animals were sacrificed and RT-PCR was performed on various CNS tissues. Expression levels of AIF1 were determined as a measure of inflammation and expression levels of GFAP were determined as a measure of glial cell activation. Results were normalized to GADPH and are presented relative to PBS control (1.0) in the table below for ISIS No. 801287, 802459, and 806679.

TABLE 17

Tolerability of antisense oligonucleotides targeting C9ORF72 in cynomolgus monkeys						
Brain Region	AIF1			GFAP		
	801287	802459	806679	801287	802459	806679
Cervical spinal cord	1.0	0.9	0.9	1.1	1.0	1.1
Thoracic spinal cord	1.1	0.9	0.9	1.2	1.2	1.2
Temporal cortex	1.0	1.0	1.5	1.2	1.1	1.7
Motor cortex	1.0	1.0	1.2	1.2	0.7	1.3
Lumbar spinal cord	1.2	0.9	0.9	1.3	0.9	1.1
Hippocampus	1.0	1.0	1.3	1.9	1.9	1.7
Frontal cortex	1.3	0.9	1.3	1.0	0.7	1.3

Example 13

Tolerability of Antisense Oligonucleotides Targeting Human C9Orf72 in Rats

Sprague Dawley rats were separated into groups of 4 or 6 rats. Each rat in each group of rats was administered a single 3 mg intrathecal (IT) dose of the oligonucleotide indicated in the table below. At 3 hours and at 8 weeks following the IT dose, the movement of 7 different parts of the body was evaluated for each rat. The 7 body parts are (1) the rat's tail; (2) the rat's posterior posture; (3) the rat's hind limbs; (4) the rat's hind paws; (5) the rat's forepaws; (6) the rat's anterior posture; (7) the rat's head. For each of the 7 different body parts, each rat was given a sub-score of 0 if the body part was moving or 1 if the body part was paralyzed. After each of the 7 body parts were evaluated, the sub-scores were summed for each rat and then averaged for each group. For example, if a rat's tail, head, and all other evaluated body parts were moving 3 hours after the 3 mg IT dose, it would get a summed score of 0. If another rat was not moving its tail 3 hours after the 3 mg IT dose but all other evaluated body

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 60

<210> SEQ ID NO 1
<211> LENGTH: 3339
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 1

acgtaaccta cgggtgtcccg ctaggaaaga gaggtgcgtc aaacagcgac aagttccgcc	60
cacgtaaaag atgacgcttg gtgtgtcagc cgtccctgct gcccggttgc ttctcttttg	120
ggggcggggt ctagcaagag caggtgtggg tttaggagat atctccggag catttgata	180
atgtgacagt tggaatgcag tgatgtcgac tctttgcca ccgccatctc cagctgttgc	240
caagacagag attgctttaa gtggcaaadc acctttatta gcagctactt ttgcttactg	300
ggacaatatt cttggtccta gagtaaggca catttgggct ccaaagacag aacagggtact	360
tctcagtgat ggagaaataa cttttcttgc caaccacact ctaaatggag aaatccttcg	420
aaatgcagag agtgggtgcta tagatgtaaa gttttttgtc ttgtctgaaa agggagtgat	480
tattgtttca ttaatctttg atggaaactg gaatggggat cgcagcacat atggactatc	540
aattatactt ccacagacag aacttagttt ctacctcca cttcatagag tgtgtgttga	600
tagattaaca catataatcc ggaaaggaag aatatggatg cataaggaaa gacaagaaaa	660
tgtccagaag attatcttag aaggcacaga gagaatggaa gatcagggtc agagtattat	720
tccaatgctt actggagaag tgattcctgt aatggaactg ctttcatcta tgaaatcaca	780
cagtgttctt gaagaaatag atatagctga tacagtactc aatgatgatg atattgggtga	840
cagctgtcat gaaggctttc ttctcaatgc catcagctca cacttgcaaa cctgtggctg	900
ttccgttgta gtaggtagca gtgcagagaa agtaataaag atagtcagaa cattatgcct	960
ttttctgact ccagcagaga gaaaatgctc caggttatgt gaagcagaat catcatttaa	1020
atatgagtca gggctctttg tacaaggcct gctaaaggat tcaactggaa gctttgtgct	1080
gcctttccgg caagtcatgt atgtccata tcccaccaca cacatagatg tggatgtcaa	1140
tactgtgaag cagatgccac cctgtcatga acatatttat aatcagcgta gatacatgag	1200
atccgagctg acagccttct ggagagccac ttcagaagaa gacatggctc aggatacgat	1260
catctacact gacgaaagct ttactcctga tttgaatatt tttcaagatg tcttacacag	1320
agacactcta gtgaaagcct tcctggatca ggtctttcag ctgaaacctg gcttatctct	1380
cagaagtact ttccttgcac agtttctact tgtccttcac agaaaagcct tgacactaat	1440
aaaatatata gaagacgata cgcagaaggg aaaaaagccc tttaaatctc ttcggaacct	1500
gaagatagac cttgatttaa cagcagaggg cgatcttaac ataataatgg ctctgggtga	1560
gaaaattaaa ccaggcctac actcttttat ctttggaaga cctttctaca ctagtgtgca	1620
agaacgagat gttctaata ctttttaaat gtgtaactta ataagcctat tccatcacia	1680
tcatgatcgc tggtaaagta gctcagtggt gtggggaaac gtccccctgg atcatactcc	1740
agaattctgc tctcagcaat tgcagttaag taagttacac tacagttctc acaagagcct	1800
gtgaggggat gtcaggtgca tcattacatt ggggtgtctct tttcctagat ttatgctttt	1860
gggatacaga cctatgttta caatataata aatattattg ctatctttta aagatataat	1920
aataggatgt aaacttgacc acaactactg tttttttgaa atacatgatt catggtttac	1980
atgtgtcaag gtgaaatctg agttggcttt tacagatagt tgactttcta tcttttgga	2040
ttctttggtg tgtagaatta ctgtaatact tctgcaatca actgaaaact agagccttta	2100

-continued

aatgatttca attccacaga aagaaagtga gcttgaacat aggatgagct ttagaaagaa	2160
aattgatcaa gcagatgttt aattggaatt gattattaga tctacttttg tggatttagt	2220
ccctgggatt cagtctgtag aaatgtctaa tagttctcta tagtccttgt tccctggtgaa	2280
ccacagttag ggtgttttgt ttattttatt gttcttgcta ttgttgatat tctatgtagt	2340
tgagctctgt aaaaggaaat tgtattttat gttttagtaa ttgttgccaa ctttttaa	2400
taattttcat tatttttgag ccaaattgaa atgtgcacct cctgtgcctt ttttctcctt	2460
agaaaatcta attacttgga acaagttcag atttcactgg tcagtcattt tcatcttggt	2520
ttcttcttgc taagtcttac catgtacctg ctttggcaat cattgcaact ctgagattat	2580
aaaatgcctt agagaatata ctaactaata agatcttttt ttcagaaaca gaaaatagtt	2640
ccttgagtac ttccttcttg catttctgcc tatgtttttg aagttgttgc tgtttgcctg	2700
caataggcta taaggaatag caggagaaat tttactgaag tgctgttttc ctaggtgcta	2760
ctttggcaga gctaagttat cttttgtttt cttaatgcgt ttggaccatt ttgctggcta	2820
taaaataact gattaatata attctaacac aatgttgaca ttgtagttac acaaacacaa	2880
ataaatatth tatttaaaat tctggaagta atataaaagg gaaaatatat ttataagaaa	2940
gggataaagg taatagagcc cttctgcccc ccacccacca aatttacaca acaaaatgac	3000
atgttcgaat gtgaaaggtc ataatagctt tcccatcatg aatcagaaag atgtggacag	3060
cttgatgttt tagacaacca ctgaactaga tgactgttgt actgtagctc agtcatttaa	3120
aaaatatata aatactacct tgtagtgtcc catactgtgt tttttacatg gtagattctt	3180
atttaagtgc taactggtta ttttctttgg ctggtttatt gtactgttat acagaatgta	3240
agttgtacag tgaaataagt tattaaagca tgtgtaaaca ttgttatata tcttttctcc	3300
taaatggaga attttgaata aaatatatth gaaatthtg	3339
<210> SEQ ID NO 2	
<211> LENGTH: 30001	
<212> TYPE: DNA	
<213> ORGANISM: homo sapiens	
<400> SEQUENCE: 2	
caaagaaaag ggggaggttt tgtaaaaaa gagaaatgtt acatagtgtt ctttgagaaa	60
attcattggc actattaagg atctgaggag ctggtgagtt tcaactggtg agtgatggtg	120
gtagataaaa ttagagctgc agcaggcat tttagcaact attagataaa actgggtctca	180
ggtcacaacg ggcagttgca gcagctggac ttggagagaa ttacactgtg ggagcagtgt	240
catttgtcct aagtgccttt ctaccccccta cccccactat tttagttggg tataaaaaga	300
atgacccaat ttgtatgatc aactttcaca aagcatagaa cagtaggaaa aggggtctgtt	360
tctgcagaag gtgtagacgt tgagagccat tttgtgtatt tattcctccc tttcttctc	420
ggtgaatgat taaaacgttc tgtgtgattt ttagtgatga aaaagattaa atgctactca	480
ctgtagtaag tgccatctca cacttgcala tcaaaaggca cacagtttaa aaaacctttg	540
tttttttaca catctgagtg gtgtaaatgc tactcatctg tagtaagtgg aatctatata	600
cctgcagacc aaaagacgca aggtttcaaa aatctttgtg ttttttacac atcaaacaga	660
atggtacgtt tttcaaaagt taaaaaaaa caactcatcc acatattgca actagcaaaa	720
atgacattcc ccagtgtgaa aatcatgctt gagagaatth ttacatgtaa aggcaaaatt	780
gcgatgactt tgcaggggac cgtgggattc ccgcccgcag tgccggagct gtcccctacc	840

-continued

agggtttgca	gtggagtttt	gaatgcactt	aacagtgtct	tacggtaaaa	acaaaatttc	900
atccaccaat	tatgtgttga	gcgcccactg	cctaccaagc	acaaacaaaa	ccattcaaaa	960
ccacgaaatc	gtcttcactt	tctccagatc	cagcagcctc	ccctattaag	gttcgcacac	1020
gctattgcgc	caacgctcct	ccagagcggg	tcttaagata	aaagaacagg	acaagttgcc	1080
ccgccccatt	tcgctagcct	cgtgagaaaa	cgtcatcgca	catagaaaac	agacagacgt	1140
aacctacggg	gtcccgctag	gaaagagagg	tgcgtcaaac	agcgacaagt	tcgccccacg	1200
taaaagatga	cgcttggtgt	gtcagccgtc	cctgctgccc	ggttgcttct	cttttggggg	1260
cggggtctag	caagagcagg	tgtgggttta	ggaggtgtgt	gtttttgttt	ttcccaccct	1320
ctctccccac	tacttgctct	cacagtactc	gctgaggggtg	aacaagaaaa	gacctgataa	1380
agattaacca	gaagaaaaa	aggagggaaa	caaccgcagc	ctgtagcaag	ctctggaact	1440
caggagtcgc	gcgctagggg	cgggggcggg	ggcggggcg	tggtcggggc	gggcccgggg	1500
gcggggccgg	ggcggggctg	cggttgcggg	gcctgcgccc	gcggcggcg	aggcgcaggc	1560
ggtggcgagt	gggtgagtga	ggaggcggca	tcttgccggg	tggctgtttg	gggttcggct	1620
gccgggaaga	ggcgcgggta	gaagcggggg	ctctcctcag	agctcgacgc	atttttactt	1680
tcctctcat	ttctctgacc	gaagctgggt	gtcgggcttt	cgcctctagc	gactgggtga	1740
attgcctgca	tccgggcccc	gggcttcccc	gcggcgccgg	cggcgccggc	ggcgcaggga	1800
caagggatgg	ggatctggcc	tcttccttgc	tttcccgcgc	tcagtaccgc	agctgtctcc	1860
ttcccgggga	cccgtctggg	gcgtgcccgc	tgcgggctcg	agaaaaggga	gcctcgggta	1920
ctgagaggcc	tcgcctgggg	gaaggccgga	gggtgggcgg	cgcgcggctt	ctgcggacca	1980
agtcgggggt	cgctaggaac	ccgagacggg	ccctgccggc	gaggagatca	tgcgggatga	2040
gatgggggtg	tggagacgcc	tgcacaattt	cagcccaagc	ttctagagag	tggtagtgac	2100
ttgcatatga	gggcagcaat	gcaagtcggg	gtgctcccca	ttctgtggga	catgacctgg	2160
ttgcttcaca	gctccgagat	gacacagact	tgcttaaagg	aagtgactat	tgtgacttgg	2220
gcatcacttg	actgatggta	atcagttgtc	taaagaagtg	cacagattac	atgtccgtgt	2280
gctcattggg	tctatctggc	cgcgttgaac	accaccaggc	tttgtattca	gaaacaggag	2340
ggaggtcctg	cactttccca	ggaggggtgg	ccctttcaga	tgcaatcgag	attgttaggc	2400
tctgggagag	tagttgcctg	gttgtggcag	ttggtaaatt	tctattcaaa	cagttgccat	2460
gcaccagttg	ttcacaacaa	gggtacgtaa	tctgtctggc	attacttcta	cttttgtaca	2520
aaggatcaaa	aaaaaaaaag	atactgttaa	gatatgattt	ttctcagact	ttgggaaact	2580
tttaacataa	tctgtgaata	tcacagaaac	aagactatca	tataggggat	attaataacc	2640
tggagtcaga	atacttgaac	tacgggtgtc	tttgacacgg	gcattgttgt	caccacctct	2700
gccaaggcct	gccactttag	gaaaaccctg	aatcagttgg	aaactgctac	atgctgatag	2760
tacatctgaa	acaagaacga	gagtaattac	cacattccag	attgttcact	aagccagcat	2820
ttacctgctc	caggaaaaaa	ttacaagcac	cttatgaagt	tgataaaaata	ttttgtttgg	2880
ctatgttggc	actccacaat	ttgctttcag	agaaacaaag	taaaccaagg	aggacttctg	2940
tttttcaagt	ctgccctcgg	gttctattct	acgttaatta	gatagttccc	aggaggacta	3000
ggttagccta	cctattgtct	gagaaacttg	gaactgtgag	aaatggccag	atagtgatat	3060
gaacttcacc	ttccagtctt	ccctgatgtt	gaagattgag	aaagtgttgt	gaactttctg	3120
gtactgtaaa	cagttcactg	tccttgaagt	ggcctggggc	agctcctgtt	gtggaaagtg	3180
gacggtttag	gatcctgctt	ctctttgggc	tgggagaaaa	taaacagcat	ggttacaagt	3240

-continued

attgagagcc	aggttggaga	aggtaggctta	cacctgtaat	gccagagctt	tgaggaggcgg	3300
aggcaagagg	atcacttgaa	gccaggagtt	caagctcaac	ctgggcaacg	tagaccctgt	3360
ctctacaaaa	aattaaaaac	ttagccgggc	gtggtgatgt	gcacctgtag	tcctagctac	3420
ttgggaggct	gaggcaggag	ggtcatttga	gccaagagt	ttgaagttac	cgagagctat	3480
gatcctgcc	gtgcattcca	gcctggatga	caaaacgaga	ccctgtctct	aaaaaacaag	3540
aagtgagggc	tttatgattg	tagaattttc	actacaatag	cagtggacca	accacctttc	3600
taaataccaa	tcagggaaga	gatggttgat	tttttaacag	acgtttaaag	aaaagcaaa	3660
acctcaaact	tagcactcta	ctaacagttt	tagcagatgt	taattaatgt	aatcatgtct	3720
gcatgtatgg	gattatttcc	agaaagtgt	ttgggaaacc	tctcatgaac	cctgtgagca	3780
agccacogtc	tcactcaatt	tgaatcttgg	cttccctcaa	aagactggct	aatgtttggt	3840
aactctctgg	agtagacagc	actacatgta	cgtaagatag	gtacataaac	aactattggt	3900
tttgagctga	tttttttcag	ctgcatttgc	atgtatggat	ttttctcacc	aaagacgatg	3960
acttcaagta	ttagtataat	aattgtacag	ctctcctgat	tatacttctc	tgtgacattt	4020
catttcccag	gctatttctt	ttggtaggat	ttaaaactaa	gcaattcagt	atgatctttg	4080
tccttcattt	tctttcttat	tctttttgtt	tgtttgtttg	tttgtttttt	tcttgaggca	4140
gagtctctct	ctgtcgccca	ggctggagtg	cagtggcgcc	atctcagctc	attgcaacct	4200
ctgccacctc	cgggttcaag	agattctcct	gcctcagcct	cccagtagc	tgaggattaca	4260
ggtgtccacc	accacacccg	gctaattttt	tgtattttta	gtagaggtgg	ggtttcacca	4320
tgttggccag	gctggtcttg	agctcctgac	ctcaggtgat	ccacctgcct	cggcctacca	4380
aagagctggg	ataacaggtg	tgaccaccca	tgcccgcccc	atTTTTTTTT	tcttattctg	4440
ttaggagtga	gagtgttaact	agcagtataa	tagttcaatt	ttcacaacgt	ggtaaaagtt	4500
tccctataat	tcaatcagat	tttgctccag	ggttcagttc	tgttttagga	aatactttta	4560
ttttcagttt	aatgatgaaa	tattagagtt	gtaatatgtc	ctttatgatt	atccaccttt	4620
ttaacctaaa	agaatgaaag	aaaaatatgt	ttgcaatata	atTTTtatggt	tgtatgttaa	4680
cttaattcat	tatgttggcc	tccagtttgc	tgtgtttagt	tatgacagca	gtagtgtcat	4740
taccatttca	attcagatta	cattcctata	ttgatcatt	gtaaactgac	tgettacatt	4800
gtattaaaaa	cagtggatat	tttaaagaag	ctgtacggct	tatatctagt	gctgtctctt	4860
aagactatta	aattgatata	acataattta	aagtaaatat	tacctaaatg	aatttttgaa	4920
attacaaata	cacgtgttaa	aactgtcggt	gtgttcaacc	atTTctgtac	atacttagag	4980
ttactgtttt	tgccaggctc	tgtatgccta	ctcataatat	gataaaagca	ctcatcta	5040
gctctgtaaa	tagaagtcag	tgttttccat	cagactgaac	tctcttgaca	agatgtggat	5100
gaaattcttt	aagtaaaatt	gtttactttg	tcatacattt	acagatcaaa	tgttagctcc	5160
caaagcaatc	atatggcaaa	gataggtata	tcatagtttg	cctattagct	gctttgtatt	5220
gctattatta	taaatagact	tcacagtttt	agacttgctt	aggtgaaatt	gcaattcttt	5280
ttactttcag	tcttagataa	caagtcttca	attatagtac	aatcacacat	tgttaggaa	5340
tgcataatta	ggcgattttg	tcattatgca	aacatcatag	agtgtactta	cacaaaccta	5400
gatagtatag	cctttatgta	cctaggccgt	atggtatagt	ctgttgctcc	taggccacaa	5460
acctgtacaa	ctgttactgt	actgaatact	atagacagtt	gtaacacagt	ggtaaatatt	5520
tatctaaata	tatgcaaaca	gagaaaaggt	acagttaaag	tatggtataa	aagataatgg	5580

-continued

tatacctgtg	taggccactt	accacgaatg	gagcttgcag	gactagaagt	tgctctgggt	5640
gagtcagtga	gtgagtgggtg	aattaatgtg	aaggcctaga	acactgtaca	ccactgtaga	5700
ctataaacac	agtacgctga	agctacacca	aatttatctt	aacagttttt	cttcaataaa	5760
aaattataac	tttttaactt	tgtaaacctt	ttaatttttt	aacttttaaa	atacttagct	5820
tgaaacacaa	atacattgta	tagctataca	aaaatatctt	ttctttgtat	ccttattcta	5880
gaagcttttt	tctatcttct	atcttaaat	ttttttttta	cttggttagtc	gtttttgtta	5940
aaaactaaaa	cacacacact	ttcacctagg	catagacagg	attaggatca	tcagtatcac	6000
tcccttccac	ctcactgcct	tccacctcca	catcttgctc	cactggaagg	tttttagggg	6060
caataacaca	catgtagctg	tcacctatga	taacagtgtc	ttctgttgaa	tacctcctga	6120
aggacttgcc	tgaggctgtt	ttacatttaa	cttaaaaaaa	aaaaaagtag	aaggagtgca	6180
ctctaaaata	acaataaaaag	gcatagtata	gtgaatacat	aaaccagcaa	tgtagtagtt	6240
tattatcaag	tggtgtacac	tgtaataatt	gtatgtgcta	tactttaaat	aacttgcaaa	6300
atagtactaa	gaccttatga	tggttacagt	gtcactaagg	caatagcata	ttttcaggct	6360
cattgtaatc	taatgggact	accatcatat	atgcagtcta	ccattgactg	aaacggtaca	6420
tggcacataa	ctgtatttgc	aagaatgatt	tgttttacat	taatatcaca	taggatgtac	6480
cttttttagag	tggtatgttt	atgtggatta	agatgtacaa	gttgagcaag	gggaccaaga	6540
gccctggggt	ctgtcttggg	tgtgagcggt	tatgttcttc	tctcatgtc	tgttttctca	6600
ttaaattcaa	aggcttgaac	gggccctatt	tagcccttct	gttttctacg	tgttctaaat	6660
aactaaagct	tttaattct	agccatttag	tgtagaactc	tctttgcagt	gatgaaatgc	6720
tgtattgggt	tcttggttag	catattaaat	atctttatct	ttgtcttgat	acttcaatgt	6780
cgttttaaac	atcaggatcg	ggcttcagta	ttctcataac	cagagagttc	actgaggata	6840
caggactggt	tgcccatctt	ttgttatggc	tccagacttg	tggtatttcc	atgtcttttt	6900
tttttttttt	tttttgacc	ttttagcggc	tttaaagtat	ttctgttggt	aggtgttgta	6960
ttacttttct	aagattactt	aacaaagcac	cacaaactga	gtggctttta	acaacagcaa	7020
tttattctct	cacaattcta	gaagctagaa	gtccgaaatc	aaagtgttga	caggggcatg	7080
atcttcaaga	gagaagactc	tttccttgcc	tcttcctggc	ttctggtggt	taccagcaat	7140
cctgagtgtt	cctttcttgc	cttgtagttt	caacaatcca	gtatctgcct	tttgtcttca	7200
catggctgtc	taccatttgt	ctctgtgtct	ccaaatctct	ctccttataa	acacagcagt	7260
tattggatta	ggccccactc	taatccagta	tgaccccat	ttaacatgat	tacacttatt	7320
tctagataag	gtcacattca	cgtacaccaa	gggttaggaa	ttgaacatat	ctttttgggg	7380
gacacaatc	aaccacaag	tgtcagtctc	tagctgagcc	ttcccttcc	tgtttttctc	7440
cttttttagtt	gctatgggtt	aggggccaaa	tctccagtca	tactagaatt	gcacatggac	7500
tggatatattg	ggaatactgc	gggtctattc	tatgagcttt	agtatgtaac	atttaatatc	7560
agtgtaaaga	agcccttttt	taagttatct	ctttgaattt	ctaaatgtat	gccctgaata	7620
taagtaacaa	gttaccatgt	cttgtaaaat	gatcatatca	acaaacattt	aatgtgcacc	7680
tactgtgcta	gttgaatgtc	tttatcctga	taggagataa	caggattcca	catctttgac	7740
ttaagaggac	aaaccaaata	tgtctaaatc	atctgggggt	ttgatggata	tctttaaatt	7800
gctgaacctc	atcattgggt	tcatatgtca	ttgttttagat	atctccggag	catttggtata	7860
atgtgacagt	tggaatgcag	tgatgtcgac	tctttgccc	ccgccatctc	cagctgttgc	7920
caagacagag	attgctttta	gtggcaaatc	acctttatta	gcagctactt	ttgcttactg	7980

-continued

ggacaatatt	cttggtccta	gagtaaggca	catttgggct	ccaaagacag	aacagggtact	8040
tctcagtgat	ggagaaataa	cttttcttgc	caaccacact	ctaaatggag	aaatccttcg	8100
aaatgcagag	agtgggtgcta	tagatgtaaa	gttttttgtc	ttgtctgaaa	agggagtgat	8160
tattgtttca	ttaatctttg	atggaaactg	gaatggggat	cgcagcacat	atggactatc	8220
aattatactt	ccacagacag	aacttagttt	ctacctccca	cttcatagag	tgtgtgttga	8280
tagattaaca	catataatcc	ggaaaggaag	aatatggatg	cataaggtaa	gtgatttttc	8340
agcttattaa	tcatgttaac	ctatctgttg	aaagcttatt	ttctggtaca	tataaatctt	8400
atTTTTTTaa	ttatatgcag	tgaacatcaa	acaataaatg	ttatttattt	tgcatttacc	8460
ctattagata	caaatacatc	tggctctgata	cctgtcatct	tcatattaac	tgtggaaggt	8520
acgaaatggg	agctccacat	tatagatgaa	aagctaaagc	ttagacaaat	aaagaaactt	8580
ttagaccctg	gattcttctt	gggagccttt	gactctaata	ccttttgttt	ccctttcatt	8640
gcacaattct	gtcttttgc	tactactatg	tgtaagtata	acagttcaaa	gtaatagt	8700
cataagctgt	tggctcatgta	gcctttgggc	tctttaacct	ctttgccaa	ttcccagggt	8760
cataaaatga	ggaggttgaa	tggaatgggt	cccaagagaa	ttccttttaa	tcttacagaa	8820
attattgttt	tcctaaatcc	tgtagttgaa	tatataatgc	tatttacatt	tcagtatagt	8880
tttgatgtat	ctaaagaaca	cattgaattc	tccttcctgt	gttccagttt	gatactaacc	8940
tgaaagtcca	ttaagcatta	ccagttttaa	aaggcttttg	cccaatagta	aggaaaaata	9000
atatctttta	aaagaataat	tttttactat	gtttgcaggc	ttacttcctt	ttttctcaca	9060
ttatgaaact	cttaaaatca	ggagaatctt	ttaaacaaca	tcataatgtt	taatttgaaa	9120
agtgcaagtc	attcttttcc	tttttgaaac	tatgcagatg	ttacattgac	tgttttctgt	9180
gaagttatct	ttttttcact	gcagaataaa	ggttgttttg	attttatttt	gtattgttta	9240
tgagaacatg	catttgttgg	gttaatttcc	taccctgcc	cccatttttt	ccctaaagta	9300
gaaagtat	ttcttgta	ctaaattact	acacaagaac	atgtctattg	aaaaataagc	9360
aagtatcaaa	atggtgtggg	ttgttttttt	aaataaat	tctcttgctc	aggaaagaca	9420
agaaaatgtc	cagaagatta	tcttagaagg	cacagagaga	atggaagatc	aggtatatgc	9480
aaattgcata	ctgtcaaatg	tttttctcac	agcatgtatc	tgtataaggt	tgatggctac	9540
atttgtcaag	gccttgga	catacgaata	agcctttaat	ggagctttta	tggaggtgta	9600
cagaataaac	tggaggaaga	ttccatatac	ttaaacccaa	agagttaaat	cagtaaacaa	9660
aggaaaatag	taattgcatc	tacaaattaa	tatttgctcc	cttttttttt	ctgtttgccc	9720
agaataaatt	ttggataact	tgttcatagt	aaaaataaaa	aaaattgtct	ctgatatgtt	9780
ctttaaggta	ctacttctcg	aacctttccc	tagaagtagc	tgtaacagaa	ggagagcata	9840
tgtaccctg	aggtatctgt	ctgggggtgta	ggcccaggtc	cacacaatat	ttcttctaag	9900
tcttatgttg	tatcgtaag	actcatgcaa	ttacat	attccataac	tatttttagta	9960
ttaaaatttg	tcagtgatat	ttcttaccct	ctcctctagg	aaaatgtgcc	atgtttatcc	10020
cttggttttg	aatgcccctc	aggaacagac	actaagagtt	tgagaagcat	ggttacaagg	10080
gtgtggcttc	ccctgcggaa	actaagtaca	gactatttca	ctgtaaagca	gagaagttct	10140
tttgaaggag	aatctccagt	gaagaaagag	ttcttcactt	ttacttccat	ttcctcttgt	10200
gggtgacct	caatgctcct	tgtaaaactc	caatatttta	aacatggctg	ttttgccttt	10260
ctttgcttct	ttttagcatg	aatgagacag	atgatacttt	aaaaaagtaa	ttaaaaaaaa	10320

-continued

aaacttgtga	aaatacatgg	ccataataca	gaacccaata	caatgatctc	ctttacccaa	10380
ttgttatggt	tgtacttttg	tagatagctt	tccaattcag	agacagttat	tctgtgtaaa	10440
ggtctgactt	aacaagaaaa	gatttccctt	tacccaaaga	atcccagtcc	ttatttgctg	10500
gtcaataagc	aggggtcccca	ggaatggggt	aactttcagc	accctctaac	ccactagtta	10560
ttagtagact	aattaagtaa	acttatcgca	agttgaggaa	acttagaacc	aactaaaatt	10620
ctgcttttac	tgggattttg	ttttttcaaa	ccagaaacct	ttacttaagt	tgactactat	10680
taatgaattt	tgggtctctt	tttaagtgtt	cttcttaaaa	atgttatctt	actgctgaga	10740
agttcaagtt	tgggaagtac	aaggaggaat	agaaacttaa	gagattttct	tttagagcct	10800
cttctgtatt	tagccctgta	ggattttttt	tttttttttt	ttttttgggt	ttgttgagct	10860
tcagtgaggc	tattcattca	cttatactga	taatgtctga	gatactgtga	atgaaatact	10920
atgtatgctt	aaacctaaga	ggaaatatat	tcccaaaaatt	attcttcccg	aaaaggagga	10980
gttgcccttt	gattgagttc	ttgcaaactt	cacaacgact	ttattttgaa	caatactgtt	11040
tggggatgat	gcattagttt	gaaacaactt	cagttgtagc	tgtcatctga	taaaattgct	11100
tcacagggaa	ggaaatttaa	cacggatcta	gtcattattc	ttgttagatt	gaatgtgtga	11160
attgtaattg	taaacaggca	tgataattat	tacttttaaa	actaaaaaca	gtgaatagtt	11220
agttgtggag	gttactaaag	gatgggtttt	ttttaataaa	aactttcagc	attatgcaaa	11280
tgggcatatg	gcttaggata	aaacttccag	aagtagcatc	acatttaaat	tctcaagcaa	11340
cttaataata	tggggctctg	aaaaactggg	taaggttact	ccaaaaatgg	ccctgggtct	11400
gacaaagatt	ctaacttaaa	gatgcttatg	aagactttga	gtaaaatcat	ttcataaaat	11460
aagtgaggaa	aaacaactag	tattaaattc	atcttaataa	atgtatgatt	taaaaaatat	11520
gtttagctaa	aatgcatag	tcatttgaca	atttcattta	tatctcaaaa	aatttactta	11580
accaagttgg	tcacaaaact	gatgagactg	gtggtggtag	tgaataaatg	agggaccatc	11640
catatttgag	acactttaca	tttgtgatgt	gttatactga	attttcagtt	tgattctata	11700
gactacaaat	ttcaaaatta	caatttcaag	atgtaataag	tagtaatatc	ttgaaatagc	11760
tctaaagggg	atttttctgt	tttattgatt	cttaaaatat	atgtgctgat	tttgatttgc	11820
atttgggtag	attatacttt	tatgagtatg	gagggttaggt	attgattcaa	gttttcctta	11880
cctatttggt	aaggatttca	aagtcttttt	gtgcttggtt	ttcctcattt	ttaaatatga	11940
aatatattga	tgacctttta	caaatttttt	ttatctcaaa	ttttaagga	gatcttttct	12000
aaaagaggca	tgatgactta	atcattgcat	gtaacagtaa	acgataaacc	aatgattcca	12060
tactctctaa	agaataaaaag	tgagcttttag	ggccgggcat	ggtcagaaat	ttgacaccaa	12120
cctggccaac	atggcgaaac	cccgctctta	ctaaaaatac	aaaaatcagc	cgggcatggg	12180
ggcggcacct	atagtcccag	ctacttggga	ggatgagaca	ggagagtcac	ttgaacctgg	12240
gaggagaggt	tgcaagtgag	tgagatcacg	ccattgcact	ccagcctgag	caatgaaagc	12300
aaaactccat	ctcaaaaaaa	aaaaaagaaa	agaaagaata	aaagtgagct	ttggattgca	12360
tataaatcct	ttagacatgt	agtagacttg	tttgatactg	tgtttgaaca	aattacgaag	12420
tattttcatc	aaagaatgtt	attgtttgat	gttattttta	ttttttattg	cccagcttct	12480
ctcatattac	gtgattttct	tcacttcatg	tcactttatt	gtgcagggtc	agagtattat	12540
tccaatgctt	actggagaag	tgattcctgt	aatggaaactg	ctttcatcta	tgaaatcaca	12600
cagtgttcc	gaagaaatag	atgtaagttt	aatgagagc	aattatacac	tttatgagtt	12660
ttttgggggt	atagtattat	tatgtatatt	attaatatc	taattttaat	agtaaggact	12720

-continued

ttgtcataca	tactattcac	atacagtatt	agccacttta	gcaaataagc	acacacaaaa	12780
tcctggattt	tatggcaaaa	cagaggcatt	tttgatcagt	gatgacaaaa	ttaaattcat	12840
tttgtttatt	tcattacttt	tataattcct	aaaagtggga	ggatcccagc	tcttatagga	12900
gcaattaata	tttaatgtag	tgtcttttga	aacaaaactg	tgtgccaaag	tagtaaccat	12960
taatggaagt	ttacttgtag	tcacaaattt	agtttcctta	atcatttggt	gaggacgttt	13020
tgaatcacac	actatgagtg	ttaagagata	cctttaggaa	actattcttg	ttgttttctg	13080
attttgtcat	ttaggttagt	ctcctgattc	tgacagctca	gaagaggaag	ttgttcttgt	13140
aaaaattggt	taacctgctt	gaccagcttt	cacatttggt	cttctgaagt	ttatggtagt	13200
gcacagagat	tgttttttgg	ggagtcttga	ttctcggaaa	tgaaggcagt	gtgttatatt	13260
gaatccagac	ttccgaaaac	ttgtatatta	aaagtgttat	ttcaacacta	tggtacagcc	13320
agactaattt	ttttattttt	tgatgcattt	tagatagctg	atacagtact	caatgatgat	13380
gatattggtg	acagctgtca	tgaaggcttt	cttctcaagt	agaatttttt	cttttcataa	13440
aagctggatg	aagcagatac	catcttatgc	tcacctatga	caagatttgg	aagaaagaaa	13500
ataacagact	gtctacttag	attgttctag	ggacattacg	tatttgaact	gttgcttaaa	13560
tttgtgttat	ttttcactca	ttatatctct	atatatat	gggtgttattc	catttgctat	13620
ttaaagaaac	cgagtttcca	tcccagacaa	gaaatcatgg	ccccttgctt	gattctgggt	13680
tcttgtttta	cttctcatta	aagctaacag	aatcctttca	tattaagttg	tactgtagat	13740
gaacttaagt	tatttaggcg	tagaacaaaa	ttattcatat	ttatactgat	ctttttccat	13800
ccagcagtg	agtttagtac	ttaagagttt	gtgcccttaa	accagactcc	ctggattaat	13860
gctgtgtacc	cgtgggcaag	gtgcctgaat	tctctataca	cctatttcct	catctgtaaa	13920
atggcaataa	tagtaatagt	acctaagtgt	tagggttggt	ataagcattg	agtaagataa	13980
ataatataaa	gcacttagaa	cagtgcctgg	aacataaaaa	cacttaataa	tagctcatag	14040
ctaacatttc	ctatttacat	ttcttctaga	aatagccagt	atttggtgag	tgccacatg	14100
ttagttcctt	tactagttgc	tttacatgta	ttatcttata	ttctgtttta	aagtttcttc	14160
acagttacag	attttcatga	aattttactt	ttaataaaag	agaagtaaaa	gtataaagta	14220
ttcactttta	tgttcacagt	cttttccttt	aggctcatga	tggagtatca	gaggcatgag	14280
tgtgtttaac	ctaagagcct	taatggcttg	aatcagaagc	actttagtcc	tgtatctggt	14340
cagtgtcagc	ctttcataca	tcattttaaa	tcccatttga	ctttaagtaa	gtcacttaat	14400
ctctctacat	gtcaatttct	tcagctataa	aatgatggta	tttcaataaa	taaatacatt	14460
aattaaatga	tattatactg	actaattggg	ctgttttaag	gctcaataag	aaaatttctg	14520
tgaaaggctc	ctagaaaatg	taggttccta	tacaaaataa	agataacatt	gtgcttatag	14580
cttcgggtgt	tatcatataa	agctattctg	agttatttga	agagctcacc	tacttttttt	14640
tgttttttagt	ttgttaaatt	gttttatagg	caatgttttt	aatctgtttt	ctttaactta	14700
cagtgccatc	agctcacact	tgcaaacctg	tggtgttcc	gttgtagtag	gtagcagtgc	14760
agagaaagta	aataaggtag	tttattttat	aatctagcaa	atgatttgac	tctttaagac	14820
tgatgatata	tcattgattg	tcattttaa	ggtaggttgc	aattaaatg	atctagtagt	14880
ataaggaggc	aatgtaatct	catcaaattg	ctaagacacc	ttgtggcaac	agtgagtttg	14940
aaataaactg	agtaagaatc	atztatcagt	ttattttgat	agctcggaaa	taccagtgtc	15000
agtagtgtat	aatgggtttt	gagaatatat	taaaatcaga	tatataaaaa	aaattactct	15060

-continued

tctatttccc	aatgttatct	ttaacaaatc	tgaagatagt	catgtacttt	tggtagtagt	15120
tccaaagaaa	tgttatttgt	ttattcatct	tgatttcatt	gtcttcgctt	tccttctaaa	15180
tctgtccctt	ctagggagct	attgggatta	agtggtcatt	gattattata	ctttattcag	15240
taatgtttct	gaccctttcc	ttcagtgcta	cttgagttaa	ttaaggatta	atgaacagtt	15300
acatttccaa	gcattagcta	ataaactaaa	ggattttgca	cttttcttca	ctgaccatta	15360
gtagaaaga	gttcagagat	aagtatgtgt	atctttcaat	ttcagcaaac	ctaatttttt	15420
aaaaaaagtt	ttacatagga	aatatgttgg	aaatgatact	ttacaaagat	attcataatt	15480
tttttttgta	atcagctact	ttgtatat	acatgagcct	taatttatat	ttctcatata	15540
accatttatg	agagcttagt	atacctgtgt	catttatattg	catctacgaa	ctagtgcact	15600
tattccttct	gttacctcaa	acaggtggct	ttccatctgt	gatctccaaa	gccttagggt	15660
gcacagagtg	actgccgagc	tgctttatga	aggagagaaag	gctccatagt	tgtagtgttt	15720
tttttttttt	ttttaaacat	ttttcccatc	ctccatcctc	ttgagggaga	atagcttacc	15780
ttttatcttg	ttttaatttg	agaaagaagt	tgccaccact	ctaggttgaa	aaccactcct	15840
ttaacataat	aactgtggat	atggtttgaa	tttcaagata	gttacatgcc	tttttatatt	15900
tcctaataka	gctgtaggtc	aaatattatt	agaatcagat	ttctaaatcc	cacccaatga	15960
cctgcttatt	ttaaatcaaa	ttcaataatt	aattctcttc	tttttgagg	atctggacat	16020
tctttgatat	ttcttacaac	gaatttcatg	tgtagacca	ctaaacagaa	gctataaaag	16080
ttgcatggtc	aaataagtct	gagaaagtct	gcagatgata	taattcacct	gaagagtcac	16140
agtatgtagc	caaatgttaa	aggttttgag	atgccatata	gtaaatttac	caagcatttt	16200
ctaaatttat	ttgaccacag	aatccctatt	ttaagcaaca	actgttacat	cccatggatt	16260
ccaggtgact	aaagaatact	tatttcttag	gatatgtttt	attgataata	acaattaaaa	16320
tttcagatat	ctttcataag	caaatcagtg	gtctttttac	ttcatgtttt	aatgctaaaa	16380
tattttcttt	tatagatagt	cagaacatta	tgcttttttc	tgactccagc	agagagaaaa	16440
tgctccaggt	tatgtgaagc	agaatcatca	tttaaatatg	agtcagggtc	ctttgtacaa	16500
ggcctgctaa	aggtatagtt	tctagtattc	acaagtga	ccacttttct	aaaatcattt	16560
ttgagactct	ttatagacaa	atcttaaata	ttagcattta	atgtatctca	tattgacatg	16620
cccagagact	gacttccttt	acacagttct	gcacatagac	tatatgtctt	atggatttat	16680
agttagtatc	atcagtga	caccatagaa	taccctttgt	gttccaggtg	ggccctgtt	16740
cctacatgtc	tagcctcagg	actttttttt	ttttaacaca	tgcttaaata	aggttgcaca	16800
tcaaaaataa	gatcatttct	ttttaactaa	atagatttga	attttattga	aaaaaaattt	16860
taaacatctt	taagaagctt	ataggattta	agcaattcct	atgtatgtgt	actaaaatat	16920
atatatttct	atatataata	tatattagaa	aaaaattgta	tttttctttt	atttgagtct	16980
actgtcaagg	agcaaaacag	agaaatgtaa	attagcaatt	atttataata	cttaaaggga	17040
agaaaagttg	tcaccttggt	gaatctatta	ttgttatctc	aattatagtc	ccaagacgtg	17100
aagaaatagc	tttctaatg	gttatgtgat	tgtctcatag	tgactacttt	cttgaggatg	17160
tagccacggc	aaaatgaaat	aaaaaaattt	aaaaattggt	gcaaatataa	gttatattag	17220
gcttttggtg	atcttcaata	atgtgctgct	atgaactcag	aatgatagta	tttaaatata	17280
gaaactagtt	aaaggaaacg	tagtttctat	ttgagttata	catatctgta	aattagaact	17340
tctcctgtta	aaggcataat	aaagtgccta	atacttttgt	ttcctcagca	ccctctcatt	17400
taattatata	atcttagttc	tgaaagggac	ctataaccaga	tgcttagagg	aaatttcaaa	17460

-continued

actatgatct	aatgaaaaaa	tatttaaatag	ttctccatgc	aaatacaaat	catatagttt	17520
tccagaaaaat	acctttgaca	ttatacaaaag	atgattatca	cagcattata	atagtaaaaa	17580
aatggaaata	gcctctttct	tctgttctgt	tcatagcaca	gtgcctcata	cgcagtaggt	17640
tattattaca	tggtaactgg	ctaccccaac	tgattaggaa	agaagtaa	at	17700
aaaatacata	ctcattgagg	tgcatagaat	aattaagaaa	ttaaaagaca	cttgtaattt	17760
tgaatccagt	gaatacccac	tgttaatat	tggtatatct	ctttctagtc	tttttttccc	17820
ttttgcatgt	attttcttta	agactccac	ccccactgga	tcatctctgc	atgttcta	17880
ctgctttttt	cacagcagat	tctaagcctc	tttgaatatc	aacacaaact	tcaacaactt	17940
catctataga	tgccaaataa	taaattcatt	tttatctact	taaccacttc	ctttggatgc	18000
ttaggtcatt	ctgatgtttt	gctattgaaa	ccaatgctat	actgaacact	tctgtcacta	18060
aaactttgca	cacactcatg	aatagcttct	taggataaat	tttttagagat	ggatttgcta	18120
aatcagagac	cattttttta	aattaaaaaa	caattattca	tatcgtttgg	catgtaagac	18180
agtaaatttt	ccttttattt	tgacaggatt	caactggaag	ctttgtgctg	cctttccggc	18240
aagtcatgta	tgctccatat	cccaccacac	acatagatgt	ggatgtcaat	actgtgaagc	18300
agatgccacc	ctgtcatgaa	catatttata	atcagcgtag	atacatgaga	tccgagctga	18360
cagccttctg	gagagccact	tcagaagaag	acatggctca	ggatacgatc	atctacactg	18420
acgaaagctt	tactcctgat	ttgtacgtaa	tgctctgcct	gctggtactg	tagtcaagca	18480
atatgaaatt	gtgtctttta	cgaataaaaa	caaaacagaa	gttgcattta	aaaagaaaga	18540
aatattacca	gcagaattat	gcttgaagaa	acatttaatc	aagcattttt	ttcttaa	18600
ttcttctttt	tccatacaat	tgtgtttacc	ctaaaatagg	taagattaac	ccttaaagta	18660
aatatttaac	tatttgttta	ataaatatat	attgagctcc	taggcactgt	tctaggtacc	18720
gggcttaata	gtggccaacc	agacagcccc	agccccagcc	cctacattgt	gtatagtcta	18780
ttatgtaaca	gttattgaat	ggacttatta	acaaaaccaa	agaagtaatt	ctaagtcttt	18840
tttttcttga	catatgaata	taaaatacag	caaaactggt	aaaatatatt	aatggaacat	18900
ttttttactt	tgcattttat	attgttat	acttcttatt	tttttttaaa	aaaaaaagcc	18960
tgaacagtaa	attcaaaagg	aaaagtaatg	ataattaatt	gttgagcatg	gacccaactt	19020
gaaaaaaaaa	atgatgatga	taaatctata	atcctaaaac	cctaagtaaa	cacttaaaag	19080
atgttctgaa	atcaggaaaa	gaattatagt	atacttttgt	gtttctcttt	tatcagttga	19140
aaaaaggcac	agtagctcat	gcctgtaaga	acagagcttt	gggagtgcaa	ggcaggcgga	19200
tcacttgagg	ccaggagttc	cagaccagcc	tgggcaacat	agtgaaaccc	catctctaca	19260
aaaaataaaa	aagaattatt	ggaatgtgtt	tctgtgtgcc	tgtaatccta	gctattccga	19320
aagctgaggc	aggaggatct	tttgagccca	ggagtttgag	gttacaggga	gttatgatgt	19380
gccagtgtac	tccagcctgg	ggaacaccga	gactctgtct	tatttaaaaa	aaaaaaaaaa	19440
aaaatgcttg	caataatgcc	tggcacatag	aaggtaacag	taagtgttaa	ctgtaataac	19500
ccaggtctaa	gtgtgtaagg	caatagaaaa	attggggcaa	ataagcctga	cctatgtatc	19560
tacagaatca	gtttgagctt	aggtaacaga	cctgtggagc	accagtaatt	acacagtaag	19620
tgtaaaccaa	aagcatagaa	taggaatatc	ttgttcaagg	gacccccagc	cttatacatc	19680
tcaagggtgca	gaaagatgac	ttaatatagg	acccattttt	tcctagttct	ccagagtttt	19740
tattggttct	tgagaaagta	gtaggggaat	gttttagaaa	atgaattggt	ccaactgaaa	19800

-continued

ttacatgtca	gtaagttttt	atatattggt	aaatttttagt	agacatgtag	aagttttcta	19860
attaatctgt	gccttgaaac	atcttctttt	ttcctaaagt	gcttagtatt	ttttccgttt	19920
tttgattggt	tacttgggag	cttttttgag	gaaatttagt	gaactgcaga	atgggtttgc	19980
aaccatttgg	tatttttggt	ttgtttttta	gaggatgtat	gtgtatttta	acattttctta	20040
atcattttta	gccagctatg	tttgttttgc	tgatttgaca	aactacagtt	agacagctat	20100
tctcattttg	ctgatcatga	caaaataata	tctgaattt	ttaaattttg	catccagctc	20160
taaattttct	aaacataaaa	ttgtccaaaa	aatagtattt	tcagccacta	gatttgtgtg	20220
taagtctatt	gtcacagagt	cattttactt	ttaagtatat	gtttttacat	gttaattatg	20280
tttgttattt	ttaattttaa	ctttttaaaa	taattccagt	cactgccaat	acatgaaaaa	20340
ttggtcactg	gaattttttt	tttgactttt	attttagggt	catgtgtaca	tgtgcagggtg	20400
tgttatacag	gtaaattgcg	tgtcatgagg	gttggtgta	cagggtgattt	cattaccag	20460
gtaataagca	tagtacccaa	taggtagttt	tttgatcttc	acccttctcc	cacctcaag	20520
taggcoctgg	tgttgctgtt	tccttctttg	tgccatgta	tactcagtgt	ttagctccca	20580
cttagaagtg	agaacatgcg	gtagttgggt	ttctgttctt	ggattagtgc	acttaggata	20640
atgacctcta	gctccatctg	gtttttatgg	ctgcatagta	ttccatgggtg	tatatgtatc	20700
acattttctt	tatccagtct	accattgata	ggcatttagg	ttgattccct	gtctttgtta	20760
tcatgaatag	tgctgtgatg	aacatacaca	tgcatgtgtc	tttatggtag	aaaaatttgt	20820
attccttttag	gtacatatag	aataatgggg	ttgctagggt	gaatggtagt	tctattttca	20880
gttatttgag	aaatcttcaa	actgcttttc	ataatagcta	aactaattta	cagtcccgcc	20940
agcagtgtat	aagtgttccc	ttttctccac	aaccttgcca	acatctgtga	ttttttgact	21000
ttttaataat	agccattcct	agagaattga	tttgcaattc	tctattagtg	atattaagca	21060
ttttttcata	tgcttttttag	ctgtctgtat	atattcttct	gaaaaatttt	catgtccttt	21120
gccagtttg	tagtgggggtg	ggttggtttt	tgcttgtaa	ttagttttta	gttccttcca	21180
gattctgcat	atccctttgt	tggatacatg	gtttgcagat	atctttctcc	cattgtgtag	21240
gttgctcttt	actctgttga	tagtttcttt	tgccatgcag	gagctcgta	ggteccattt	21300
gtgtttgttt	ttgttgaggt	tgcttttggc	gtcttcatca	taaaatctgt	gccagggcct	21360
atgtccagaa	tggtatttcc	taggttgtct	tccagggttt	ttacaatttt	agattttacg	21420
tttatgtctt	taatccatct	tgagttgatt	tttgatatag	gcacaaggaa	ggggtccagt	21480
ttcactccaa	ttcctatggc	tagcaattat	cccagcacca	tttattgaat	acggagtcct	21540
ttccccattg	cttggttttt	gtcaactttg	ttgaagatca	gatggttgta	agtgtgtggc	21600
tttatttctt	ggctctctat	tctccattgg	tctatgtgtc	tgtttttata	acagtaccct	21660
gctgttcagg	ttcctatagc	cttttagtat	aaaatcggt	aatgtgatgc	ctccagcttt	21720
gttctttttg	cttaggattg	ctttggctat	ttgggctcct	ttttgggtcc	atattaattt	21780
taaaacagtt	ttttctgggt	ttgtgaagga	tatcattgggt	agtttatagg	aatagcattg	21840
aatctgtaga	ttgctttggg	cagtatggcc	attttaacaa	tattaattct	tcctatctat	21900
gaatatggaa	tgtttttcca	tgtgtttgtg	tcatctcttt	atacctgatg	tataaagaaa	21960
agctgggtatt	attcctactc	aatctgttcc	aaaaaattga	ggaggaggaa	ctcttcccta	22020
atgaggccag	catcattctg	ataccaaaac	ctggcagaga	cacaacagaa	aaaagaaaac	22080
ttcaggccaa	tatccttgat	gaatatagat	gcaaaaatcc	tcaacaaaat	actagcaaac	22140
caaatccagc	agcacatcaa	aaagctgatc	tactttgatc	aagtaggctt	tatccctggg	22200

-continued

atgcaagggt	ggttcaacat	acacaaatca	ataagtgtga	ttcatcacat	aaacagagct	22260
aaaaacaaaa	accacaagat	tatctcaata	ggtagagaaa	aggttgtcaa	taaaatttaa	22320
catcctccat	gttaaaaacc	ttcagtaggt	caggtgtagt	gactcacacc	tgtaatccca	22380
gcactttggg	aggccaaggc	gggcataatc	cttaagccca	ggagttcaag	acgagcctag	22440
gcagcatggt	gaaaccccat	ctctacaaaa	aaaaaaaaaa	aaaaaatta	gcttggtatg	22500
gtgacatgca	cctatagtcc	cagctattca	ggaggttgag	gtgggaggat	tgtttgagcc	22560
cgggaggcag	aggttggcag	cgagctgaga	tcatgccacc	gcactccagc	ctgggcaacg	22620
gagtgagacc	ctgtctcaaa	aaagaaaaat	cacaaacaat	cctaaacaaa	ctaggcattg	22680
aaggaacatg	cctcaaaaaa	ataagaacca	tctatgacag	acccatagcc	aatatcttac	22740
caaatgggca	aaagctggaa	gtattctcct	tgagaaccgt	aacaagacaa	ggatgtccac	22800
tctcaccact	ccttttcagc	atagttctgg	aagtccctagc	cagagcaatc	aggaaagaga	22860
aagaaagaaa	gacattcaga	taggaagaga	agaagtcaaa	ctatttctgt	ttgcaggcag	22920
tataattctg	tacctagaaa	atctcatagt	ctctgccag	aaactcctaa	atctgttaaa	22980
aatttcagca	aagttttggc	attctctata	ctccaacacc	ttccaaagtg	agagcaaaat	23040
caagaacaca	gtccattca	caatagccgc	aaaacgaata	aaatacctag	gaatccagct	23100
aaccagggag	gtgaaagatc	tctatgagaa	ttacaaaaca	ctgctgaaag	aatcagaga	23160
tgacacaaac	aatggaaat	gttctttttt	aacaccttgc	tttatcta	tcacttatga	23220
tgaagatact	cattcagtgg	aacaggtata	ataagtccac	tcgattaaat	ataagcctta	23280
ttctctttcc	agagcccaag	aaggggcact	atcagtcccc	agtcaataat	gacgaaatgc	23340
taatattttt	cccctttacg	gtttctttct	tctgtagtgt	ggtacactcg	tttcttaaga	23400
taaggaaact	tgaactacct	tcctgtttgc	ttctacacat	accattctc	tttttttgcc	23460
actctggtca	ggtataggat	gatccctacc	actttcagtt	aaaaactcct	cctcttacta	23520
aatgttctct	taccctctgg	cctgagtaga	acctagggaa	aatggaagag	aaaaagatga	23580
aagggagggtg	gggcctggga	agggaataag	tagtccctgt	tgtttgtgtg	tttgctttag	23640
cacctgctat	atcctaggtg	ctgtgttagg	cacacattat	tttaagtggc	cattatatta	23700
ctactactca	ctctggtcgt	tgccaaggta	ggtagtactt	tcttgtagat	ttggttcatg	23760
ttacttacag	atggtgggct	tggtgaggca	aaccagtggt	ataatcatcg	gagtgtgttc	23820
tctaactctca	ctcaaatttt	tcttcacatt	ttttggtttg	ttttggtttt	tgatggtagt	23880
ggcttatttt	tggtgctggg	ttgttttttg	tttttttttg	agatggcaag	aattggtagt	23940
tttattttatt	aattgcctaa	gggtctctac	ttttttttaa	agatgagagt	agtaaaatag	24000
attgatagat	acatacatac	ccttactggg	gactgcttat	attctttaga	gaaaaaatta	24060
catattagcc	tgacaaacac	cagtaaaatg	taaatatatc	cttgagtaaa	taaatgaatg	24120
tatatattgt	gtctccaaat	atatatatct	atattcttac	aaatgtgttt	atatgtaata	24180
tcaatttata	agaacttaaa	atgttggtc	aagtgaggga	ttgtggaagg	tagcattata	24240
tggccatttc	aacatttgaa	cttttttctt	ttcttcattt	tcttcttttc	ttcaggaata	24300
tttttcaaga	tgtcttacac	agagacactc	tagtgaaagc	cttcctggat	caggtaaattg	24360
ttgaacttga	gattgtcaga	gtgaatgata	tgacatgttt	tcttttttaa	tatatcctac	24420
aatgcctggt	ctatatattt	atattcccct	ggatcatgcc	ccagagttct	gtcagcaat	24480
tgcagttaag	ttagttacac	tacagttctc	agaagagtct	gtgagggcac	gtcaagtgca	24540

-continued

tcattacatt	ggttgcctct	tgtcctagat	ttatgcttcg	ggaattcaga	cctttgttta	24600
caatataata	aatattattg	ctatctttta	aagatataat	aataagatat	aaagttgacc	24660
acaactactg	ttttttgaaa	catagaattc	ctgggtttaca	tgtatcaaag	tgaaatctga	24720
cttagcctttt	acagatataa	tatatacata	tatatatcct	gcaatgcttg	tactatatat	24780
gtagtacaag	tatatatata	tgtttgtgtg	tgtatatata	tatagtacga	gcatatatac	24840
atattaccag	cattgtagga	tatatatatg	tttatatatt	aaaaaaaaagt	tataaactta	24900
aaaccctatt	atgttatgta	gagtatatgt	tatatatgat	atgtaaaata	tataacatat	24960
actctatgat	agagtgtaat	atattttttta	tatatatttt	aacattttata	aatgataga	25020
attaagaatt	gagtccta	ctgttttatt	aggtgctttt	tgtagtgtct	ggctctttcta	25080
aagtgtctaa	atgatttttc	cttttgactt	attaatgggg	aagagcctgt	atattaacaa	25140
ttaagagtgc	agcattccat	acgtcaaaca	acaaacattt	taattcaagc	attaacctat	25200
aacaagtaag	tttttttttt	ttttttgaga	aaggagggtt	gtttatttgc	ctgaaatgac	25260
tcaaaaatat	ttttgaaaca	tagtgtactt	atttaaataa	catctttatt	gtttcattct	25320
tttaaaaaat	atctacttaa	ttacacagtt	gaaggaaatc	gtagattata	tggaacttat	25380
ttcttaatat	attacagttt	gttataataa	cattctgggg	atcaggccag	gaaactgtgt	25440
catagataaa	gctttgaaat	aatgagatcc	ttatgtttac	tagaaatttt	ggattgagat	25500
ctatgaggtc	tgtgacatat	tgogaagttc	aaggaaaatt	cgtaggcctg	gaatttcatg	25560
cttctcaagc	tgacataaaa	tccctccac	tctccacctc	atcatatgca	cacattctac	25620
tctacccac	ccactccacc	ccttgcaaaa	gtacagggtat	atgaatgtct	caaaaccata	25680
ggctcatctt	ctaggagctt	caatgttatt	tgaagatttg	ggcagaaaaa	attaagtaat	25740
acgaaataac	ttatgtatga	gttttaaaag	tgaagtaaac	atggatgtat	tctgaagtag	25800
aatgcaaaat	ttgaatgcat	ttttaaagat	aaattagaaa	acttctaaaa	actgtcagat	25860
tgtctgggcc	tggtggctta	tgctgtaat	cccagcactt	tgggagtcctg	aggtgggtgg	25920
atcacaaggt	caggagatcg	agaccatcct	gccaacatgg	tgaaaccccg	tctctactaa	25980
gtatacaaaa	attagctggg	cgtggcagcg	tgtgcctgta	atcccagcta	cctgggaggc	26040
tgaggcagga	gaatcgcttg	aaccaggag	gtgtagggtg	cagtgagtca	agatcgcgcc	26100
actgcacttt	agcctgggtga	cagagctaga	ctccgtctca	aaaaaaaaaa	aaaatatcag	26160
attgttccta	cacctagtgc	ttctatacca	cactcctggt	agggggcatc	agtggaaatg	26220
gttaaggaga	tgtttagtgt	gtattgtctg	ccaagcactg	tcaacactgt	catagaaact	26280
tctgtacgag	tagaatgtga	gcaaattatg	tgttgaaatg	gttcctctcc	ctgcaggtct	26340
ttcagctgaa	acctggctta	tctctcagaa	gtactttcct	tgcacagttt	ctacttgtcc	26400
ttcacagaaa	agccttgaca	ctaataaaat	atatagaaga	cgatacgtga	gtaaaactcc	26460
tacacggaag	aaaaaccttt	gtacattgtt	tttttgtttt	gtttcctttg	tacattttct	26520
atatcataat	ttttgcgctt	cttttttttt	tttttttttt	tttttttcca	ttatttttag	26580
gcagaaggga	aaaaagccct	ttaaatctct	tcggaacctg	aagatagacc	ttgatttaac	26640
agcagagggc	gatcttaaca	taataatggc	tctggctgag	aaaattaaac	caggcctaca	26700
ctcttttato	tttggaagac	ctttctacac	tagtgtgcaa	gaacgagatg	ttctaataac	26760
tttttaaatg	tgtaacttaa	taagcctatt	ccatcacaat	catgatcgct	ggtaaagtag	26820
ctcagtgggtg	tggggaaacg	ttcccctgga	tcatactcca	gaattctgct	ctcagcaatt	26880
gcagttaagt	aagttacact	acagttctca	caagagcctg	tgaggggatg	tcaggtgcat	26940

-continued

cattacattg	ggtgtctctt	ttcctagatt	tatgcttttg	ggatacagac	ctatgtttac	27000
aatataataa	atattattgc	tatcttttaa	agatataata	ataggatgta	aacttgacca	27060
caactactgt	ttttttgaaa	tacatgattc	atggttttaca	tgtgtcaagg	tgaaatctga	27120
gttggttttt	acagatagtt	gactttctat	cttttgcat	tctttggtgt	gtagaattac	27180
tgtaaacttt	ctgcaatcaa	ctgaaaacta	gagcctttta	atgatttcaa	ttccacagaa	27240
agaaagtgag	cttgaacata	ggatgagctt	tagaaagaaa	attgatcaag	cagatgttta	27300
attggaattg	attattagat	cctactttgt	ggatttagtc	cctgggattc	agtctgtaga	27360
aatgtctaat	agttctctat	agtccttggt	cctggtgaac	cacagttagg	gtgttttggt	27420
tattttattg	ttcttgctat	tgttgatatt	ctatgtagtt	gagctctgta	aaaggaaatt	27480
gtattttatg	ttttagtaat	tgttgccaac	tttttaaatt	aattttcatt	atttttgagc	27540
caaattgaaa	tgtgcacctc	ctgtgccttt	tttctcctta	gaaaatctaa	ttacttgga	27600
caagttcaga	tttactgggt	cagtcatttt	catcttggtt	tcttcttgct	aagtcttacc	27660
atgtacctgc	tttggaatc	attgcaactc	tgagattata	aaatgcctta	gagaatatac	27720
taactaataa	gatctttttt	tcagaaacag	aaaatagttc	cttgagtact	tccttcttgc	27780
atttctgcct	atgtttttga	agttggtgct	gttgccctgc	aataggctat	aaggaatagc	27840
aggagaaatt	ttactgaagt	gctgttttcc	taggtgctac	tttggcagag	ctaagttatc	27900
ttttgttttc	ttaatgcgtt	tggaccatth	tgctggctat	aaaataactg	attaatataa	27960
ttctaacaca	atgttgacat	tgtagttaca	caaacacaaa	taaatatttt	atttaaaatt	28020
ctggaagtaa	tataaaaggg	aaaatatatt	tataagaaag	ggataaaggt	aatagagccc	28080
ttctgcccc	caccaccaa	atttacacaa	caaatgaca	tgttcgaatg	tgaaagggtca	28140
taatagcttt	cccatcatga	atcagaaaga	tgtggacagc	ttgatgtttt	agacaaccac	28200
tgaactagat	gactgttgta	ctgtagctca	gtcatttaaa	aaatatataa	atactacctt	28260
gtagtgtccc	atactgtgtt	ttttacatgg	tagattctta	tttaagtgtc	aactgggttat	28320
tttctttggc	tggtttattg	tactgttata	cagaatgtaa	gttgtacagt	gaaataagtt	28380
attaaagcat	gtgtaaacad	tgttatatat	cttttctcct	aaatggagaa	ttttgaataa	28440
aatatatattg	aaattttgcc	tctttcagtt	gttcattcag	aaaaaaatac	tatgatattt	28500
gaagactgat	cagcttctgt	tcagctgaca	gtcatgctgg	atctaaactt	tttttaaaat	28560
taattttgtc	ttttcaaaga	aaaaatatth	aaagaagctt	tataatataa	tcttatgtta	28620
aaaaaaacttt	ctgcttaact	ctctggattt	cattttgatt	tttcaaatta	tatattaata	28680
tttcaaagt	aaaatactat	ttagataaat	tgtttttaaa	cattcttatt	attataatat	28740
taatataacc	taaactgaag	ttattcatcc	caggatatcta	atacatgtat	ccaaagtaaa	28800
aatccaagga	atctgaacac	tttcatctgc	aaagctagga	ataggtttga	cattttcact	28860
ccaagaaaaa	gttttttttt	gaaaatagaa	tagttgggat	gagaggtttc	tttaaaagaa	28920
gactaactga	tcacattact	atgattctca	aagaagaaac	caaaacttca	tataatacta	28980
taaagtaaat	ataaaatagt	tccttctata	gtatatttct	ataatgctac	agtttaaaaca	29040
gatcactctt	atataatact	attttgattt	tgatgtagaa	ttgcacaaat	tgatatttct	29100
cctatgatct	gcagggtata	gcttaaagta	acaaaaacag	tcaaccacct	ccatttaaca	29160
cacagtaaca	ctatgggact	agttttatta	cttccatttt	acaaatgagg	aaactaaagc	29220
ttaaagatgt	gtaatacacc	gccaaggtc	acacagctgg	taaaggtgga	tttcatccca	29280

-continued

gacagttaca gtcattgcc	atgggcacagc tcctaactta	gtaactccat gtaactggta	29340
ctcagtgtag ctgaattgaa	aggagagtaa ggaagcagg	tttacaggtc tacttgca	29400
attcagagcc cgagtgtgaa	tccctgctgt gctgcttga	gaagttactt aacctatgc	29460
aggttcattt tgtaaatatt	ggaaatggag tgataatac	g tacttcacca gaggattta	29520
tgagacotta tacgatcct	agttcagtac ctgactagt	g cttcataaat gctttttcat	29580
ccaatctgac aatctccag	c ttgtaattgg ggcatttag	a acatttaata tgattattg	29640
catggtaggt taaagctgt	c atcttgctgt tttctattt	g ttctttttgt tttctccta	29700
cttttggaatt tttttattc	t actatgtctt ttctattgt	c ttattaacta tactctttg	29760
tttatttttag tggttgttt	t agggttatac ctctttcta	a tttaccagtt tataaccag	29820
ttatatacta cttgacatat	a gcttaagaa acttactgt	t gttgtctttt tgctgttat	29880
gtcttaacgt tttttattc	t acaaacatta taaactcca	c actttattgt tttttaatt	29940
tacttataca gtcaattat	c ttttaaagat atttaaata	a aaacattcaa aacacccca	30000
t			30001

<210> SEQ ID NO 3
<211> LENGTH: 1031
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 3

attcccggga tacgtaacct	acgggtgtccc gctaggaaag	agaggtgcgt caaacagcga	60
caagttccgc ccacgtaaaa	gatgacgctt ggtgtgtcag	ccgtccctgc tgcccgggtg	120
cttctctttt gggggcgggg	tctagcaaga gcaggtgtgg	gtttaggaga tatctccgga	180
gcatttggat aatgtgacag	ttggaatgca gtgatgtcga	ctctttgccc accgccatct	240
ccagctgttg ccaagacaga	gattgcttta agtggcaa	aat cacctttatt agcagctact	300
tttgcttact gggacaatat	tcttggtcct agagtaagg	c acatttgggc tccaaagaca	360
gaacaggta	c ttctcagtga tggagaaata	acttttcttg ccaaccacac	tctaaatgga 420
gaaatccttc gaaatgcaga	gagtggtgct atagatgtaa	agttttttgt cttgtctgaa	480
aaggggagtga ttattgttt	c attaatcttt gatggaaact	ggaatgggga tcgcagcaca	540
tatggactat caattatact	tccacagaca gaacttagt	t tctacctccc acttcataga	600
gtgtgtgttg atagattaac	acatataatc cggaaaggaa	gaatatggat gcataaggaa	660
agacaagaaa aatgtccaga	agattatctt agaaggcaca	gagagaatgg aagatcaggg	720
tcagagtatt attccaatgc	ttactggaga agtgattcct	gtaatggaaa ctgcttccct	780
ctatgaaatt ccccgggtt	cctggaggaa atagatatag	gctgatacag ttacceaatg	840
atggatgaat attgggggac	cgcctggtca ttgaaaggct	ttcttttctc caggaaagaa	900
atttttttcc ttttccataa	aaagcttggg aatggaagac	aacaattccc attctttttt	960
tgcgttccac ccctatgtga	caacagaaat ttttggggaa	acaacaacga aaaaatttta	1020
tcccgcgcgc a			1031

<210> SEQ ID NO 4
<211> LENGTH: 3244
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 4

gggcggggct gcggttgcg	gtgctgcgcc cgcggcggcg	gaggcgcagg cggtggcgag	60
----------------------	-----------------------	-----------------------	----

-continued

tggatatctc	cggagcattt	ggataatgtg	acagttggaa	tgcagtgatg	tcgactcttt	120
gcccaccgcc	atctccagct	gttgccaaga	cagagattgc	tttaagtggc	aaatcacctt	180
tattagcagc	tacttttgct	tactgggaca	atattcttgg	tcctagagta	aggcacattt	240
gggctccaaa	gacagaacag	gtactttctc	gtgatggaga	aataactttt	cttgccaacc	300
acactctaaa	tggagaaatc	cttcgaaatg	cagagagtgg	tgctatagat	gtaaagtttt	360
ttgtcttgtc	tgaaaaggga	gtgattattg	tttcattaat	ctttgatgga	aactggaatg	420
gggatcgcag	cacatatgga	ctatcaatta	tacttccaca	gacagaactt	agtttctacc	480
tcccacttca	tagagtgtgt	gttgatagat	taacacatat	aatccgaaa	ggaagaatat	540
ggatgcataa	ggaaagacaa	gaaaatgtcc	agaagattat	cttagaaggc	acagagagaa	600
tggaagatca	gggtcagagt	attattccaa	tgcttactgg	agaagtgatt	cctgtaatgg	660
aactgctttc	atctatgaaa	tcacacagtg	ttcctgaaga	aatagatata	gctgatacag	720
tactcaatga	tgatgatatt	ggtgacagct	gtcatgaagg	ctttcttctc	aatgccatca	780
gctcacactt	gcaaacctgt	ggctgttccg	ttgtagtagg	tagcagtgca	gagaaagtaa	840
ataagatagt	cagaacatta	tgcttttttc	tgactccagc	agagagaaaa	tgctccaggt	900
tatgtgaagc	agaatcatca	tttaaatatg	agtcagggct	ctttgtacaa	ggcctgctaa	960
aggattcaac	tggaagcttt	gtgctgcctt	tccggcaagt	catgtatgct	ccatatccca	1020
ccacacacat	agatgtggat	gtcaatactg	tgaagcagat	gccaccctgt	catgaacata	1080
tttataatca	gcgtagatac	atgagatccg	agctgacagc	cttctggaga	gccacttcag	1140
aagaagacat	ggctcaggat	acgatcatct	acactgacga	aagctttact	cctgatttga	1200
atatttttca	agatgtctta	cacagagaca	ctctagtga	agccttcctg	gatcaggtct	1260
ttcagctgaa	acctggctta	tctctcagaa	gtactttcct	tgcacagttt	ctacttgtcc	1320
ttcacagaaa	agccttgaca	ctaataaaat	atatagaaga	cgatacgcag	aagggaaaaa	1380
agccctttaa	atctcttcgg	aacctgaaga	tagacctga	tttaacagca	gagggcgatc	1440
ttaacataat	aatggctctg	gctgagaaaa	ttaaaccagg	cctacactct	tttatctttg	1500
gaagaccttt	ctacactagt	gtgcaagaac	gagatgttct	aatgactttt	taaatgtgta	1560
acttaataag	cctattccat	cacaatcatg	atcgctggta	aagtagctca	gtgggtgtggg	1620
gaaacgttcc	cctggatcat	actccagaat	tctgctctca	gcaattgcag	ttaagtaagt	1680
tacactacag	ttctcacaag	agcctgtgag	gggatgtcag	gtgcatcatt	acattgggtg	1740
tctcttttcc	tagatttatg	cttttgggat	acagacctat	gtttacaata	taataaatat	1800
tattgctatc	ttttaagat	ataataatag	gatgtaaact	tgaccacaac	tactgttttt	1860
ttgaaataca	tgattcatgg	tttacetgtg	tcaaggtgaa	atctgagttg	gcttttacag	1920
atagttgact	ttctatcttt	tggcattctt	tggtgtgtag	aattactgta	atacttctgc	1980
aatcaactga	aaactagagc	ctttaaatga	tttcaattcc	acagaaagaa	agtgagcttg	2040
aacataggat	gagctttaga	aagaaaattg	atcaagcaga	tgtttaattg	gaattgatta	2100
ttagatccta	ctttgtggat	ttagtccttg	ggattcagtc	tgtagaaatg	tctaatagtt	2160
ctctatagtc	cttgttcctg	gtgaaccaca	gttaggggtg	tttgtttatt	ttattgttct	2220
tgctattggt	gatattctat	gtagttgagc	tctgtaaaag	gaaattgtat	tttatgtttt	2280
agtaattggt	gccaactttt	taaattaatt	ttcattattt	ttgagccaaa	ttgaaatgtg	2340
cacctcctgt	gccttttttc	tccttagaaa	atctaattac	ttggaacaag	ttcagatttc	2400

-continued

actggtcagt cattttcatc ttgttttctt cttgctaagt cttaccatgt acctgctttg	2460
gcaatcattg caactctgag attataaaaat gccttagaga atataactaac taataagatc	2520
tttttttcag aaacagaaaa tagttccttg agtacttctt tcttgcatth ctgcctatgt	2580
ttttgaagtt gttgctgttt gcctgcaata ggctataagg aatagcagga gaaatthtac	2640
tgaagtgtg ttttctagg tgctactttg gcagagctaa gttatcttht gttthcttaa	2700
tgcgtttgga ccattttgct ggctataaaa taactgatta atataattct aacacaatgt	2760
tgacattgta gttacacaaa cacaaataaa tatthttatt aaaattctgg aagtaatata	2820
aaagggaaaa tatatthata agaaagggat aaaggtaata gagcccttct gccccccacc	2880
caccaaattt acacaacaaa atgacatgtt cgaatgtgaa aggtcataat agctthccca	2940
tcatgaatca gaaagatgtg gacagcttga tgtthtagac aaccactgaa ctagatgact	3000
gttgactgt agctcagtca thtaaaaaat atataaatac taccttgtag tgtccatac	3060
tgtgtthttt acatggtaga thctthatta agtgctaact ggthatttht thtggtgtgt	3120
thattgtact gttatacaga atgtaagttg tacagtgaat taagthatta aagcatgtgt	3180
aaacattgtt atatatctth tctctaaat ggagaattth gaataaaaata thttgaaat	3240
thtg	3244

<210> SEQ ID NO 5
<211> LENGTH: 761
<212> TYPE: DNA
<213> ORGANISM: homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (693)..(693)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (722)..(722)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 5	
cacgaggctt tgatatttct tacaacgaat ttcattgtga gaccactaa acagaagcta	60
taaaagttgc atggtcaaat aagtctgaga aagtctgcag atgatataat tcacctgaag	120
agtcacagta tntagccaaa tgttaaaggt thtgagatgc catacagtaa atthaccaag	180
cattthctaa atthatttga ccacagaatc cctattthta gcaacaactg thacatccca	240
tggattccag gtgactaaag aatacttatt tcttaggata tgtthtattg ataataacaa	300
thaaaatttc agatatctth cataagcaaa tcagtgggtt thttacttca tgtthtaatg	360
ctaaaatatt thctthtata gatagtcaga acattatgcc thtttctgac tccagcagag	420
agaaaatgct ccaggttatg tgaagcagaa tcatcattta aatatgagtc agggctctth	480
gtacaaggcc tgctaaagga thcaactgga agctthgtgc tgctthtccg gcaagtcatg	540
tatgtccat atcccaccac acacatagat gtggatgtca atactgtgaa gcagatgcca	600
ccctgtcatg aacatattth taatcagcgt agatacatga gatccgagct gacagcttc	660
tggagagcca thtcagaaga agacatggct cangatacga tcattctacac tgacgaaagc	720
thtactctg atthgaatat thttcaagat gtcttacaca g	761

<210> SEQ ID NO 6
<211> LENGTH: 1901
<212> TYPE: DNA
<213> ORGANISM: homo sapiens

<400> SEQUENCE: 6

-continued

acgtaaccta	cgggtgtccc	ctaggaaaga	gaggtgcgtc	aaacagcgac	aagttccgcc	60
cacgtaaaag	atgacgcttg	atatctccgg	agcattttgga	taatgtgaca	gttggaatgc	120
agtgatgtcg	actctttgcc	caccgccatc	tccagctggt	gccaagacag	agattgcttt	180
aagtggcaaa	tcacctttat	tagcagctac	ttttgcttac	tgggacaata	ttcttggtcc	240
tagagtaagg	cacatttggg	ctccaaagac	agaacaggta	cttctcagtg	atggagaaat	300
aacttttctt	gccaaccaca	ctctaaatgg	agaaatcctt	cgaaatgcag	agagtgggtgc	360
tatagatgta	aagttttttg	tcttgtctga	aaagggagtg	attattgttt	cattaatctt	420
tgatggaaac	tggaatgggg	atcgcagcac	atatggacta	tcaattatac	ttccacagac	480
agaacttagt	ttctacctcc	cacttcatag	agtgtgtggt	gatagattaa	cacatataat	540
ccggaaagga	agaatatgga	tgcataagga	aagacaagaa	aatgtccaga	agattatctt	600
agaaggcaca	gagagaatgg	aagatcaggg	tcagagtatt	attccaatgc	ttactggaga	660
agtgattcct	gtaatggaac	tgctttcatc	tatgaaatca	cacagtgttc	ctgaagaaat	720
agatatagct	gatacagtac	tcaatgatga	tgatattggg	gacagctgtc	atgaaggctt	780
tcttctcaag	taagaatttt	tcttttcata	aaagctggat	gaagcagata	ccatcttatg	840
ctcacctatg	acaagatttg	gaagaaagaa	aataacagac	tgtctactta	gattgttcta	900
gggacattac	gtatttgaac	tgttgcttaa	atttgtgtta	tttttcactc	attatatctt	960
tatatatatt	tggtgttatt	ccatttgcta	tttaaagaaa	ccgagtttcc	atcccagaca	1020
agaaatcatg	gccccttgct	tgattctggg	ttcttgtttt	acttctcatt	aaagctaaca	1080
gaatcctttc	atattaagtt	gtactgtaga	tgaacttaag	ttatttaggc	gtagaacaaa	1140
attattcata	tttatactga	tctttttcca	tccagcagtg	gagtttagta	cttaagagtt	1200
tgtgccctta	aaccagactc	cctggattaa	tgctgtgtac	ccgtgggcaa	ggtgccctgaa	1260
ttctctatac	acctatttcc	tcatctgtaa	aatggcaata	atagtaatag	tacctaattgt	1320
gtagggttgt	tataagcatt	gagtaagata	aataatataa	agcacttaga	acagtgcctg	1380
gaacataaaa	acacttaata	atagctcata	gctaacatct	cctatttaca	tttcttctag	1440
aaatagccag	tatttggtga	gtgcctacat	gttagttcct	ttactagttg	ctttacatgt	1500
attatcttat	attctgtttt	aaagtttctt	cacagttaca	gattttcatg	aaatcttact	1560
tttaataaaa	gagaagtaaa	agtataaagt	attcactttt	atgttcacag	tcttttcctt	1620
taggctcatg	atggagtatc	agaggcatga	gtgtgtttta	cctaagagcc	ttaatggctt	1680
gaatcagaag	cacttttagt	ctgtatctgt	tcagtgtcag	cctttcatac	atcattttta	1740
atcccatttg	actttaagta	agtcacttaa	tctctctaca	tgtcaatttc	ttcagctata	1800
aaatgatggg	atttcaataa	ataaatacat	taattaaatg	atattatact	gactaattgg	1860
gctgttttaa	ggcaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	a		1901
<210> SEQ ID NO 7						
<211> LENGTH: 562						
<212> TYPE: DNA						
<213> ORGANISM: homo sapiens						
<220> FEATURE:						
<221> NAME/KEY: misc_feature						
<222> LOCATION: (166)..(166)						
<223> OTHER INFORMATION: n is a, c, g, or t						
<400> SEQUENCE: 7						
agacgtaacc	tacgggtgtcc	cgctaggaaa	gagagatatc	tccggagcat	ttggataatg	60

-continued

tgacagttgg aatgcagtga tgtcgactct ttgcccaccg ccatctccag ctgttgccaa	120
gacagagatt gctttaagtg gcaaatacacc tttattagca gctacntttt gcttactggg	180
acaatattct tggtcctaga gtaaggcaca tttgggctcc aaagacagaa caggtacttc	240
tcagtgatgg agaaataact tttcttgcca accacactct aaatggagaa atccttcgaa	300
atgcagagag tggtgctata gatgtaaagt tttttgtctt gtctgaaaag ggagtgatta	360
ttgtttcatt aatctttgat ggaaactgga atggggatcg cagcacatat ggactatcaa	420
ttatacttcc acagacagaa cttagtttct acctccact tcatagagtg tgtgttgata	480
gattaacaca tataatccgg aaaggaagaa tatggatgca taaggaaaga caagaaaatg	540
tccagaagat tatcttagaa gg	562
<210> SEQ ID NO 8	
<211> LENGTH: 798	
<212> TYPE: DNA	
<213> ORGANISM: homo sapiens	
<400> SEQUENCE: 8	
gggctctctt ttgggggcgg ggtctagcaa gacagatat ctccggagca tttggataat	60
gtgacagttg gaatgcagtg atgtcgactc tttgcccacc gccatctcca gctgttgcca	120
agacagagat tgctttaagt ggcaaatac ctttattagc agctactttt gcttactggg	180
acaatattct tggtcctaga gtaaggcaca tttgggctcc aaagacagaa caggtacttc	240
tcagtgatgg agaaataact tttcttgcca accacactct aaatggagaa atccttcgaa	300
atgcagagag tggtgctata gatgtaaagt tttttgtctt gtctgaaaag ggagtgatta	360
ttgtttcatt aatctttgat ggaaactgga atggggatcg cagcacatat ggactatcaa	420
ttatacttcc acagacagaa cttagtttct acctccact tcatagagtg tgtgttgata	480
gattaacaca tataatccgg aaaggaagaa tatggatgca taaggaaaga caagaaaatg	540
tccagaagat tatcttagaa ggcacagaga gaatggaaga tcagggtcag agtattattc	600
caatgcttac tggagaagtg attcctgtaa tgggactgct ttcactatg aaatcacaca	660
gtgttctctga agaaatagat atagctgata cagtactcca tgatgatgat atttggtgac	720
agctgtcatg aaaggctttc ttctcaagta ggaatttttt cttttcataa aagctgggat	780
gaagccagat tcccatct	798
<210> SEQ ID NO 9	
<211> LENGTH: 169	
<212> TYPE: DNA	
<213> ORGANISM: homo sapiens	
<400> SEQUENCE: 9	
aaacagcgac aagttccgcc cacgtaaaag atgatgcttg gtgtgtcagc cgteccctgct	60
gcccggttgc ttctcttttg ggggcggggg ctagcaagag cagatatctc cggagcattt	120
ggataatgtg acagttggaa tgcggtgatg tcgactcttt gccaccgc	169
<210> SEQ ID NO 10	
<211> LENGTH: 176	
<212> TYPE: DNA	
<213> ORGANISM: homo sapiens	
<400> SEQUENCE: 10	
aaaacgtcat cgcacataga aaacagacag acgtaacctc cgggtgtccc ctaggaaaga	60
gaggtgcgtc aaacagcgac aagttccgcc cacgtaaaag atgacgcttg atatctccgg	120

-continued

agcatttgga taatgtgaca gttggaatgc agtgatgtcg actctttgcc caccgc	176
<210> SEQ ID NO 11 <211> LENGTH: 30001 <212> TYPE: DNA <213> ORGANISM: macaca mulatta <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (26113)..(26155) <223> OTHER INFORMATION: n is a, c, g, or t <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (28797)..(29186) <223> OTHER INFORMATION: n is a, c, g, or t <400> SEQUENCE: 11	
aatctctaag caattttttg ggggaagaaag aattgcaatt agggcatacg tgtagatcag	60
atgggtcttcg gtatatccaa cgacaaagaa aagggtgggag gtttcgttaa aaaagagaaa	120
tgttacatag tactttttaga gaaaattcac tggcactatt aagggtctga ggagctggta	180
agtttcaatt ggtgagtgat ggtggtagat aaaattagag ctgcagcagg tcatttcagc	240
aactatcaga taaaactggc ctcaggtcac aacgggcagt ttcagcagct agacttgaaa	300
gaattacact gcgggagcaa tgtcatttgt cctgcattgct tttctacccc ctacccccac	360
tttttttagtt ggggtataaca agaacgaccc aaattgtatg atcaactttc acaaagcata	420
gaacagtagg aaaagggctc gtttctgcag aagatgtaga cgttgagagc cattttatgt	480
atttatttct ccctttcttc atcgggtgaat gattaaaatg ttctgtatga tttttagtga	540
tgagaaaggt taaacgccac tcatctgtag taagtgtaat ctacacactt gcagacccaa	600
aggcataagg tttaaaaaac ctttggtttt ttacacatca aacagagtgg tataaatgct	660
actcatctgt agtaagtga atctatacac ctgcagacca acgacgcaag gtttcaaaaa	720
tctttgtgtt ttttacacat caaacagaat ggtacatttt tcaaaagttt aaaaaaaaaa	780
aaaatccaca tatcacaact agcaaaaatg acattcccca gtgtgaaaat catgcttgag	840
agaattctta catgtaaagg caaaattgca gtgactttac aagggaacctg gggattcccc	900
cccacagtgt ggagctgtcc cctaccaggg tttgcggcgg agttttgaat gtacttaaca	960
gtgtctcacg gtaaaaaaca aacttcatcc accaaatatt tgttgagcgc ccaactgcctg	1020
ccaagcaca acaaaaccat tcaaaaccac gaaatcgtct gcactttctc cggatccagc	1080
agcctctgcg attaaggttt gcacacgcta ttgcgccaac gtcctccag agcgcgctct	1140
aagataaaag aatgggacaa gttgcccctc cccctttcac gggcctcgtg cgtcaacgtc	1200
atcgcatata gaaaacacac agacgtaacc tacggtgtcc cgctaggaaa gagaggcgcg	1260
tcaaacagcg acaagttccg cccacgtaaa agatgacgct tgggtgcgtca gccgtccctg	1320
ctgcccgggt ccttctctct gggggcgggg cctggctaga gcaggtgtgg gtttaggagg	1380
tgtgtgtttt tgtttttct accctctccc ctctacttgc tctcacagta ctcgctgagg	1440
gtgaacaaga aaagacctga taaagattaa ccagaagaaa acaaagaggg aaacaactgc	1500
agcctgtagc gggctctgga gcttaagaga ggcgcgctag gcgccggggc gtgggcgtgg	1560
tcggggcggg gtcggggccag gggcggggct gcggttgcgg tccctgcgcc cgcggcgggc	1620
gcggcgggcg cagcggaggc gcaggcgggt gcgagtgggt gagtgaagag gcggcgctct	1680
ggcgggtgtc tgtttgccgt ccggttgccg ggaagagacg cgggtagcag ccggggctct	1740
cctcagagct cgacacattt ttactttccc tctcgtttct ctgaccgaag tcgggtgtcc	1800

-continued

ggctttcgcc	tctagcgact	ggtggaattg	cctgcatctg	ggccccgggc	ttcgcggcgg	1860
cgcagggacg	agggatggga	atctggcctc	ttctcgctt	ccccgccgc	agtgcgctgc	1920
cccagctgtc	tccttcccgg	ggacctgtg	ggagcgctgc	cgctacagac	tcgagagaaa	1980
ggagcctcgg	gcactgagag	gcctcgcccc	ggggaaggcc	ggagggcggg	cggcgggcgg	2040
cgagcggctc	ctgcggacca	agtctgggtt	ctctgggaac	ccgagacggt	ccctgatggc	2100
gaggagatca	tgcggggtgc	tatgggggtg	tggagacgtc	tgcagaattt	tagcccaagc	2160
ttctaaggag	tgctgatgac	ttgcatatga	gggcagcaat	gccagtcggt	gtactcccta	2220
ttctgtggga	catgatgtgg	ttgcttcaca	gctccgagat	gacacagact	tgcttaaagg	2280
aagtgaccat	tgtgacttgg	gcataccttg	actgatggta	atcagttgca	gacagaagtg	2340
cacagattac	atgtctgtgt	ccacactgga	tcagtctggc	cacgaggaac	accacaggct	2400
ttgtattgag	aaacaggagg	gaggtcctgc	actttcccag	gaggggtggc	cctttcagat	2460
gcaatcgaga	ttgttaggct	ctggtagagt	ggttgccctg	ttgtggcagt	tggcaaattc	2520
ctattcaaac	tggtgccgtg	cgtcaccagt	taacaacaag	ggtacacgat	ctgtctggca	2580
ttactttctac	tttgtacaaa	ggatcaaaaa	tactgttaga	tatgattttt	ctcagacttt	2640
gggaaacttt	taacgtaatc	tgtgaatatc	acagaagcaa	gactgtcata	tagaggatat	2700
taataacctg	gagtcagaat	acttgaaata	tgggtgtcatt	tgacacgggc	tctgttatca	2760
ccacctttgc	caagcccttt	cacttgagga	aaacctcaa	tcagttggaa	actgcctcat	2820
gctgacagta	catctgaaac	aaaaacgaga	gtagttacca	cattccagat	tgttcactaa	2880
ggcagcattt	atctgctcca	ggaaaacatt	acaagcaact	tatgaagttg	ataaaatatt	2940
ttgtttggct	atgttggtac	tccaaaagtt	gctttcagag	aaacaaagta	aaccaaggag	3000
gactttctgtt	gttcacgtct	gcccttgggc	tctattctac	gttaattagg	tagttcccag	3060
gaggactaga	ttagcctacc	tattgtctga	gaaacttgga	tctgtgagaa	atggccagat	3120
agtgatacga	acttcacctc	ccagtctttc	ctgatgttta	agattgagaa	agtgttgtga	3180
actttctggg	gctgtaagca	gttcactgtc	cttaaagtgg	tcctgggcag	ctcctgttgt	3240
ggaaaagtga	ccgatttagg	attctgcttg	gctttggact	gggagaaaaat	aaactgcatg	3300
gttacaagta	ttgagagcca	agttggagaa	ggtggcttac	acctataatg	ccagagcctt	3360
aggaggcagg	ggcaagagga	tcactggaag	tcaggagttc	aagcccaacc	tgggcagcct	3420
agaccctgtc	tctacaaaaa	attaaaaact	tagccgggcg	cggtggtgtg	cacctgtagt	3480
cctagctact	tgggaggctg	aggcaggagg	gtcttttgag	cccaggagtt	tgaagttaca	3540
gggagctatg	atcctgccag	tgactccag	cctggatggc	aaaacgagac	cctgtctcta	3600
aaaaacaaga	agtgagggtc	ttatgatcgt	agaaattttg	cttacaatag	cagtggacca	3660
accacctttc	taaataccaa	tcagggaaga	catagttgat	ttttaacaaa	catttaaaga	3720
aaaagcaaaa	cctcaaactt	agcactctac	taacagtttt	agccgatgct	aattaaggta	3780
atcatgtctg	catatatggg	attactttca	gaaagtgtat	tgggaaacct	ctcatgaacc	3840
ctgtgcaacc	ctgagcaagc	caccgtctca	ctcagtttga	atcttggcct	ccctcaaaag	3900
actctgtggc	taatgtttgg	taactctctg	gagtagccag	cactgcatgt	acataggata	3960
ggtacataaa	acaattattg	gttttgagct	gatttttttc	agctgcattt	gcgtgtatgg	4020
atttttctca	ccaaagacaa	tgacttcaag	tgtaataaaa	ataattgtac	agctctccta	4080
attatacttc	tctgtaacat	ttcattttct	agactatttc	ttttggtagg	atttaaaact	4140
aaacaattca	gtatgatctt	tgttcttcat	tttctttctt	attctttttt	ttttcgagac	4200

-continued

agagtctccc	tctgttgtgc	cactctcagcc	cattgcaacc	tccgccacct	gggttcaagt	4260
gattctcctg	cctcagcctc	ctgagtagct	gggattacag	gtgcccgcga	ccacacctac	4320
ctaatttttt	gtatttttag	tagaggcggg	gtttcaccat	gttggttagg	ctggtcttaa	4380
actcctgacc	tcagatgac	cacctgcctc	ggcctcccaa	agagctggga	tgataggcgt	4440
gaccacccat	gcccgcacca	ttttttttct	tattctgtta	ggagtgagag	tgtaactagc	4500
agtctataat	agttcaattt	tcacaacgtg	gtaaaaattt	tccctgtaat	tcaacgagat	4560
tttgcttcag	ggctcagttc	tgtttttagga	aatactttta	ttttcagttt	gatgatgaaa	4620
tattagagtt	gtgatattgc	ctttatgatt	acctaccttt	ttaacctaaa	agaatgaaag	4680
aaaaatatgt	ttacagtata	attgtatggg	tgcgtgttaa	cttaattcat	tatgttggcc	4740
tccagtttgc	tggtgttcgt	tatgacagca	gtagtgtcat	taccatttca	attcagatta	4800
cattcctgta	tttgatcatt	gtaaactgat	tgcttaaatt	gtattaaaaa	cagtggatat	4860
tttaaacaag	ctgtactgct	tatatccagt	gctgtctcct	aagactatta	aattgatata	4920
acatatTTaa	aagtaaatat	ttcctaaatg	aatttttgaa	attaaaaata	cacgtgttaa	4980
aactgtcttt	gtgttcaacc	atttctgtac	gtacttagag	ttaactgttt	tgccaggctc	5040
tgtatgccta	ctcataatgt	gataaaagca	ctcatcta	gctctataaa	tagaagtcag	5100
tgctttccat	cagactgaac	actcttggca	agatgtggat	aaaattatTT	aagtaaaatt	5160
gtttactttg	tcatacattt	acagatcaaa	tgtagctcc	caaagcaatc	atatggcaaa	5220
gataggcata	tcataatttg	cctattagct	gctttgtatt	gctattatga	tagatttcac	5280
agtttttagat	ctgcttagat	gaaaatgtaa	ttctttttac	tgtcagtcct	agatataagt	5340
cttcaattat	agtacagtca	cacattgctt	aggaatgcat	cattaggcga	ttttgtcatt	5400
atgcaaacat	catagagtat	acttacataa	acatatatag	tacagccttt	acgtacgtag	5460
gcatatgggt	atagtctatt	gctcctaggc	tacaaatctg	tacagctgtt	actgtactga	5520
atactataga	cagttgtaac	acagtgggat	ttatttatct	aaatatatcc	aaacatagaa	5580
aagggtacagt	taaagtatgg	tataaaaaat	aatgatatac	ctatataggc	cacttacgt	5640
gaatggagct	tgcaggacta	gaagttgctc	tgggtgagtc	agtaagtaag	tggtgaatga	5700
atgtgaaggc	ctagaacatt	actgtacaca	ctgtagactt	tataaacaca	gtatgcttaa	5760
gctacaccaa	atttatcttt	acagtttttc	ttcaataaaa	aattaatgtg	aacctactat	5820
aactttttta	ctttgtaaac	tttttaattt	tttaactttt	aaaatactta	gcttgaaaca	5880
caaacacgca	tagctataca	aaaatatTTT	ttcttttatat	ccttatttcta	gaagctTTTT	5940
cctatttttta	actTTTTTTT	ttttacttgt	tagtcgtttt	tgTTaaaaac	taaaacacac	6000
acactttcac	ctaagcatag	acaggattag	gatcatcagt	ttcactccct	tccacctcac	6060
tgcttccac	ctccacatct	tgtccactg	gaacgtTTTT	agggggaata	acacacatgt	6120
agctgtcacc	tgctatgata	acagtgcttt	ctgttgaata	cctcctgaag	gacctgcctg	6180
aggctgtttt	acatttaact	taaaaaaaaa	aataagtaga	aggagtacac	tctaaaataa	6240
caataaaagg	tatagtctag	tgaatacata	aaccagcaac	atagtagttt	attatcaagt	6300
gttgataact	gtaataattg	tatgtgctat	actttaaatg	acttgaaaaa	ttgtactaag	6360
accttatgat	ggttacagtg	tcactaaggc	gatagcatat	tttcaggctc	attgtaatct	6420
aatgggacca	ccatcatata	tgcagtccac	cattgactga	aatgttacat	ggtacgtaac	6480
tgtatttgca	agaatgattt	gttttacatt	aatatcacat	aggatgtacc	tttttagagt	6540

-continued

gatatgttta	tgtggattaa	gatgtacaag	tggagcaagg	ggacaagagc	ccttggttct	6600
gtcttgga	tgagctttta	tgtctctctc	atcatgtctg	ttttcttatt	aaattcaaag	6660
gcttgga	gccctattta	gcccttggtt	tctatgtgtt	ctaaataact	aaagctttta	6720
aattctagcc	atttagtgga	gaactctctt	tgcaatggta	aaatgctgta	ttggtttctt	6780
gactagcata	ttaaataat	ttatctttgt	cttgatattt	caatgtcatt	ttaaacaatc	6840
ggattgggct	ttagtattct	cataccaga	gagttcactg	aggatacagg	actgtttgcc	6900
cattttttgt	tatggctcca	gacttggtt	atttcgatgt	cttttttttt	tttttttttt	6960
tttaaccttt	tagcagcttt	aaagtatttc	tgttggttagg	tgttggtatta	cttttctaag	7020
attactgtaa	caaagcacca	caaactgagt	ggctttaaac	aacagcaatt	tattctctca	7080
caattctaga	agctagaagt	ccgaaatgga	agtgttgatg	gggcatgatc	ctcaaaagag	7140
agaagactct	ttccttgctt	cttcctggct	tctgggtggt	accagcaatc	ctgagcggtc	7200
ctttcttgct	tcgtagtctt	agcagtcctg	tatctgcctt	ttgtcttcac	atggatgtct	7260
acctcttgct	tctgtgtctc	cagatctctc	tccttataaa	cacagaagtt	actggattag	7320
gccccactct	aatccagtat	gaccccatct	taacacgatt	acacctattt	ctaaataagg	7380
tcacattcac	atataccaag	ggtaggaat	tgagcatatc	ttttgcaggg	acacaattca	7440
accacaagt	gtcagctctc	agctgagcct	ttcccttctt	ggttttctcc	tttttagttg	7500
ctgtgggtta	ggggccaaat	ctccagtcct	actagacttg	cacatggact	ggagatttgg	7560
gaatactgct	ggctctattc	atgagcttta	gtatgtaaca	tttaatatca	gtgtaaagaa	7620
gccatttttt	cagttcacta	tttctttgaa	tttcttaatg	tatgccctga	atataagtaa	7680
caagttacta	tgtctcataa	aatgatcata	tcaacaaaca	tttaatgtgc	acctactgtg	7740
ctagttgaat	gtctttatcc	tgataggaga	taacaggcct	ccgcatcttt	gacttaagag	7800
gacaaaccaa	gtatgtctga	atcatttggg	gttttgatgg	atatctttaa	attgctgaac	7860
ctaactcattg	gttttatatg	tcattgttta	gatatctcag	gagcatttgg	ataatgtgac	7920
agttggaatg	cagtgtgtgc	gactctttgc	ccaccgccat	ctccagctgt	tgccaagaca	7980
gagattgctt	taagtgggtg	atcaccttta	ttagcagcta	cttttgctta	ctgggacaat	8040
attcttggtc	ctagagtaag	gcacatttgg	gctccaaaga	cagaacaggt	acttctcagt	8100
gatggagaaa	taacttttct	tgccaaccac	actctaaatg	gagaaatcct	tcgaaatgca	8160
gagagtgggtg	ctatagatgt	aaagtttttt	gtcttgtctg	aaaaggagg	gattattgtt	8220
tcattaatct	ttgatggaaa	ctggaatggg	gatgcagca	catacggact	atcaattata	8280
ctccacaga	cagaacttag	tttctacctc	ccacttcata	gagtgtgtgt	tgatagatta	8340
acacatataa	tccggaaagg	aagaatatgg	atgcataagg	taagtgattt	ttcagcttat	8400
taatcatgtt	aacctatctt	ttgaaagctt	attttctgat	acataataat	cttattttta	8460
aatttatatg	agtgaacatc	aaacaataga	tattatttat	tttgcattta	tcctgttaga	8520
tacaaataca	tctgggtctga	tgctgtcat	cttcatatta	actgtggaag	gtaggaaatg	8580
gtagctccac	attacagatg	aaaagctaaa	gcttaaaca	atgcagaaac	ttttagatcc	8640
tggattcttc	ttgggagcct	ttgactctaa	taccttttgt	ttccctttca	ttgcacaatc	8700
ctgtctttcg	cttactacta	tgtgtaagta	taacagttca	aaaaaatagt	ttcataagct	8760
gttggttatg	tagcctttgg	tctctttaac	ctctttgcca	agttcccagg	ttcataaaat	8820
gaggagggtg	aaccgcatgg	ttcccaagag	aattcctttt	aattttacag	aaattattgt	8880
ttccccgaa	gtcctatagt	tcaatatata	atgatattta	catttcagta	tagttttggc	8940

-continued

atatctaaag	aacacattaa	gttctccttc	ctgtgttcca	gtttgatact	aacctggaag	9000
tccattaagc	attaccaatt	ttaaaaggct	tttgcccaat	agtaaggaaa	aataatatct	9060
tttaaaagaa	taatttttta	ctatgtttgc	aggcttactt	ccttttttct	cacattatga	9120
aactcttaaa	atcaggagaa	tcttttaaac	atcataatgt	ttaatttgaa	aagtgcaagt	9180
cattcttttc	ctttttgaaa	ctatgcagat	gttacattga	ctattttctg	tgaagttatc	9240
ttttttttcc	ctgcagaata	aagggtgttt	tgattttatt	ttgtgttgtt	tataagaaca	9300
tacattcggt	gggttaattt	cctgcccctg	cccccgtttt	ttccctaaag	tagaaagtat	9360
ttttcttgtg	aactaaatta	ctacacaaga	acatgtctat	tgaaaaataa	gtatcaaaat	9420
gttgtggggt	gtttttttta	ataaattctt	tcttgctcag	gaaagacaag	aaaatgtcca	9480
gaagattatc	ttagaaggca	cagagagaat	ggaagatcag	gtatatgcag	attgcatact	9540
gtcaaatatt	attctcatgg	catgtatctg	tgtaaagttg	atggctacat	ttgtgaaggc	9600
cttggggaca	tacagagtaa	gccttaatgg	agcttttatg	gaggtgtaca	gaataaacta	9660
gaggaagatt	tccatatctt	agacctgaag	agttaaataa	gtaaacaaag	gaaaatagta	9720
attgcatcta	caaattaata	tttgctccct	ttttttttct	gtttgaacag	aataaatttt	9780
ggataacttg	ttactagtaa	aaaatttaaa	aattgtctgt	gatatgttct	ttaagggtact	9840
acttctcgaa	ctttttccta	gaagtagctg	taacaggagg	agagcatatg	tacccttaag	9900
gtatctgggg	tataggccca	tgtccaaaca	atattttctt	taagtcttgt	gttgatatct	9960
taagactcat	gcaatttaca	ttttattcca	tgatataact	attttaatat	taaaatttgt	10020
cagtgatatt	tcttaccctc	tcctctagga	aaatgtgcc	tgtttatact	ttggctttga	10080
gtgcccctga	ggaacagaca	ctagagtttg	agaagcatgg	ttacacaggc	gtggcttccc	10140
ctgcagaaat	taagtacaga	ctatttcagt	gtaaagcaga	gaagttcttt	tgaaggggaa	10200
tctccagtga	agaaagggtt	cttcactttt	acttccatct	cctcttgagg	gtgacctca	10260
ttgctccttg	taaaactccg	atatttttaa	catggctgtt	ttgctttcct	ctggttcttt	10320
ttaacatgag	tgagacagat	gatactttta	aaagtaattt	taaaaaaaaa	tgtaaaaata	10380
tatggccata	atgcagaacc	ctatgctgtg	atctccttta	ccaaattgtt	gtgtttgtac	10440
ttttgtagat	agctttccag	tccagagaca	gttattctgt	gtaaagggtc	gactcaacaa	10500
gaaaagattt	ccctttacc	aaagaatgcc	agtctttatt	tgctgggtcaa	taagcagggt	10560
ccccaggaaa	ggggtaactt	tcaccaccct	ctaaccact	ggttattagt	aaactaatta	10620
agtagactta	tctcaagatg	aggaaactta	aaaccaagta	aaattctgct	tttactggga	10680
ttttattttt	tgaaaccaga	aacgtttact	taagttgact	actattaatg	aattttgggtc	10740
tctcttttaa	gtactcttct	taaaaatgtt	atcctactgc	tgagaagttc	aagtttgaga	10800
agtacaagga	ggaatagaaa	cttgagagat	tttctttttc	ttttagagcc	tcttctgtat	10860
ttagccctgt	aggaattttt	tttttcccc	aagattcttc	ttcgtgaaaa	ggaggagtgt	10920
ccttttgatt	gagttcttgc	aaatctcaca	acgactttat	tttgaacaat	actgtttggg	10980
gatgatgcat	gagtctgaaa	caacttcagt	tgtagctgtc	atctgataaa	attgcttcac	11040
aggaagggaa	atttagcacg	gatctagtca	ttattcttgt	tagattgaat	gtgttaatca	11100
taattgtaaa	caggcatgat	aattattact	ttaaaaactg	aaaacagtga	atagttagtt	11160
gtggagggtta	ctaaagcatg	atttttttta	aataaaaact	tcagcatttt	gcaaatatgc	11220
atatggttta	ggatagaact	tccagaggta	gcatcacatt	taaattctca	agcaacttag	11280

-continued

taatacgagg	ctctgaaaaa	ctggttaaag	ttactccaga	aatggccctg	ggtctgacag	11340
acatttctaac	ttaaagatgc	atatgaagac	tttgaataaa	atcatttcat	atgaagacat	11400
tgaataaaat	catttcataa	aataagtgag	gaaaaacaac	tactattgaa	ttcatcttaa	11460
tgtatgattt	taaaaatatg	tttagctaaa	aattcataga	catttgacaa	tttcgtttat	11520
atctcaaaaa	gttgacttac	ccaagttgat	cacaaaactg	atgagactgg	tggtggtagt	11580
gaataaatga	gggaccaccc	atatttgaga	cactttacat	ttgtgatgtg	ttatactgaa	11640
ttttcagttt	gattctataa	actaccaatt	tcaaaattac	aatttcaagg	tgtaataagt	11700
agtggtatta	tcttgaaata	ggtctaaagg	gaacttttct	gttttaaaat	attcttaaac	11760
tatatgtgct	gattttgatt	tgcatttggg	tagattatac	tcttatgaat	cggggggctg	11820
ggtattgatt	caggttttcc	ttacctat	ggtaaggatt	tcaaagtctt	tttgtgcttg	11880
attttctctg	tttttaata	tgaaacatat	tgatgacttt	taattaacaa	atgtttttat	11940
ctcgaataaa	ttttaaagga	gatcttttct	aaaagaggta	tgatgactta	attattgcat	12000
ataacaataa	atgagaaacc	agtgattcca	tactctctaa	agaataaaaag	tgagcttttag	12060
gccagggcat	ggtggctcat	gcctgtaatc	ccagcacttt	ggaaggccga	ggcaggcgga	12120
tcacctgagg	tcaggaattc	gacaccagcc	tggccaaatg	gcaaaaccct	gtctctacta	12180
caaatacaaa	aattagctgg	gcatgggtggc	agccctata	gtcccagcta	cttgaagac	12240
tgagacagga	gagtcactcg	aacccgagag	gcagagggtg	cagtaagctg	aatcacacc	12300
attgcactcc	agcctgggca	acaagagcaa	aactccgtct	aaaaaaaaa	aaaaaaaaa	12360
aaaaagaata	aaagtgagct	ttggattgcg	tataaatcct	ttagacaagt	agtagacttg	12420
tttgatactg	tgtttgaaca	aattacaaag	tattttcact	aaagaatgtt	attgtttgct	12480
gttattttta	ttttttattg	cccagcttct	ctcatattca	ttatgtgatt	ttcttcactt	12540
catgttactt	tattgtgcag	ggtcagagta	ttattccaat	gcttactgga	gaagtgattc	12600
ctgtaatgga	actgctttca	tctatgaaat	cacacagtgt	tcctgaagaa	atagatgtaa	12660
gtttttatat	ttttaaatga	gagcaattat	accctttatc	agtttttttg	ggttatatta	12720
ttattatgta	tattattaat	attctaattt	taatactaag	cacttcgctg	tacgtactat	12780
ccacatgcag	tattagccac	ttgaacagat	aagcacacac	aaaatcctgg	attttatggc	12840
ataacagagg	catttttgat	cagtgatgac	aaaactaaat	ttattttgtt	tatttcacta	12900
cttttataat	tcctaaaagt	gggaggatcc	cagctcttat	aggagcaatt	aatatttaat	12960
gcagtacctt	ttgaaacaaa	actgtgtgcc	aaagcagtaa	ccattaatgg	aagttgactt	13020
atagtcacaa	atttagtttc	cttaatcatt	tggtgaggat	gttttgaatc	acacactatg	13080
agtggttaaga	gatatcttta	ggacactatt	cttggtgttt	tattgtcatt	taggttagtc	13140
tcctgtctga	cagctcagaa	gaggaagttg	ttcttgtaaa	aattgtttac	acaacctgat	13200
tgaccagctt	tcacatttgt	tcttctgaaa	gctgatggta	gtgcacagat	tgttttatgg	13260
ggagtcttga	ttctcagaaa	tgaaggcagt	gtgttatatt	gaatccagac	ttcagaaaac	13320
ttgtatatta	aaagtgtttt	ttcaacacta	tgttatagcc	agactaattt	ttttattttt	13380
ttgatgcatt	ttagatagct	gatacagtac	tcaatgatga	tgatattggg	gacagttgtc	13440
atgaaggctt	tcttctcaag	taagaatttt	tcttttcata	aaacctggat	gaagcatatg	13500
ttcacctatg	acaagatttg	gaaggaagaa	aataacagac	tgtctactta	gattgttcta	13560
gggacaacat	tgcatatttg	aattgttgct	taaatttggt	ttatttttca	ttcgttatat	13620
ttctataata	tatttgatgt	tattccattt	gctatttaaa	gaaactgagt	ttccatattt	13680

-continued

```

cccagacaag aaatcatggc cccttgcttg attctgggttt cttgtttttac ttctcattaa 13740
agctaaaaga accctttcaa attaagttgt actgtagatg aacttaagtt atttaggcct 13800
agaaaaaaaa aattcatatt tatactgata tttttccatc cagcagtga gtttagtact 13860
taagagtttg tgccttaaa ccagactccc tgggttaatg ctgtgtacct gtgggcaagg 13920
tcctgaatt ctctatacac ctatttcctc atctgtaaaa tggcaataat aataatagta 13980
cctaattgat agagttgtta taagcattga gtaagataaa taatataaag cacttagaac 14040
agtgcctgga acataagaac acttaataat agctaacatt ttctatttac atttcttcta 14100
aggaaaagg taaacagaaat agccaatatt tggtcagtgc ctacatgtta gtccctatac 14160
taagtgcctt acatgtatta tcttatattc tattttaatg tttcttcaca gttgcagatt 14220
atcatgaaat tttatTTTTT aaaaaagaga agtaaaagga taaagtattc acttttatgt 14280
ccacagtctt ttccttttagg ctcatgatgg agtatcagag gcatgaatgt gtttaaccta 14340
agagccttaa tggcttgaat cagaagcact ttagtcctgt atctgttcag tgtcagcctt 14400
tcaaacatca ttttaaattc catttgactt taagtaaata acttaatctc tctacatgtc 14460
aatttcttca gctataaaat gatgggtattt caataaataa atacattaat taaatgatat 14520
tttacaaact aattgggctg ttttaaggct caataagaaa atttctgtga aaggctctta 14580
gcaaatgtag ggttctatac aaataaaaga taacattatg cttatatctt cgggtgtttat 14640
catgcaaagc tcttctgagt tttttgaaga gtcacctac tattttttgt ttttagtttg 14700
ttaaattggt ttataggcaa tgtttttaat ctgttttctt taacttacag tgccatcagc 14760
tcacacttgc aaacctgtgg ctgttccgtt gtagtaggta gcagtgcaga gaaagtaaat 14820
aaggtagttt attttataat ctagcaaatg atttgactct ttaagactga tgatatatca 14880
tggattgtca tttaaatggg aggttgcaat taaaatgatc taatagtata aggaggcaat 14940
gtaatctcat cgaattgctg agacaacttg tggcaacagt gagtttgaaa taaagtgaat 15000
aggagtcatt tatcagttta ttttgataac ttgtaaatac cagtgtcaga tgtgtataaa 15060
tggttttgag aatatattaa aatcagggtat ttaaaaaaac actattcttc tatttcccaa 15120
tgtaatcttt aacaaatctg aaggtagtca tgtactttcg gtactagttc tgaagaaatg 15180
ttatttggtt attcatcttg atttcattgt ctgggtttc cttctaaatc tatcccttct 15240
tgggagctat tgggattaag tggtcattga tgattatact ttattcagta atgtttctga 15300
ccctttcctt cagtgtact tgagttaata aaggattaat gaacagttac atttccaagc 15360
attagctaat aaactaaagg attttgcact tttcttccact gaccattagt taaaaacagt 15420
tcagagataa gtacatgtat ctttcaattc tagcaaacct aatttttttaa agaagtttt 15480
acataggaaa tatgttgga atgattattt actttacaaa gatattcata atttattttt 15540
tctgtaacta gctactttgt atatttacat gacctaata ttatcaaaat tatatttctc 15600
atataaccat ttatgagagc ttagtattcc tctgtcatta tattgcgtct acggactagt 15660
gatcttacta cttctgttac ctgaacaag tggcttcccg tctgtgacct ccaaagccgt 15720
aggttccaca gagtgactgc tgagctgctt tatgaaggga gaaaggctcc atagttgggt 15780
ttttggtttt gcttttggtt ttgtttttaa catttttcct atcctccatc ctcttgaggc 15840
agagtagctt accttttata ttgttttaat ttgagaaaga agttgccact gctctagatt 15900
gaaaaccact gctttaacat aataactctg aatatgggtt gaatttcaag atagtacat 15960
gcctttttat ttttactaat agagctgtag gtcgaatatt attagatttc taaacccac 16020

```

-continued

ccaatgacct	ccttatTTtTa	aatcaaattt	aataattaat	tatcttctta	ttggaggatc	16080
tggacattct	ttgatgtttc	ttacaatgaa	tttcacatgt	agaccacta	aacagaagtt	16140
ataaagggtc	catggtcaaa	taagtctgag	aaagtctgca	tattatataa	ttcacctaaa	16200
gagtcacagt	atgtacccaa	atgttaaagg	ttttgagatg	ccatacagta	aatttaccaa	16260
gcattttcta	aatttatTTtg	accacagaat	ccctatTTtTa	agcaacaact	gttatatccc	16320
ataggttcca	ggtgactaaa	gaatacttat	tgcttaggat	atgttttatt	gataataaca	16380
attaaaatgt	cagatatctt	tcataagcag	atcagtggtc	tttttaaaac	tttgtatTTt	16440
aatgctaaaa	tcttttcttt	tgtagatagt	cagaacatta	tgcttttttc	tgactgcagc	16500
agagagaaaa	tgctccaggt	tatgtgaagc	agaatcatca	tttaaataTG	agtcagggct	16560
ctttgtacag	ggcctgctaa	aggtatagtt	tctacttatc	acaagggaaa	ccaattttct	16620
aaaatcattt	ttgagactct	ttgtagacaa	atattaaata	ttagcattta	atgtatctca	16680
tattgacatg	cccagtgact	gacttccttt	gcacagttct	gcgcataGac	tatatgtctt	16740
atggatttat	agttagtatc	atcagtgtaa	caccatagaa	taccctttgt	tttccagggtg	16800
ggtccttgta	cctacatgtc	tagcatcagg	tgttgTTTTt	TTTTTTTTtT	tttaaaacat	16860
atgcttaaat	caggttgcac	atctaaaata	agatcatttc	tttttaacta	aatagatttg	16920
aattttattg	aaaaaaattt	taaaacatct	ttaagaagca	tataggattt	aagcagttac	16980
tatgtatgtg	tactaaaata	tatatatatt	cctaaatata	tattcctata	tataatatat	17040
gtatttctat	atataatata	tattagaaaa	aacttagagt	tttctttcat	ttgagtctac	17100
tgttcaagga	gcaaaacaga	gaaatgtaaa	ttagcaatta	tttacaataa	ttaaagggaa	17160
gaaagtgtgt	cacctgttg	gatctattat	tgttgTTTTa	attatagtcc	caagacgtga	17220
agaaatagct	ttcctaattg	ttatgtgatt	gtctcatagt	gactactttc	ttgaggatgt	17280
agccacagca	aatgaaatt	taaaaaattt	aaaaattggt	gcaaataaaa	gttatattag	17340
gcttttgtgc	aatttcaata	atgtgctgct	atgaactcag	aatgatagta	tttaaataa	17400
gaaactagtt	aaaggaaaca	cagtttctat	ttgagttata	caaactctgta	aattagaact	17460
tctcctgtta	aggcattata	aagtgcTTaa	tacttttggt	tcctcagcac	cctctcattt	17520
aattatataa	ttttagctct	gaaagggacc	tataccagat	gtgtagagga	aatttcaaaa	17580
ctatgatcta	atgaaaaaat	atttaatagt	tctccatgca	aatacaaatt	atatagtttt	17640
ctggaaaata	cctttgacat	tatacaaaga	tgattatcac	agcattataa	tagtgaaaaa	17700
atggaaaatag	cctctttctt	ctgttctggt	cacagcatat	ggcacagtac	ctcatatgca	17760
gtaggttatt	atgacctggt	aactggctcc	cccaactgat	taggaaagaa	gtaaatttgt	17820
tatttataaa	aatacgtgtt	cattgagatg	catagaataa	ttaaGaaatt	aaaagacact	17880
tgtaatTTca	aatccagtga	ataccactg	ttaatatttg	gcatactctc	ttctagtctt	17940
tttttccctt	ttgcatgtat	tttctTTaag	actcccaccc	ccactggatc	atctctgcat	18000
attctaactc	gcttttttca	cagcagattc	taagcctttt	tgcatatcaa	cacaaacttc	18060
aacaacttca	tcttttagatg	ctaaataatg	aattcatttt	tatttactta	accactttct	18120
ttggatgctc	aggttattct	gatgttttgc	cattaaaacc	aatgctatac	tgaacacttc	18180
tgtcactaaa	acttgaacac	actcatgaat	aatttcttag	gataaatttt	tagagatgga	18240
tttgctaaat	caaagaccat	tttttaaaaa	ttgaaaaaca	attatatcgt	ttggcatgta	18300
agacagtaca	ttttcctttt	atTTtgacag	gattcaactg	gaagctttgt	gctgcctttc	18360
cggcaagtca	tgtatgctcc	atatcccacc	acacacatag	atgtggatgt	caatactgtg	18420

-continued

aagcagatgc	caccctgtca	tgaacatatt	tataatcagc	gtagatacat	gagatccgag	18480
ctgacagcct	tctggagagc	cacttcagaa	gaagacatgg	ctcaggatac	gatcatctac	18540
actgacgaaa	gctttactcc	tgatttgtac	gtaatgctct	gcgtgctggt	actgtagtca	18600
agcaatatga	aactgtgtct	tttatgaata	aaaacaaaac	agaagttgca	ttcaaaaaga	18660
aagaaatatt	actagcagaa	ttatgcttga	agaaacattt	aatcaagcat	ttttttctta	18720
aatgttcttc	tttttccata	cgattgtggt	taccctaaaa	taagtaagat	taacccttaa	18780
agtgaatatt	taactatttg	tttaataaat	atatattgag	ctcctaagca	ctgttctagg	18840
tactgggctt	aatagtggct	aaccacacag	ctccagcccc	tacattgcat	atagtctatt	18900
gtataagtta	ctgaatggac	ttactaacaa	aaccagagaa	gtaattctaa	gtcttttttt	18960
tcttgacata	tgaatataaa	atacaacaaa	actggtaaaa	tatattaata	gagcattctt	19020
ttactttgca	ttttatattg	ttactcactt	cgtattttaag	aaaaacagtc	tgatcaggaa	19080
attcaaaagg	aaaagtaatg	ataattaatt	gagcatagac	ccaacttgaa	aagaaaaaaa	19140
aggatgatga	taaatctata	atcctaaaac	cctaagtaaa	cacttaaattg	atgttctgaa	19200
atcaggaaaa	gaattatagt	atatttttgt	ggttctcttt	tattagttga	aaaaaggcac	19260
agtagctcat	gcctataaga	acagagcttt	gggattccaa	ggcaggcaga	tcacttgagg	19320
ccaggagttc	cagaccagcc	tgggcaacat	agtgaacccc	catctctaca	aaaataaaaa	19380
agaattagtt	gaatgtgttt	ctgtgtgcct	ataatcctag	ctattcagaa	agctgaggca	19440
ggaggatctc	ttgagcccag	gagtttgagg	ttacatggag	ttatgatgtg	ccagtgtact	19500
ccagcctgcg	ggacaatgag	actctgtctt	gttaaaaaaa	aaagtgcttg	gaataatgtt	19560
tggcatatag	aaggtaacaa	cagtaaatgt	taactgtaat	aaccaggtta	taagtgtgta	19620
aggtgataga	aaaattgggg	caaacaaccc	tgacctgtgt	ctctacagaa	taagtttgag	19680
ttgaggcaac	agacatgtgg	agcaccagta	attacacact	aaatgttaac	caaagcggtt	19740
gaatagtaac	atcttattca	agggaccccc	agccttatat	atctcaaggt	gcagaaagat	19800
gacttaatat	aggaccatt	ttttccgagt	tctccagagt	ttttattggt	tcttgagaaa	19860
gtagtggggg	aattgtttta	gaaaatgaat	tgggtcaaact	gaaattccat	gtcagtaagt	19920
ttttacatat	tggtaaattt	tgatagacat	gtagaagttt	tctaattaat	ctgcgccttg	19980
aaacattttc	ctttttccta	aagtgccttag	tattttttcc	cttttttgat	tggttgcttg	20040
ggagcttttt	tgaggaaatt	tagtgaactg	cagaatgagt	ttgcaaccat	ttagtatttt	20100
tgttttgtgt	tttagaggag	gtatgtgtat	tttaacattt	cttaatcatt	tttagccagc	20160
tatgtttggt	ttgctgattg	acaaactata	attaaacagc	tattctcatt	ttgctgatca	20220
tgacaaagta	atatcctgaa	tttttaaatt	ttgcatccag	ctctaaattt	tctaaatttt	20280
ctaaacataa	aattgttcaa	aaaatagtat	ttttagccac	tagattgtgt	gttggttaagt	20340
ctgttgtcac	agactcattt	tacttttcag	tgtgtgtttt	tacatgttaa	ttatgtttgt	20400
catttttaat	tttaactttt	taaaataatt	ccagtcactg	ccaaaacatg	aaaaattggt	20460
cactggaaat	ttttttttta	acttttattt	taggttcatg	tgtacatgtg	caggtttggt	20520
atacaggtaa	attgcgtgtc	gtgaggggtt	ggtgtacca	ggtaataagg	gtagtacca	20580
ataggtagtt	ttttgatcct	tacccttctc	ccacccttct	ttcaccctcg	agtaggcctt	20640
ggtgttgctg	tttcttctt	tgtgtccatg	tgtactcaat	ggtagctcc	tacttagaag	20700
tgagaacatg	cggatatttg	ttttctgttc	ctggattagt	tcactcagga	taatggcctc	20760

-continued

tagctccatc	tgttttttat	ggctgcatag	tattccatgg	tgtatatgta	tcatgttttc	20820
tttatccagt	ctaccattga	tagacattta	gggtgattct	ctgtctttac	tatcatgaat	20880
agcgctgtga	tgaacatata	cacatgcatg	tgtccttatg	gtggaacaat	ttgtattcct	20940
ttaagtatat	acagaataat	ggggttgcta	gggtgaatgg	tagttctatt	gtaagttatt	21000
tgtgaaatct	tcaaactgct	tttcacaata	gctaaactaa	tttacagtcc	caccagcagt	21060
gtataagtgt	tcccttttct	ccacaacctt	gccaacatgt	tattttttta	cttttcaata	21120
ataggcattc	ctagagaatt	gatttgcaat	tctctaata	ttagtatat	tgagcatttt	21180
ttcgtatgct	ttttagctgt	gtgtatatat	tcttttgaaa	aatgttaatg	tcctttgccc	21240
agtttgtaat	ggggttgttt	gtttttgctt	gttaattaaa	gttccttcca	gattctggat	21300
atccctttgt	cagatgcgtg	gtttgcagat	atttttctcc	ccttgtgtag	gttgtctttt	21360
tactctgttg	atagtttctt	ttgccgggca	ggagctcatt	aggtctcatt	tgtgtttggt	21420
tttgttgcag	ttgcttttgg	cgtcttcac	ataaaatctg	tgccagggcc	tatgtccaga	21480
atgggtatttc	ctagtttgtc	ttccaggggt	tttacaattt	tagattttac	gtttatgtct	21540
ttaatccgtc	ttgagttgat	taaggaaggg	gtccagtttc	actctaattc	ctatggctaa	21600
caattatccc	agcaccattt	attgaatacg	gagtcctttc	cccattgctt	gtttttgtca	21660
attttgttga	agatctgatg	gttgtagggt	ctatgtggct	ttatttcttg	gctctctatt	21720
ctccactggg	ctgtctgttt	ttataccagt	accctgctgt	taaggttcct	atagcctttt	21780
agtataaggt	cggctaagt	gatgcctcca	gctttgttct	ttttgcttag	gattgctttg	21840
gctatttggg	ctcctttttg	gttccatatt	aattttaaaa	tagttttttc	tagttttgtg	21900
aagaatgtca	ttgatagttt	agaggaatag	cgttgaatct	gtagattgct	ttgggcaaat	21960
ggccatttta	acaatattga	ttcttcctat	ctatgaacat	ggaatgtttt	tccatgtgtt	22020
tgtgtcatct	ctttatacct	gatgtataaa	gaaaaaccag	tattattgct	actcaatctg	22080
ttccaaaaaa	ttgaggagga	ggaactcttc	cctaatagaga	ccggcttcct	tctgatacca	22140
aaacctggca	gagatacaac	agaaaaaga	aaacttcagg	ccaatatcct	tgatgaatat	22200
agatgcaaaa	atcctcaaca	aaatactagc	aaccaaattc	agcagtacgt	caaaaagcta	22260
atctacttta	agtaggcttt	atccctggga	tgcaagggtg	gttcaacata	cacaaatcaa	22320
taagtgtgat	tcatcacata	aacagagcta	aaaacaaaaa	ccacaagatt	atctcaatag	22380
gtgcagaaaa	ggctttcaat	acaatttaac	atccttcatg	ttaaaaacct	tcagtaggtc	22440
aggtgcagtg	actcacacct	gtaatcccag	cactttggga	ggccaaggcg	gacgtatatc	22500
ttaagcccag	gagttcaaga	ccagcctagg	cagcatgggtg	aaaccccatc	tctacaggaa	22560
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	agcttaatat	ggcggcatgc	acctatagtc	22620
ccagctactc	aggaggttga	ggtgggagga	ttgcttgagc	ccaggaggca	gaggttgagc	22680
cgagctgaga	tcgtgccact	gcactccaac	ctgggcaata	aagtgagacc	ctgtctcaaa	22740
aagaaaaaca	aaaataatcc	taaaccaact	aggcattgaa	ggaatatgcc	tcaaaaaaat	22800
aagaaccatc	tatgacagac	ccacagccaa	tatcttacca	aatgggcaaa	agctggaagt	22860
attctccttg	agaaccgtaa	caagacaagg	atacacactc	tcatccctcc	ttttcagcat	22920
agttctggaa	gtcctcgcca	gagcagtcag	gaaagagaaa	gaaagaaaag	gcattcagat	22980
aggaagagaa	gaagtcaaac	tatttctgtt	tgcaaggcagt	ataattctat	acctagaaaa	23040
tgccatagtt	tctgcccaga	agtcctaca	tctgttaaaa	atttcagcaa	agtttttagca	23100
ttgtctgtat	tccaacagct	tccaggggtga	gagtgaaatc	aggaacacag	tcccgttcac	23160

-continued

aatagccgca	aaaagaataa	aataccttgg	aatccagcta	accaggagg	tgaaacatct	23220
ctacgagaat	tacaaaacgc	tgctgaaaga	aatcagagat	gacacaaaca	aatggaaatg	23280
ttgtttttta	caccttgctt	tatctaattc	acttataact	aagatattca	ttcagtggaa	23340
caggtataat	aagaccactc	gacttaaata	taagccttat	tctctttcca	gagcccaaga	23400
aggggcacta	tcagtgccca	gtcaataatg	ataaaatgct	gatatttttc	ccctttactg	23460
tttctttctt	ctgtagtggt	gtacactcat	ttcttaagat	tagaaaactt	gacctacctt	23520
cctgtttgct	tctacacacc	cccattctct	ttttttgcca	ctccggtcag	gtataggatg	23580
atccctacca	cttttagtta	aaacctcctt	cccttattaa	atgttctctt	accactctgg	23640
cctgagtaga	acctagggaa	aatggaagag	aaaagatgaa	agggaggtgg	gggctgggaa	23700
gggaatagtc	ttgtttgtgt	gtttgcttta	gcacctacta	tatcctaggt	gctgtgttag	23760
gcacacatta	ttttaagtgg	ccattatatt	gctacatctc	actctgggtca	ttgccaaagg	23820
aggtagtact	ttcttgata	gttggttcat	gttacttata	ggtggtggac	ttgttgaggc	23880
aacccaatg	gataatcatc	tgagtgtgtt	ctctaactc	agatttttct	tcatattttt	23940
tggtttgttt	tggtttttga	tggtggtggt	tgtgtgctta	tttttggtgc	tggttggttt	24000
ttttgttttg	tttttgatat	ggcaagaatt	ggtagtttta	tttattaatt	gcctaagggt	24060
ctctactttt	tttaaaagat	gagagtacta	aaatagattg	ataggtacat	acataccttt	24120
atgggggact	gcttatattc	cttagagaaa	aaaattactt	attagcctga	caaacaccag	24180
taaaatgtaa	atatatccgt	gagtaaataa	atgaatgtat	gctttgtatc	tccaaatata	24240
tacatctata	ttcttacaaa	tatgttttta	tgtaatacca	atttataaga	acttaaaatg	24300
ttggctcaag	tgagggatgg	tggaaagtag	cattatatag	ccatttcaac	atttgaactt	24360
ttttcttcat	tttcttcttt	tcttcaggaa	tatttttcaa	gatgtcttac	acagagacac	24420
tctagtgaag	gccttcctgg	atcaggtaaa	tgttgaactt	gagattgtca	gagtgaatga	24480
tatgacatgt	tttctttttt	aatatatctt	acaatgcctg	ttctctctct	ctatatatat	24540
atatttatat	atttccttgg	atcatgcccc	agagttctgc	tgagcaattg	cagttaagtt	24600
agttacacta	cagttctcac	aagagtctgt	gaggggatgt	caggtgcac	attacattgg	24660
atgcctcttg	tcctagattt	atgtttcggg	aattcagacc	tatgtttaca	atataataaa	24720
tattgttgct	gccttttaca	gataaaataa	taagatataa	acttgaccac	aactactgtt	24780
ttttgaaaca	tagagttcat	ggtttacatg	tatcaaagtg	aaatctgagt	tagcttttac	24840
agatataata	tatacatata	tatatcctac	aatgcttgta	ctatatatgt	agtacaagta	24900
tatatatgtg	tgtgtgtgtg	tgtatatata	ttatggcact	gtagtatata	tatgtttata	24960
tgttaaaaaa	tatataaata	tatgtttacat	atttaacata	aacatatata	catatatgtt	25020
aaatatataa	catatactct	atatatgaca	aatagagtat	aatatatatt	tttatttttt	25080
atatatatat	aaaacatgat	agaattaaga	attaagtcct	aatctgtttt	attaggtgct	25140
ttttgtagtg	ttcagtcttt	ctaaagtgtc	taaatgattt	ttccttttga	cttattaatg	25200
gggaagagcc	tctatattaa	caattaaggc	tgcagcattg	attacttcaa	acaacaaaca	25260
ttttaattca	agcattaacc	tataactcaa	gtaagttttt	tttttttttt	ttttgagaaa	25320
gggagggtgt	ttatttgctt	gaattgagtc	aaaaatattt	ttgaaacatc	atgtactcat	25380
ttaaagtata	acatctttat	tgtttcattc	ttttaaaaaa	tatctactta	attacacagt	25440
tgaaggaaat	tgtagattat	atggaactta	tttcttaata	tattacagtt	ttgttataat	25500

-continued

aacattctgg	ggatcaggcc	aggaaactgt	gtcatagata	aagctttgaa	ataatgagat	25560
ccttatgttt	actagaaatt	ttggattgag	atctatgtgg	tctgtgacat	attgcaaagt	25620
tcaaggaaaa	ttcgtaggca	tggaatttct	caaactgaaa	atccctccca	ctgtccacct	25680
catcacatgc	acacattcta	ctcttaccba	cccactccac	cccttgcaaa	agtacagata	25740
tatgaatgtc	tcaaaaccat	gggctcatct	tctagaagct	tcaatgttat	ttgaagattt	25800
gggcagagga	agttaagaaa	tatgaaatag	cttacatatg	agttttaata	gtgaaacaaa	25860
catggatgta	ttctgaagta	gaatgcaaaa	tttgagtgc	tttttttttt	tttgagactg	25920
agtctggctc	tgtcgcccag	gctggagtgc	agtggccgga	tctcagctca	ctgcaagctc	25980
cacctcccgg	gtttacgcca	ttctcctgcc	tcagcctccc	gagtagctgg	gaccacaggc	26040
gcccgccact	tcgcccggct	agtttgtttg	tatttttttag	tagagatggg	gtttcaccgt	26100
gtagccagg	gannnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnccatt	26160
gagtgcattt	ttaaagataa	atcagaaaac	ttcgaaaaac	tatcagattg	gccggacatg	26220
gtggcttatg	cctgtaatcc	tagcactttg	ggaggctgag	gtgggtggat	cacgaggtca	26280
ggagatcgag	accatcctgc	caacatgggtg	aaaccccatc	tctactaagt	atacaaaaat	26340
tagctgggcg	tgacagcacg	tgctgtaat	cccagctact	tgaggaggctg	aggcaggaga	26400
atcgcttgaa	cccgggaggt	ggagggttgc	gtgagtcaag	atcacaccac	tgcacttcag	26460
cttggtgaca	gagctagact	ccatatcaaa	aaaaaaaaaa	aaaaaaagaa	gtcagattgt	26520
tcctacaccc	agtgtctcta	taccacactc	ctactagggg	gcatcagtgg	aatgggttaa	26580
ggagatgttt	agtgtgtatt	gtctgccaa	gactgttaac	actgtcctag	aaacattgct	26640
gtacaagtag	aatgtgagca	aattatgtat	tgaatgggtt	cctctccctg	cagggtctttc	26700
agctgaaacc	tggtttatct	ctcaggagta	ctttccttgc	acagttttta	cttgtccttc	26760
acagaaaagc	cttgacacta	ataaaatata	tagaagatga	tacgtgagta	caactcctac	26820
atggaggaaa	aaccttttgt	acgttgtttt	ttgttttatt	tcctttgtac	attttctgta	26880
tcataatttt	tgcttttttt	tttttttttt	ttttctccat	tactttcagg	cagaagggaa	26940
aaaagccctt	taaatctctt	cggaacctga	agatagacct	tgatttaaca	gcagagggcg	27000
atcttaacat	aataatggct	ctggctgaga	aaattaaacc	aggcctacac	tcttttatct	27060
ttggaagacc	tttctacact	agtgtacaag	aacgagatgt	tctaataact	ttttaaatgt	27120
gtaacttaat	aagcctattc	catcacaatc	gtgatcgctg	ctaaagtagc	tcgggtgggtg	27180
ggggaaacat	tcccctggat	catactccag	agctctgctc	ggcagttgca	gttaagttag	27240
ttacactaca	gttctcacia	gagtctgtga	ggggatgtca	ggtgcatcat	tacattggat	27300
gtctcttttc	ctagatttat	gcttttgagg	tacagacctc	tgtttacaat	ataataggta	27360
ttattgctgt	cttttaataa	tataataata	ggatataaac	ttgaccacaa	ctgctgtttt	27420
tttgaaatat	atgattcatg	gtttacatgt	attaagggtga	aatccgagtt	cgctttttaca	27480
gatattagtt	gactttctat	cttttggtat	tctttgggtg	gtggaattac	tgtaataactt	27540
ctgcaatcaa	ctgaaaatta	gagcctttta	atgatttcag	ttccacagaa	agaaagtgag	27600
cttcaacata	ggataagctt	tagaaagaga	attgatcaag	cagatgttta	attggaattg	27660
attattagat	cctgctttgt	ggatttagcc	ctcgggatcc	agtctgtaga	aatgtctgat	27720
agttctctat	agtccttgc	catgggtgaac	cacagttagg	atgttttggt	tgttttattg	27780
ttgttgctat	tggtgatgtt	ctatatagtt	gagctctata	aaaggaaatt	gtattttatg	27840
ttttagtagt	tggtgccaac	tttttaaat	aattttcatt	atttttgagc	caaattgaaa	27900

-continued

tgtgcacctc	ctgtgccttt	tttttccttg	gaaaatcgaa	ttacttggaa	gaagttcaga	27960
tttcaactgg	cagtcgtttt	catcttggtt	tcttcttgca	gagtcctacc	atgtacctgc	28020
tttggcaatc	attgtaactc	tgagattata	aaatgcatta	gagaatatat	taactaataa	28080
gatctttttt	ttcaggaaca	gaaaatagtt	ccttgagtac	ttccttctta	cattttctgcc	28140
catgtttttg	aagttgttgc	catttgcctg	caataggcta	taaggaatag	caggagaaat	28200
tttactgaag	tgctattttt	ctaggtgcta	ctttggcaga	gctaagtggg	ctgtttcttt	28260
tgtttcctta	atgcgtttgg	accattttgc	tggtctgtaa	ataactgatt	aatataattc	28320
taacacaata	ttgacattgt	agtgtacaca	aacacaaata	ttttatttaa	aactggaagt	28380
aacataaaag	ggaaaatata	tttataagaa	aggaataaag	gtaatagagc	tcttctgtcc	28440
cccagccacc	aaatttacac	aacaaaatca	tatgttctaa	tgtgaaaggt	cataatagct	28500
ttcccatcat	taatcagaaa	gatgtggcag	cttgattttt	tagacaaccc	ctgaactaga	28560
tgactgttgt	actgtagctc	agtcatttaa	aaaatatata	aatactatct	cgtagtgtcc	28620
catactatgt	tttttacatg	atagattctt	atttaagtgc	taactggtta	ttttctttgg	28680
ctggtttatt	gtactgttat	atagaatgta	agttgtacag	tgaaataagt	tattaaagca	28740
tgtgtaaaca	ttgttatata	tcttttctcc	tagatggaga	attttgaata	aaatatnnnn	28800
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	28860
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	28920
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	28980
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	29040
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	29100
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	29160
nnnnnnnnnn	nnnnnnnnnn	nnnnnnagaa	gactaattga	tcatatcact	atgattctca	29220
aagaagaacc	aaaacttcat	ataatactac	aaatatgaga	tagttacttc	tgtagtatat	29280
ttctgtaatg	ctacaggtta	aacaggtcac	tcttatataa	cactattttg	attttgatgt	29340
agaattgcac	aaattgatat	ttcttctatg	atctgtaggg	tatagcttaa	agtagcaaaa	29400
acagtccacc	acctccagtt	aacacacagt	aacactatgg	gactagtatt	attatttcca	29460
ttttacaaag	gaggaaacta	aagcttaaag	atgtgtaata	tacagcccaa	ggtcacacag	29520
ctggtaaagg	tagatttcat	cccagacagt	tacagtcatt	gccgtgggca	cagctcctaa	29580
cttattaact	ccatgtaact	ggtactcagt	ttagttgaat	tgaaaggaga	gtagggaagc	29640
aggtctgttt	gcactattca	gagcccaagt	gtgaatccct	gctgtgctgc	ttggagaagt	29700
tacttaacct	atgcaagggt	cattttttta	atatttgaaa	cggaatgata	atacatactt	29760
caccagtggg	tttaatgaga	ccttataaga	tcgttagttc	agtacctgac	cagtgcctca	29820
taaatgcctt	ttcatccaat	ctgacaatct	ctagcttgta	attggggcat	ttagaacatt	29880
taatatgatt	attggcatgg	taggttaaag	ttgtcatctt	gctgttttct	ctttgttctt	29940
ttttctcctt	tcttttgat	ttttttttta	ttttactgtg	tcttctctgt	tgtcttatta	30000
a						30001

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Primer		
<400> SEQUENCE: 12		
gggtctagca agagcaggtg	20	
<210> SEQ ID NO 13		
<211> LENGTH: 21		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Primer		
<400> SEQUENCE: 13		
gtcttggaac cagctggaga t	21	
<210> SEQ ID NO 14		
<211> LENGTH: 24		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Probe		
<400> SEQUENCE: 14		
tgatgtcgac tctttgccca ccgc	24	
<210> SEQ ID NO 15		
<211> LENGTH: 22		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Primer		
<400> SEQUENCE: 15		
tgtgacagtt ggaatgcagt ga	22	
<210> SEQ ID NO 16		
<211> LENGTH: 25		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Primer		
<400> SEQUENCE: 16		
gccacttaaa gcaatctctg tcttg	25	
<210> SEQ ID NO 17		
<211> LENGTH: 21		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Probe		
<400> SEQUENCE: 17		
tcgactcttt gccaccgcc a	21	
<210> SEQ ID NO 18		
<211> LENGTH: 783		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
<400> SEQUENCE: 18		
gcctctcagt acccgaggct cctttttctc gagcccgag cggcagcgct cccagcgggt	60	
ccccgggaag gagacagctc gggtactgag ggcgggaaag caaggaagag gccagatccc	120	
catcccttgt ccctgcgcg cgcgcgcgcg cgccgcgcgc gggaagcccg gggcccgat	180	

-continued

gcaggcaatt ccaccagtcg ctagaggcga aagcccgaca cccagcttcg gtcagagaaa	240
tgagaggggaa agtaaaaatg cgtcgagctc tgaggagagc ccccgcttct acccgcgctt	300
cttcccgga gccgaacccc aaacagccac ccgcccaggat gccgcctcct cactcaccca	360
ctcgccaccg cctgcgcctc cgccgcccgc ggcgaggca ccgcaaccgc agccccgccc	420
cgggcccgc cccggggccc ccccgaccac gccccggccc cggccccggc cccggccccg	480
gccccctagc cgcgactcct gagttccaga gcttgctaca ggctgcggtt gtttccctcc	540
ttgttttctt ctggttaatc tttatcaggt cttttcttgt tcacctcag cgagtactgt	600
gagagcaagt agtggggaga gaggggtggga aaaacaaaaa cacacacctc ctaaaccac	660
acctgctctt gctagacccc gccccaaaaa gagaagcaac cgggcagcag ggacggctga	720
cacaccaagc gtcactcttt acgtgggcgg aacttgtcgc tgtttgacgc acctctcttt	780
cct	783
<210> SEQ ID NO 19	
<400> SEQUENCE: 19	
000	
<210> SEQ ID NO 20	
<400> SEQUENCE: 20	
000	
<210> SEQ ID NO 21	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 21	
gccttactct aggaccaaga	20
<210> SEQ ID NO 22	
<211> LENGTH: 20	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 22	
ccggccccta gcgcgcgact	20
<210> SEQ ID NO 23	
<211> LENGTH: 19	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 23	
cggcccctag gcgcgcgact	19
<210> SEQ ID NO 24	
<211> LENGTH: 19	
<212> TYPE: DNA	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	

-continued

<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 24		
ccggccccta gcgcgcgac	19	
<210> SEQ ID NO 25		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 25		
ccctagcgcg cgactcctga	20	
<210> SEQ ID NO 26		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 26		
cccctagcgc gcgactcctg	20	
<210> SEQ ID NO 27		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 27		
gcccctagcg cgcgactcct	20	
<210> SEQ ID NO 28		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 28		
ggcccctagc gcgcgactcc	20	
<210> SEQ ID NO 29		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 29		
cggccccctag cgcgcgactc	20	
<210> SEQ ID NO 30		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 30		
cctagcgcgcg gactcctg	18	

-continued

<210> SEQ ID NO 31 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 31		
ccctagcgcg cgactcct	18	
<210> SEQ ID NO 32 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 32		
cccctagcgc gcgactcc	18	
<210> SEQ ID NO 33 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 33		
gcccctagcg cgcgactc	18	
<210> SEQ ID NO 34 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 34		
ggcccctagc gcgcgact	18	
<210> SEQ ID NO 35 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 35		
cggcccctag cgcgcgac	18	
<210> SEQ ID NO 36 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 36		
aggctgcggt tgtttcctc	20	
<210> SEQ ID NO 37 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide		

-continued

<400> SEQUENCE: 37		
caggctgcgg ttgtttccct		20
<210> SEQ ID NO 38		
<211> LENGTH: 19		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 38		
ggctgcggtt gtttcctc		19
<210> SEQ ID NO 39		
<211> LENGTH: 19		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 39		
aggctgcggt tgtttccct		19
<210> SEQ ID NO 40		
<211> LENGTH: 19		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 40		
caggctgcgg ttgtttccc		19
<210> SEQ ID NO 41		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 41		
gctgcggttg tttccctc		18
<210> SEQ ID NO 42		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 42		
ggctgcggtt gtttcct		18
<210> SEQ ID NO 43		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 43		
aggctgcggt tgtttccc		18
<210> SEQ ID NO 44		
<211> LENGTH: 18		

-continued

<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 44		
caggctgcgg	ttgtttcc	18
<210> SEQ ID NO 45		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 45		
tctgtctttg	gagcccaa	20
<210> SEQ ID NO 46		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 46		
ctgcgatccc	cattccagtt	20
<210> SEQ ID NO 47		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 47		
gccttactct	aggaccaa	18
<210> SEQ ID NO 48		
<211> LENGTH: 18		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 48		
tctgtctttg	gagcccaa	18
<210> SEQ ID NO 49		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 49		
ggttaatctt	tatcaggtct	20
<210> SEQ ID NO 50		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 50		

-continued


tggttaatct ttatcaggtc	20
<210> SEQ ID NO 51 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 51	
ctgggtaatc tttatcaggt	20
<210> SEQ ID NO 52 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 52	
gttaatcttt atcaggtc	18
<210> SEQ ID NO 53 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 53	
ggttaatctt tatcaggt	18
<210> SEQ ID NO 54 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 54	
tggttaatct ttatcagg	18
<210> SEQ ID NO 55 <211> LENGTH: 18 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 55	
ctagcgcgcg actcctga	18
<210> SEQ ID NO 56 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic oligonucleotide	
<400> SEQUENCE: 56	
gggacactac aaggtagtat	20
<210> SEQ ID NO 57 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence	

-continued


<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 57		
tacaggctgc ggttgtttcc	20	
<210> SEQ ID NO 58		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 58		
cccggeccct agcgcgcgac	20	
<210> SEQ ID NO 59		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 59		
ggtaacttca aactcttggg	20	
<210> SEQ ID NO 60		
<211> LENGTH: 20		
<212> TYPE: DNA		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic oligonucleotide		
<400> SEQUENCE: 60		
aatctttatc aggtcttttc	20	

What is claimed is:
1. A modified oligonucleotide according to the following 40
formula:


(SEQ ID NO:33)



185



-continued




186

or a salt thereof.


45

2. A modified oligonucleotide according to the following formula:


(SEQ ID NO: 33)



187



-continued



188

3. The modified oligonucleotide of claim 1, which is a sodium salt of the formula.

4. A pharmaceutical composition comprising the modified oligonucleotide of claim 1 and a pharmaceutically acceptable diluent or carrier.

5. The pharmaceutical composition of claim 4, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

6. The pharmaceutical composition of claim 4, wherein the pharmaceutical composition consists essentially of the modified oligonucleotide and phosphate-buffered saline (PBS).

7. A pharmaceutical composition comprising the modified oligonucleotide of claim 3 and a pharmaceutically acceptable diluent or carrier.

8. The pharmaceutical composition of claim 7, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

9. The pharmaceutical composition of claim 7, wherein the pharmaceutical composition consists essentially of the modified oligonucleotide and phosphate-buffered saline (PBS).

10. A pharmaceutical composition comprising the modified oligonucleotide of claim 2 and a pharmaceutically acceptable diluent or carrier.

11. The pharmaceutical composition of claim 10, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

12. The pharmaceutical composition of claim 10, wherein the pharmaceutical composition consists essentially of the modified oligonucleotide and phosphate-buffered saline (PBS).

13. A compound comprising a modified oligonucleotide, wherein the modified oligonucleotide is a gapmer consisting of a 5' wing segment, a central gap segment, and a 3' wing segment, wherein:

the 5' wing segment consists of four 2'-O-methoxyethyl nucleosides,

the central gap segment consists of eight β -D-deoxyribo-nucleosides, and

the 3' wing segment consists of six 2'-O-methoxyethyl nucleosides;

wherein the modified oligonucleotide has the nucleobase sequence 5'-GCCCCCTAGCGCGCGACTC-3' (SEQ ID NO: 33), wherein each cytosine is a 5-methylcytosine; and wherein the internucleoside linkages of the modified oligonucleotide are, from 5' to 3', soossssssssoos, wherein each s is a phosphorothioate linkage and each o is a phosphodiester linkage.

14. The compound of claim 13, comprising the modified oligonucleotide covalently linked to a conjugate group.

15. A pharmaceutical composition comprising the compound of claim 13 and a pharmaceutically acceptable diluent or carrier.

16. The pharmaceutical composition of claim 15, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS).

17. The pharmaceutical composition of claim 15, wherein the pharmaceutical composition consists essentially of the compound and phosphate-buffered saline (PBS). 5

18. A pharmaceutical composition comprising the compound of claim 14 and a pharmaceutically acceptable diluent or carrier.

19. The pharmaceutical composition of claim 18, wherein the pharmaceutically acceptable diluent is phosphate-buffered saline (PBS). 10

20. The pharmaceutical composition of claim 18, wherein the pharmaceutical composition consists essentially of the compound and phosphate-buffered saline (PBS). 15

* * * * *


UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 9,605,263 B2
APPLICATION NO. : 15/130818
DATED : March 28, 2017
INVENTOR(S) : Frank Rigo

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In Claim 1, the structure shown at Columns 183-186 is replaced with the following structure:



Signed and Sealed this
Sixth Day of November, 2018

Andrei Iancu

Andrei Iancu
Director of the United States Patent and Trademark Office