


Lab - 10

NAVEEN KUMAR A G

210123075





- 1) First 25 values of the Van der Corput sequence using the radical function with base $b = 2$:
[0, 0.5, 0.25, 0.75, 0.125, 0.625, 0.375, 0.875, 0.0625, 0.5625, 0.3125, 0.8125, 0.1875, 0.6875, 0.4375, 0.9375, 0.03125, 0.53125, 0.28125, 0.78125, 0.15625, 0.65625, 0.40625, 0.90625, 0.09375]

(x_i, x_{i+1}) plot of 1000 values generated with this sequence.



Observation: A linear pattern can be observed. This means that successive points are linearly related to their previous points.


2) The values used for generating LCG are: $x_0 = 100$, $a = 1597$, $b = 0$, $m = 244944$.



Observations:

- 1) As n (sample size) increases both LCG and VDC are more uniform distribution.
- 2) VDC has a better uniform distribution than LCG for both $n = 100$, $n = 100000$.

3) Halton sequence plots for both $n = 100$, 100000 in \mathbb{R}^2 .



Observations:

- 1) For both values of n , the points are scattered uniformly. This suggests that Halton sequence has nearly uniform distribution.
- 2) For $n = 100000$, the plot is much denser than the one for $n = 100$.