

---

%HW-3 Prb-4  
%Navneet Singh (nsingh1@andrew.cmu.edu)

The equation in simplified form is

$$\frac{d^2\theta}{dr^2} = \left(\frac{hA}{k(1-\epsilon)}\right)\theta - \frac{1}{r} \frac{d\theta}{dr}$$

Above problem is BVP. First we have converted it to system of FODEs. Then we have used shooting method to calculate variation of  $\theta$  with  $r$ .

```
function problem4
clc          %clear screen
clear all   %clearing all stored variables
close all   %close previous plots

%Given data
h = 0.001; %cal/cm^2 s
e = 0.36;  %porosity
A = 15;    %cm^-1
k = 0.0034; %cal/cm s

%calculating values of constant to consise equations
a = (h*A)/(k*(1-e));

%radius of the reactor
rspan = [0 1.3];

%Dirichlet BC at r = R
bc = 1;

%using fsolve to guess initial conditions
%guess for fsolve
guess = 0.1;

options = optimset('Display','off');
sol = fsolve(@func, guess, options);
fprintf('u1(r=0) = %f\n', sol)

%function used to calculate Initial values
function y = func(guess)
    [r, u] = ode45(@dudt, rspan, [guess, 0]);
    y = bc - u(end,1); %checking whether u1(r=1) matches our BC.
end

%using initla values obtained to calculate theta as fucntion of r.
initial = [sol, 0];
[r, u] = ode45(@dudt, rspan, initial);

%The second order ODE converted to two FODEs.
function f = dudt(r, u)
```

---


---

```

f = zeros(2,1);
f(1) = u(2);
if r == 0 %as equation contain 1/r it will get undefined at r=0.
    f(2) = a*u(1);
else
    f(2) = a*u(1) - (1/r)*u(2);
end
end
%plotting
plot(r, u(:,1))
title('\theta vs radius of the reactor')
xlabel('Radius of the reactor (cm)')
ylabel('\theta')
end

u1(r=0) = 0.145772

```



*Published with MATLAB® R2016a*