



## Review

# 18F-FDG positron emission tomography for the assessment of histological response to neoadjuvant chemotherapy in osteosarcomas: A meta-analysis

Li Hongtao<sup>a,1</sup>, Zhao Hui<sup>a</sup>, Wang Bingshun<sup>b</sup>, Wang Xiaojin<sup>b</sup>, Wang Zhiyu<sup>a</sup>, Zheng Shuier<sup>a</sup>, He Aina<sup>a</sup>, Sun Yuanjue<sup>a</sup>, Min Daliu<sup>a</sup>, Shen Zan<sup>a,\*,1</sup>, Yao Yang<sup>a,\*,1</sup>

<sup>a</sup> Department of Oncology, Affiliated Sixth People's Hospital, Shanghai Jiaotong University, Yishan Road 600#, 200233 Shanghai, People's Republic of China

<sup>b</sup> Department of Biostatistics, Shanghai Jiao Tong University School of Medicine, 200025 Shanghai, People's Republic of China

## ARTICLE INFO

## Article history:

Accepted 18 July 2012

## Keywords:

18F-FDG positron emission tomography

Response evaluation

Necrosis

## ABSTRACT

**Purpose:** The purpose of this meta-analysis was to evaluate the predicting value of fluorine-18-fluorodeoxyglucose positron emission tomography with computed tomography (18F-FDG PET-CT) in the assessment of histological response to neoadjuvant chemotherapy in patients with osteosarcomas. **Methods:** A detailed search was made in MEDLINE, EMBASE and the Web of Knowledge for relevant original articles published in English; methodological quality of the included studies were also assessed. Two reviewers extracted data independently. Sufficient data was presented to construct a  $2 \times 2$  contingency table. Pooled sensitivity and specificity, positive and negative likelihood ratios were estimated. A summary receiver operating characteristic curve (SROC) was constructed with the Moses' constant of linear model. A  $\chi^2$  test was performed to test for heterogeneity.

**Results:** Eight studies comprising 178 patients met the inclusion criteria. The pooled sensitivity and specificity for standardized uptake values (SUV) after chemotherapy ( $SUV_2 \leq 2.5$ ) were 0.734 (95% CI, 0.537–0.867) and 0.864 (95% CI, 0.510–0.975), for the ratio of standardized uptake values after ( $SUV_2$ ) to before ( $SUV_1$ ) chemotherapy  $SUV_2:1 \leq 0.5$  were 0.690 (95% CI, 0.497–0.833) and 0.653 (95% CI, 0.492–0.786), the positive and negative likelihood ratio ( $LR+/LR-$ ) for  $SUV_2 \leq 2.5$  were 5.397 (95% CI, 1.169–24.920) and 0.308 (95% CI, 0.165–0.577), for  $SUV_2:1 \leq 0.5$  were 1.989 (95% CI, 1.145–3.457) and 0.475 (95% CI, 0.247–0.915). There was no significant difference between-study heterogeneity for either  $LR+$  or  $LR-$  in any of these analyses. The area under the SROC curve for  $SUV_2 \leq 2.5$  and  $SUV_2:1 \leq 0.5$  were 0.81 and 0.72, respectively.

**Conclusions:** The present meta-analysis showed that 18F-FDG PET-CT scan, as measured by the SUV before and after treatment,  $SUV_2 \leq 2.5$  and  $SUV_2:1 \leq 0.5$  are valuable for predicting the histological response to chemotherapy.  $SUV_2 \leq 2.5$  have better predicting performance than  $SUV_2:1 \leq 0.5$ .

© 2012 Elsevier Ltd. All rights reserved.

## Contents

Introduction .....	e166
Methods .....	e166
Search strategy .....	e166
Selection criteria .....	e166
Data extraction .....	e166
Study character .....	e166
Statistical analysis .....	e166
Results .....	e166

**Abbreviations:** OS, Osteosarcoma; 18F-FDG PET-CT, Fluorine-18-fluorodeoxyglucose positron emission tomography with computed tomography; SUV, standardized uptake value; TBR, tumor to background ratios; ROI, region-of-interest; SROC, summary receiver operating characteristic;  $LR+/-$ , positive/negative likelihood ratios; AUC, area under curve.

\* Corresponding author. Tel.: +86 021 24058408.

E-mail addresses: [lhtmed@yahoo.com.cn](mailto:lhtmed@yahoo.com.cn) (L. Hongtao), [shenzanok@yahoo.com](mailto:shenzanok@yahoo.com) (S. Zan), [yangyao\\_6@hotmail.com](mailto:yangyao_6@hotmail.com) (Y. Yang).

<sup>1</sup> Li Hongtao is the first author, Shen Zan and Yao Yang are co-corresponding authors.

Study selection and characteristics analysis .....	e166
Predictive value of SUV2 .....	e167
Predictive value of SUV2:1 .....	e167
Discussion .....	e167
Author contribution .....	e168
Conflict of interest statement .....	e169
Acknowledgments .....	e169
References .....	e169

## Introduction

Osteosarcoma (OS) is the most common bone cancer in children and young adults [1,2]. In the past decades, the combination of multi-agent chemotherapy has improved long-term survival in patients with OS [3,4]. Histological response or tumor necrosis rate to neoadjuvant chemotherapy has been shown to be associated with survival [5]. Tumor response to neoadjuvant chemotherapy has important implications in subsequent patient management. However, histopathological examination of tumor response can be assessed only after resection. Therefore, an accurate and noninvasive preoperative marker of response would be ideal for planning surgical margins and as a prognostic tool.

Fluorine-18-fluorodeoxyglucose (18F-FDG) positron emission tomography with computed tomography (18F-FDG PET-CT) is now widely used with promising results in the initial diagnosis, staging, and detection of recurrence in many kinds of cancer [6]. Furthermore, some studies revealed that a decreased 18F-FDG uptake after chemotherapy or radiotherapy is associated with good pathological response [7–14]. However, there have been conflicting results. The other studies have not found a significant association between 18F-FDG uptake and tumor necrosis [15–17]. Therefore, the results of predictive value of 18F-FDG uptake remained undetermined.

In the present study, we conducted a comprehensive review of relevant studies and a meta-analysis of data to determine the predictive value of 18F-FDG uptake in determining histological response to treatment in patient with osteosarcoma.

## Methods

### Search strategy

Medline, PubMed, Embase, and the Web of Science were searched (last search was updated on 24 May 2011, using the search terms: “positron emission tomography”, “prognosis” and “sarcoma”). All searched studies were retrieved, and their bibliographies were checked for other relevant publications. References of selected articles and reviews were also searched manually for additional relevant studies. When more than one of the same patient populations was included in several publications, only the most recent or complete study was used to avoid duplication of information.

### Selection criteria

Two independent reviewers assessed the eligibility of studies by reviewing titles and abstracts identified by the search, differences were resolved by discussion. The inclusion criteria required that the specified articles were studies that included: at least 10 patients, 18F-FDG PET-CT scan before and/or after chemotherapy, 18F-FDG PET-CT performed with the intention of monitoring the histological response to neoadjuvant chemotherapy, attenuation-corrected 18F-FDG PET-CT studies. Duplicate studies on the same patients,

studies written in a language other than English, and reviews and abstracts were excluded. No unpublished data or data from abstracts were used. “Good” histopathologic response was defined as tumor necrosis of more than 90% according to Salzer-Kuntschik’s grading systems [18], whereas the presence of 10% or more viable tumor after chemotherapy was defined a “poor” response.

### Data extraction

Two investigators extracted data from eligible studies independently, discussed discrepancies and reached consensus for all items. The following data were collected from each study: first author’s name, year of publication, study design, total number of patients, age, sex, and 18F-FDG PET-CT parameters such as standardized uptake value (SUV). We considered the description of 18F-FDG PET-CT as adequate if the publication reported details on the scanner type, the timing of scanning after injection, a clear description of quantitative procedures including region-of-interest (ROI) methodology, and the performance of attenuation correction.

### Study character

The QUADAS quality assessment tool, an evidence-based quality assessment tool for diagnostic accuracy studies, was used to extract relevant study design characteristics from each study. This tool and the definitions of the characteristics have been fully described by Whiting et al. [19].

### Statistical analysis

Data on the diagnostic performance of the SUV after chemotherapy (SUV2), the ratio of SUV2 to SUV before chemotherapy (SUV1) (SUV 2:1) for determining histological response to chemotherapy were analyzed separately. Pooled sensitivity and specificity for each study were calculated by extracting data into  $2 \times 2$  tables including the numbers of true positive (TP), true negative (TN), false positive (FP) and false negative (FN) results in each study. A value of 0.5 was added to all cells of studies that contained a count of zero to avoid potential problems in odds calculations for studies with sensitivities or specificities of 100%, and a summary receiver operating characteristic (SROC) curve was constructed. Combined positive and negative likelihood ratios (LR+ and LR–, respectively) were also estimated. Heterogeneity was assessed with the  $\chi^2$ -test using a random effects model. All analyses were performed by using Rev Manager V5.1, and STATA [19].

## Results

### Study selection and characteristics analysis

We initially identified 26 articles including discussion of the role of 18F-FDG PET-CT scan in patients with OS. Of those, 18 reports were excluded (5 studies due to reviews [20–24], 2 studies aiming

at staging or grading [25,26], and 3 focused on survival [13,27,28], four for not providing detailed data of enrolled patients [25,29–31], 2 for discussing Ewing sarcoma [8,16], one had less than 10 patients [32], and one used T:NT (tumor to non-tumor background) as parameter [33]. In all, 8 independent eligible studies, which had detailed data of 18F-FDG PET-CT scan and pathological grade score for each patient, were included in the quantitative synthesis [9,10,12,15,17,34–36]. Different methods to quantify 18F-FDG uptake were used: the SUV approach was used in 10 studies (maximum SUV in 9 studies; mean SUV in 1 study), tumor to background ratios (TBR) was used in 3 studies. All analyzed patients were treated with combination chemotherapy regimens. These 18F-FDG PET-CT characteristics are listed in Table 1. The QUADAS quality assessment result is presented in Table 2. The mean time intervals of second 18F-FDG PET-CT scan were provided in 7 studies: 2–3 weeks after treatment and before surgery in brief, while it was not mentioned in the other 4 studies. Five studies mentioned that 18F-FDG PET-CT scan was evaluated blindly to histopathological results, whereas detailed observation methods were not provided in 6 studies.

### Predictive value of SUV2

Eight studies investigated the prediction value of SUV after therapy (SUV2) to tumor necrosis rate. As showed in Fig. 1b, the results of estimation of 8 studies involving 138 patients were included in this comparison. There was no significance between-study heterogeneity for these analyses ( $I^2 = 34.39\%$ ). At a cut-off level at  $\leq 2.5$ , the pooled sensitivity and specificity estimated for  $SUV2 \leq 2.5$  were 0.734 (95% CI, 0.537–0.867) and 0.864 (95% CI, 0.510, 0.975) respectively, and pooled LR+ and LR– were 5.397 (95% CI, 1.169–24.920) and 0.308 (95% CI, 0.165–0.577) respectively (Fig. 1a,b). The SROC curve for  $SUV \leq 2.5$  showed that area under curve (AUC) was 0.81 (95% CI, 0.77–0.84), indicating a lower diagnostic accuracy (Fig. 2a).

### Predictive value of SUV2:1

The diagnostic performances for the ratio of SUV after treatment (SUV2) to SUV before treatment (SUV1) were assessed in nine eligible studies. These studies involving 178 patients were analyzed in comparison. There was no significance between-study heterogeneity for these analyses ( $I^2 = 0.00\%$ ). The results were showed in Fig. 1c. In this meta-analysis, when the cut-off value were set at  $\leq 0.5$ , the pooled sensitivity and specificity estimated were 0.690 (95% CI, 0.497–0.833) and 0.653 (95% CI, 0.492–0.786) respectively, and pooled LR+ and LR– estimated were 1.989 (95% CI, 1.145–3.457) and 0.475 (95% CI, 0.247–0.915) respectively. The

**Table 2**

Results of the distribution of study design characteristics in 8 studies.

	Question about study design characteristic	Response		
		Yes	No	Unclear
1	Was the spectrum of patients representative of the patients who will receive the test in practice?	8	0	0
2	Were selection criteria clearly described?	8	0	0
3	Is the reference standard likely to correctly classify the target condition?	8	0	0
4	Is the time period between reference standard and index test short enough to be reasonable?	6	0	2
5	Did the whole sample or a random selection of the sample, receive verification using a reference standard of diagnosis?	8	0	0
6	Did patients receive the same reference standard regardless of the index test result?	8	0	0
7	Was the reference standard independent of the index test?	8	0	0
8	Were the reference standard results interpreted without knowledge of the results of the index test?	8	0	0
9	Were the index test results interpreted without knowledge of the results of the reference standard?	4	0	4
10	Was the execution of the reference standard described in sufficient detail to permit its replication?	8	0	0
11	Were the same clinical data available when test results were interpreted as would be available when the test is used in practice?	8	0	0
12	Was the execution of the index test described in sufficient detail to permit its replication?	8	0	0
13	Were uninterpretable/intermediate results reported?	8	0	0
14	Were withdrawals from the study explained?	8	0	0

Data were the numbers of responses from the QUADAS tool. The numbers indicated how many articles were assigned a score of “Yes”, “No” or “Unclear”.

SROC curve for SUV2:1 showed a median diagnostic accuracy. The AUC was 0.72 (95% CI, 0.68–0.76) (Fig. 2b).

## Discussion


18F-FDG PET-CT imaging is a useful, noninvasive method for staging tumors and detecting disease recurrence. Lines of studies have evidenced that the decrease of SUV2 is significantly correlated to percent of necrosis. Quarles et al. demonstrated that the change in 18F-FDG uptake was the strongest indicator for histological response to neoadjuvant chemotherapy through a meta-analysis of 19 studies [6], indicating that 18F-FDG PET-CT scan might be plausible for predicting tumor necrosis after treatment. The present meta-analysis explored the association between the 18F-FDG uptake and histological response after neoadjuvant chemotherapy focusing on a population with osteosarcoma. Since previous studies used an SUV2 cutoff value of 2.5 and an SUV2:1 cutoff value at 0.5 to discriminating histological response, we also selected the same

**Table 1**

Comparison of PET parameters in patients with osteosarcoma.

No.	Study	R/N*	PET	Design	Age(mean, median)	Sex(M/F)	Time interval after treatment(d)	Time interval before surgery(d)	Observer /Blinded	Reference
1	Cheon 2009	33/37	SUV2/SUV2:1	Prospective	14(5–59)	48/22	14	14	1/nm	[35]
2	Denecke 2010	17/10	SUV2	Prospective	13(3–18)	15/12	21	nm	2/blinded	[12]
3	Gaston 2011	16/15	SUV2	Retrospective	26.75 (18–62) 24.11 (14–38)	13/18	nm	nm	1/blinded	[10]
4	Hamada 2009	5/6	SUV2	Retrospective	27(10–68)	7/4	nm	nm	nm/nm	[34]
5	Hawkins 2002	5/9	SUV2	Unclear	13(6.2–19.2)	22/11	nm	18.2(2–84)	nm/nm	[9]
6	Huang 2006	2/8	SUV2	Prospective	19(4–47)	8/2	7	nm	nm/nm	[17]
7	Ye 2008	8/7	SUV2	Prospective	17(7–31)	9/6	8 (4–14)	12 (2–22)	2/blinded	[15]
8	Bajpai 2011	10/21	SUV2/SUV2:1	Prospective	17(5–66)	25/6	21–28	nm	2/blinded	[36]

N: numbers of patients that included, R/N: number of patients with necrosis  $\geq 90\%$ /number of patients with necrosis  $< 90\%$ , SUV: standardized uptake value, SUV2: standardized uptake value after chemotherapy, SUV 2:1: the ratio of standardized uptake values after chemotherapy (SUV2) to before (SUV1) chemotherapy, \*:for patient with pathological evaluation, nm: not mentioned.



**Figure 1.** The forest plots of sensitivity and specificity for SUV2  $\leq$  2.5 (a, b), SUV2:1  $\leq$  0.5 (c, d). Each solid represents each study in the meta-analysis. The pooled sensitivity and specificity for SUV2  $\leq$  2.5 were 0.734 and 0.864, for SUV2:1  $\leq$  0.5 were 0.690 and 0.653.

cutoff point [9,34]. The results showed that 18F-FDG PET-CT parameters, SUV2  $\leq$  2.5 and SUV2:1  $\leq$  0.5, were of predictive value for tumor response to chemotherapy and/or radiotherapy, and SUV2  $\leq$  2.5 exhibited better performance than SUV2:1  $\leq$  0.5 according to AUC of SROC (0.81–0.72).

According to widely accepted interpretation of AUC values, an AUC of 0.5 is classified as no value for diagnosis; 0.6–0.69 provides poor discrimination; 0.7–0.79 provides acceptable discrimination; 0.8–0.89 provides excellent discrimination and 0.9–1.0 provides outstanding discrimination. The AUC estimates for SUV2  $\leq$  2.5 was 0.81, while for SUV2:1  $\leq$  0.5 was 0.72, indicating a relatively high level of accuracy of SUV2  $\leq$  2.5 than SUV2:1  $\leq$  0.5 in predicting treatment response.

Since likelihood ratios are considered to be more clinically meaningful than SROC [37], LR+/- were also analyzed in this review. A good test should have LR+ above 5 and LR- below 0.2, even there is no absolute cut-off. Our results showed the value of LR+ for SUV2  $\leq$  2.5 and SUV2:1  $\leq$  0.5 were 5.397 and 1.989, suggesting that SUV2  $\leq$  2.5 is more valuable than SUV2:1  $\leq$  0.5 in discriminating tumor necrosis rate >90% after treatment. In the other hand, the value of LR- for SUV2  $\leq$  2.5 and SUV2:1  $\leq$  0.5 were 0.308 and 0.475, indicating that a negative examination result could not to rule out good response of neoadjuvant therapy.

Some limitations of this meta-analysis should be acknowledged. Firstly, although eleven studies were included in present analysis, the numbers of patients enrolled in these studies were relatively small, with a mean of 25 patients per study. Therefore, large number of patient enrollment would be more powerful to improve this statistical result. Secondly, there was no single reference

standard strategy for 18F-FDG PET-CT imaging. For example, the time intervals between neoadjuvant therapy and 18F-FDG PET-CT scan, and the scan to surgery were varied at most of the studies, different time window would inevitably make result conflicting between studies. Thirdly, since tumor volumes and time of 18F-FDG PET-CT scan are showed to influence scan results, these parameters should be adjusted in estimation. However, they are not taken in this analysis.

In conclusion, this meta-analysis suggests that 18F-FDG PET-CT parameters SUV2  $\leq$  2.5 and SUV2:1  $\leq$  0.5 are valuable predictor to assess the chemotherapy-induced tumor necrosis. However, more patients should be enrolled to improve statistical power and avoid clinical heterogeneity; moreover it will be necessary to define the prognostic assessment system in combination with other factors to develop the more precise strategies to improve discriminating ability for better treatment of patients.

#### Author contribution

Guarantor of the integrity of the study: Li Hongtao, Zhao Hui, Yao Yang, Shen Zan.

Study concepts: Li Hongtao, Shen Zan.

Study design: Yao Yang, Shen Zan, Zhao Hui.

Definition of intellectual content: Li Hongtao.

Literature research: Li Hongtao, Wang Zhiyu, Zheng Shuier.


Clinical studies: Li Hongtao, Zhao Hui.

Experimental studies: Li Hongtao.

Data acquisition: Li Hongtao, He Aina.

Data analysis: Li Hongtao, Wang Bingshun, Wang Xiaojin.





**Figure 2.** The summary receiver operating characteristic (SROC) curves for  $SUV2 \leq 2.5$  (a),  $SUV2:1 \leq 0.5$  (b) on a per-patient basis. Each cross represents each study in the meta-analysis. The size of the circle indicates the study size. The AUC estimate for  $SUV2 \leq 2.5$  and  $SUV2:1 \leq 0.5$  were 0.81 and 0.72 respectively. AUC: area under the curve.

Statistical analysis: Li Hongtao, Wang Xiaojin.

Manuscript preparation: Li Hongtao.

Manuscript editing: Li Hongtao, Sun Yuanjue.

Manuscript review: Li Hongtao, Shen Zan, Min Dalui.

### Conflict of interest statement

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the position presented in, or the review of, the manuscript entitled “18F-FDG positron emission tomography for the assessment of neoadjuvant therapy response in osteosarcomas: A meta-analysis”.

### Acknowledgments

This work was supported by NSFC (NO: 81102038, 81172105).

We thank Dr Ren Shengxiang from Shanghai Pulmonary Hospital at Tongji University, for his personal advice for this paper.

### References

- [1] Picci P. Osteosarcoma (osteogenic sarcoma). *Orphanet Journal of Rare Diseases* 2007;2:6.
- [2] Meyers PA, Gorlick R. Osteosarcoma. *Pediatric Clinics of North America* 1997; 44:973–89.
- [3] Bacci G, Longhi A, Versari M, Mercuri M, Briccoli A, Picci P. Prognostic factors for osteosarcoma of the extremity treated with neoadjuvant chemotherapy:

- 15-year experience in 789 patients treated at a single institution. *Cancer* 2006;106:1154–61.
- [4] Link MP, Goorin AM, Miser AW, Green AA, Pratt CB, Belasco JB, et al. The effect of adjuvant chemotherapy on relapse-free survival in patients with osteosarcoma of the extremity. *The New England Journal of Medicine* 1986;314:1600–6.
- [5] Davis AM, Bell RS, Goodwin PJ. Prognostic factors in osteosarcoma: a critical review. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* 1994;12:423–31.
- [6] Quarles van Ufford HM, van Tinteren H, Stroobants SG, Riphagen II, Hoekstra OS. Added value of baseline 18F-FDG uptake in serial 18F-FDG PET for evaluation of response of solid extracerebral tumors to systemic cytotoxic neoadjuvant treatment: a meta-analysis. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine* 2010;51:1507–16.
- [7] Schulte M, Brecht-Krauss D, Werner M, Hartwig E, Sarkar MR, Keppler P, et al. Evaluation of neoadjuvant therapy response of osteogenic sarcoma using FDG PET. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine* 1999;40:1637–43.
- [8] Hawkins DS, Schuetze SM, Butrynski JE, Rajendran JG, Vernon CB, Conrad 3rd EU, et al. [18F]Fluorodeoxyglucose positron emission tomography predicts outcome for Ewing sarcoma family of tumors. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* 2005; 23:8828–34.
- [9] Hawkins DS, Rajendran JG, Conrad 3rd EU, Bruckner JD, Eary JF. Evaluation of chemotherapy response in pediatric bone sarcomas by [F-18]-fluorodeoxy-D-glucose positron emission tomography. *Cancer* 2002;94:3277–84.
- [10] Gaston LL, Di Bella C, Slavin J, Hicks RJ, Choong PF. 18F-FDG PET response to neoadjuvant chemotherapy for Ewing sarcoma and osteosarcoma are different. *Skeletal Radiology* 2011;40:1007–15.
- [11] Franzius C, Sciuk J, Brinkschmidt C, Jurgens H, Schober O. Evaluation of chemotherapy response in primary bone tumors with F-18 FDG positron emission tomography compared with histologically assessed tumor necrosis. *Clinical Nuclear Medicine* 2000;25:874–81.
- [12] Denecke T, Hundsdoerfer P, Misch D, Steffen IG, Schonberger S, Furth C, et al. Assessment of histological response of paediatric bone sarcomas using FDG PET in comparison to morphological volume measurement and standardized MRI parameters. *European Journal of Nuclear Medicine and Molecular Imaging* 2010;37:1842–53.
- [13] Costelloe CM, Macapinlac HA, Madewell JE, Fitzgerald NE, Mawlawi OR, Rohren EM, et al. 18F-FDG PET/CT as an indicator of progression-free and overall survival in osteosarcoma. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine* 2009;50:340–7.
- [14] Nair N, Ali A, Green AA, Lamonica G, Alibazoglu H, Alibazoglu B, et al. Response of osteosarcoma to chemotherapy. Evaluation with F-18 FDG-PET Scans. *Clinical Positron Imaging: Official Journal of the Institute for Clinical P.E.T* 2000;3:79–83.
- [15] Ye Z, Zhu J, Tian M, Zhang H, Zhan H, Zhao C, et al. Response of osteogenic sarcoma to neoadjuvant therapy: evaluated by 18F-FDG-PET. *Annals of Nuclear Medicine* 2008;22:475–80.
- [16] Igaru A, Masamed R, Chawla SP, Menendez LR, Fedenko A, Conti PS. F-18 FDG PET and PET/CT evaluation of response to chemotherapy in bone and soft tissue sarcomas. *Clinical Nuclear Medicine* 2008;33:8–13.
- [17] Huang TL, Liu RS, Chen TH, Chen WY, Hsu HC, Hsu YC. Comparison between F-18-FDG positron emission tomography and histology for the assessment of tumor necrosis rates in primary osteosarcoma. *Journal of the Chinese Medical Association: JCMA* 2006;69:372–6.
- [18] Salzer-Kuntschik M, Delling G, Beron G, Sigmund R. Morphological grades of regression in osteosarcoma after polychemotherapy - study COSS 80. *Journal of Cancer Research and Clinical Oncology* 1983;106(Suppl.):21–4.
- [19] Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Medical Research Methodology* 2003;3:25.
- [20] Young H, Baum R, Cremerius U, Herholz K, Hoekstra O, Lammertsma AA, et al. Measurement of clinical and subclinical tumour response using [18F]-fluorodeoxyglucose and positron emission tomography: review and 1999 EORTC recommendations. European Organization for Research and Treatment of Cancer (EORTC) PET Study Group. *European Journal of Cancer* 1999;35: 1773–82.
- [21] Tewfik JN, Greene GS. Fluorine-18-deoxyglucose-positron emission tomography imaging with magnetic resonance and computed tomographic correlation in the evaluation of bone and soft-tissue sarcomas: a pictorial essay. *Current Problems in Diagnostic Radiology* 2008;37:178–88.
- [22] Reiman RE. PET tracers for osteosarcoma. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine* 2004;45:1424. author reply 1424–5.
- [23] Nanni C, Marzola MC, Rubello D, Fanti S. Positron emission tomography for the evaluation of soft-tissue sarcomas and bone sarcomas. *European Journal of Nuclear Medicine and Molecular Imaging* 2009;36:1940–3.
- [24] Brenner W, Bohuslavizki KH, Eary JF. PET imaging of osteosarcoma. *Journal of nuclear medicine: official publication. Society of Nuclear Medicine* 2003;44: 930–42.
- [25] Folpe AL, Lyles RH, Sprouse JT, Conrad 3rd EU, Eary JF. (F-18) fluorodeoxyglucose positron emission tomography as a predictor of pathologic grade and other prognostic variables in bone and soft tissue sarcoma. *Clinical Cancer Research: An Official Journal of the American Association for Cancer Research* 2000;6:1279–87.

- [26] Charest M, Hickeson M, Lisbona R, Novales-Diaz JA, Derbekyan V, Turcotte RE. FDG PET/CT imaging in primary osseous and soft tissue sarcomas: a retrospective review of 212 cases. *European Journal of Nuclear Medicine and Molecular Imaging* 2009;36:1944–51.
- [27] Kim MS, Lee SY, Cho WH, Song WS, Koh JS, Lee JA, et al. Tumor necrosis rate adjusted by tumor volume change is a better predictor of survival of localized osteosarcoma patients. *Annals of Surgical Oncology* 2008;15:906–14.
- [28] Costelloe CM, Raymond AK, Fitzgerald NE, Mawlawi OR, Nunez RF, Madewell JE, et al. Tumor necrosis in osteosarcoma: inclusion of the point of greatest metabolic activity from F-18 FDG PET/CT in the histopathologic analysis. *Skeletal Radiology* 2010;39:131–40.
- [29] Volker T, Denecke T, Steffen I, Misch D, Schonberger S, Plotkin M, et al. Positron emission tomography for staging of pediatric sarcoma patients: results of a prospective multicenter trial. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* 2007;25:5435–41.
- [30] Sato J, Yanagawa T, Dobashi Y, Yamaji T, Takagishi K, Watanabe H. Prognostic significance of 18F-FDG uptake in primary osteosarcoma after but not before chemotherapy: a possible association with autocrine motility factor/phosphoglucose isomerase expression. *Clinical Experimental Metastasis* 2008;25:427–35.
- [31] Benz MR, Evilevitch V, Allen-Auerbach MS, Eilber FC, Phelps ME, Czernin J, et al. Treatment monitoring by 18F-FDG PET/CT in patients with sarcomas: interobserver variability of quantitative parameters in treatment-induced changes in histopathologically responding and nonresponding tumors. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine* 2008;49:1038–46.
- [32] Jones DN, McCowage GB, Sostman HD, Brizel DM, Layfield L, Charles HC, et al. Monitoring of neoadjuvant therapy response of soft-tissue and musculoskeletal sarcoma using fluorine-18-FDG PET. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine* 1996;37:1438–44.
- [33] Franzius C, Bielack S, Flege S, Sciuk J, Jurgens H, Schober O. Prognostic significance of (18)F-FDG and (99m)Tc-methylene diphosphonate uptake in primary osteosarcoma. *Journal of nuclear medicine: official publication, Society of Nuclear Medicine* 2002;43:1012–7.
- [34] Hamada K, Tomita Y, Inoue A, Fujimoto T, Hashimoto N, Myoui A, et al. Evaluation of chemotherapy response in osteosarcoma with FDG-PET. *Annals of Nuclear Medicine* 2009;23:89–95.
- [35] Cheon GJ, Kim MS, Lee JA, Lee SY, Cho WH, Song WS, et al. Prediction model of chemotherapy response in osteosarcoma by 18F-FDG PET and MRI. *Journal of nuclear medicine: official publication, Society of Nuclear Medicine* 2009;50:1435–40.
- [36] Bajpai J, Kumar R, Sreenivas V, Sharma MC, Khan SA, Rastogi S, et al. Prediction of chemotherapy response by PET-CT in osteosarcoma: correlation with histologic necrosis. *Journal of Pediatric Hematology Oncology* 2011;33:e271–8.
- [37] Deeks JJ. Systematic reviews in health care: systematic reviews of evaluations of diagnostic and screening tests. *BMJ* 2001;323:157–62.