

WITH GRAPH PAPER

भारतीय गणनात्मक शिक्षा बोर्ड, दिल्ली  
गणित साहित्य केन्द्र परीक्षा (कक्षा १०वीं)  
परीक्षा प्रश्न-पत्र के अङ्गक २

Mathematics

041

I

Wednesday 18.03.2015

English

Code Number

65/2/c

Set Number

① ● ③ ④

Yes / No

40

B D H S C A

Yes / No

40

Each letter be written in one box and one box be left blank between each part of the answer. In case Candidates' Name exceeds 24 letters, write first 24 letters.

341-0892 4414974

## Section - 1

Q: 20

$$(1+x^2) \frac{dy}{dx} = e^{\tan^{-1}x} - y$$

$$(1+x^2) \frac{dy}{dx} + y = e^{\tan^{-1}x}$$

$$\frac{dy}{dx} + \frac{y}{1+x^2} = \frac{e^{\tan^{-1}x}}{1+x^2}$$

comparing it with

$$\frac{dy}{dx} + Py = Q \rightarrow \text{linear differential equation}$$

$$P = \frac{1}{1+x^2}$$

$$I \cdot O F = e^{\int P dx}$$

$$= e^{\int \frac{1}{1+x^2} dx}$$

$$I \cdot O F = e^{\tan^{-1}x}$$

$$\left\{ \int \frac{1}{x^2+1} dx = \tan^{-1}x \right\}$$

the diff equation becomes

$$y \text{ I.O.F} = \int Q \times \text{I.O.F} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{m \tan^{-1}x}}{1+x^2} \times e^{\tan^{-1}x} \, dx + c$$

$$y \times e^{\tan^{-1}x} = \int \frac{e^{(m+1)\tan^{-1}x}}{1+x^2} \, dx + c$$

but  $\tan^{-1}x = t$

$$\frac{1}{1+x^2} \, dx = dt$$

$$y e^{\tan^{-1}x} = \int e^{(m+1)t} \times dt + c$$

$$y e^{\tan^{-1}x} = \frac{e^{(m+1)t}}{(m+1)} + c$$

$$\text{after } y e^{\tan^{-1}x} = \frac{e^{(m+1)\tan^{-1}x}}{m+1} + c$$

when  $x=0$   $y=1$



$$y e^{\tan^{-1} x} = \frac{e^{(m+1)\tan^{-1} x}}{(m+1)} + c$$

when  $x=0$   $y=1$

$$1 \times e^{\tan^{-1} 0} = \frac{e^{(m+1)\tan^{-1} 0}}{(m+1)} + c$$

$$1 \times e^0 = \frac{e^{(m+1)0}}{m+1} + c \quad e^0 = 1$$

$$1 = \frac{1}{m+1} + c$$

$$1 - \frac{1}{m+1} = c$$

$$\frac{m+1-1}{m+1} = c$$

$$c = \frac{m}{m+1}$$

the equation is

$$\left[ y e^{\tan^{-1} x} = \frac{e^{(m+1)\tan^{-1} x}}{(1+m)} + \frac{m}{m+1} \right]$$

Q: 21.

$$f(x) = \sin^2 x - \cos x \quad x \in [0, \pi]$$

$$f'(x) = 2 \sin x \cos x - (-\sin x)$$

$$f'(x) = 2 \sin x \cos x + \sin x$$

$$\text{put } f'(x) = 0$$

$$\text{then } (2 \sin x \cos x + \sin x) = 0$$

$$\sin x (2 \cos x + 1) = 0$$

$$\sin x = 0 \quad \text{or} \quad \cos x = -\frac{1}{2}$$

$$x = 0, \pi \quad \text{or}$$

$$x = \frac{2\pi}{3}$$

$$\text{also } f(0) = [\sin(0)]^2 - \cos 0$$

$$= 0 - 1$$

$$= -1$$

$$f(\pi) = \sin^2 \pi - \cos \pi$$

$$= 0 - (-1)$$

$$= 1$$

$$f\left(\frac{2\pi}{3}\right) = \sin^2 \frac{2\pi}{3} - \cos \frac{2\pi}{3}$$

$$= \left(\frac{\sqrt{3}}{2}\right)^2 - \left(-\frac{1}{2}\right)$$

$$= \frac{3}{4} + \frac{1}{2} = \frac{10}{8} = \frac{5}{4}$$

both the extreme value are automatically included



$$f(0) = -1$$

$$f(1) = 1$$

$$f\left(\frac{20}{3}\right) = \frac{5}{4}$$

So Absolute maxima is  $\frac{5}{4}$  at  $x = \frac{20}{3}$

Absolute minima is  $-1$  at  $x = 0$

Q: 22

$$R = 1 + j + k + \lambda(1 - j + k)$$

$$R = 4j + 2k + \mu(2i - j + 3k)$$

These line are coplaner if they are parallel or they are intersecting

But these lines are not parallel.

So they are coplaner if shortest distance between them is zero.

$$a_1 = (1, 1, 1)$$

$$(a_2 - a_1) = (-1, 3, 1)$$

$$a_2 = (0, 4, 2)$$

$$b_1 = \langle 1, -1, 1 \rangle$$

$$b_2 = \langle 2, -1, 3 \rangle$$

$$\text{Shortest distance} = \frac{|(a_2 - a_1) \cdot (b_1 \times b_2)|}{|b_1 \times b_2|}$$

$$\text{Let us find } |(a_2 - a_1) \cdot (b_1 \times b_2)|$$

$$(a_2 - a_1) \cdot (b_1 \times b_2) = \begin{vmatrix} -1 & 3 & 1 \\ 1 & -1 & 1 \\ 2 & -1 & 3 \end{vmatrix}$$

(expanding by 1st row)

$$= -1(-3+1) - 3(3-2) + 1(-1+2)$$

$$= -1(-2) - 3(1) + 1$$

$$= 2 + 1 - 3$$

$$= 0$$

So shortest distance between the lines is zero  
So they are coplanar.



Let  $\langle a, b, c \rangle$  be the direction ratio of normal to the plane.

As the dot product of direction ratio of normal to plane and the direction ratio of line is 0.

$$\text{So } a - b + c = 0$$

$$2a - b + 3c = 0$$

$$\begin{array}{ccc|ccc} a & b & c & & & \\ -1 & 1 & 1 & & & -1 \\ -1 & 3 & 2 & & & -1 \end{array}$$

$$\frac{a}{-3+1} = \frac{b}{2-3} = \frac{c}{-1+2}$$

$$\text{So } \langle a, b, c \rangle = \langle -2, -1, 1 \rangle$$

So passing point of line is also the passing point of plane

$$\text{So passing point} = (1, 1, 1)$$



also equation becomes

$$-2(x-1) - 1(y-1) + 1(z-1) = 0$$

$$-2x + 2 - y + 1 + z - 1 = 0$$

$$\boxed{-2x - y + z + 2 = 0}$$

equation of plane.

Q: 23

$$f: W \rightarrow W$$

$$f(n) = \begin{cases} n-1 & \text{if } n \text{ is odd} \\ n+1 & \text{if } n \text{ is even} \end{cases}$$

prove that it is one-one.

let us suppose that  $n_1, n_2 \in W$

$$n_1 \neq n_2 \text{ but } f(n_1) = f(n_2)$$

(i) 1st case : if  $n_1$  &  $n_2$  are odd

$$f(n_1) = f(n_2)$$

$$n_1 - 1 = n_2 - 1$$

which is contradiction.

(ii) and case :  $n_1, n_2$  are even

$$f(n_1) = f(n_2)$$

$$n_1 + 1 = n_2 + 1$$

$$n_1 = n_2$$

which is contradiction.

(iii) and case if  $n_1$  is even &  $n_2$  is odd

$$f(n_1) = f(n_2)$$

$$n_1 + 1 = n_2 - 1$$

$$n_1 + 2 = n_2$$



even

odd

if we add 2 in an even no. then we get an even no. but  $n_2$  is odd which is again contradiction

So from these 3 point we see that

$\Rightarrow f(n_1) = f(n_2)$  only if  $n_1 = n_2$   
 $\Rightarrow f$  is one-one.



To prove that  $f$  is onto:

Let  $y \in \mathbb{N}$

and  $y$  is even

then  $y+1$  is odd and it belongs in vehicle no.

$$f(y+1) = y+1-1$$

$$= y$$

So for every  $y \in \mathbb{N}$  which is even there exist a preimage  $y+1$  which is odd and  $(y+1) \in \mathbb{N}$

Let  $y \in \mathbb{N}$

$y$  is odd

then  $y-1$  is even and  $y-1 \in \mathbb{N}$

$$f(y-1) = y-1+1$$

$$= y$$

So for every  $y \in \mathbb{N}$  which is odd there exist a preimage  $y-1 \in \mathbb{N}$  which is even.

So for all  $y \in \mathbb{N}$  there exist preimage in vehicle no. So  $f$  is onto.

$f$  is one-one and onto so it is invertible

To find  $f^{-1}(x)$

$$y = x - 1 \quad \text{if } x \text{ is odd}$$

$$y + 1 = x$$

$$\text{so } x = y + 1 \quad \text{if } y \text{ is even}$$

$$f^{-1}(x) = x + 1 \quad \text{if } x \text{ is even}$$

$$y = x + 1 \quad \text{if } x \text{ is even}$$

$$y - 1 = x \quad \text{if } y \text{ is odd here}$$


$$\text{so } f^{-1}(x) = x - 1 \quad \text{if } x \text{ is odd}$$

$$\text{so } f^{-1}(x) = \begin{cases} x - 1 & \text{if } x \text{ is odd} \\ x + 1 & \text{if } x \text{ is even} \end{cases}$$

$$[x \in W]$$

$$\text{so } [f^{-1} = f]$$



240

intersection point :

$$y = \sqrt{5-x^2}$$

$$y = x-1$$

$$\text{so } \sqrt{5-x^2} = (x-1)$$

$$5-x^2 = x^2 + 1 - 2x$$

$$2x^2 - 2x - 4 = 0$$

$$x^2 - x - 2 = 0$$

$$x^2 - 2x + x - 2 = 0$$

$$x(x-2) + 1(x-2) = 0$$

$$\text{so } x=2 \text{ or } x=-1$$

$$x=2$$

rejected as  $x \geq 1$

$$y = \sqrt{5-x^2}$$

$$y = -x+1$$

$$y \quad x < 1$$

$$5-x^2 = x^2 + 1 - 2x$$

$$2x \quad x = 2, 9 - 1$$

$$x = 2$$

$$[x = -1]$$

Rejected

Required Area

$$(A) = 2 \int_{-1}^2 \sqrt{5-x^2} dx - \left[ \int_{-1}^2 (-x+1) dx + \int_1^2 (x-1) dx \right]$$

$$A = \frac{x\sqrt{5-x^2}}{2} + \frac{5}{2} \sin^{-1} \frac{x}{\sqrt{5}} \Big|_{-1}^2 - \left[ \frac{-x^2+x}{2} \Big|_{-1}^2 + \frac{x^2-x}{2} \Big|_1^2 \right]$$

$$A = 1 \times 1 + \frac{5}{2} \sin^{-1} \frac{2}{\sqrt{5}} - \left( \frac{-1 \times 2}{2} + \frac{5}{2} \sin^{-1} \left( \frac{-1}{\sqrt{5}} \right) \right) - \left[ \frac{-1}{2} [2-1] + 2 + \frac{1}{2} [4-1] - 1 \right]$$



$$A = \frac{1+5 \sin^{-1} 2}{2} + \frac{1+5 \sin^{-1} 1}{2} - \left[ 0 + 2 + \frac{3}{2} - 1 \right]$$

$$A = \frac{2+5 \left[ \sin^{-1} 2 + \sin^{-1} 1 \right]}{2} - \frac{5}{2}$$

$$A = \frac{5 \times \frac{\pi}{2} + 2 - 5}{2}$$

$$A = \frac{5\pi}{4} - \frac{1}{2} \text{ } \cancel{\text{89 units}}$$

$$\left\{ \begin{aligned} & \frac{\sin^{-1} 2}{\sqrt{5}} + \frac{\sin^{-1} 1}{\sqrt{5}} \\ &= \frac{\sin^{-1} 2 + \sin^{-1} 1}{\sqrt{5}} \\ &= \frac{\pi}{2} \end{aligned} \right\}$$

$$\int_0^1 dx$$

25°

Let 6 positive integers = (1, 2, 3, 4, 5, 6)

Random Variable =  $X$  = larger of 2 numbers.

$$x \left| \begin{array}{c} 2 \\ 1 \end{array} \right.$$

2

$$\{4-1\}-1\}$$

sample space  $S = \{$

$$\begin{aligned} & (1,2) (1,3) (1,4) (1,5) (1,6) \\ & (2,1) (3,1) (4,1) (5,1) (6,1) \\ & (2,3) (2,4) (2,5) (2,6) (3,2) (4,2) (5,2) \\ & (6,2) (3,4) (3,5) (3,6) (4,3) (5,3) (6,3) \\ & (4,5) (4,6) (5,4) (6,4) (5,6) (6,5) \end{aligned}$$

X	P(X)	$X \cdot P_c$	$X^2 \cdot P_c$	Range
2	$\frac{2}{30} = \frac{1}{15}$	$\frac{2}{15}$	$\frac{4}{15}$	2+6
3	$\frac{4}{30} = \frac{2}{15}$	$\frac{6}{15}$	$\frac{18}{15}$	
4	$\frac{6}{30} = \frac{1}{5}$	$\frac{12}{15}$	$\frac{48}{15}$	
5	$\frac{8}{30} = \frac{4}{15}$	$\frac{20}{15}$	$\frac{100}{15}$	
6	$\frac{10}{30} = \frac{1}{3}$	$\frac{30}{15}$	$\frac{180}{15}$	

$$\sum X \cdot P_c = 70 \quad \sum X^2 \cdot P_c = 350$$

$$\text{Mean} = \frac{\sum X \cdot P_c}{n} = \frac{70}{15} = 4.66$$

$$\text{Variance} = \frac{\sum X^2 \cdot P_c - (\mu)^2}{n}$$

$$= \frac{350 - 70 \times 70}{15}$$

Handwritten calculations and corrections at the bottom of the page, including a circled '3' and various numerical scribbles.



$$\sigma^2 = \frac{70}{15} \left[ 5 - \frac{70}{15} \right]$$

$$= \frac{70}{15} \left[ \frac{75-70}{15} \right]$$

$$= \frac{70 \times 5}{15 \times 15}$$

$$= \frac{35}{15} = \frac{7}{3}$$


$$= \frac{14}{9}$$

$$\sigma^2 = 1.55$$

$$\boxed{\text{Variance} = 1.55}$$

### Section - B

Q.19



$$x^x + x^y + y^x = a^b$$

$\downarrow$     $\downarrow$     $\downarrow$   
 p   q   r

$$\frac{d}{dx} [p + q + r] = 0$$

$$\left[ \frac{dp}{dx} + \frac{dq}{dx} + \frac{dr}{dx} = 0 \right]$$

$$P = x^x$$

$$\log P = x \log x$$

$$\frac{1}{P} \frac{dP}{dx} = x \left[ \frac{1}{x} \right] + \log x$$

$$\frac{dP}{dx} = x^x (1 + \log x) \quad \text{--- (1)}$$

$$Q = x^y$$

$$\log Q = y \log x$$

$$\frac{1}{Q} \frac{dQ}{dx} = \left( \frac{y}{x} + \log x \frac{dy}{dx} \right)$$

$$\frac{dQ}{dx} = x^y \left[ \frac{y}{x} + \log x \frac{dy}{dx} \right] \quad \text{--- (2)}$$

$$Q = y^x$$

$$R = y^x$$

$$\log R = x \log y$$

$$\frac{1}{R} \frac{dR}{dx} = \frac{x}{y} \frac{dy}{dx} + \log y$$



$$\frac{dx}{dy} = y^x \left[ \frac{x}{y} \frac{dy}{dy} + \log y \right] \quad \text{--- (3)}$$

adding ① & ② & ③ we get

$$x^x + x^x \log x + x^{y-1} y + x^y \log x \frac{dy}{dx} + y^{x-1} x \frac{dy}{dx} + y^x \log y = 0$$

$$\frac{dy}{dx} = - \left[ x^x (1 + \log x) + x^{y-1} y + y^x \log y \right] \\ x^y \log x + y^{x-1} x$$

Q:18  $y = e^{ax} \cosh x$

$$\frac{dy}{dx} = e^{ax} (-\sinh x) b + \cosh x \times e^{ax} \times a$$

$$\frac{dy}{dx} = -b e^{ax} \sinh x + a y \Rightarrow \left[ \frac{-1}{b} \left( \frac{dy}{dx} - ay \right) = e^{ax} \sinh x \right] \quad \text{--- (1)}$$

$$\frac{d^2 y}{dx^2} = -b \left[ e^{ax} \cosh x \times b + \sinh x \times e^{ax} \times a \right] + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} = -b \left[ by - \frac{a}{b} \left( \frac{dy}{dx} - ay \right) \right] + a \frac{dy}{dx} \quad \text{from ①}$$

$$\frac{d^2y}{dx^2} = -b^2y + a \frac{dy}{dx} - a^2y + a \frac{dy}{dx}$$

$$\frac{d^2y}{dx^2} + (a^2 + b^2)y - 2a \frac{dy}{dx} = 0$$

hence proved.

$$\underline{\underline{26^o}} \quad Z = 5x + 2y$$

↓  
objective function to be made maximum and minimum

$$x - 2y \leq 2$$

$$3x + 2y \leq 12$$

$$-3x + 2y \leq 3$$

$$x \geq 0, y \geq 0$$



corresponding equations

$$x - 2y = 2$$

put  $(0, 0)$  in it

$$0 - 0 \leq 2$$

which is true so the region is towards

origin

$$3x + 2y \leq 12 \Rightarrow 3x + 2y = 12$$

put  $(0, 0)$

$$0 \leq 12 \text{ which is true}$$

so region towards origin

$$-3x + 2y = 3$$

put  $(0, 0)$

$$0 \leq 3$$

which is true

so region is towards origin

$x \geq 0$   $y \geq 0$  so it means I st quadrant.

at A (2,

at B

at C

at

intersection point of intersection point

$$-3x + 2y = 3$$

$$3x + 2y = 12$$

$$4y = 15$$

$$y = \frac{15}{4}$$

$$-3x + 2 \times \frac{15}{4} = 3$$

$$-3x = 3 - \frac{15}{2}$$

$$-x = 1 - \frac{5}{2}$$

$$-x = -\frac{3}{2}$$

$$x = \frac{3}{2}$$

$$Z = 5x + 2y$$

$$\text{at A } (2, 0) \quad Z = 10$$

$$\text{at B } (0, 0) \quad Z = 0$$

$$\text{at C } (0, \frac{3}{2}) \quad Z = 3$$

$$\text{at D } (\frac{3}{2}, \frac{15}{4}) \quad Z = \frac{15}{2} + \frac{15}{2} = 15$$

of:

$$x - 2y = 2$$

$$3x + 2y = 12$$

$$4x = 14$$

$$x = \frac{7}{2}$$

$$\frac{7}{2} - 2y = 2$$

$$\frac{7}{2} - 2 = 2y$$

$$\frac{3}{2} = 2y$$

$$y = \frac{3}{4}$$

at E

$$(\frac{7}{2}, \frac{3}{4})$$


$$Z = \frac{15}{2} + \frac{3}{2}$$

$$Z = 9$$



Q126

0 small box = 1 unit





As  $Z$  is maximum  
at  $D(3, 15)$

$Z = 15$

$Z$  is minimum at  $(0, 0)$

$[Z \leq 0]$

170

A =	Jams Mats Toys			
	30	12	70	
40	15	55		→ school Y
35	20	75		→ school Z
				3x2

B =	25	→ cost of Jams
	100	→ cost of Mats
	50	→ cost of Toys

2010th hand 200500 1000000



$$AB = \begin{bmatrix} 30 & 12 & 70 \\ 40 & 15 & 55 \\ 30 & 20 & 75 \end{bmatrix} \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix}$$

$$X = AB = \begin{bmatrix} 750 + 1200 + 3500 \\ 1000 + 1500 + 2750 \\ 875 + 2000 + 3750 \end{bmatrix}$$

$$X = \begin{bmatrix} 5450 \\ 5250 \\ 6625 \end{bmatrix} \begin{array}{l} \rightarrow \text{fund collected by school X} \\ \rightarrow \text{fund collected by school Y} \\ \rightarrow \text{fund collected by school Z} \end{array}$$

$$\begin{array}{l} \text{fund by } X = \text{Rs. } 5450 \\ \text{fund by } Y = \text{Rs. } 5250 \\ \text{fund by } Z = \text{Rs. } 6625 \end{array}$$

$$\text{Total fund} = \text{Rs. } 17325$$

They are helping victims and hence

the value of helping others is generated.

16°

$$I = \int \frac{x+3}{(x+5)^3} e^x dx$$

$$I = \int \frac{x+5}{(x+5)^3} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$I = \int \frac{1}{(x+5)^2} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

↓  
integrating this by parts is taken as 1st function

$$I = \frac{1}{(x+5)^2} e^x - \int \frac{d}{dx} \frac{1}{(x+5)^2} e^x dx - \int \frac{2}{(x+5)^3} e^x dx$$

$$I = \frac{1}{(x+5)^2} e^x + \frac{2}{(x+5)^3} e^x dx - \frac{2}{(x+5)^3} e^x dx$$



$$\text{So } I = \frac{e^x}{(x+5)^2} + C$$

15°

$$x = a \sin \omega t (1 + \cos 2t)$$

$$\frac{dx}{dt} = a [\dot{\sin \omega t} (-\sin \omega t)(2) + (1 + \cos 2t) \cos 2t \times 2]$$

$$\frac{dx}{dt} = 2a [\cos^2 \omega t + \cos 2t - \sin^2 2t]$$

$$\frac{dx}{dt} = 2a [\cos 4t + \cos 2t] \quad [\cos^2 x - \sin^2 x = \cos 2x]$$

$$y = b \cos \omega t (1 + \cos 2t)$$

$$y = b \cos \omega t - b \cos 2t$$

$$\frac{dy}{dt} = -b \sin \omega t \times 2 + b \times 2 \cos 2t \sin \omega t \times 2$$

$$\frac{dy}{dt} = 2b [-\sin \omega t + \sin 4t]$$

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx}$$

$$\frac{dy}{dx} = \frac{2b}{2a} \left[ \frac{\sin 4t - \sin 2t}{\cos 2t + \cos 4t} \right]$$

$$\text{at } t = \frac{\pi}{4}$$

$$\frac{dy}{dx} = \frac{b}{a} \left[ \frac{\sin \pi - \sin \frac{\pi}{2}}{\cos \frac{\pi}{2} + \cos \pi} \right]$$

$$= \frac{b}{a} \left[ \frac{0 - 1}{0 - 1} \right]$$

$$\left. \frac{dy}{dx} \right|_{t=\frac{\pi}{4}} = \frac{b}{a}$$



147

$$I = \frac{1}{2} \int_0^a \frac{x \sin^2 x}{\sin x + \cos x} dx \quad \text{--- (1)}$$

$$\left[ \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$I = \frac{1}{2} \int_0^a \frac{x \sin^2(a-x)}{\sin(a-x) + \cos(a-x)} dx$$

$$I = \frac{1}{2} \int_0^a \frac{\cos^2 x}{\cos x + \sin x} dx \quad \text{--- (2)}$$

adding (1) & (2) we get

$$2I = \frac{1}{2} \int_0^a \frac{x \sin^2 x + \cos^2 x}{\sin x + \cos x} dx$$

$$2I = \frac{1}{2} \int_0^a \frac{1}{(\sin x + \cos x)} dx$$

$$2I = \int_0^1 \frac{1 + \tan^2 x}{2 \tan x + 1 - \tan^2 x} dx$$

$$2I = \int_0^1 \frac{\sec^2 x}{2 \tan x + 1 - \tan^2 x} dx \quad \left\{ \begin{array}{l} \text{put } \sin = 2 \tan x \\ \text{and } \cos x = \frac{1 + \tan^2 x}{1 + \tan^2 x} \end{array} \right.$$

$$\text{put } \tan x = t$$

$$\sec^2 x \times \frac{1}{2} dx = dt$$

$$\begin{array}{l} x \rightarrow 0 \quad t \rightarrow 0 \\ \text{when } x \rightarrow \frac{\pi}{4} \quad t \rightarrow 1 \end{array}$$

$$2I = \int_0^1 \frac{dt}{2t + 1 - t^2}$$

$$I = \int_0^1 \frac{dt}{2t + 1 - t^2} \Rightarrow I = \int_0^1 \frac{dt}{2(t-1)^2}$$



$$I = \int_0^1 \frac{dx}{(\sqrt{x})^2 - (x-1)^2} \quad \int \frac{1}{a^2 - x^2} = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} + \frac{x-1}{\sqrt{x}}}{\sqrt{x} - \frac{x-1}{\sqrt{x}}} \right| \Bigg|_0^1$$

$$I = \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} + \frac{x-1}{\sqrt{x}}}{\sqrt{x} - \frac{x-1}{\sqrt{x}}} \right| - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} - 1}{\sqrt{x} + 1} \right|$$

$$I = \frac{1}{2\sqrt{2}} \log 1 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} - 1}{\sqrt{x} + 1} \right|$$

$$= 0 - \frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} - 1}{\sqrt{x} + 1} \right|$$

$$I = -\frac{1}{2\sqrt{2}} \log \left| \frac{\sqrt{x} - 1}{\sqrt{x} + 1} \right|$$

13°

$$\Delta = \begin{vmatrix} x+2 & x+6 & x-1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$R_1 \rightarrow R_1 + R_2 + R_3$$

$$\Delta = \begin{vmatrix} 3x+7 & 3x+7 & 3x+7 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$\Delta = 3x+7 \begin{vmatrix} 1 & 1 & 1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix}$$

$$C_1 \rightarrow C_1 - C_2 \quad C_2 \rightarrow C_2 - C_3$$

$$\Delta = (3x+7) \begin{vmatrix} 0 & 0 & 1 \\ 7 & -3 & x+2 \\ -3 & -4 & x+6 \end{vmatrix}$$



expanding by  $R_1$

$$\Delta = (3x+7) \begin{bmatrix} -28 & 9 \end{bmatrix}$$

$$\Delta = 0$$

$$\Rightarrow 3x+7=0$$

$$x = -\frac{7}{3}$$

120

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = A \cdot A =$$

$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 1+4+4 & 2+2+4 & 2+4+2 \\ 2+2+4 & 4+1+4 & 4+2+2 \\ 2+4+2 & 4+2+2 & 4+4+1 \end{bmatrix}$$

$$A^2 = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix}$$

$$A^2 - 4A - 5I = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 9 \\ 8 & 8 & 9 \end{bmatrix} + \begin{bmatrix} -4 & -8 & -8 \\ -8 & -4 & -8 \\ -8 & -8 & -4 \end{bmatrix} + \begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{bmatrix}$$

$$A^2 - 4A - 5I =$$

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = 0$$

$$\text{Hence } A^2 - 4A - 5I = 0$$

Hence proved.



$$A^2 - 4A - 5I = 0$$

pre-multiplying by  $A^{-1}$

$$A^{-1}AA - 4A^{-1}A - 5A^{-1}I = 0$$

$$A^{-1}A = I$$

$$IA - 4I - 5A^{-1} = 0$$

$$A - 4I - 5A^{-1} = 0$$

$$IA = A$$

$$A - 4I = 5A^{-1}$$

$$A^{-1} = \frac{A - 4I}{5}$$

5

$$A^{-1} = \frac{1}{5} [A - 4I]$$

$$5A^{-1} =$$

$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} + \begin{bmatrix} -4 & 0 & 0 \\ 0 & -4 & 0 \\ 0 & 0 & -4 \end{bmatrix}$$

$$5A^{-1} =$$

$$\begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

$$\det A^{-1} = \frac{1}{5} \begin{vmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{vmatrix}$$

11.

$$\sin^{-1}(1-x) - 2 \sin^{-1} x = \frac{\pi}{2}$$

$$\sin^{-1}(1-x) = \frac{\pi}{2} + 2 \sin^{-1} x$$

$$(1-x) = \sin\left(\frac{\pi}{2} + 2 \sin^{-1} x\right)$$

$$(1-x) = \cos(2 \sin^{-1} x)$$

$$2 \sin^{-1} x = 0$$

$$x = \sin 0$$

$$\cos 2\theta = 1 - 2 \sin^2 \theta$$

$$\cos 2\theta = 1 - 2x^2$$

$\Rightarrow$

$$1-x = \cos 2\theta$$

$$1-x = 1-2x^2$$

$$\Rightarrow 2x^2 - x = 0$$

$$x(2x-1) = 0$$



$$x=0 \text{ or } x=\frac{1}{2}$$

put  $x=\frac{1}{2}$  in equation

$$x \sin \frac{1}{2} - 2x \sin \frac{1}{2}$$

$$= \frac{1}{6} - 2 \times \frac{1}{6}$$

$$\neq \frac{1}{2}$$

$$\text{So } x \neq \frac{1}{2}$$

$$\text{So } \boxed{x=0}$$

passing point of line =  $(492, 2)$

since it is 11 to line

direction ratio of line  $\langle 2, 13, 6 \rangle$


the equation of line

$$\frac{x-4}{2} = \frac{y-2}{3} = \frac{z-2}{6} = \lambda$$

general point on line

$$(2\lambda+4, 3\lambda+2, 6\lambda+2)$$

$$P(1, 2, 3)$$



Q be the foot of  $LP$

$PQ$  is  $\perp$  to line

the dot product of direction ratios of line and  $PQ$  is 0

$$\text{direction ratios of } Q = \langle 2\lambda+3, 3\lambda, 6\lambda-1 \rangle$$



also according to question:

$$2(2\lambda + 3) + (3\lambda)3 + 6(6\lambda - 1) = 0$$

$$4\lambda + 6 + 9\lambda + 36\lambda - 6 = 0$$

$$\boxed{\lambda = 0}$$

also point  $Q = (4, 2, 9, 2)$

also for distance

$$PQ = \sqrt{(4-1)^2 + (2-2)^2 + (2-3)^2}$$

$$= \sqrt{(3)^2 + (0)^2 + (-1)^2}$$

$$= \sqrt{9+1}$$

$$\boxed{\text{length of } PQ = \sqrt{10} \text{ unit}}$$

Qo  
~~AB~~  
 Are A, B, C, D are coplanar?

Ans.  $\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = 0$   
 Triple product is 0.

$$\vec{AB} = 1\hat{i} + (x-1)\hat{j} + 4\hat{k}$$

$$\vec{BC} = 0\hat{i} + (1-x)\hat{j} - 7\hat{k}$$

$$\vec{CD} = 2\hat{i} + 3\hat{j} + \hat{k}$$

$$\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = \begin{vmatrix} 1 & x-1 & 4 \\ 0 & 1-x & -7 \\ 2 & 3 & 1 \end{vmatrix} = 0$$

Expanding by  $R_1$

$$1(1-x+21) - (x-1)14 + 4(2(x-1)) = 0$$

$$22-x - 14x + 14 + 8x - 8 = 0$$

$$-7x = -28 \quad \boxed{x=4}$$



8°

$P(\text{probability of success}) = \frac{1}{2}$   
 i.e. that head comes

$Q(\text{probability of failure}) = \frac{1}{2}$   
 i.e. that tail comes

Let the coin be tossed  $n$  times

this event follows the conditions of Bernoulli trial

$X$  be the random variable = no. of heads

$$P(X \geq 1) = P(1) + \dots + P(n)$$

$$P(X \geq 1) = 1 - P(0) \\ = 1 - {}^nC_0 \times \left(\frac{1}{2}\right)^0 \left(\frac{1}{2}\right)^n$$

$P(X \geq 8)$  should be more than 80%.

$$P(X \geq 1) > \frac{8}{10} \\ \frac{86}{100} < 1 - \left(\frac{1}{2}\right)^n$$

$$1 - \left(\frac{1}{2}\right)^n > \frac{8}{10}$$

$$\left(1 - \frac{8}{10}\right)^n > \left(\frac{1}{2}\right)^n$$

$$\frac{18}{5} > \left(\frac{1}{2}\right)^n$$

$$5 < 2^n$$

$$n=3$$

follow the condition  
So the coin should be tossed  
at least 3 times.

Ex.

$$I = \int \frac{x^2}{x^4 + x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^4 + 2x^2 - x^2 - 2} dx$$

$$I = \int \frac{x^2}{x^2(x^2 + 2) - 1(x^2 + 2)} dx$$



$$I = \int \frac{x^2}{(x^2-1)(x^2+2)} dx$$

put  $x^2 = t$

then  $\frac{t}{(t-1)(t+2)} = \frac{A}{t-1} + \frac{B}{t+2}$

$$t = A(t+2) + B(t-1)$$

put  $t=1$

$$1 = A \times 3$$

$$\boxed{A = \frac{1}{3}}$$

put  $t=-2$

$$-2 = -3B$$

$$\boxed{B = \frac{2}{3}}$$

$$I = \int \frac{1}{3} \frac{dx}{(t-1)} + \frac{2}{3} \frac{dx}{(t+2)}$$

$$I = \int \frac{1}{3} \frac{dx}{(x^2-1)} + \frac{2}{3} \int \frac{dx}{(x^2+2)}$$

$$\int \frac{1}{x^2-a^2} \frac{dx}{2a} \log \left| \frac{x-a}{x+a} \right|$$

$$I = \frac{1}{3} \times \frac{1}{2} \log \left| \frac{x-1}{x+1} \right| + \frac{2}{3} \times \frac{1}{\sqrt{2}} \tan^{-1} \frac{x}{\sqrt{2}} + C$$

$$\int \frac{1}{x^2+a^2} \frac{dx}{a} \tan^{-1} \frac{x}{a}$$

$$I = \frac{1}{6} \log \left| \frac{x-1}{x+1} \right| + \frac{\sqrt{2}}{3} \tan^{-1} \frac{x}{\sqrt{2}} + C$$



## Section - A

Q: 6

equation of plane

$$6x - 3y + 2z - 4 = 0$$

$$\text{distance} = \frac{|6 \times 2 - 3 \times 5 + 2(-3) - 4|}{\sqrt{36 + 9 + 4}}$$

$$= \frac{|12 - 15 - 6 - 4|}{\sqrt{49}}$$

$$= \frac{|12 - 15 - 10|}{7}$$

$$\text{distance} = \frac{13}{7} \text{ units}$$

Q: 5

$$\vec{a} = \hat{i} - \hat{j}$$

$$|\vec{a}| = \sqrt{2}$$

$$\vec{b} = \hat{j} - \hat{k}$$

$$|\vec{b}| = \sqrt{2}$$

0902

Fictitious Roll No.  
(To be entered by Board)

4474874

आपका उत्तर लिखें  
वे न लिखें

Please do not write your  
Roll Number on this Answer-book

आपका उत्तर लिखें  
वे न लिखें

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$$

$$(\hat{i} - \hat{j}) \cdot (\hat{j} - \hat{k}) = \sqrt{2} \times \sqrt{2} \cos \theta$$

$$\frac{-1}{2} = \cos \theta$$

$$\cos \theta = \frac{2\pi}{3}$$

angle between vectors =  $\frac{2\pi}{3}$

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 3 \\ 3 & 5 & -2 \end{vmatrix}$$

$$\vec{a} \times \vec{b} = \hat{i}(-2-15) - \hat{j}(-4-9) + \hat{k}(10-3)$$

$$\vec{a} \times \vec{b} = -17\hat{i} + 13\hat{j} + 7\hat{k}$$



$$|a \times b| = \sqrt{(-17)^2 + (13)^2 + (7)^2}$$

$$= \sqrt{289 + 169 + 49}$$

$$= \sqrt{507}$$

$$|a \times b| = 13\sqrt{3}$$

Q. 30

$$x \log x \frac{dy}{dx} + y = 2 \log x$$

$$\frac{dy}{dx} + \frac{y}{x \log x} = 2 \log x$$

compare it with  $\frac{dy}{dx} + P_y = Q$

$$I.F. =$$

$$e^{\int P dx}$$

$$= e^{\int \frac{1}{x \log x} dx}$$

$$\text{put } \log x = t$$

$$\frac{1}{x} dx = dt$$

$$I \circ F = e^{\int \frac{dt}{t}}$$

$$= e^{\log |t|}$$

$$= t$$

$$I \circ F = \log x$$

20

general equation of family of lines passing through origin

$$y = mx$$

$$m = \frac{y}{x}$$

$$\frac{dy}{dx} = m$$

$$\frac{dy}{dx} = \frac{y}{x}$$

$$x \frac{dy}{dx} - y = 0$$



$$A_{12} = e^{2 \times 1 \times 2} \sin 2x$$

$$A_{12} = e^{2x} \sin 2x$$

Done as per  
 CRSE  
 Monday  
 11/07/2022

Calculation

XII - U