

Some ideas about Tomography

Todd Quinto

Tufts University

Todd.Quinto@tufts.edu
<http://www.tufts.edu/~equinto>

Key ideas:

1. *Algorithms to reconstruct features of objects can be effective and easier than inverting the forward operator.*

Key ideas:

1. *Algorithms to reconstruct features of objects can be effective and easier than inverting the forward operator.*
 - ▶ Reconstruction algorithms can be simpler (Lambda CT).

Key ideas:

1. *Algorithms to reconstruct features of objects can be effective and easier than inverting the forward operator.*
 - ▶ Reconstruction algorithms can be simpler (Lambda CT).
 - ▶ Limited data X-ray CT inversion is highly ill-posed, but singularity reconstruction (for visible singularities) is easy and stable [FFRS, KLM, NC, Q1993, Q2005].

Key ideas:

1. *Algorithms to reconstruct features of objects can be effective and easier than inverting the forward operator.*
 - ▶ Reconstruction algorithms can be simpler (Lambda CT).
 - ▶ Limited data X-ray CT inversion is highly ill-posed, but singularity reconstruction (for visible singularities) is easy and stable [FFRS, KLM, NC, Q1993, Q2005].
2. *Microlocal analysis (pure math) can show what features are stably visible from tomographic data.*


Key ideas:

1. *Algorithms to reconstruct features of objects can be effective and easier than inverting the forward operator.*
 - ▶ Reconstruction algorithms can be simpler (Lambda CT).
 - ▶ Limited data X-ray CT inversion is highly ill-posed, but singularity reconstruction (for visible singularities) is easy and stable [FFRS, KLM, NC, Q1993, Q2005].
2. *Microlocal analysis (pure math) can show what features are stably visible from tomographic data.*
 - ▶ It provides the relation between singularities of an object and those of its tomographic data ([Q 1993], [NC]).

Radon Transform:

$$L(\theta, p) = \{x \in \mathbb{R}^2 \mid x \cdot \bar{\theta} = p\} \quad \bar{\theta} = (\cos \theta, \sin \theta).$$

$$x \in L(\theta, x \cdot \bar{\theta}).$$



$$Rf(\theta, p) = \int_{x \in L(\theta, p)} f(x) ds$$

•
•
•

Notation: $f = f(x), \quad g = g(\theta, p)$

Dual Radon Transform: $R^*g(x) = \int_0^{2\pi} g(\theta, x \cdot \bar{\theta}) d\theta$

Notation: $f = f(x)$, $g = g(\theta, p)$

Dual Radon Transform: $R^*g(x) = \int_0^{2\pi} g(\theta, x \cdot \bar{\theta}) d\theta$

Fourier Transform: $\mathcal{F}f(y) = \frac{1}{2\pi} \int_{x \in \mathbb{R}^2} e^{-ix \cdot y} f(x) dx$

1-D Fourier Transform: $\mathcal{F}_p g(\theta, \tau) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-ip\tau} g(\theta, p) dp$

Notation: $f = f(x)$, $g = g(\theta, p)$

Dual Radon Transform: $R^*g(x) = \int_0^{2\pi} g(\theta, x \cdot \bar{\theta}) d\theta$

Fourier Transform: $\mathcal{F}f(y) = \frac{1}{2\pi} \int_{x \in \mathbb{R}^2} e^{-ix \cdot y} f(x) dx$

1-D Fourier Transform: $\mathcal{F}_p g(\theta, \tau) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-ip\tau} g(\theta, p) dp$

Lambda Operators:

$$\Lambda_p g(\theta, p) = \sqrt{-\frac{d^2}{dp^2}} g = \frac{1}{\sqrt{2\pi}} \int_{\tau=-\infty}^{\infty} e^{ip\tau} |\tau| (\mathcal{F}_p g)(\theta, \tau) d\tau,$$

$$\Lambda_x f = \sqrt{-\Delta} f = \frac{1}{2\pi} \int_{\xi \in \mathbb{R}^2} e^{ix \cdot \xi} |\xi| \hat{f}(\xi) d\xi.$$

Filtered Back Projection (FBP):

Filtered Back Projection Inversion formula:

$$f = \frac{1}{4\pi} R^* \Lambda_p R f$$

Filtered Back Projection (FBP):

Filtered Back Projection Inversion formula:

$$f = \frac{1}{4\pi} R^* \Lambda_p R f$$

- ▶ FBP reconstructs f .

Filtered Back Projection (FBP):

Filtered Back Projection Inversion formula:

$$f = \frac{1}{4\pi} R^* \Lambda_p R f$$

- ▶ FBP reconstructs f .
- ▶ Reconstruction **is not** local

Filtered Back Projection (FBP):

Filtered Back Projection Inversion formula:

$$f = \frac{1}{4\pi} R^* \Lambda_p R f \approx \frac{1}{4\pi} R^* (\varphi(p) *_p R f)$$

- ▶ FBP reconstructs f .
- ▶ Reconstruction **is not** local as the approximate to Λ_p , $\phi(p)$, is nonzero “everywhere.”

Lambda CT [FFRS]:

Lambda reconstruction:

$$\Lambda_x f = \frac{1}{4\pi} R^* \left(-d^2 / dp^2 \right) R f$$

Lambda CT [FFRS]:

Lambda reconstruction:

$$\Lambda_x f = \frac{1}{4\pi} R^* \left(-d^2 / dp^2 \right) R f$$

- ▶ The algorithm replaces Λ_p by $-d^2 / dp^2$ and it reconstructs $\Lambda_x f$ **not** f .

Lambda CT [FFRS]:

Lambda reconstruction:

$$\Lambda_x f = \frac{1}{4\pi} R^* \left(-d^2 / dp^2 \right) R f$$

- ▶ The algorithm replaces Λ_p by $-d^2 / dp^2$ and it reconstructs $\Lambda_x f$ **not** f .
- ▶ Reconstruction **is** local.

Lambda CT [FFRS]:

Lambda reconstruction:

$$\Lambda_x f = \frac{1}{4\pi} R^* \left(-d^2 / dp^2 \right) R f$$

- ▶ The algorithm replaces Λ_p by $-d^2 / dp^2$ and it reconstructs $\Lambda_x f$ **not** f .
- ▶ Reconstruction **is** local.
- ▶ Reconstructions look great (limited data, too)!

Lambda CT [FFRS]:

Lambda reconstruction:

$$\Lambda_x f = \frac{1}{4\pi} R^* \left(-d^2 / dp^2 \right) R f = \frac{1}{2\pi} \int_{\xi \in \mathbb{R}^2} e^{ix \cdot \xi} |\xi| \hat{f}(\xi) d\xi$$

- ▶ The algorithm replaces Λ_p by $-d^2 / dp^2$ and it reconstructs $\Lambda_x f$ **not** f .
- ▶ Reconstruction **is** local.
- ▶ Reconstructions look great (limited data, too)!
- ▶ **Key point:** Λ_x emphasizes singularities of f

(Λ_x is an elliptic Ψ DO.)