

Contents

1	Introduction	1
1.1	Curve Fitting	1
1.2	Probability Theory	2

Chapter 1

Introduction

1.1 Curve Fitting

Problem 1

This can be solved by substituting the definition of:

$$y(x, \mathbf{w}) = \sum_{j=0}^M w_j x^j$$

into the error function and then taking the derivative.

$$\begin{aligned} E(\mathbf{w}) &= \frac{1}{2} \sum_{n=1}^N (y(x, \mathbf{w}) - t_n)^2 \\ &= \frac{1}{2} \sum_{n=1}^N \left(\sum_{j=0}^M w_j x^j - t_n \right)^2 && \text{Substitute} \\ \frac{dE(\mathbf{w})}{dw_i} &= \sum_{n=1}^N \left(\left(\sum_{j=0}^M w_j x^j - t_n \right) x^i \right) && \text{Take the derivative} \\ 0 &= \sum_{n=1}^N \left(\left(\sum_{j=0}^M w_j x^j - t_n \right) x^i \right) && \text{Set derivative to 0} \\ 0 &= \sum_{n=1}^N \left(\sum_{j=0}^M w_j x^j x^i - t_n x^i \right) && \text{Set derivative to 0} \\ \sum_{n=1}^N t_n x^i &= \sum_{n=1}^N \sum_{j=0}^M w_j x^j x^i \\ \sum_{n=1}^N t_n x^i &= \sum_{n=1}^N \sum_{j=0}^M w_j x^{i+j} \end{aligned}$$

Problem 2

This is solved in almost the same way we just have one additional term for the regularization so:

$$\begin{aligned}
E(\mathbf{w}) &= \frac{1}{2} \sum_{n=1}^N (y(x, \mathbf{w}) - t_n)^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2 \\
&= \frac{1}{2} \sum_{n=1}^N \left(\sum_{j=0}^M w_j x^j - t_n \right)^2 + \frac{\lambda}{2} \|\mathbf{w}\|^2 \\
\frac{dE(\mathbf{w})}{dw_i} &= \sum_{n=1}^N \left(\left(\sum_{j=0}^M w_j x^j - t_n \right) x^i \right) + \lambda w_i \\
-\lambda w_i &= \sum_{n=1}^N \left(\left(\sum_{j=0}^M w_j x^j - t_n \right) x^i \right) + \lambda w_i \\
-\lambda w_i &= \sum_{n=1}^N \left(\sum_{j=0}^M w_j x^j x^i - t_n x^i \right) \\
\sum_{n=1}^N t_n x^i - \lambda w_i &= \sum_{n=1}^N \sum_{j=0}^M w_j x^j x^i \\
\sum_{n=1}^N t_n x^i - \lambda w_i &= \sum_{n=1}^N \sum_{j=0}^M w_j x^{i+j}
\end{aligned}$$

1.2 Probability Theory

Problem 3 The Probability of Selecting an Apple can be decomposed as:

$$\begin{aligned}
\mathcal{P}(\text{apple}) &= \mathcal{P}(\text{apple}, \text{red}) + \mathcal{P}(\text{apple}, \text{blue}) + \mathcal{P}(\text{apple}, \text{green}) \\
&= \mathcal{P}(\text{apple}|\text{red})\mathcal{P}(\text{red}) + \mathcal{P}(\text{apple}|\text{blue})\mathcal{P}(\text{blue}) + \mathcal{P}(\text{apple}|\text{green})\mathcal{P}(\text{green}) \\
&= (.3)(.2) + (.5)(.2) + (.3)(.6) \\
&= .34
\end{aligned}$$

The probability that observing an orange came from the green box

can be solved using Bayes rule:

$$\begin{aligned}\mathcal{P}(\text{green}|\text{orange}) &= \frac{\mathcal{P}(\text{orange}|\text{green})\mathcal{P}(\text{green})}{\mathcal{P}(\text{orange})} \\ &= \frac{(.3)(.6)}{.66} \\ &= .27\end{aligned}$$

Problem 5

$$\begin{aligned}\text{Var}[X] &= \mathbb{E}[(X - \mathbb{E}[X])^2] \\ &= \mathbb{E}[X^2 - 2X\mathbb{E}[X] + \mathbb{E}[X]^2] && \text{Distributive Law} \\ &= \mathbb{E}[X^2] + \mathbb{E}[-2X\mathbb{E}[X]] + \mathbb{E}[\mathbb{E}[X]^2] && \text{Linearity of } \mathbb{E}[X] \\ &= \mathbb{E}[X^2] + -2\mathbb{E}[X\mathbb{E}[X]] + \mathbb{E}[X]^2 && \mathbb{E}[\alpha X] = \alpha\mathbb{E}[X] \\ &= \mathbb{E}[X^2] + -2\mathbb{E}[X]^2 + \mathbb{E}[X]^2 && \mathbb{E}[X] \text{ is just another constant} \\ &= \mathbb{E}[X^2] - \mathbb{E}[X]^2\end{aligned}$$

Problem 6

$\text{Cov}[X, Y] = \mathbb{E}[X, Y] - \mathbb{E}[X]\mathbb{E}[Y]$ But because
 $X \perp Y \Rightarrow \mathbb{E}[X, Y] = \mathbb{E}[X]\mathbb{E}[Y] \Rightarrow \text{Cov}[X, Y] = 0$

Problem 7

$$\begin{aligned}
I^2 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp \left(-\frac{1}{2\sigma^2}x^2 - \frac{1}{2\sigma^2}y^2 \right) dx dy \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \exp \left(-\frac{1}{2\sigma^2}(x^2 + y^2) \right) dx dy \\
&= \int_0^{\infty} \int_0^{2\pi} \exp \left(-\frac{1}{2\sigma^2} [r^2 \cos^2(\theta) + r^2 \sin^2(\theta)] \right) r d\theta dr \\
&= \int_0^{\infty} \int_0^{2\pi} \exp \left(-\frac{r^2}{2\sigma^2} \right) r d\theta dr \\
&= 2\pi \int_0^{\infty} \exp \left(-\frac{r^2}{2\sigma^2} \right) r dr \\
&= 2\pi \int_0^{\infty} \exp \left(-\frac{u}{2\sigma^2} \right) \frac{1}{2} du \\
&= -2\pi\sigma^2 \exp \left(-\frac{u}{2\sigma^2} \right) \Big|_0^{\infty} \\
&= 2\pi\sigma^2
\end{aligned}$$

Now we just need to show that this normalizes the Gaussian. Take $y = x - \mu$ then

$$\begin{aligned}
\mathcal{N}(x|\mu, \sigma^2) &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left(-\frac{1}{2\sigma^2}(x - \mu)^2 \right) dx \\
&= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left(-\frac{1}{2\sigma^2}y^2 \right) dy
\end{aligned}$$

Problem 8

$$\begin{aligned}
\mathbb{E}[x] &= \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma) x dx \\
&= \int_{-\infty}^{\infty} \frac{x}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) dx \\
&= \frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{\infty} (y+\mu) \exp\left(-\frac{1}{2\sigma^2}y^2\right) dy \quad y = x - \mu
\end{aligned}$$

Now split this into the sum of two integrals. The second integral has the form:

$$\frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{\infty} \mu \exp\left(-\frac{1}{2\sigma^2}y^2\right) dy$$

Which is just a normalized Gaussian times μ so this is just μ . Now since we are trying to prove that this equals μ I'm fairly confident that the left integral will go to zero. Let's try and prove this.

$$\frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{\infty} y \exp\left(-\frac{1}{2\sigma^2}y^2\right) dy$$

Well if we visualize this function it looks odd which would imply that the integral from $[-\infty, \infty]$ is 0. Let's prove that it is odd quickly. We want to show that $f(x) = -f(-x)$ where $f(x) = y \exp\left(-\frac{1}{2\sigma^2}y^2\right)$

$$-(-y) \exp\left(-\frac{1}{2\sigma^2}(-y)^2\right) = y \exp\left(-\frac{1}{2\sigma^2}y^2\right)$$

Thus our integral in question is odd and evaluates to 0 yielding μ as the desired result.

Next we need to show that:

$$\mathbb{E} [x^2] = \int_{-\infty}^{\infty} \mathcal{N}(x|\mu, \sigma) x^2 dx = \mu^2 + \sigma^2$$

To do this differentiate both sides by σ^2 as follows:

$$\begin{aligned} \frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) dx &= 1 \\ \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) dx &= \sqrt{2\pi\sigma^2} \\ \int_{-\infty}^{\infty} \frac{d}{d\sigma^2} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) dx &= \frac{d}{d\sigma^2} \sqrt{2\pi\sigma^2} \\ \frac{1}{2\sigma^4} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) (x-\mu)^2 dx &= \frac{\pi}{\sqrt{(2\pi\sigma^2)}} \\ \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) (x-\mu)^2 dx &= \frac{2\pi\sigma^4}{\sqrt{(2\pi\sigma^2)}} \\ \frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2\sigma^2}(x-\mu)^2\right) (x-\mu)^2 dx &= \sigma^2 \\ \text{Var} [x] &= \sigma^2 \end{aligned}$$

The left hand side of this equation is just the definition of variance. Now to complete the proof of (1.50) just use the alternate formulation of variance:

$$\begin{aligned}
\mathbb{E}[(x - \mu)^2] &= \mathbb{E}[x^2] - \mathbb{E}[x]^2 \\
&= \mathbb{E}[x^2] - \mu^2 && \text{By part 1} \\
\sigma^2 &= \mathbb{E}[x^2] - \mu^2 && \text{By above} \\
\mathbb{E}[x^2] &= \sigma^2 + \mu^2
\end{aligned}$$

Problem 9

$$\begin{aligned}
\mathcal{N}(x|\mu, \sigma^2) &= \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(x - \mu)^2\right) \\
\frac{d\mathcal{N}(x|\mu, \sigma^2)}{dx} &= -\frac{1}{\sigma^2} \mathcal{N}(x|\mu, \sigma^2)(x - \mu) \\
0 &= \mathcal{N}(x|\mu, \sigma^2)(x - \mu)
\end{aligned}$$

But since $\mathcal{N}(x|\mu, \sigma^2)(x - \mu) > 0$ the only term that matters is $(x - \mu)$, which goes to 0 at $x = \mu$

The proof for the multivariate case is almost identical and is omitted

Problem 10

$$\begin{aligned}
\log p(\mathbf{x}|\mu, \sigma^2) &= -\frac{1}{\sqrt{2\pi\sigma^2}} \sum_{n=1}^N (x_n - \mu)^2 - \frac{N}{2} \log \sigma^2 - \frac{N}{2} \log(2\pi) \\
\frac{d \log p(\mathbf{x}|\mu, \sigma^2)}{d\mu} &= -\frac{1}{\sqrt{\pi\sigma^2}} \sum_{n=1}^N (x_n - \mu) \\
0 &= \sum_{n=1}^N (x_n - \mu) \\
&= \sum_{n=1}^N x_n - \sum_{n=1}^N \mu \\
\mu &= \sum_{n=1}^N x_n
\end{aligned}$$

Problem 11