

Over a GCD Domain – Part I

One of our big questions is to what extent the structure of R is reflected in the structure of $R[x]$; if R has more “technology” available, perhaps this can be used to say interesting things about the polynomials over R . In this section we will see that this is indeed the case if R is a GCD domain.

Definition 1 (Content of a polynomial). *Let R be a GCD domain and let $p(x) \in R[x]$ be a polynomial with coefficients a_i . We define the content of $p(x)$ to be*

$$\text{content}(p) = \begin{cases} 0 & \text{if } p(x) = 0 \\ \gcd(a_0, a_1, \dots, a_d) & \text{if } p(x) \neq 0, \text{ where } d = \deg p. \end{cases}$$

If $\text{content}(p) = 1$, we say that $p(x)$ is primitive.

For example, \mathbb{Z} is a GCD Domain, and $\text{content}(2x^3 + 4x - 6) = 2$. Every field F is a GCD domain, and every nonzero polynomial over F is primitive.

Proposition 1. *Let R be a GCD domain.*

1. *Every polynomial $a(x) \in R[x]$ can be written as $a(x) = \text{content}(a)\bar{a}(x)$, where $\bar{a}(x) \in R[x]$ is primitive.*
2. *Let $d \in R$ and $a(x) \in R[x]$. Then the constant polynomial d divides $a(x)$ in $R[x]$ if and only if d divides $\text{content}(a)$ in R .*
3. *Let $d \in R$ and $a(x), b(x) \in R[x]$. If $d \mid \text{content}(a+b)$ and $d \mid \text{content}(a)$, then $d \mid \text{content}(b)$.*
4. *If $d \in R$ and $a(x) \in R[x]$, then $\text{content}(da) = d\text{content}(a)$.*
5. *Let F be the field of fractions of R and let $q(x) \in F[x]$. Then there is a fraction $\frac{u}{v} \in F$ such that $p(x) = \frac{u}{v}q(x)$ is in $R[x]$ and is primitive there.*
6. $\text{content}(x^n a(x)) = \text{content}(a(x))$.

Proof.

1. If $a(x) = 0$, set $\bar{a}(x) = 1$. Suppose $a(x) \neq 0$. Now $\text{content}(a) = \gcd(a_0, a_1, \dots, a_n)$, and in particular for each i we have $a_i = \text{content}(a)\bar{a}_i$ for some \bar{a}_i , and $\gcd(\bar{a}_0, \bar{a}_1, \dots, \bar{a}_n) = 1$. Let $\bar{a}(x) = \sum_{i=0}^n \bar{a}_i x^i$.
2. (write these)

□

Proposition 2 (Gauss’ Lemma – Part I). *Let R be a GCD Domain with $a(x), b(x) \in R[x]$. Then we have the following.*

1. *If $a(x)$ and $b(x)$ are primitive, then $a(x)b(x)$ is primitive.*

2. $\text{content}(ab) = \text{content}(a)\text{content}(b)$
3. If $a(x)|b(x)$ in $R[x]$, then $\text{content}(a)|\text{content}(b)$ in R .

Proof.

1. We proceed by induction on the number k of nonzero terms of a and b together.

- (a) **Base Case** ($k = 0$): If a and b together have no nonzero terms, then $a(x) = b(x) = 0$; neither is primitive.
- (b) **Base Case** ($k = 1$): If a and b together have exactly one nonzero term, then either $a(x) = 0$ or $b(x) = 0$; one is not primitive.
- (c) **Base Case** ($k = 2$): If $a(x)$ and $b(x)$ together have exactly two nonzero terms, then each must have exactly one. (Otherwise one is zero and thus not primitive.) Say $a(x) = a_n x^n$ and $b(x) = b_m x^m$. If both $a(x)$ and $b(x)$ are primitive, then $a_n = \text{content}(a)$ and $b_m = \text{content}(b)$ are units, so that $\text{content}(ab) = a_n b_m$ is a unit; hence $a(x)b(x)$ is primitive.

- (d) **Inductive Step:** Suppose the result holds for all pairs of primitive polynomials having less than $n > 2$ nonzero terms together, and suppose that $a(x)$ and $b(x)$ are primitive with exactly n nonzero terms together. Say $\deg a = n$ and $\deg b = m$, so that the leading coefficients of a , b , and ab are a_n , b_m , and $a_n b_m$, respectively. Now let $c = \text{content}(ab)$, and suppose BWOC that c is *not* a unit. Note that $c|a_n b_m$. Now $\gcd(c, a_n)$ and $\gcd(c, b_m)$ cannot both be units in R . (If $\gcd(c, a_n) = 1$, then by Euclid's lemma we have $c|\gcd(c, b_m)$.) So suppose WLOG that $\gcd(c, a_n) = d$ is not a unit.

Now $d|\text{content}(ab)$ in R , so that $d|a(x)b(x)$ in $R[x]$. Since $d|a_n$, we also have $d|a_n x^n$ in $R[x]$. Thus $d|b(x)(a(x) - a_n x^n)$ in $R[x]$, and thus

$$d|\text{content}(b(x)(a(x) - a_n x^n)) = \text{content}(a(x) - a_n x^n)\text{content}(b(x)p(x)),$$

where $p(x) \in R[x]$ is primitive such that $a(x) - a_n x^n = \text{content}(a(x) - a_n x^n)p(x)$. In particular, note that $p(x)$ and $a(x) - a_n x^n$ have the same number of nonzero terms which is one fewer than the number of nonzero terms of $a(x)$. Thus b and p have fewer than n nonzero terms. Since b and p are both primitive, by the inductive hypothesis, $\text{content}(bp) = 1$. Thus we have $d|\text{content}(a(x) - a_n x^n)$. Since $d|\text{content}(a_n x^n)$, by the lemma we have $d|\text{content}(a)$. But a is primitive, so that d is a unit, a contradiction. So $a(x)b(x)$ must be primitive.

2. We have

$$\begin{aligned} \text{content}(a(x)b(x)) &= \text{content}(\text{content}(a)\bar{a}\text{content}(b)\bar{b}) \\ &= \text{content}(a)\text{content}(b)\text{content}(\bar{a}\bar{b}) \\ &= \text{content}(a)\text{content}(b) \end{aligned}$$

3. Say $a(x)c(x) = b(x)$; then $\text{content}(a)\text{content}(c) = \text{content}(b)$. □

Lemma 3. *Let R be a GCD domain with field of fractions F .*

1. *If $p(x) \in R[x]$ is primitive, $r \in R$, and $a(x) \in R$ such that $p(x)|a(x)$ and $r|a(x)$ in $R[x]$, then $rp(x)|a(x)$ in $R[x]$.*
2. *If $q(x) \in F[x]$ and $p(x) \in R[x]$ such that $p(x)$ is primitive and $p(x)q(x) \in R[x]$, then in fact $q(x) \in R[x]$.*

Proof. (type this) □

Proposition 4 (Gilmer-Parker). *If R is a GCD Domain, then $R[x]$ is a GCD Domain.*

Proof. (type this) □

Exercises

1. Let R be a GCD domain with $p(x), q(x) \in R[x]$ so that q is irreducible (hence prime), and let k be a natural number. Show that q^{k+1} divides p in $R[x]$ iff $q|p$ and $q^k|p'$ in $R[x]$. In particular, show that p is squarefree iff $\gcd(p, p') = 1$.