

Localization

In a general ring with 1, or even a general domain, elements typically do not have multiplicative inverses. Those which do are called units and are very special. In this section we will see how a domain can be “extended” to a larger ring so that any given element can be made into a unit.

Definition 1 (Multiplicative subset). *Let R be a domain and $S \subseteq R$. We say that S is a multiplicative subset of R if $0 \notin S$ and if S is closed under multiplication.*

Domains have plenty of multiplicative sets. For instance, the set of all nonzero elements is multiplicative. If $a \in R$ is not zero, then the set $S = \{1, a, a^2, a^3, \dots\}$ of powers of a is multiplicative.

Here is the punch line of this section.

If $S \subseteq R$ is a multiplicative subset, then we can construct a new ring, T , which contains R as a subset, but in which the elements of S are units.

Proposition 1. *Let R be a domain and $S \subseteq R$ a multiplicative subset. We define a relation Φ on the cartesian product $S \times R$ as follows:*

$$(s_1, r_1)\Phi(s_2, r_2) \quad \text{iff} \quad r_1 s_2 = r_2 s_1.$$

This relation Φ is an equivalence.

Proof.

- $rs = rs$ for all $r \in R$ and $s \in S$, so that $(s, r)\Phi(s, r)$.
- Suppose $(s_1, r_1)\Phi(s_2, r_2)$. Then $r_1 s_2 = r_2 s_1$, so that $r_2 s_1 = r_1 s_2$, and thus $(s_2, r_1)\Phi(s_1, r_1)$.
- Suppose $(s_1, r_1)\Phi(s_2, r_2)$ and $(s_2, r_2)\Phi(s_3, r_3)$. Now $r_1 s_2 = r_2 s_1$ and $r_2 s_3 = r_3 s_2$. We then have $r_1 s_2 r_2 s_3 = r_2 s_1 r_3 s_2$; rearranging (since R is commutative) and using cancellation, we have $r_1 s_3 = r_3 s_1$. So $(s_1, r_1)\Phi(s_3, r_3)$ as needed. \square

Since Φ is an equivalence, it induces a partition on the set $S \times R$. We will denote this quotient set $S^{-1}R = (S \times R)/\Phi$, and denote the equivalence class of (s, r) by $\frac{r}{s}$.

Proposition 2. *Let R be a domain with $S \subseteq R$ a multiplicative subset. Define operations $+$ and \cdot on $S^{-1}R$ as follows.*

$$\frac{r_1}{s_1} + \frac{r_2}{s_2} = \frac{r_1 s_2 + r_2 s_1}{s_1 s_2} \quad \text{and} \quad \frac{r_1}{s_1} \cdot \frac{r_2}{s_2} = \frac{r_1 r_2}{s_1 s_2}.$$

Then we have the following.

1. $+$ and \cdot are well-defined.
2. $S^{-1}R$, with these operations, is an integral domain, which we call the localization of R at S .
3. If $t \in S$, then the mapping $\iota : R \rightarrow S^{-1}R$ given by $\iota(r) = \frac{rt}{t}$ is an injective ring homomorphism, and $\iota(t)$ is a unit in $S^{-1}R$.

Proof. (super tedious) □

So $S^{-1}R$ is a new ring which contains a “copy” (homomorphic image) of R , within which the elements of S become units.

Definition 2. Let R be a domain and let $D = \{x \in R \mid x \neq 0\}$ be the multiplicative subset of all nonzero elements of R . Then the localization $D^{-1}R$ is a field, called the field of fractions of R .

For example, \mathbb{Q} is properly defined as the field of fractions of \mathbb{Z} .

Special things

Proposition 3. If R is a UFD then $S^{-1}R$ is also a UFD.

Proof. (type this) □

Proposition 4. If R is a Euclidean domain then $S^{-1}R$ is a Euclidean domain.

Proof. (type this) □

Exercises

1. (do stuff in $\mathbb{Z}[\frac{1}{2}]$)
2. (universal property of localization)