

Decomposing parabolic eigenvalue problem

J. Xin and N. B. Murphy

Abstract

A decomposition of a time periodic parabolic eigenvalue problem and an application to our effective diffusivity problem.

1 An Example

Consider the parabolic eigenvalue problem:

$$\psi_t - \Delta_x \psi + \cos(t) b(x) \psi = \lambda \psi, \quad (1.1)$$

subject to 2π periodic boundary conditions in x and t .

Substituting $\cos t = (e^{it} + e^{-it})/2$, and $\psi = \sum_{\ell} \psi_{\ell}(x) e^{i\ell t}$ in (1.1), we have:

$$\sum_{\ell} e^{i\ell t} (i\ell \psi_{\ell} - \Delta_x \psi_{\ell}) + \frac{1}{2} b(x) \sum_{\ell} (e^{i(\ell+1)t} + e^{i(\ell-1)t}) \psi_{\ell} = \lambda \sum_{\ell} \psi_{\ell} e^{i\ell t},$$

or:

$$\sum_{\ell} e^{i\ell t} (i\ell \psi_{\ell} - \Delta_x \psi_{\ell}) + \frac{1}{2} b(x) \sum_{\ell} (\psi_{\ell-1} + \psi_{\ell+1}) e^{i\ell t} = \lambda \sum_{\ell} \psi_{\ell} e^{i\ell t}.$$

Extracting the ℓ -th mode on both sides gives:

$$(-\Delta_x + i\ell) \psi_{\ell} + \frac{1}{2} b(x) (\psi_{\ell-1} + \psi_{\ell+1}) = \lambda \psi_{\ell}, \quad \ell \in \mathbb{Z}, \quad (1.2)$$

which can be put in a tri-diagonal matrix acting on $[\cdots, \psi_{\ell-1}, \psi_{\ell}, \psi_{\ell+1}, \cdots]'$, with $b(x)/2$ on the off-diagonals, $-\Delta_x + i\ell$ on the diagonals.

Let us find a similar derivation for the eigenvalue problem of the advection-diffusion operator, with advection field being the time periodic cell flow:

$$\vec{u}(t, \vec{x}) = (\cos y, \cos x) + \delta \cos t (\sin y, \sin x) := \vec{u}_1(\vec{x}) + \delta \cos t \vec{u}_2(\vec{x}). \quad (1.3)$$

Question: Is it possible to carry out a similar decomposition in x and y and fully reduce the differential system like (1.2) into an algebraic system ?

1.1 An Application to our effective diffusivity problem

Consider the eigenvalue problem $A\psi_l = \imath\lambda_l\psi_l$, $\imath = \sqrt{-1}$, $\lambda_l \in \mathbb{R}$, $l = 1, 2, 3, \dots$, involving the integro-differential operator $A = (-\Delta)^{-1}(\partial_t + \vec{u} \cdot \vec{\nabla})$, introduced in equation (45) of our (attached) effective-diffusivity paper, with $\vec{u} \mapsto -\vec{u}$. Here A is an anti-symmetric (normal) operator and the incompressible velocity field $\vec{u}(t, \vec{x})$ is given in equation (1.3) above. The equation $A\psi_l = \imath\lambda_l\psi_l$ may be rewritten as

$$(\partial_t + \vec{u} \cdot \vec{\nabla})\psi_l = -\imath\lambda_l\Delta\psi_l. \quad (1.4)$$

The eigenfunctions ψ_l satisfy the following orthogonality condition

$$\langle \psi_l, \psi_i \rangle_1 = \langle \vec{\nabla}\psi_l \cdot \vec{\nabla}\psi_i \rangle = \delta_{li}, \quad (1.5)$$

where δ_{li} is the Kronecker delta and $\langle \cdot \rangle$ denotes space-time averaging over the period cell $\mathcal{T} \times \mathcal{V}$, with $\mathcal{T} = [0, 2\pi]$ and $\mathcal{V} = [0, 2\pi] \times [0, 2\pi]$.

The eigenfunction ψ_l is $\mathcal{T} \times \mathcal{V}$ periodic, mean-zero, and $\psi_l \in \mathcal{A}(\mathcal{T}) \otimes \mathcal{H}^1(\mathcal{V})$, i.e. it is absolutely continuous in time for $t \in \mathcal{T}$, and is in the Sobolev space $\mathcal{H}^1(\mathcal{V})$ for $\vec{x} \in \mathcal{V}$. We denote the class of such functions by \mathcal{F}

$$\mathcal{F} = \{f \in \mathcal{A}(\mathcal{T}) \otimes \mathcal{H}^1(\mathcal{V}) \mid \langle f \rangle = 0 \text{ and is periodic on } \mathcal{T} \times \mathcal{V}\}. \quad (1.6)$$

Since the orthogonal set $\{e^{i\ell t}\}$, $\ell \in \mathbb{Z}$, is dense in $\mathcal{A}(\mathcal{T})$, we may represent ψ_l by

$$\psi_l(t, \vec{x}) = \sum_{\ell} \psi_{\ell}^l(\vec{x}) e^{i\ell t}, \quad (1.7)$$

where $\psi_{\ell}^l \in \mathcal{H}^1(\mathcal{V})$. Write $\cos t = (e^{it} + e^{-it})/2$ and insert this and (1.7) into equation (1.4), yielding

$$\sum_{\ell} (i\ell + \vec{u}_1 \cdot \vec{\nabla} + i\lambda_l \Delta) \psi_{\ell}^l(\vec{x}) e^{i\ell t} + \frac{\delta}{2} \sum_{\ell} (e^{i(\ell+1)t} + e^{i(\ell-1)t}) \vec{u}_2 \cdot \vec{\nabla} \psi_{\ell}^l(\vec{x}) = 0, \quad (1.8)$$

or:

$$\sum_{\ell} \left[(i\ell + \vec{u}_1 \cdot \vec{\nabla} + i\lambda_l \Delta) \psi_{\ell}^l(\vec{x}) + \frac{\delta}{2} \vec{u}_2 \cdot \vec{\nabla} (\psi_{\ell-1}^l(\vec{x}) + \psi_{\ell+1}^l(\vec{x})) \right] e^{i\ell t} = 0. \quad (1.9)$$

By the completeness in $L^2(\mathcal{T})$ of the orthogonal set $\{e^{i\ell t}\}$ we have, for each $\ell \in \mathbb{Z}$,

$$(i\ell + \vec{u}_1 \cdot \vec{\nabla}) \psi_{\ell}^l(\vec{x}) + \frac{\delta}{2} \vec{u}_2 \cdot \vec{\nabla} (\psi_{\ell-1}^l(\vec{x}) + \psi_{\ell+1}^l(\vec{x})) = -i\lambda_l \Delta \psi_{\ell}^l(\vec{x}). \quad (1.10)$$

The system of partial differential equations in (1.10) can be reduced to a system of algebraic equations as follows. Recall that $\vec{u}_1(\vec{x}) = (\cos y, \cos x)$ and $\vec{u}_2(\vec{x}) = (\sin y, \sin x)$, which implies

$$\begin{aligned} (\vec{u}_1 \cdot \vec{\nabla}) \psi_{\ell}^l(\vec{x}) &= \cos y \partial_x \psi_{\ell}^l(\vec{x}) + \cos x \partial_y \psi_{\ell}^l(\vec{x}) \\ (\vec{u}_2 \cdot \vec{\nabla}) \psi_{\ell}^l(\vec{x}) &= \sin y \partial_x \psi_{\ell}^l(\vec{x}) + \sin x \partial_y \psi_{\ell}^l(\vec{x}) \end{aligned} \quad (1.11)$$

Since $\psi_{\ell}^l \in \mathcal{H}^1(\mathcal{V})$ and the orthogonal set $\{e^{i(mx+ny)}\}$, $m, n \in \mathbb{Z}$, is dense in this space, we can represent $\psi_{\ell}^l(\vec{x})$ by

$$\psi_{\ell}^l(\vec{x}) = \sum_{m,n} a_{\ell,m,n}^l e^{i(mx+ny)} \quad (1.12)$$

Write $\cos x = (e^{ix} + e^{-ix})/2$ and $\sin x = (e^{ix} - e^{-ix})/(2i)$, for example, and insert this and (1.12) into equation (1.11), yielding

$$\begin{aligned} (\vec{u}_1 \cdot \vec{\nabla}) \psi_{\ell}^l &= \frac{1}{2} \sum_{m,n} a_{\ell,m,n}^l \left[im e^{imx} (e^{i(n+1)y} + e^{i(n-1)y}) + in e^{iny} (e^{i(m+1)x} + e^{i(m-1)x}) \right] \\ (\vec{u}_2 \cdot \vec{\nabla}) \psi_{\ell}^l &= \frac{1}{2i} \sum_{m,n} a_{\ell,m,n}^l \left[im e^{imx} (e^{i(n+1)y} - e^{i(n-1)y}) + in e^{iny} (e^{i(m+1)x} - e^{i(m-1)x}) \right] \end{aligned} \quad (1.13)$$

or:

$$\begin{aligned} (\vec{u}_1 \cdot \vec{\nabla}) \psi_{\ell}^l &= \frac{i}{2} \sum_{m,n} [m(a_{\ell,m,n-1}^l + a_{\ell,m,n+1}^l) + n(a_{\ell,m-1,n}^l + a_{\ell,m+1,n}^l)] e^{i(mx+ny)} \\ (\vec{u}_2 \cdot \vec{\nabla}) \psi_{\ell}^l &= \frac{1}{2} \sum_{m,n} [m(a_{\ell,m,n-1}^l - a_{\ell,m,n+1}^l) + n(a_{\ell,m-1,n}^l - a_{\ell,m+1,n}^l)] e^{i(mx+ny)}. \end{aligned} \quad (1.14)$$

We also have

$$-\Delta\psi_\ell^l = \sum_{m,n} a_{\ell,m,n}^l (m^2 + n^2) e^{i(mx+ny)} \quad (1.15)$$

By the completeness of the orthogonal set $\{e^{i(mx+ny)}\}$, inserting equations (1.14) and (1.15) into equation (1.10) yields

$$\begin{aligned} & i\ell a_{\ell,m,n}^l + \frac{\iota}{2}[m(a_{\ell,m,n-1}^l + a_{\ell,m,n+1}^l) + n(a_{\ell,m-1,n}^l + a_{\ell,m+1,n}^l)] \\ & + \frac{\delta}{4}[m(a_{\ell-1,m,n-1}^l - a_{\ell-1,m,n+1}^l) + n(a_{\ell-1,m-1,n}^l - a_{\ell-1,m+1,n}^l) \\ & + m(a_{\ell+1,m,n-1}^l - a_{\ell+1,m,n+1}^l) + n(a_{\ell+1,m-1,n}^l - a_{\ell+1,m+1,n}^l)] \\ & = i\lambda_\ell(m^2 + n^2)a_{\ell,m,n}^l, \end{aligned} \quad (1.16)$$

which is an infinite system of algebraic equations for the unknown Fourier coefficients $a_{\ell,m,n}^l$ associated with the eigenfunctions $\psi_l(t, \vec{x})$ and eigenvalues $i\lambda_\ell$, $l \in \mathbb{N}$ and $\ell, m, n \in \mathbb{Z}$. Recalling that ψ_l is mean-zero $\langle \psi_l \rangle = 0$, we have that $\ell^2 + m^2 + n^2 > 0$.

We now discuss how the orthogonality condition $\langle \psi_l, \psi_i \rangle_1 = \delta_{li}$ in (1.5) is transformed by the Fourier expansion of the eigenfunctions $\psi_l(t, \vec{x})$. This expansion of $\psi_l(t, \vec{x})$ implies that for $\vec{\nabla}\psi_l(t, \vec{x})$ as follows

$$\psi_l(t, \vec{x}) = \sum_{\ell,m,n} a_{\ell,m,n}^l e^{i(\ell t + mx + ny)} \Rightarrow \vec{\nabla}\psi_l(t, \vec{x}) = \sum_{\ell,m,n} a_{\ell,m,n}^l (m, n) e^{i(\ell t + mx + ny)}. \quad (1.17)$$

Recall the orthogonality relation

$$\left\langle e^{i(\ell t + mx + ny)}, e^{i(\ell' t + m' x + n' y)} \right\rangle_2 = (2\pi)^3 \delta_{\ell,\ell'} \delta_{m,m'} \delta_{n,n'}, \quad (1.18)$$

where $\langle \cdot, \cdot \rangle_2$ denotes the $L^2(\mathcal{T} \times \mathcal{V})$ inner-product, and that $\langle \cdot \rangle$ denotes space-time averaging. Consequently, we have that the orthogonality relation in (1.5) is transformed to

$$\delta_{li} = \langle \vec{\nabla}\psi_l \cdot \vec{\nabla}\psi_i \rangle = \sum_{\ell,m,n} (m^2 + n^2) \overline{a_{\ell,m,n}^l} a_{\ell,m,n}^i \quad (1.19)$$

Recall that $\sum_i i^{-p}$ converges for all $p > 1$. From this and equation (1.19) we see that the square modulus of the Fourier coefficients $a_{\ell,m,n}^l$ must have the asymptotic behavior $|a_{\ell,m,n}^l|^2 \sim o((m^2 + n^2)^{-3/2})$ as $m, n \rightarrow \pm\infty$. Since $\psi_l(\cdot, \vec{x}) \in \mathcal{A}(\mathcal{T})$, i.e. $\partial_t \psi_l(\cdot, \vec{x}) \in L^2(\mathcal{T})$, we also have $|a_{\ell,m,n}^l|^2 \sim o(\ell^{-3})$ as $\ell \rightarrow \pm\infty$. Since $\partial_t \vec{\nabla}\psi_l \in L^2(\mathcal{T} \times \mathcal{V})$ we may generalize both of these statements by the following

$$|a_{\ell,m,n}^l|^2 \sim o(\ell^{-3}(m^2 + n^2)^{-3/2}), \text{ as } \ell, m, n \rightarrow \pm\infty. \quad (1.20)$$

We now use the special nature of the velocity field in (1.3) and the Fourier expansion of the eigenfunctions in (1.17) to show that the Radon–Stieltjes integral representation for the symmetric κ^* and anti-symmetric α^* parts of the effective diffusivity tensor \mathcal{K}^* depend only on the Fourier coefficients $a_{\ell,m,n}^l$ for $\ell, m, n \in \{-1, 0, 1\}$. Recall that the Hilbert space \mathcal{H} underlying this problem is

$$\mathcal{H} = \{f \in L^2(\mathcal{T}) \otimes \mathcal{H}^1(\mathcal{V}) \mid \langle f \rangle = 0 \text{ and is periodic on } \mathcal{T} \times \mathcal{V}\}, \quad (1.21)$$

with A maximal normal, i.e. iA self-adjoint, on $\mathcal{F} \subset \mathcal{H}$ defined in (1.6). Since the eigenfunctions ψ_l and ψ_i of A associated with distinct eigenvalues, $\lambda_l \neq \lambda_i$, are orthonormal, $\langle \psi_l, \psi_i \rangle_1 = \langle \vec{\nabla}\psi_l \cdot \vec{\nabla}\psi_i \rangle = \delta_{li}$, the span of all eigenfunctions constitutes a closed linear manifold \mathcal{E} in \mathcal{F} . Let \mathcal{E}^\perp be its (closed) orthogonal complement in \mathcal{H} so that [Stone]

$$\mathcal{H} = \mathcal{E} \oplus \mathcal{E}^\perp. \quad (1.22)$$

Recall the cell problem

$$(\varepsilon - A)\chi_j = g_j, \quad g_j = -\Delta^{-1}u_j, \quad (1.23)$$

where u_j is the j^{th} component of the velocity field \vec{u} . Since $\chi_j, g_j \in \mathcal{F}$ and $\mathcal{F} \subset \mathcal{E} \oplus \mathcal{E}^\perp$ they have the following representations

$$\chi_j = \sum_l \langle \psi_l, \chi_j \rangle_1 \psi_l + \chi_j^\perp, \quad g_j = \sum_l \langle \psi_l, g_j \rangle_1 \psi_l + g_j^\perp, \quad (1.24)$$

where $\psi_l \in \mathcal{E}$ and $\chi_j^\perp, g_j^\perp \in \mathcal{E}^\perp$. Using $A\psi_l = \imath\lambda_l\psi_l$, $\lambda_l \in \mathbb{R}$, and the orthonormality of the set $\{\psi_l\}$, inserting (1.24) into the cell problem (1.23) yields

$$\sum_l [(\varepsilon - \imath\lambda_l)\langle \psi_l, \chi_j \rangle_1 - \langle \psi_l, g_j \rangle_1] \psi_l + (\varepsilon + A)\chi_j^\perp - g_j^\perp = 0. \quad (1.25)$$

By the orthonormality of the set $\{\psi_l\}$ and since $\langle A\chi_j^\perp, \psi_l \rangle_1 = -\langle \chi_j^\perp, A\psi_l \rangle_1 = -\imath\lambda_l \langle \chi_j^\perp, \psi_l \rangle_1 = 0$, taking the inner-product of both sides of (1.25) with ψ_l yields

$$\langle \psi_l, \chi_j \rangle_1 = \frac{\langle \psi_l, g_j \rangle_1}{(\varepsilon - \imath\lambda_l)}, \quad 0 < \varepsilon < \infty. \quad (1.26)$$

Recall that the components κ_{jk}^* and α_{jk}^* of $\boldsymbol{\kappa}^*$ and $\boldsymbol{\alpha}^*$ are given by

$$\kappa_{jk}^* = \varepsilon(\delta_{jk} + \langle \chi_j, \chi_k \rangle_1), \quad \alpha_{jk}^* = \langle A\chi_j, \chi_k \rangle_1. \quad (1.27)$$

From equations (1.24) and (1.26) and the orthonormality of the set $\{\psi_l\}$, we have

$$\begin{aligned} \langle \chi_j, \chi_k \rangle_1 - \langle \chi_j^\perp, \chi_k^\perp \rangle_1 &= \sum_l \overline{\langle \psi_l, \chi_j \rangle_1} \langle \psi_l, \chi_k \rangle_1 = \sum_l \frac{\overline{\langle \psi_l, g_j \rangle_1} \langle \psi_l, g_k \rangle_1}{\varepsilon^2 + \lambda_l^2} \\ \langle A\chi_j, \chi_k \rangle_1 - \langle A\chi_j^\perp, \chi_k^\perp \rangle_1 &= \sum_l (-\imath\lambda_l) \overline{\langle \psi_l, \chi_j \rangle_1} \langle \psi_l, \chi_k \rangle_1 = \sum_l \frac{(-\imath\lambda_l) \overline{\langle \psi_l, g_j \rangle_1} \langle \psi_l, g_k \rangle_1}{\varepsilon^2 + \lambda_l^2} \end{aligned} \quad (1.28)$$

The right hand sides of the formulas in equation (1.28) are Radon–Stieltjes integrals associated with a *discrete* measure. The terms $\langle \chi_j^\perp, \chi_k^\perp \rangle_1$ and $\langle A\chi_j^\perp, \chi_k^\perp \rangle_1$ also have Radon–Stieltjes integral representations with respect to *continuous* measures, and provides the standard decomposition of the *spectral measure* into its discrete and continuous components, in the general setting [Stone].

We now show that the special nature of the velocity field in (1.3) and the Fourier expansion of the eigenfunctions ψ_l in (1.17) allow the spectral weights $\langle \psi_l, g_j \rangle_1$ in equation (1.28) to be given in terms of the Fourier coefficients $a_{\ell,m,n}^l$ for the reduced index set $\ell, m, n \in \{-1, 0, 1\}$. First note that, since $u_j(t, \cdot) \in \mathcal{H}^1(\mathcal{V}) \subset L^2(\mathcal{V})$,

$$\langle \psi_l, g_j \rangle_1 = \langle \vec{\nabla} \psi_l \cdot \vec{\nabla} (-\Delta)^{-1} u_j \rangle = \langle \psi_l, (-\Delta)(-\Delta)^{-1} u_j \rangle_2 = \langle \psi_l, u_j \rangle_2. \quad (1.29)$$

Writing $\cos x = (e^{ix} + e^{-ix})/2$ and $\sin x = (e^{ix} - e^{-ix})/(2\imath)$, for example, from equation (1.3) we have that

$$u_1(t, x, y) = \cos y + \cos t \sin y \quad (1.30)$$

$$\begin{aligned} &= \frac{1}{2} (e^{iy} + e^{-iy}) + \frac{1}{4\imath} (e^{it} + e^{-it})(e^{iy} - e^{-iy}) \\ &= \frac{1}{2} (e^{iy} + e^{-iy}) + \frac{1}{4\imath} (e^{i(t+y)} - e^{i(t-y)} + e^{i(-t+y)} - e^{i(-t-y)}), \end{aligned} \quad (1.31)$$

and $u_2(t, x, y) = u_1(t, y, x)$. This, equation (1.29), and the orthogonality relation in (1.18) imply that

$$\begin{aligned} \langle \psi_l, g_1 \rangle_1 &= \frac{1}{2} (a_{0,0,1}^l + a_{0,0,-1}^l) + \frac{1}{4\imath} (a_{1,0,1}^l - a_{1,0,-1}^l + a_{-1,0,1}^l - a_{-1,0,-1}^l) \\ \langle \psi_l, g_2 \rangle_1 &= \frac{1}{2} (a_{0,1,0}^l + a_{0,-1,0}^l) + \frac{1}{4\imath} (a_{1,1,0}^l - a_{1,-1,0}^l + a_{-1,1,0}^l - a_{-1,-1,0}^l) \end{aligned} \quad (1.32)$$

Since \vec{u}_i is incompressible, there exists an anti-symmetric matrix \mathbf{H}_i such that $\vec{u}_i = \vec{\nabla} \cdot \mathbf{H}_i$. This allows us to write $\vec{u}_i \cdot \vec{\nabla} = \vec{\nabla} \cdot \mathbf{H}_i \vec{\nabla}$, which is an anti-symmetric operator. When $\delta = 0$, the velocity field \vec{u} is time-independent and the operator A , which arises from the cell problem, becomes $A = \Delta^{-1}(\vec{u}_1 \cdot \vec{\nabla})$. In this case, the eigenvalue problem in (1.4) becomes

$$\vec{\nabla} \cdot \mathbf{H}_1 \vec{\nabla} \psi = \lambda \Delta \psi. \quad (1.33)$$

Discretizing this equation leads to a generalized eigenvalue problem involving *sparse* matrices. This matrix formulation has all the desired properties of the associated abstract Hilbert space formulation. (I will be adding the details of this to our paper soon.) From this matrix problem, we obtain a discrete approximation of the Radon–Stieltjes integral representation for the symmetric κ^* and anti-symmetric α^* parts of the effective diffusivity tensor \mathcal{K}^* , displayed in equation (35) of our (attached) paper.

1.1.1 Matrix representations of the eigenvalue problem

In the *time-independent case*, where $\delta = 0$ in the velocity field of equation (1.3), the system of equations in (1.16) corresponding to the eigenvalue problem becomes

$$m(a_{m,n-1} + a_{m,n+1}) + n(a_{m-1,n} + a_{m+1,n}) = 2\lambda(m^2 + n^2)a_{m,n}, \quad m, n \in \mathbb{Z}, \quad (1.34)$$

where, for simplicity, we have dropped the super-script and sub-script, and have written $a_{m,n} = a_{m,n}^l$ and $\lambda = \lambda_l$. When the indices in equation (1.34) are restricted to be finite, $-M \leq m, n \leq M$ say, and suitable boundary conditions are imposed, it can be written in matrix form

$$B\vec{a}_l = 2\lambda_l C\vec{a}_l, \quad (1.35)$$

where B and C are $(2M+1)^2 \times (2M+1)^2$ symmetric matrices and $l = 1, \dots, (2M+1)^2$. More specifically, B is real-symmetric and C is real-diagonal. Equation (1.35) is a generalized eigenvalue problem. Since B and C are symmetric matrices, the generalized eigenvalues are real $\lambda_l \in \mathbb{R}$ and the eigen-vectors \vec{a}_l – consisting of the Fourier coefficients for ψ_l – satisfy the orthogonality condition

$$\vec{a}_j^T C \vec{a}_k = \delta_{jk}. \quad (1.36)$$

The rows and columns of B and C , corresponding to the $a_{0,0}$ component of \vec{a} , consist entirely of zero elements. Therefore, without loss of generality, they can be removed from the generalized eigenvalue problem, making the matrix C positive-definite.

In the *time-dependent case*, where $\delta \neq 0$ in the velocity field of equation (1.3), slightly manipulating equation (1.16) yields

$$\begin{aligned} & 4\ell a_{\ell,m,n} + 2[m(a_{\ell,m,n-1} + a_{\ell,m,n+1}) + n(a_{\ell,m-1,n} + a_{\ell,m+1,n})] \\ & - i\delta[m(a_{\ell-1,m,n-1} - a_{\ell-1,m,n+1} + a_{\ell+1,m,n-1} - a_{\ell+1,m,n+1}) \\ & + n(a_{\ell-1,m-1,n} - a_{\ell-1,m+1,n} + a_{\ell+1,m-1,n} - a_{\ell+1,m+1,n})] \\ & = 4\lambda(m^2 + n^2)a_{\ell,m,n}. \end{aligned} \quad (1.37)$$

By restricting the indices, $-M \leq l, m, n \leq M$, in equation (1.37) and imposing suitable boundary conditions, it can also be written as the generalized eigenvalue problem in (1.35) and (1.36), where B and C are $(2M+1)^3 \times (2M+1)^3$ symmetric matrices. More specifically, B is Hermitian-symmetric and C is real-diagonal. The rows and columns of B and C , corresponding to the $a_{0,0,0}$ component of \vec{a} , consist entirely of zero elements. Therefore, without loss of generality, they can be removed from the generalized eigenvalue problem, making the matrix C positive-definite.

2 Three-Dim Steady Cellular Flows

The 3 dimensional (3D) steady cellular flows are:

$$B = (\Phi_x(x, y)W'(z), \Phi_y(x, y)W'(z), k\Phi(x, y)W(z)), \quad (2.38)$$

with $-\Delta\Phi = k\Phi$.

A special case is $k = 2$, then

$$\vec{u}(x, y, z) = (-\sin x \cos y \cos z, -\cos x \sin y \cos z, 2 \cos x \cos y \sin z). \quad (2.39)$$

We now derive a system of algebraic equations, analogous to that of equation (1.16), corresponding to the velocity field \vec{u} displayed in (2.39). The steady state eigenvalue problem in equation (1.4) is given by

$$\vec{u} \cdot \vec{\nabla} \psi_l = -\imath \lambda_l \Delta \psi_l. \quad (2.40)$$

The analog of equation (1.11) is

$$(\vec{u} \cdot \vec{\nabla}) \psi_l(\vec{x}) = -\sin x \cos y \cos z \partial_x \psi_l(\vec{x}) - \cos x \sin y \cos z \partial_y \psi_l(\vec{x}) + 2 \cos x \cos y \sin z \partial_z \psi_l(\vec{x}). \quad (2.41)$$

For this three dimensional flow, we use the following notation for the Fourier expansion of $\psi_l(\vec{x})$:

$$\psi_l(\vec{x}) = \sum_{\ell, m, n} a_{\ell, m, n}^l e^{\imath(\ell x + m y + n z)}. \quad (2.42)$$

Using the identities

$$\begin{aligned} u_1(\vec{x}) &= -\sin x \cos y \cos z = -\frac{1}{4}(\sin(x-y-z) + \sin(x-y+z) + \sin(x+y-z) + \sin(x+y+z)) \\ u_2(\vec{x}) &= -\cos x \sin y \cos z = \frac{1}{4}(\sin(x-y-z) + \sin(x-y+z) - \sin(x+y-z) - \sin(x+y+z)) \\ u_3(\vec{x}) &= 2 \cos x \cos y \sin z = \frac{1}{2}(-\sin(x-y-z) + \sin(x-y+z) - \sin(x+y-z) + \sin(x+y+z)), \end{aligned} \quad (2.43)$$

and writing $\sin x = (e^{\imath x} - e^{-\imath x})/(2\imath)$, for example, one can more easily see that the analogue of equation (1.16) for the velocity field in (2.39) is given by

$$\begin{aligned} & -\frac{\imath(\ell-1)}{8\imath} a_{\ell-1, m+1, n+1} + \frac{\imath(\ell+1)}{8\imath} a_{\ell+1, m-1, n-1} - \frac{\imath(\ell-1)}{8\imath} a_{\ell-1, m+1, n-1} + \frac{\imath(\ell+1)}{8\imath} a_{\ell+1, m-1, n+1} \\ & -\frac{\imath(\ell-1)}{8\imath} a_{\ell-1, m-1, n+1} + \frac{\imath(\ell+1)}{8\imath} a_{\ell+1, m+1, n-1} - \frac{\imath(\ell-1)}{8\imath} a_{\ell-1, m-1, n-1} + \frac{\imath(\ell+1)}{8\imath} a_{\ell+1, m+1, n+1} \\ & -\frac{\imath(m+1)}{8\imath} a_{\ell-1, m+1, n+1} - \frac{\imath(m-1)}{8\imath} a_{\ell+1, m-1, n-1} + \frac{\imath(m+1)}{8\imath} a_{\ell-1, m+1, n-1} - \frac{\imath(m-1)}{8\imath} a_{\ell+1, m-1, n+1} \\ & -\frac{\imath(m-1)}{8\imath} a_{\ell-1, m-1, n+1} + \frac{\imath(m+1)}{8\imath} a_{\ell+1, m+1, n-1} - \frac{\imath(m-1)}{8\imath} a_{\ell-1, m-1, n-1} + \frac{\imath(m+1)}{8\imath} a_{\ell+1, m+1, n+1} \\ & -\frac{\imath(n+1)}{4\imath} a_{\ell-1, m+1, n+1} + \frac{\imath(n-1)}{4\imath} a_{\ell+1, m-1, n-1} + \frac{\imath(n-1)}{4\imath} a_{\ell-1, m+1, n-1} - \frac{\imath(n+1)}{4\imath} a_{\ell+1, m-1, n+1} \\ & -\frac{\imath(n+1)}{4\imath} a_{\ell-1, m-1, n+1} + \frac{\imath(n-1)}{4\imath} a_{\ell+1, m+1, n-1} + \frac{\imath(n-1)}{4\imath} a_{\ell-1, m-1, n-1} - \frac{\imath(n+1)}{4\imath} a_{\ell+1, m+1, n+1} \\ & = \imath \lambda_l (m^2 + n^2) a_{\ell, m, n}, \end{aligned} \quad (2.44)$$

where we have suppressed the super-script notation and written $a_{\ell, m, n}^l = a_{\ell, m, n}$. Simplifying equation (2.44) yields

$$\begin{aligned} & \imath((\ell-1) - (m+1) + 2(n+1)) a_{\ell-1, m+1, n+1} + \imath(-(\ell+1) + (m-1) - 2(n-1)) a_{\ell+1, m-1, n-1} \\ & + \imath((\ell-1) - (m+1) - 2(n-1)) a_{\ell-1, m+1, n-1} + \imath(-(\ell+1) + (m-1) + 2(n+1)) a_{\ell+1, m-1, n+1} \\ & + \imath((\ell-1) + (m-1) + 2(n+1)) a_{\ell-1, m-1, n+1} + \imath(-(\ell+1) - (m+1) - 2(n-1)) a_{\ell+1, m+1, n-1} \\ & + \imath((\ell-1) + (m-1) - 2(n-1)) a_{\ell-1, m-1, n-1} + \imath(-(\ell+1) - (m+1) + 2(n+1)) a_{\ell+1, m+1, n+1} \\ & = 8\lambda_l (m^2 + n^2) a_{\ell, m, n}, \end{aligned} \quad (2.45)$$

From equations (1.29) and (2.43), one can see that the analogue of equation (1.32) is given by

$$\begin{aligned}
\langle \psi_l, g_1 \rangle_1 &= -\frac{1}{8l} (+ (a_{1,-1,-1} - a_{-1,1,1}) + (a_{1,-1,1} - a_{-1,1,-1}) + (a_{1,1,-1} - a_{-1,-1,1}) + (a_{1,1,1} - a_{-1,-1,-1})) \\
\langle \psi_l, g_2 \rangle_1 &= \frac{1}{8l} (+ (a_{1,-1,-1} - a_{-1,1,1}) + (a_{1,-1,1} - a_{-1,1,-1}) - (a_{1,1,-1} - a_{-1,-1,1}) - (a_{1,1,1} - a_{-1,-1,-1})) \\
\langle \psi_l, g_3 \rangle_1 &= \frac{1}{4l} (- (a_{1,-1,-1} - a_{-1,1,1}) + (a_{1,-1,1} - a_{-1,1,-1}) - (a_{1,1,-1} - a_{-1,-1,1}) + (a_{1,1,1} - a_{-1,-1,-1}))
\end{aligned} \tag{2.46}$$

By restricting the indices, $-M \leq l, m, n \leq M$, in equation (2.45) and imposing suitable boundary conditions, it can also be written as the generalized eigenvalue problem in (1.35) and (1.36), where B and C are $(2M+1)^3 \times (2M+1)^3$ symmetric matrices. More specifically, B is Hermitian-symmetric and C is real-diagonal positive-semi-definite. The rows and columns of B and C , corresponding to the $a_{0,0,0}$ component of \vec{a} , consist entirely of zero elements. Therefore, without loss of generality, they can be removed from the generalized eigenvalue problem, making the matrix C positive-definite.

References

- [1] L. Biferale, A. Crisanti, M. Vergassola, A. Vulpiani, *Eddy diffusivities in scalar transport*, Phys. Fluids 7(11), pp. 2725–2734, 1995.
- [2] A. Fannjiang and G. Papanicolaou, *Convection enhanced diffusion for periodic flows*, SIAM J. Appl. Math., **54** (1992), pp. 333-408.
- [3] L. Ryzhik and A. Zlatos, *KPP pulsating front speed-up by flows*, Comm. Math. Sci., **5** (2007), pp. 575-593.
- [4] L. Shen, J. Xin and A. Zhou, *Finite Element Computation of KPP Front Speeds in 3D Cellular and ABC Flows*, Math Model. Natural Phenom., 8(3), 2013, pp. 182-197.