



Name and section: _____

ID number: _____

E-mail: _____

1. Use the known values of the function $f(x) = \sin(x)$ (with $y = f(x)$) at $x = 0, \pi/6, \pi/4, \pi/3$ and $\pi/2$ in order to derive an interpolating polynomial $p(x)$ using a **monomial basis**. What is the degree of your polynomial? What is the interpolation error magnitude $|p(1.2) - \sin(1.2)|$? Plot your data points and the underlying interpolating polynomial for $x \in [0, \pi/2]$ in the same figure. Attach any codes/scripts producing your figure and the figure itself.
2. Assume the data pairs $\{(x_i, y_i)\}_{i=0}^n$ together with the functions:

$$\rho_j = \prod_{i \neq j} (x_j - x_i), \quad j = 0, 1, \dots, n,$$

$$\psi(x) = \prod_{i=0}^n (x - x_i).$$

- (a) Show that:

$$\rho_j = \psi'(x_j).$$

- (b) Show that the interpolating polynomial of degree at most n is given by

$$p_n(x) = \psi(x) \sum_{j=0}^n \frac{y_j}{(x - x_j) \psi'(x_j)}.$$

3. Let $f(x) = 1/x$ and data points $x_0 = 2$, $x_1 = 3$ and $x_2 = 4$. Note that you can use the abscissae to find the corresponding ordinates.
 - (a) Find **by hand** the **Lagrange form**, the **standard form**, and the **Newton form** of the interpolating polynomial $p_2(x)$ of $f(x)$ at the given points. **State which is which!** Then, expand out the Newton and Lagrange form to verify that they agree with the standard form of p_2 that you obtained (this is true due to the uniqueness of polynomial interpolation). Also, verify that $p_2(x_i) = f(x_i)$ for $i = 0, 1, 2$.
 - (b) Use the Polynomial Interpolation Error theorem to find an upper bound for the error

$$\|f - p_2\|_{\infty} = \max_{2 \leq x \leq 4} |f(x) - p_2(x)|.$$

- (c) Find the exact value of $\|f - p_2\|_\infty$ to at least 5 decimal places of accuracy. Of course, the answer should be less than or equal to the upper bound you found in part (b).
4. Consider the data set $\{x_i\}_{i=0}^n$ containing $(n+1)$ **distinct points** and the corresponding Lagrange basis functions $\{L_i(x)\}_{i=0}^n$. Then, prove that

$$\sum_{j=0}^n L_j(x) = 1.$$

Hint: Consider interpolating the function $f(x) = 1$ at the points given!

5. For some function f , the divided difference table is given:

i	x_i	$f[\cdot]$	$f[\cdot, \cdot]$	$f[\cdot, \cdot, \cdot]$	$f[\cdot, \cdot, \cdot, \cdot]$
0	1	$f[x_0]$	—	—	—
1	5	$f[x_1]$	$f[x_0, x_1]$	—	—
2	6	4	0	$-1/4$	—
3	4	2	$f[x_2, x_3]$	$f[x_1, x_2, x_3]$	$f[x_0, x_1, x_2, x_3]$

Fill in the unknown entries in the table.