

Nausea and Vomiting Related to Cancer Treatment (PDQ®)–Health Professional Version

Overview

Prevention and control of nausea and vomiting (N&V) are paramount in the treatment of patients with cancer. Chemotherapy-induced N&V is one of the most common and distressing acute side effects of cancer treatment. It occurs in up to 80% of patients and can have a significant impact on a patient's quality of life. N&V can also result in the following:

- Serious metabolic derangements.

- Nutritional depletion and anorexia.
- Deterioration of the patient's physical and mental status.
- Esophageal tears.
- Fractures.
- Wound dehiscence.
- Withdrawal from potentially useful and curative antineoplastic treatment.
- Degeneration of self-care and functional ability.

In this summary, unless otherwise stated, evidence and practice issues as they relate to adults are discussed. The evidence and application to practice related to children may differ significantly from information related to adults. When specific information about the care of children is available, it is summarized under its own heading.

Pathophysiology of N&V

Nausea is the subjective experience of an unpleasant, wavelike sensation in the back of the throat and/or the epigastrium that may culminate in vomiting. **Vomiting** (emesis) is the forceful expulsion of the contents of the stomach, duodenum, or jejunum through the oral cavity. Retching (dry heaves) involves the gastric and esophageal movements of vomiting without expulsion of vomitus.

Progress has been made in understanding the neurophysiological mechanisms that control nausea and vomiting (N&V). Both are controlled or mediated by the central nervous system but by different mechanisms. Nausea is mediated through the autonomic nervous system. Vomiting results from stimulation of a complex reflex that includes a convergence of afferent stimulation from the following:[1,2]

- A chemoreceptor trigger zone (CTZ, area postrema).
- The cerebral cortex and the limbic system in response to sensory stimulation (particularly smell and taste), psychological distress, and pain.
- The vestibular-labyrinthine apparatus of the inner ear in response to body motion.
- Peripheral stimuli from visceral organs and vasculature (via vagal and spinal sympathetic nerves) as a result of exogenous chemicals and endogenous substances that accumulate during inflammation, ischemia, and irritation.

Neurotransmitters (including serotonin, substance P, and dopamine) found in the CTZ, the vomiting center (thought to be located in the nucleus tractus solitarius), and enterochromaffin cells in the gastrointestinal tract release efferent impulses. These impulses are transmitted to the abdominal musculature, salivation center, and

respiratory center. The relative contribution from these multiple pathways, culminating in N&V symptoms, is complex. It is postulated to account for agents' variable emetogenicity (intrinsic emetogenicity and mitigating factors such as dosage, administration route, and exposure duration) and emetogenic profile (i.e., time to onset, symptom severity, and duration).[3,4]

References

1. Wickham R: Evolving treatment paradigms for chemotherapy-induced nausea and vomiting. Cancer Control 19 (2 Suppl): 3-9, 2012. [\[PUBMED Abstract\]](#)
2. Navari RM: Antiemetic control: toward a new standard of care for emetogenic chemotherapy. Expert Opin Pharmacother 10 (4): 629-44, 2009. [\[PUBMED Abstract\]](#)
3. Cefalo MG, Ruggiero A, Maurizi P, et al.: Pharmacological management of chemotherapy-induced nausea and vomiting

in children with cancer. J Chemother 21 (6): 605-10, 2009. [\[PUBMED Abstract\]](#)

4. Darmani NA, Crim JL, Janoyan JJ, et al.: A re-evaluation of the neurotransmitter basis of chemotherapy-induced immediate and delayed vomiting: evidence from the least shrew. Brain Res 1248: 40-58, 2009. [\[PUBMED Abstract\]](#)

General Risk Factors and Etiologies

Although most patients receiving chemotherapy are at risk of nausea and vomiting (N&V), the onset, severity, triggers, and duration vary. Factors related to the tumor, treatment, and patient all contribute to N&V, including tumor location, chemotherapy agents used, and radiation exposure.[\[1-3\]](#)

Patient-related factors may include the following:

- Incidence and severity of N&V during past courses of chemotherapy. Patients with poor control of N&V during past chemotherapy cycles are likely to experience N&V in subsequent cycles.
- History of chronic alcohol use. Patients with a history of chronic high intake of alcohol are less likely to experience cisplatin-induced N&V. [4]
- Age. N&V is more likely to occur in patients younger than 50 years.[5]
- Gender. N&V is more likely to occur in women. [5,6]
- History of morning sickness or emesis during pregnancy.

Additional causal factors may include the following:

- Fluid and electrolyte imbalances, such as hypercalcemia, volume depletion, or water intoxication.
- Tumor invasion or growth in the gastrointestinal tract, liver, or central nervous system, especially the posterior fossa.
- Constipation.
- Certain drugs, such as opioids.
- Infection or septicemia.
- Uremia.

Clinicians must be aware of all potential causes and factors of N&V, especially in patients with cancer who may receive several treatments and medications. For more information about opioid-induced N&V, see the [Adverse effects](#) section in Cancer Pain.

Classifications

N&V have been classified as acute, delayed, anticipatory, breakthrough, refractory, and chronic, as outlined below:[7-9]

- **Acute N&V:** N&V experienced during the first 24 hours after chemotherapy administration. [10]
- **Delayed (or late) N&V:** N&V that occurs more than 24 hours after chemotherapy administration. Delayed N&V is associated with cisplatin, cyclophosphamide, and other drugs (e.g., doxorubicin and ifosfamide) given at high doses or on 2 or more consecutive days.
- **Anticipatory N&V (ANV):** N&V that occurs before a new cycle of chemotherapy in response to conditioned stimuli such as the smells, sights, and sounds of the treatment room. ANV is a classically conditioned response that typically occurs after three or four chemotherapy treatments that led to acute or delayed N&V.

- **Breakthrough N&V:** Vomiting that occurs within 5 days of prophylactic use of antiemetics and requires rescue.
- **Refractory N&V:** N&V that does not respond to treatment.
- **Chronic N&V in patients with advanced cancer:** N&V associated with a variety of potential etiologies. These etiologies are neither well known nor well researched, but potential causal factors include gastrointestinal, cranial, metabolic, drug-induced (e.g., morphine), cytotoxic chemotherapy-induced, and radiation-induced mechanisms.[11]

The National Cancer Institute has published a descriptive terminology for adverse event reporting (see Table 1). A grading (severity) scale is provided for each term.

Table 1. National Cancer Institute’s
Common Terminology Criteria for
Adverse Events: N&V^a

Adverse Event	Grade	Description
Nausea ^b	1	Loss of appetite without alteration in eating habits
	2	Oral intake decreased without significant weight loss, dehydration, or malnutrition
	3	Inadequate oral caloric or

Adverse Event	Grade	Description
		fluid intake; tube feeding, TPN, or hospitalization indicated
	4	Grade not assigned
	5	Grade not assigned
Vomiting ^c	1	Intervention not indicated
	2	Outpatient IV hydration; medical intervention indicated

Adverse Event	Grade	Description
	3	Tube feeding, TPN, or hospitalization indicated
	4	Life-threatening consequences; urgent intervention indicated
	5	Death

Adverse Event	Grade	Description
<p>IV = intravenous; N&V = nausea and vomiting (emesis); TPN = total parenteral nutrition.</p> <p>^aAdapted from National Cancer Institute.[12]</p> <p>^bDefinition: A disorder characterized by a queasy sensation and/or the urge to vomit.</p> <p>^cDefinition: A disorder characterized by the reflexive act of ejecting the contents of the stomach through the mouth.</p>		

References

1. Farrell C, Brearley SG, Pilling M, et al.: The impact of chemotherapy-related nausea on patients' nutritional status, psychological distress and quality of life. Support Care Cancer 21 (1): 59-66, 2013. [\[PUBMED Abstract\]](#)

2. Dranitsaris G, Bouganim N, Milano C, et al.: Prospective validation of a prediction tool for identifying patients at high risk for chemotherapy-induced nausea and vomiting. J Support Oncol 11 (1): 14-21, 2013. [\[PUBMED Abstract\]](#)
3. Bouganim N, Dranitsaris G, Hopkins S, et al.: Prospective validation of risk prediction indexes for acute and delayed chemotherapy-induced nausea and vomiting. Curr Oncol 19 (6): e414-21, 2012. [\[PUBMED Abstract\]](#)
4. Sullivan JR, Leyden MJ, Bell R: Decreased cisplatin-induced nausea and vomiting with chronic alcohol ingestion. N Engl J Med 309 (13): 796, 1983. [\[PUBMED Abstract\]](#)
5. Tonato M, Roila F, Del Favero A: Methodology of antiemetic trials: a review. Ann Oncol 2 (2): 107-14, 1991. [\[PUBMED Abstract\]](#)
6. Roila F, Tonato M, Basurto C, et al.: Antiemetic activity of high doses of metoclopramide combined with methylprednisolone versus

- metoclopramide alone in cisplatin-treated cancer patients: a randomized double-blind trial of the Italian Oncology Group for Clinical Research. J Clin Oncol 5 (1): 141-9, 1987. [\[PUBMED Abstract\]](#)
7. Kris MG, Urba SG, Schwartzberg LS: Clinical roundtable monograph. Treatment of chemotherapy-induced nausea and vomiting: a post-MASCC 2010 discussion. Clin Adv Hematol Oncol 9 (1): suppl 1-15, 2011. [\[PUBMED Abstract\]](#)
 8. Hesketh PJ: Chemotherapy-induced nausea and vomiting. N Engl J Med 358 (23): 2482-94, 2008. [\[PUBMED Abstract\]](#)
 9. Grunberg SM, Osoba D, Hesketh PJ, et al.: Evaluation of new antiemetic agents and definition of antineoplastic agent emetogenicity--an update. Support Care Cancer 13 (2): 80-4, 2005. [\[PUBMED Abstract\]](#)
 10. Wickham R: Nausea and vomiting. In: Yarbo CH, Frogge MH, Goodman M, eds.: Cancer

Symptom Management. 2nd ed. Jones and Bartlett Publishers, 1999, pp 228-263.

11. Schwartzberg L: Chemotherapy-induced nausea and vomiting: state of the art in 2006. J Support Oncol 4 (2 Suppl 1): 3-8, 2006. [\[PUBMED Abstract\]](#)
12. National Cancer Institute: Common Terminology Criteria for Adverse Events (CTCAE), Version 5.0. Bethesda, Md: U.S. Department of Health and Human Services, National Institutes of Health, 2017. [Available online](#). Last accessed Dec. 18, 2024.

Anticipatory Nausea and Vomiting

Prevalence

The prevalence of anticipatory nausea and vomiting (ANV) has varied because of changing

definitions and assessment methods.[1] Anticipatory nausea appears to occur in approximately 29% of patients receiving chemotherapy, while anticipatory vomiting appears to occur in 11% of patients.[2] With the introduction of pharmacological agents such as 5-hydroxytryptamine-3 (5-HT₃) receptor antagonists, the prevalence of ANV was expected to decline, but studies have shown mixed results. One study found a lower incidence of ANV,[3] and three studies found comparable incidence rates.[2,4,5] It appears that the 5-HT₃ agents reduce postchemotherapy vomiting but not postchemotherapy nausea,[2,5] and the resulting impact on ANV is unclear.

Classical Conditioning

Although other theoretical mechanisms have been proposed,[6] ANV appears to be best explained by classical conditioning, also known as Pavlovian or respondent conditioning.[7] In classical conditioning, a previously neutral stimulus (e.g.,

smells of the chemotherapy environment) elicits a conditioned response (e.g., ANV) after a number of pairings or learning trials. In cancer chemotherapy, the first few chemotherapy infusions are the learning trials. The chemotherapy drugs are the unconditioned stimuli that elicit postchemotherapy nausea and vomiting (N&V) in some patients. The drugs are paired with a variety of other neutral, environmental stimuli (e.g., smells of the setting, presence of the oncology nurse, chemotherapy room). These previously neutral stimuli then become conditioned stimuli and elicit ANV in future chemotherapy cycles. ANV is not an indication of psychopathology. It is a learned response that, in other life situations (e.g., food poisoning), results in adaptive avoidance.

A variety of correlational studies provide empirical support for classical conditioning. For example, the prevalence of ANV before treatment with any chemotherapy is rare, and few patients ever experience ANV without previous

postchemotherapy nausea.[8] Also, most studies have found (1) a higher probability of ANV with an increasing number of chemotherapy infusions and (2) the intensity of ANV increasing as patients get closer to the time of their infusion.[9] In one experimental study, it was shown that a novel beverage could become a conditioned stimulus to nausea when paired with several chemotherapy treatments.[10]

Variables Correlated with ANV

Many variables have been investigated as potential risk factors that correlate with the incidence of ANV. There is no agreement on which factors predict ANV. However, a patient with fewer than three of the first eight characteristics listed below is unlikely to develop ANV. Screening after the first chemotherapy infusion could identify patients at increased risk.[11]

Variables Found to Correlate With ANV

1. Age younger than 50 years.
2. N&V after the last chemotherapy session.
3. Posttreatment nausea described as moderate, severe, or intolerable.
4. Posttreatment vomiting described as moderate, severe, or intolerable.
5. Feeling warm or hot all over after the last chemotherapy session.
6. Susceptibility to motion sickness.
7. Female gender.
8. High-state anxiety (anxiety reactive to specific situations).[12,13]
9. Greater reactivity of the autonomic nervous system and slower reaction time.[14]
10. Patient expectations of chemotherapy-related nausea before beginning treatment.
[15,16]
11. Percentage of chemotherapy infusions followed by nausea.[17]

12. Postchemotherapy dizziness.
13. Longer latency of onset of posttreatment N&V.[18]
14. Emetogenic potential of various chemotherapeutic agents. Patients receiving drugs with a moderate to severe potential for posttreatment N&V are more likely to develop ANV.[12]
15. History of morning sickness during pregnancy.

Treatment of ANV

Antiemetic drugs do not seem to control ANV once it has developed;[2] however, a variety of behavioral interventions has been investigated. [19] These interventions include the following:

- Progressive muscle relaxation with guided imagery.[20]
- Hypnosis.[21]

- Systematic desensitization.[22]
- Electromyography and thermal biofeedback.
[23]
- Distraction via the use of video games.[24,25]

Progressive muscle relaxation with guided imagery, hypnosis, and systematic desensitization has been studied the most and should be considered as treatment. Referral to a psychologist or other mental health professional with specific training and experience in working with cancer patients should be considered when ANV is identified. The earlier ANV is identified, the more likely treatment will be effective, so early screening and referral are essential. However, physicians and nurses often underestimate the incidence of chemotherapy-induced N&V.[26][[Level of evidence: II](#)]

Clearly, the most important aspect of ANV is prevention of acute and delayed N&V associated with chemotherapy. Most antiemetics have not

shown benefit for the treatment of ANV, but their use during chemotherapy may markedly decrease the incidence of ANV. The only class of medication that has shown benefit in some studies is benzodiazepines, most commonly lorazepam.[27]
[Level of evidence: IV]

References

1. Andrykowski MA: Defining anticipatory nausea and vomiting: differences among cancer chemotherapy patients who report pretreatment nausea. J Behav Med 11 (1): 59-69, 1988. [\[PUBMED Abstract\]](#)
2. Morrow GR, Roscoe JA, Kirshner JJ, et al.: Anticipatory nausea and vomiting in the era of 5-HT3 antiemetics. Support Care Cancer 6 (3): 244-7, 1998. [\[PUBMED Abstract\]](#)
3. Aapro MS, Kirchner V, Terrey JP: The incidence of anticipatory nausea and vomiting after repeat cycle chemotherapy: the effect of granisetron. Br J Cancer 69 (5): 957-60, 1994. [\[PUBMED Abstract\]](#)

4. Fernández-Marcos A, Martín M, Sanchez JJ, et al.: Acute and anticipatory emesis in breast cancer patients. Support Care Cancer 4 (5): 370-7, 1996. [\[PUBMED Abstract\]](#)
5. Roscoe JA, Morrow GR, Hickok JT, et al.: Nausea and vomiting remain a significant clinical problem: trends over time in controlling chemotherapy-induced nausea and vomiting in 1413 patients treated in community clinical practices. J Pain Symptom Manage 20 (2): 113-21, 2000. [\[PUBMED Abstract\]](#)
6. Reesal RT, Bajramovic H, Mai F: Anticipatory nausea and vomiting: a form of chemotherapy phobia? Can J Psychiatry 35 (1): 80-2, 1990. [\[PUBMED Abstract\]](#)
7. Stockhorst U, Klosterhalfen S, Steingruber HJ: Conditioned nausea and further side-effects in cancer chemotherapy: a review. Journal of Psychophysiology 12 (suppl 1): 14-33, 1998.

8. Morrow GR, Rosenthal SN: Models, mechanisms and management of anticipatory nausea and emesis. *Oncology* 53 (Suppl 1): 4-7, 1996. [\[PUBMED Abstract\]](#)
9. Montgomery GH, Bovbjerg DH: The development of anticipatory nausea in patients receiving adjuvant chemotherapy for breast cancer. *Physiol Behav* 61 (5): 737-41, 1997. [\[PUBMED Abstract\]](#)
10. Bovbjerg DH, Redd WH, Jacobsen PB, et al.: An experimental analysis of classically conditioned nausea during cancer chemotherapy. *Psychosom Med* 54 (6): 623-37, 1992 Nov-Dec. [\[PUBMED Abstract\]](#)
11. Morrow GR, Roscoe JA, Hickok JT: Nausea and vomiting. In: Holland JC, Breitbart W, Jacobsen PB, et al., eds.: *Psycho-oncology*. Oxford University Press, 1998, pp 476-484.
12. Andrykowski MA, Redd WH, Hatfield AK: Development of anticipatory nausea: a

- prospective analysis. J Consult Clin Psychol 53 (4): 447-54, 1985. [\[PUBMED Abstract\]](#)
13. Roscoe JA, Morrow GR, Hickok JT, et al.: Biobehavioral factors in chemotherapy-induced nausea and vomiting. J Natl Compr Canc Netw 2 (5): 501-8, 2004. [\[PUBMED Abstract\]](#)
 14. Kvale G, Psychol C, Hugdahl K: Cardiovascular conditioning and anticipatory nausea and vomiting in cancer patients. Behav Med 20 (2): 78-83, 1994 Summer. [\[PUBMED Abstract\]](#)
 15. Montgomery GH, Tomoyasu N, Bovbjerg DH, et al.: Patients' pretreatment expectations of chemotherapy-related nausea are an independent predictor of anticipatory nausea. Ann Behav Med 20 (2): 104-9, 1998 Spring. [\[PUBMED Abstract\]](#)
 16. Shelke AR, Roscoe JA, Morrow GR, et al.: Effect of a nausea expectancy manipulation on chemotherapy-induced nausea: a university of Rochester cancer center community

- clinical oncology program study. J Pain Symptom Manage 35 (4): 381-7, 2008. [\[PUBMED Abstract\]](#)
17. Tomoyasu N, Bovbjerg DH, Jacobsen PB: Conditioned reactions to cancer chemotherapy: percent reinforcement predicts anticipatory nausea. Physiol Behav 59 (2): 273-6, 1996. [\[PUBMED Abstract\]](#)
 18. Chin SB, Kucuk O, Peterson R, et al.: Variables contributing to anticipatory nausea and vomiting in cancer chemotherapy. Am J Clin Oncol 15 (3): 262-7, 1992. [\[PUBMED Abstract\]](#)
 19. Carey MP, Burish TG: Etiology and treatment of the psychological side effects associated with cancer chemotherapy: a critical review and discussion. Psychol Bull 104 (3): 307-25, 1988. [\[PUBMED Abstract\]](#)
 20. Lyles JN, Burish TG, Krozely MG, et al.: Efficacy of relaxation training and guided imagery in reducing the aversiveness of cancer

- chemotherapy. J Consult Clin Psychol 50 (4): 509-24, 1982. [\[PUBMED Abstract\]](#)
21. Redd WH, Andresen GV, Minagawa RY: Hypnotic control of anticipatory emesis in patients receiving cancer chemotherapy. J Consult Clin Psychol 50 (1): 14-9, 1982. [\[PUBMED Abstract\]](#)
22. Morrow GR, Morrell C: Behavioral treatment for the anticipatory nausea and vomiting induced by cancer chemotherapy. N Engl J Med 307 (24): 1476-80, 1982. [\[PUBMED Abstract\]](#)
23. Burish TG, Shartner CD, Lyles JN: Effectiveness of multiple muscle-site EMG biofeedback and relaxation training in reducing the aversiveness of cancer chemotherapy. Biofeedback Self Regul 6 (4): 523-35, 1981. [\[PUBMED Abstract\]](#)
24. Kolko DJ, Rickard-Figueroa JL: Effects of video games on the adverse corollaries of chemotherapy in pediatric oncology patients:

- a single-case analysis. *J Consult Clin Psychol* 53 (2): 223-8, 1985. [\[PUBMED Abstract\]](#)
25. Vasterling J, Jenkins RA, Tope DM, et al.: Cognitive distraction and relaxation training for the control of side effects due to cancer chemotherapy. *J Behav Med* 16 (1): 65-80, 1993. [\[PUBMED Abstract\]](#)
26. Chan CW, Cheng KK, Lam LW, et al.: Psycho-educational intervention for chemotherapy-associated nausea and vomiting in paediatric oncology patients: a pilot study. *Hong Kong Med J* 14 (5 Suppl): 32-5, 2008. [\[PUBMED Abstract\]](#)
27. Rock EM, Limebeer CL, Parker LA: Anticipatory nausea in animal models: a review of potential novel therapeutic treatments. *Exp Brain Res* 232 (8): 2511-34, 2014. [\[PUBMED Abstract\]](#)

Etiology of Acute or Delayed Chemotherapy-Induced Nausea and Vomiting

Acute Nausea and Vomiting (N&V)

The incidence of acute N&V with moderate- or high-risk chemotherapy ranges from 30% to 90%. [1-3] It can result in significant morbidity and can negatively affect quality of life. However, in recent years many new antiemetic medications and combinations have become available, dramatically decreasing the incidence and severity of this dreaded complication. Risk factors include the following:

- The emetogenic potential of the specific drug.
- The dose used.
- The treatment schedule.
- How chemotherapy agents are combined.

A drug with a low emetogenic potential given in high doses may cause a dramatic increase in the potential to induce N&V.[4] For example, standard doses of cytarabine rarely produce N&V, but high doses often do. Another influence is the use of drug combinations. Because most patients receive combination chemotherapy, the emetogenic potential of all of the drugs combined and individual drug doses needs to be considered.[5-9]

Other risk factors include the following:[10]

- Poor control with previous chemotherapy.
- Female gender.
- Age younger than 50 years.
- Experience with previous chemotherapy.
- History of motion sickness.
- History of morning sickness during pregnancy.
- Dehydration.
- Malnutrition.

- Recent surgery.
- Radiation therapy.

The American Society of Clinical Oncology (ASCO) provides a summary of intravenous chemotherapeutic agents and their respective risk of acute and delayed emesis.[10] For more information, see Table 2.

Table 2. Intravenous Chemotherapeutic Agents and Their Risk of Acute and Delayed Emesis^a

High Risk	Moderate Risk
Emesis has been documented in >90% of patients.	Emesis has been documented in 30%–90% of patients.
Anthracycline/cyclophosphamide combination	Alemtuzumab
Carmustine	Azacitidine
Cisplatin	Bendamustine
Cyclophosphamide ($\geq 1,500$ mg/m ²)	Carboplatin
Dacarbazine	Clofarabine
^a From Hesketh et al.[10]	

High Risk	Moderate Risk
Dactinomycin	Cyclophosphamide ($<1,500 \text{ mg/m}^2$)
Mechlorethamine	Cytarabine ($>1,000 \text{ mg/m}^2$)
Streptozotocin	Daunorubicin
	Doxorubicin
	Epirubicin
	Idarubicin

^aFrom Hesketh et al.[10]

High Risk	Moderate Risk
	Ifosfamide
	Irinotecan
	Irinotecan liposomal inject
	Oxaliplatin
	Romidepsin
	Temozolomide
	Thiotepa
^a From Hesketh et al.[10]	

High Risk	Moderate Risk
	Trabectedin
^a From Hesketh et al.[10]	

High Risk	Moderate Ris
^a From Hesketh et al.[10]	

ASCO also provides a summary of oral chemotherapeutic agents and their respective risk of acute and delayed emesis.^[10] For more information, see Table 3.

Table 3. Oral Chemotherapeutic Agents and Their Risk of Acute and Delayed Emesis^a

High Risk	Moderate Risk	Low Risk
Emesis has been documented in >90% of patients.	Emesis has been documented in 30%–90% of patients.	Emesis has been documented in 10%–30% of patients.
Altretamine	Bosutinib	Afatinib
^a From Hesketh et al. ^[10]		

High Risk	Moderate Risk	Low Risk
Procarbazine	Cabozantinib	Alectinib
	Ceritinib	Axitinib
	Crizotinib	Capecitabine
	Cyclophosphamide	Cobimetinib
	Imatinib	Dabrafenil
	Lenvatinib	Dasatinib
	Temozolomide	Etoposide
^a From Hesketh et al.[10]		

High Risk	Moderate Risk	Low Risk
	Trifluridine-tipiracil	Everolimus
	Vinorelbine	Fludarabine
		Ibrutinib
		Idelalisib
		Ixazomib
		Lapatinib
		Lenalidomide
^a From Hesketh et al.[10]		

High Risk	Moderate Risk	Low Ris
		Olaparib
		Osimertini
		Nilotinib
		Palbociclik
		Panobinos
		Pazopanib
		Ponatinib
^a From Hesketh et al.[10]		

High Risk	Moderate Risk	Low Ris
		Regorafen
		Sonidegib
		Sunitinib
		Thalidomide
		Trametinib
		Vandetanib
		Venetoclax

^aFrom Hesketh et al.[10]

High Risk	Moderate Risk	Low Risk
		Vorinostat
^a From Hesketh et al.[10]		

Delayed N&V

Delayed (or late) N&V occurs more than 24 hours after chemotherapy administration. Delayed N&V is associated with cisplatin, cyclophosphamide, and other drugs (e.g., doxorubicin and ifosfamide) given at high doses or given on 2 or more consecutive days.[1,11,12]

- Etiologies: Patients who experience acute emesis with chemotherapy are significantly more likely to have delayed emesis.
- Risk factors: All predictive characteristics for acute emesis are considered risk factors for

delayed emesis.

- Emetic classifications: For more information, see the [Acute Nausea and Vomiting \(N&V\)](#) section.

References

1. Hesketh PJ, Sanz-Altamira P, Bushey J, et al.: Prospective evaluation of the incidence of delayed nausea and vomiting in patients with colorectal cancer receiving oxaliplatin-based chemotherapy. *Support Care Cancer* 20 (5): 1043-7, 2012. [\[PUBMED Abstract\]](#)
2. Schwartzberg L: Addressing the value of novel therapies in chemotherapy-induced nausea and vomiting. *Expert Rev Pharmacoecon Outcomes Res* 14 (6): 825-34, 2014. [\[PUBMED Abstract\]](#)
3. Sekine I, Segawa Y, Kubota K, et al.: Risk factors of chemotherapy-induced nausea and vomiting: index for personalized antiemetic

prophylaxis. Cancer Sci 104 (6): 711-7, 2013. [\[PUBMED Abstract\]](#)

4. Roscoe JA, Morrow GR, Hickok JT, et al.: Nausea and vomiting remain a significant clinical problem: trends over time in controlling chemotherapy-induced nausea and vomiting in 1413 patients treated in community clinical practices. J Pain Symptom Manage 20 (2): 113-21, 2000. [\[PUBMED Abstract\]](#)
5. Viale PH, Grande C, Moore S: Efficacy and cost: avoiding undertreatment of chemotherapy-induced nausea and vomiting. Clin J Oncol Nurs 16 (4): E133-41, 2012. [\[PUBMED Abstract\]](#)
6. Dranitsaris G, Bouganim N, Milano C, et al.: Prospective validation of a prediction tool for identifying patients at high risk for chemotherapy-induced nausea and vomiting. J Support Oncol 11 (1): 14-21, 2013. [\[PUBMED Abstract\]](#)

7. Kris MG, Urba SG, Schwartzberg LS: Clinical roundtable monograph. Treatment of chemotherapy-induced nausea and vomiting: a post-MASCC 2010 discussion. Clin Adv Hematol Oncol 9 (1): suppl 1-15, 2011. [\[PUBMED Abstract\]](#)
8. Phillips RS, Gopaul S, Gibson F, et al.: Antiemetic medication for prevention and treatment of chemotherapy induced nausea and vomiting in childhood. Cochrane Database Syst Rev (9): CD007786, 2010. [\[PUBMED Abstract\]](#)
9. Olver I, Clark-Snow RA, Ballatori E, et al.: Guidelines for the control of nausea and vomiting with chemotherapy of low or minimal emetic potential. Support Care Cancer 19 (Suppl 1): S33-6, 2011. [\[PUBMED Abstract\]](#)
10. Hesketh PJ, Kris MG, Basch E, et al.: Antiemetics: American Society of Clinical Oncology Clinical Practice Guideline Update. J

Clin Oncol 35 (28): 3240-3261, 2017. [\[PUBMED Abstract\]](#)

11. Geling O, Eichler HG: Should 5-hydroxytryptamine-3 receptor antagonists be administered beyond 24 hours after chemotherapy to prevent delayed emesis? Systematic re-evaluation of clinical evidence and drug cost implications. J Clin Oncol 23 (6): 1289-94, 2005. [\[PUBMED Abstract\]](#)
12. Fleishman SB, Mahajan D, Rosenwald V, et al.: Prevalence of Delayed Nausea and/or Vomiting in Patients Treated With Oxaliplatin-Based Regimens for Colorectal Cancer. J Oncol Pract 8 (3): 136-40, 2012. [\[PUBMED Abstract\]](#)

Prevention and Management of Acute or Delayed Nausea and Vomiting

Several organizations—including the American Society of Clinical Oncology, the National Comprehensive Cancer Network, and the Pediatric Oncology Group of Ontario—have published antiemetic guidelines for their members. PDQ does not endorse specific guidelines, but examples can be found in the literature.[1-4]

Antiemetic agents are the most common intervention for treatment-related nausea and vomiting (N&V). The basis for antiemetic therapy is the neurochemical control of vomiting. Although the exact mechanism is not well understood, peripheral neuroreceptors and the chemoreceptor trigger zone (CTZ) are known to contain receptors for serotonin, histamine (H1 and H2), dopamine, acetylcholine, opioids, and numerous other endogenous neurotransmitters.[5,6] Many antiemetics act by competitively blocking receptors for these substances, which inhibit stimulation of peripheral nerves at the CTZ and possibly at the vomiting center.

Current guidelines [2,7] recommend that prechemotherapy management of chemotherapy-induced N&V (CINV) be based on the emetogenic potential of the chemotherapy agent(s) selected. For patients receiving regimens with high emetogenic potential, the combination of a 5-hydroxytryptamine-3 (5-HT₃) receptor antagonist, neurokinin-1 (NK-1) receptor antagonist, and dexamethasone with or without olanzapine is recommended prechemotherapy. Aprepitant (if chosen as the NK-1 receptor antagonist prechemotherapy), olanzapine, and dexamethasone are recommended for the prevention of delayed emesis. Guidelines differ with respect to using a three- or four-drug regimen for prophylaxis for highly emetogenic chemotherapy. One guideline includes the option of omitting an NK-1 antagonist completely if dexamethasone, palonosetron, and olanzapine are used.[7]

For patients receiving moderately emetogenic chemotherapy, the combination of a 5-HT₃ receptor antagonist and dexamethasone is used prechemotherapy. Patients receiving carboplatin (area under the curve ≥ 4 mg/mL) may also receive an NK-1 receptor antagonist. Postchemotherapy, a 5-HT₃ receptor antagonist, dexamethasone, or both are recommended for the prevention of delayed emesis.

For regimens with low emetogenic potential, dexamethasone or a 5-HT₃ receptor antagonist is recommended. For regimens with minimal emetogenic risk, no prophylaxis is recommended.
[2,7]

Antiemetic guidelines [2,7] have included oral 5-HT₃ receptor antagonists as optional therapy for the prevention of delayed emesis, but the level of evidence supporting this practice is low.[8]

Studies have strongly suggested that patients experience more acute and delayed CINV than is

perceived by practitioners.[8-10] One study suggested that patients who are highly expectant of nausea appear to experience more postchemotherapy nausea.[11] In addition, the current and new agents that have been used as prophylaxis for acute and delayed CINV have not been studied for use in established CINV. One study reported the effective use of intravenous (IV) palonosetron and dexamethasone to prevent CINV in patients receiving multiple-day chemotherapy. [12]

Table 4 summarizes prechemotherapy and postchemotherapy recommendations by emetogenic potential.

Table 4. Antiemetic Recommendations by Emetic Risk Categories^{a,b}

Emetic Risk Category	ASCO Guidelines	MASCC Guidelines
High risk (>90%)	4-drug combination of NK-1 antagonist, 5-HT ₃ receptor antagonist, dexamethasone, and olanzapine recommended prechemotherapy	3-drug combination of NK-1 antagonist, 5-HT ₃ receptor antagonist, and dexamethasone recommended prechemotherapy
	Olanzapine and dexamethasone to be continued on days 2–4	

Emetic Risk Category	ASCO Guidelines	MASCC Guidelines
	<p>For anthracycline and cyclophosphamide combinations only, olanzapine to be continued on days 2-4</p>	
	<p>Note: Depending on NK-1 antagonist, dosing may be ≥ 1 day</p>	

Emetic Risk Category	ASCO Guidelines	MASCC Guide
<p>Moderate risk (30%–90%)</p>	<p>Carboplatin AUC ≥ 4 mg/mL per min; 3-drug combination of NK-1 antagonist, 5-HT₃ receptor antagonist, and dexamethasone recommended prechemotherapy</p>	<p>For carboplatin-containing regimens, 3-drug combination of NK-1 antagonist, 5-HT₃ receptor antagonist, and dexamethasone recommended prechemotherapy</p>

Emetic Risk Category	ASCO Guidelines	MASCC Guidelines
	<p>For patients receiving chemotherapies of moderate emetic risk excluding carboplatin AUC ≥ 4 mg/mL per min, 2-drug combination of 5-HT₃ receptor antagonist and dexamethasone recommended prechemotherapy</p>	<p>For patients receiving chemotherapy of moderate emetic risk excluding carboplatin, 2-drug combination of 5-HT₃ receptor antagonist and dexamethasone recommended prechemotherapy</p>
	<p>For patients receiving cyclophosphamide, doxorubicin, oxaliplatin, and other moderate-emetic-risk antineoplastic agents known to cause delayed nausea,</p>	<p>For patients receiving cyclophosphamide, doxorubicin, oxaliplatin, and other moderate-emetic-risk antineoplastic agents, dexamethasone may be offered on days 2–3 for prevention of delayed emesis</p>

Emetic Risk Category	ASCO Guidelines	MASCC Guidelines
	dexamethasone may be offered on days 2–3 for prevention of delayed emesis	
Low risk (10%–30%)	Single dose of 5-HT ₃ receptor antagonist or dexamethasone (8 mg) recommended	Single dose of 5-HT ₃ receptor antagonist or dexamethasone or dopamine antagonist recommended

Emetic Risk Category	ASCO Guidelines	MASCC Guidelines
Minimal risk (<10%)	No antiemetic administered routinely pre- or postchemotherapy	No routine prophylaxis recommended

5-HT₃ = 5-hydroxytryptamine-3; ASCO = American Society of Clinical Oncology; AUC = area under the curve; MASCC = Multinational Association of Supportive Care in Cancer; NCCN = National Comprehensive Cancer Network; NK-1 = neurokinin-1.

^aAdapted from National Comprehensive Cancer Network. Hesketh et al.[\[2\]](#)

^bOrder of listed antiemetics does not reflect preference

Most drugs with proven antiemetic activity can be categorized into one of the following groups:

- Competitive antagonists at dopaminergic (D₂ subtype) receptors:
 - Phenothiazines.
 - Butyrophenones (droperidol, haloperidol).
 - Substituted benzamides (metoclopramide).
- Competitive antagonists at serotonergic (5-hydroxytryptamine-3 or 5-HT₃ subtype) receptors.
- Substance P antagonists (NK-1 receptor antagonists).
- Corticosteroids.
- Benzodiazepines (lorazepam).
- Cannabinoids.

Although Table 5 lists all routes of administration, the intramuscular (IM) route is used only when no other access is available. IM delivery is painful, is associated with erratic absorption of drug, and

may lead to sterile abscess formation or fibrosis of the tissues. This is particularly important when more than one or two doses of a drug are to be given.

Table 5. Prevention of Acute or Delayed CINV

Drug Category	Medication	
Dopamine antagonists: phenothiazines	Chlorpromazine	10–25 6h
		25–50 4h

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manufacturer.

Drug Category	Medication	
	Prochlorperazine	25 mg
		5–10 q6–8h
	Promethazine	12.5–
Dopamine antagonists: butyrophenones	Haloperidol	0.5–5 divided
<p>5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C extrapyramidal symptoms; IM = intramuscular; IV = intravenous = sublingual; SQ = subcutaneous.</p> <p>^aDolasetron may be difficult to obtain from the manu</p>		

Drug Category	Medication	
	Droperidol	1–2.5 6h

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manufacturer.

Drug Category	Medication	
Dopamine antagonists: substituted benzamides	Metoclopramide	Prevent CINV: x1 dose prechemo then 12 then 24
		Treat CINV: PO q4h 0.5 mg

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manufacturer.

Drug Category	Medication	
	Trimethobenzamide	300 n
		200 n
Serotonin (5-HT ₃) receptor antagonists	Dolasetron ^a	100 n prech
	Granisetron	1–2 n µg/kg IV wit chem

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
		3.1 mg trans
		10 mg prech
	Ondansetron	0.15 mg min prech then repea later; 16 mg

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = contraindicated; E = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
		24 mg before emet day cl
		8 mg before emet chem follow 8 mg PO q'

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C
extrapyramidal symptoms; IM = intramuscular; IV = i
= sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
	Palonosetron	0.25 mg PO prechemo day 1
Substance P antagonists (NK- 1 receptor antagonists)	Aprepitant	125 mg PO prechemo day 1 daily
	Aprepitant, emulsion	130 mg PO prechemo day 1

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manufacturer.

Drug Category	Medication	
	Fosaprepitant	150 n prech day 1

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C
extrapyramidal symptoms; IM = intramuscular; IV = i
= sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
	Netupitant (combined with palonosetron)	Netup mg/p 0.5 m prech day 1
	Fosnetupitant (combined with palonosetron)	Fosne mg/p 0.25 r prech day 1
	Rolapitant	180 n prech day 1

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C
extrapyramidal symptoms; IM = intramuscular; IV = i
= sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
Corticosteroids	Dexamethasone	12–20 mg high-risk chemo follow 1–2 times daily
		8 mg moderate risk chemo

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = contraindicated; E = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; PO = oral; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manufacturer.

Drug Category	Medication	
		follow mg/d
	Methylprednisolone	0.5–1 min p and 8 postc
Benzodiazepines	Alprazolam	0.25–
	Lorazepam	0.5–2

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C
extrapyramidal symptoms; IM = intramuscular; IV = i
= sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
Atypical antipsychotics	Olanzapine	Prevent acute CINV comb 5-HT ₃ antagonist dexar and N antagonist PO qd
		Treat breakthrough CINV: daily

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manufacturer.

Drug Category	Medication	
Other pharmacological agents	Dronabinol	5 mg/ prech follow h by s up to
		Dose increa increa mg/r maxir mg/r

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C = contraindicated; E = extrapyramidal symptoms; IM = intramuscular; IV = intravenous; SL = sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Drug Category	Medication	
	Nabilone	1–2 r maxir in 3 d
	<i>Cannabis</i>	No cl on dc
	Ginger	0.5–2 prech

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; C
extrapyramidal symptoms; IM = intramuscular; IV = i
= sublingual; SQ = subcutaneous.

^aDolasetron may be difficult to obtain from the manu

Competitive Dopamine (D₂) Antagonists

Phenothiazines

Phenothiazines act on dopaminergic receptors at the CTZ, possibly at other central nervous system (CNS) centers, and peripherally.

In selecting phenothiazines, the primary consideration is assessing differences in adverse effect profiles, which correlate with the structural classes of the drugs. Generally, aliphatic phenothiazines (e.g., chlorpromazine) produce sedation and anticholinergic effects, while piperazines (e.g., prochlorperazine) are associated with less sedation but higher incidence of extrapyramidal symptoms (EPS) such as acute dystonias, akathisia, neuroleptic malignant syndrome (uncommon), and, rarely, akinesias and dyskinesias. Marked hypotension may also result if

IV doses are administered rapidly at high doses. The concomitant use of H1 blockers, such as diphenhydramine, can often decrease the risk and severity of EPS. Phenothiazines may be of particular value in treating patients who experience delayed N&V with cisplatin regimens. [25-29][[Level of evidence: I](#)] Given their anticholinergic properties, phenothiazines are listed in the American Geriatrics Society Beers Criteria for Potentially Inappropriate Medication Use in Older Adults.[30]

Butyrophenones

Droperidol and haloperidol represent butyrophenones, another class of dopaminergic (D₂ subtype) receptor antagonists that are structurally and pharmacologically similar to the phenothiazines. While droperidol is used primarily as an adjunct to anesthesia induction, haloperidol is indicated as a neuroleptic antipsychotic drug. However, both agents have some antiemetic activity. Results of small, uncontrolled, open-label

studies show some efficacy for haloperidol in patients receiving palliative care.[31,32] Both agents may produce EPS, akathisia, hypotension, and sedation.

Substituted benzamides

Metoclopramide is a substituted benzamide, which, before serotonin (5-HT₃) receptor antagonists were introduced, was considered the most effective antiemetic agent against highly emetogenic chemotherapy. Although metoclopramide is a competitive antagonist at dopaminergic (D₂) receptors, it is most effective against acute vomiting when given IV at high doses, probably because it is a weak competitive antagonist (relative to other serotonin antagonists) at 5-HT₃ receptors. It may act on the CTZ and the periphery. Metoclopramide also increases lower esophageal sphincter pressure and enhances the rate of gastric emptying, which may factor into its overall antiemetic effect. Metoclopramide has also been safely given by IV bolus injection at higher

single doses (up to 6 mg/kg) and by continuous IV infusion, with or without a loading bolus dose, with efficacy comparable to that of multiple intermittent dosing schedules.[33-35]

Metoclopramide is associated with akathisia and dystonic EPS. Akathisia is seen more frequently in patients older than 30 years, and dystonic EPS are seen more commonly in patients younger than 30 years. Diphenhydramine, benztropine mesylate, and trihexyphenidyl are commonly used prophylactically or therapeutically to pharmacologically antagonize EPS.[36] While cogwheeling rigidity, acute dystonia, and tremor are responsive to anticholinergic medications, akathisia is best treated by lowering the metoclopramide dose, changing to a different agent, or adding a benzodiazepine.

Trimethobenzamide is believed to act centrally on the CTZ by blocking emetic impulses. It has been studied in a limited number of oncology patients experiencing nausea from various chemotherapy

regimens. Compared with placebo, trimethobenzamide, 200 mg IM every 6 hours for 2 days, significantly reduced episodes of N&V.[17]

5-HT₃ Receptor Antagonists

Four serotonin receptor antagonists—ondansetron, granisetron, dolasetron, and palonosetron—are available in the United States. Agents in this class are thought to prevent N&V by preventing serotonin, which is released from enterochromaffin cells in the gastrointestinal (GI) mucosa, from initiating afferent transmission to the CNS via vagal and spinal sympathetic nerves. [37-39] The 5-HT₃ receptor antagonists may also block serotonin stimulation at the CTZ and other CNS structures. Major side effects of this class of medications include mild headache and constipation. Multiple studies have shown that the 5-HT₃ receptor antagonists are most effective when given in conjunction with steroids.

Comparison of agents

Studies suggest that there are no major differences in efficacy or toxicity of the three first-generation 5-HT₃ receptor antagonists (dolasetron, granisetron, and ondansetron) in the treatment of acute CINV. These three agents are equivalent in efficacy and toxicity when used in appropriate doses.[40,41]; [42][Level of evidence: I] These agents have been shown to be effective in the first 24 hours postchemotherapy (acute phase), but not on days 2 to 5 postchemotherapy (delayed phase).

Palonosetron, the second-generation 5-HT₃ receptor antagonist, has been approved for acute emesis with highly and moderately emetogenic chemotherapy and for delayed emesis in patients receiving moderately emetogenic chemotherapy. [43]; [44][Level of evidence: I]

Despite the use of both first- and second-generation 5-HT₃ receptor antagonists, the control of acute CINV, especially delayed N&V, is suboptimal. There is considerable opportunity for improvement with either the addition or

substitution of new agents in current regimens.
[8,45-47]

Ondansetron

Several studies have demonstrated that ondansetron produces an antiemetic response that is equal or superior to that of high doses of metoclopramide, but with an improved toxicity profile, compared with that of dopaminergic antagonist agents.[48-51][Level of evidence: I]; [52,53] A randomized trial of ondansetron, 8 mg and 32 mg, given prophylactically to patients receiving cisplatin found no difference between the doses.[54] A single-center, retrospective chart review has reported ondansetron-loading doses of 16 mg/m² IV (maximum, 24 mg) to be safe in infants, children, and adolescents.[55] However, data reported to the U.S. Food and Drug Administration (FDA) raise concerns about QT prolongation and potentially fatal arrhythmias with a single 32-mg IV dose. Current drug labeling calls for a maximum single 16-mg IV dose.[56]

Currently, oral and injectable ondansetron formulations are approved for use without dosage modification in patients older than 4 years, including elderly patients and patients with renal insufficiency. Oral ondansetron is given 3 times daily starting 30 minutes before chemotherapy and continuing for up to 2 days after chemotherapy is completed. Ondansetron clearance is diminished in patients with severe hepatic insufficiency; these patients receive a single injectable or oral dose no higher than 8 mg. There is currently no information evaluating the safety of repeated daily ondansetron doses in patients with hepatic insufficiency. Other effective dosing schedules, such as a continuous IV infusion (e.g., 1 mg/h for 24 h) or oral administration, have also been evaluated.[\[57\]](#)

The major adverse effects of ondansetron include the following:[\[58\]](#)

- Headache (which can be treated with mild analgesics).

- Constipation.
- Fatigue.
- Dry mouth.
- Transient asymptomatic elevations in liver function tests (alanine and aspartate transaminases), which may be related to concurrent cisplatin administration.

Ondansetron has been etiologically implicated in a few case studies involving thrombocytopenia, renal insufficiency, and thrombotic events.[\[59\]](#) Rare electrocardiogram changes in the form of QTc prolongation may occur. In addition, a few case reports have implicated ondansetron in causing EPS. However, it is not clear whether the events described were in fact EPS. In other reports, the evidence is confounded by concurrent use of other agents that are known to produce EPS. Nevertheless, the greatest advantage of serotonin receptor antagonists over dopaminergic receptor antagonists is that they have fewer adverse

effects. Despite prophylaxis with ondansetron, many patients receiving doxorubicin, cisplatin, or carboplatin will experience acute and delayed N&V.[60] Randomized, double-blind, placebo-controlled trials support the addition of aprepitant, an NK-1 receptor antagonist, for additional mitigation of N&V.[61,62][Level of evidence: I]

Granisetron

Granisetron has shown efficacy in preventing and controlling N&V at a broad range of doses. In the United States, granisetron injection, extended-release injection, transdermal patch, and oral tablets are approved for initial and repeat prophylaxis for patients receiving emetogenic chemotherapy, including high-dose cisplatin. Granisetron is pharmacologically and pharmacokinetically distinct from ondansetron. However, clinically it is equally efficacious and equally safe.[60-63][Level of evidence: I]

The subcutaneous extended-release formulation of granisetron was compared with palonosetron to prevent CINV for patients receiving moderately or highly emetogenic chemotherapy in a randomized, double-blind noninferiority phase III trial.[64]

Patients were randomly assigned to receive IV palonosetron, 0.25 mg; or subcutaneous granisetron, 5 mg or 10 mg. Patients who received palonosetron in cycle 1 were then randomly assigned to receive granisetron in cycles 2 through 4. Both subcutaneous doses of granisetron were noninferior to palonosetron in cycle 1 of moderately emetogenic chemotherapy (74.8% and 76.9% for granisetron 5 mg and 10 mg, respectively, vs. 75.0% for palonosetron) and highly emetogenic chemotherapy (77.7% and 81.3% for granisetron 5 mg and 10 mg, respectively, vs. 80.7% for palonosetron).

Subcutaneous granisetron was not superior to palonosetron in the prevention of delayed CINV after highly emetogenic chemotherapy.

Currently, granisetron is approved for use without dosage modification in patients older than 2 years, including elderly patients and patients with hepatic and renal insufficiency.

Dolasetron

Oral formulations of dolasetron are indicated for the prevention of N&V associated with moderately emetogenic cancer chemotherapy, including initial and repeat courses. However, the drug may be difficult to obtain from the manufacturer. Oral dolasetron may be dosed as 100 mg within 1 hour before chemotherapy. Dolasetron was given IV or orally at 1.8 mg/kg as a single dose approximately 30 minutes before chemotherapy. However, injection formulations are no longer approved for CINV because of the risk of QTc interval prolongation.[[65](#)]

The effectiveness of oral dolasetron in the prevention of CINV has been proven in a large randomized, double-blind, comparative trial of 399

patients.[66][Level of evidence: I] Oral dolasetron was administered in the range of 25 to 200 mg 1 hour before chemotherapy. The other study arm consisted of oral ondansetron (8 mg) administered 1.5 hours before chemotherapy and every 8 hours after chemotherapy for a total of three doses. Rates of complete response (CR), defined as no emetic episodes and no use of escape antiemetic medications, improved with increasing doses of dolasetron. Both dolasetron 200 mg and ondansetron had significantly higher CR rates than did dolasetron 25 or 50 mg.

Palonosetron

Palonosetron is a 5-HT₃ receptor antagonist (second generation) that has antiemetic activity at both central and GI sites. Palonosetron is FDA approved for the prevention of acute N&V associated with initial and repeat courses of moderately and highly emetogenic cancer chemotherapy and for the prevention of delayed N&V associated with initial and repeat courses of

moderately emetogenic cancer chemotherapy. Compared with the older 5-HT₃ receptor antagonists, palonosetron has a higher binding affinity to the 5-HT₃ receptors, a higher potency, a significantly longer half-life (approximately 40 hours, four to five times longer than that of dolasetron, granisetron, or ondansetron), and an excellent safety profile.[67][[Level of evidence: I](#)] A dose-finding study demonstrated that the effective dose was 0.25 mg or higher.[68-72]

In two large studies of patients receiving moderately emetogenic chemotherapy, CR (no emesis, no rescue) was significantly improved in the acute and delayed periods for patients who received 0.25 mg of palonosetron alone, compared with either ondansetron or dolasetron alone.[43]; [44][[Level of evidence: I](#)] Dexamethasone was not given with the 5-HT₃ receptor antagonists in these studies, and it is not yet known whether the differences in CR would persist if it were used.

In another study,[73][Level of evidence: I] 650 patients receiving highly emetogenic chemotherapy (cisplatin ≥ 60 mg/m²) also received either dexamethasone and one of two doses of palonosetron (0.25 mg or 0.75 mg) or dexamethasone and ondansetron (32 mg). Single-dose palonosetron was as effective as ondansetron in preventing acute CINV with dexamethasone pretreatment. It was significantly more effective than ondansetron throughout the 5-day postchemotherapy period. In an analysis of the patients in the above studies who received repeated cycles of chemotherapy, one study [74] reported that the CR rates for both acute and delayed CINV were maintained with single IV doses of palonosetron without concomitant corticosteroids.

NK-1 Receptor Antagonists (Substance P Antagonists)

Substance P, found in the vagal afferent neurons in the nucleus tractus solitarius, the abdominal

vagus, and the area postrema, induces vomiting. NK-1 receptor antagonists, including aprepitant, fosaprepitant, netupitant, fosnetupitant, and rolapitant, block substance P from binding to the NK-1 receptor. In combination with a 5-HT₃ receptor antagonist and a corticosteroid, NK-1 receptor antagonists are indicated for the prevention of acute and delayed N&V associated with initial and repeat courses of highly and moderately emetogenic chemotherapy. There have been no randomized trials comparing the individual NK-1 receptor antagonists. All are considered effective at their FDA-approved doses.

Aprepitant and fosaprepitant

Clinical trials [75-78] demonstrated that the addition of aprepitant to a 5-HT₃ receptor antagonist plus dexamethasone before cisplatin chemotherapy improved the control of acute emesis, compared with a 5-HT₃ receptor antagonist plus dexamethasone. This regimen also improved the control of delayed emesis, compared

with placebo. In two randomized, double-blind, parallel, controlled studies, patients received cisplatin ($\geq 70 \text{ mg/m}^2$) and were randomly assigned to receive either (1) standard therapy with ondansetron and dexamethasone prechemotherapy and dexamethasone on days 2 to 4 postchemotherapy or (2) standard therapy plus aprepitant prechemotherapy on days 2 and 3. [79,80][Level of evidence: I] The CR (no emesis, no rescue) of the aprepitant group in both studies was significantly higher in both the acute and the delayed periods. An additional study confirmed the efficacy of aprepitant in the delayed period, when it was compared with ondansetron.[81] [Level of evidence: I] Finally, aprepitant has been shown to be efficacious in preventing N&V in breast cancer patients receiving highly emetogenic chemotherapy with cyclophosphamide and doxorubicin.[82]

The benefit of aprepitant has also been demonstrated outside of highly emetogenic

chemotherapy. The addition of aprepitant to ondansetron and dexamethasone before moderately emetogenic chemotherapy versus ondansetron and dexamethasone alone resulted in improved CINV outcomes.[83-85] An alternative dosing strategy was evaluated in a randomized, double-blind, placebo-controlled, phase III crossover study in patients receiving 5-day cisplatin combination chemotherapy for germ cell tumors.[86] In addition to standard antiemetic therapy, patients received aprepitant 125 mg on day 3, followed by aprepitant 80 mg on days 4 through 7. There was a significant improvement in CINV CR with the three-drug regimen.

Fosaprepitant dimeglumine, a water-soluble, phosphorylated analog of aprepitant, is rapidly converted to aprepitant after IV administration. [87] Fosaprepitant is approved as a single dose of 150 mg before chemotherapy on day 1, as an alternative to the 3-day oral aprepitant regimen. As demonstrated in a randomized, double-blind

study of patients receiving cisplatin chemotherapy, single-dose IV fosaprepitant (150 mg) given with ondansetron and dexamethasone was noninferior to the standard 3-day dosing of oral aprepitant in preventing CINV.[87] Fosaprepitant is formulated with polysorbate 80, a solubilizing agent, which can cause rare but serious hypersensitivity reactions.[88,89] Aprepitant is also available in a parenteral emulsion form, which has a reduced risk of thrombophlebitis and hypersensitivity reactions.[90]

Netupitant and fosnetupitant

Netupitant is a competitive antagonist to the NK-1 receptor that is marketed as either an oral fixed-combination product containing 300 mg of netupitant and 0.5 mg of palonosetron (NEPA) or an IV fixed-combination product containing 235 mg of fosnetupitant and 0.25 mg of palonosetron. Of note, the IV formulation of NEPA does not contain the surfactant polysorbate 80 or any other allergenic excipients and could be considered for

patients who have had hypersensitivity reactions to fosaprepitant.[91][Level of evidence: I] It is given with dexamethasone before chemotherapy to prevent both acute and delayed CINV. This drug combination has been used successfully for prevention of CINV in a single cycle of both highly and moderately emetogenic chemotherapy regimens.[92,93]

The antiemetic benefit of NEPA was demonstrated throughout multiple cycles of chemotherapy in a randomized, double-blind, controlled trial.[94] [Level of evidence: I] Patients starting combination anthracycline/cyclophosphamide regimens were randomly assigned to receive oral fixed-dose NEPA with 12 mg of dexamethasone or 0.5 mg of oral palonosetron with 20 mg of dexamethasone. The percentage of patients with a CR was significantly greater for NEPA than for oral palonosetron for cycles 1 to 4. The most common treatment-related side effects, headache and constipation, were similar between the two arms.

A Japanese study compared single-agent fosnetupitant to fosaprepitant combined with palonosetron and dexamethasone in patients receiving highly emetic chemotherapy.[95][[Level of evidence: I](#)] Fosnetupitant was found to be noninferior to the fosaprepitant regimen. Additionally, fosnetupitant had an improved safety profile with fewer injection site reactions (11% vs. 20.6%, $P < .001$). Single-agent fosnetupitant is not currently FDA approved in the United States.

Similarly, NEPA has been compared with granisetron and aprepitant in patients receiving highly emetogenic chemotherapy. In a phase III, randomized, double-blind study, a single dose of NEPA was shown to be noninferior to a 3-day regimen of granisetron and aprepitant. Additionally, significantly more patients did not need rescue medications when they received NEPA (96.6%), compared with those who received granisetron plus aprepitant (93.5%). Toxicities

were similar between treatment arms.[96][Level of evidence: I]

Rolapitant

Rolapitant is an oral competitive NK-1 receptor inhibitor. It is approved for the prevention of delayed N&V associated with highly and moderately emetogenic chemotherapy. In addition to granisetron and dexamethasone, rolapitant significantly increases CINV CR versus standard therapy plus placebo for patients receiving both highly and moderately emetogenic chemotherapy. Unlike other drugs in its class, rolapitant has no effect on cytochrome P450 3A4 enzymes; therefore, no dose adjustment for dexamethasone is required.[97-99] The IV formulation has been associated with hypersensitivity reactions, including anaphylaxis, which have limited its use. [100]

Corticosteroids

Steroids are commonly used in combination with other antiemetics. Their antiemetic mechanism of action is not fully understood, but they may affect prostaglandin activity in the brain. Clinically, steroids quantitatively decrease or eliminate episodes of N&V and may improve patients' mood, producing a subjective sense of well-being or euphoria (although they also can cause depression and anxiety). Steroids are sometimes used as single agents against mildly emetogenic chemotherapy but are more often used in antiemetic drug combinations.[101,102][Level of evidence: I];[103]

Steroids are given orally or intravenously before chemotherapy and may be repeated. Dosages and administration schedules are selected empirically. Dexamethasone is often the treatment of choice for N&V in patients receiving radiation to the brain, as it also reduces cerebral edema. It is administered orally or intravenously in the dose range of 8 mg to 40 mg (pediatric dose, 0.25–0.5

mg/kg).[104,105] Methylprednisolone is also administered orally or intravenously at doses and schedules that vary from 40 mg to 500 mg every 6 to 12 hours for up to 20 doses.[102,106]

Dexamethasone is also used orally for delayed N&V. Long-term corticosteroid use, however, is inappropriate and may cause substantial morbidity, including the following:[107-109]

- Immunosuppression.
- Proximal muscle weakness (especially involving the thighs and upper arms).
- Aseptic necrosis of the long bones.
- Cataract formation.
- Hyperglycemia and exacerbation of preexisting diabetes or escalation of subclinical diabetes to clinical pathology.
- Adrenal suppression with hypocortisolism.
- Lethargy.

- Weight gain.
- GI irritation.
- Insomnia.
- Anxiety.
- Mood changes.
- Psychosis.

A study that examined chemotherapy in a group of patients with ovarian cancer found that short-term use of glucocorticoids as antiemetics had no negative effects on outcomes (e.g., overall survival or efficacy of chemotherapy).[110] As previously shown with metoclopramide, numerous studies have demonstrated that dexamethasone potentiates the antiemetic properties of 5-HT₃ receptor–blocking agents.[107,111] If administered intravenously, dexamethasone may be given over 10 to 15 minutes because rapid administration may cause sensations of generalized warmth, pharyngeal tingling or burning, or acute transient perineal and/or rectal pain.[112-115]

Benzodiazepines

Benzodiazepines, such as lorazepam and alprazolam, are valuable adjuncts in the prevention and treatment of anxiety and the symptoms of anticipatory N&V associated with chemotherapy, especially with the highly emetogenic regimens given to children.[107-109] Benzodiazepines have not demonstrated intrinsic antiemetic activity as single agents, so they are adjuncts to other antiemetic agents in antiemetic prophylaxis and treatment.[116] Benzodiazepines presumably act on higher CNS structures, the brainstem, and spinal cord, and they produce anxiolytic, sedative, and anterograde amnesic effects. In addition, these drugs markedly decrease the severity of EPS, especially akathisia, associated with dopaminergic receptor antagonist antiemetics.

The adverse effects of lorazepam include sedation, perceptual and vision disturbances, anterograde amnesia, confusion, ataxia, and depressed mental

acuity.[117];[118][Level of evidence: I];[119,120]

Alprazolam has been shown to be effective when given in combination with metoclopramide and methylprednisolone.[19]

Olanzapine

Olanzapine is an antipsychotic in the thienobenzodiazepine drug class that blocks multiple neurotransmitters: dopamine at D₁, D₂, D₃, and D₄ brain receptors; serotonin at 5-HT_{2a}, 5-HT_{2c}, 5-HT₃, and 5-HT₆ receptors; catecholamines at alpha-1 adrenergic receptors; acetylcholine at muscarinic receptors; and histamine at H₁ receptors.[121] Common side effects include the following:[122,123]

- Sedation.
- Dry mouth.
- Increased appetite.
- Weight gain.
- Postural hypotension.

- Dizziness.

Olanzapine's activity at multiple receptors, particularly at the D₂ and 5-HT₃ receptors that appear to be involved in N&V, suggests that it may have significant antiemetic properties.[124][Level of evidence: II] Subsequent studies have shown its effectiveness as a CINV antiemetic.[125,126][Level of evidence: II] A large study [127][Level of evidence: I] demonstrated that in patients receiving either highly emetogenic chemotherapy or moderately emetogenic chemotherapy, the addition of olanzapine to azasetron and dexamethasone improved the CR of delayed CINV.

A randomized, double-blind, phase III trial evaluated olanzapine versus placebo in addition to standard antiemetics for the prevention of CINV associated with highly emetogenic chemotherapy. [20][Level of evidence: I] Chemotherapy-naïve patients receiving either (1) cisplatin at least 70 mg/m² of body surface area (BSA) with or without

additional agents or (2) doxorubicin 60 mg/m² of BSA with cyclophosphamide 600 mg/m² of BSA were randomly assigned to receive olanzapine 10 mg orally on days 1 through 4 or matching placebo with guideline-directed antiemetics. The antiemetic regimen included an NK-1 antagonist (fosaprepitant or aprepitant), 5-HT₃ receptor antagonist (palonosetron, granisetron, or ondansetron), and dexamethasone 12 mg on day 1, followed by 8 mg orally daily on days 2 through 4. Patients were stratified by sex, chemotherapy regimen, and the specific 5-HT₃ receptor antagonist chosen. The primary endpoint, no nausea, was defined as a score of 0 on the visual analogue scale of 0 to 10 and assessed at three time points postchemotherapy:

- Early, 0 to 24 hours.
- Later, 25 to 120 hours.
- Overall, 0 to 120 hours.

The percentage of patients experiencing no nausea was significantly higher in the olanzapine group than in the placebo group at the early (74% vs. 45%; $P = .002$), later (42% vs. 25%; $P = .002$), and overall time points (37% vs. 22%; $P = .002$). CR rate and freedom from clinically significant nausea (a score lower than 3 on the visual analog scale of 0–10) were also significantly improved with the addition of olanzapine at all time points. Patients receiving olanzapine reported increased sedation from baseline on day 2, which resolved on days 3 through 5. Based on these data and additional clinical trials, olanzapine appears to be safe and effective in controlling acute and delayed CINV in patients receiving highly emetogenic and moderately emetogenic chemotherapy.[[128](#),[129](#)]

Other Pharmacological Agents

Cannabis

The plant *Cannabis* contains more than 60 different types of cannabinoids, or components

that have physiological activity. The most popular, and perhaps the most psychoactive, is delta-9-tetrahydrocannabinol (delta-9-THC).[130] There are two FDA-approved *Cannabis* products for CINV:

- **Dronabinol** (a synthetic delta-9-THC), as prophylaxis for CINV, 5 mg/m² orally 1 to 3 hours before chemotherapy and every 2 to 4 hours after chemotherapy, for a total of no more than 6 doses per day.
- **Nabilone**, for CINV that has failed to respond to other antiemetics, 1 to 2 mg orally twice a day.

With respect to CINV, *Cannabis* products probably target cannabinoid-1 and cannabinoid-2 receptors, which are in the CNS.[131]

Much of the research on agents in this class, conducted in the late 1970s and 1980s, compared nabilone, dronabinol, or levonantradol to older antiemetic agents that targeted the dopamine

receptor, such as prochlorperazine (Compazine) and metoclopramide (Reglan).[132-136] This group of studies demonstrated that cannabinoids were as effective as dopaminergic antiemetics for moderately emetogenic chemotherapy or were more effective than placebo.[130] Side effects included euphoria, dizziness, dysphoria, hallucinations, and hypotension.[130] Despite earlier reports of efficacy, in at least one study, patients did not significantly prefer nabilone because of the side effects.[132]

Since the 1990s, research in N&V has elucidated newer and more physiological targets, namely 5-HT₃ and NK-1 receptors. Subsequently, 5-HT₃ and NK-1 receptor antagonists have become standard prophylactic therapy for CINV. Few studies have investigated the role of *Cannabis* extract and cannabinoids with these newer agents, so only limited conclusions can be drawn. In published trials, however, *Cannabis* extract and cannabinoids have not demonstrated more efficacy than 5-HT₃

receptor antagonists, and synergistic or additive effects have not been fully investigated.[[137,138](#)]

In summary, *Cannabis* and cannabinoids' role in the prevention and treatment of CINV is not fully known. Discussions with patients about their use may include responses to available agents, known side effects of *Cannabis*, and an assessment of the risks versus benefits of this therapy.[[139](#)] For more information, see [Cannabis and Cannabinoids](#).

Ginger

There are conflicting data on the efficacy of ginger for prophylaxis of CINV. A phase III, randomized, dose-finding trial of 576 patients with cancer evaluated 0.5 g, 1 g, and 1.5 g of ginger versus placebo in twice-a-day dosing for the prevention of acute nausea (defined as day 1 postchemotherapy). Patients experienced some level of nausea (as measured on an 11-point scale) caused by their current chemotherapy regimen, despite standard prophylaxis with a 5-HT₃ receptor

antagonist. Patients began taking ginger or placebo capsules 3 days before each chemotherapy treatment and continued them for 6 days. For average nausea severity, 0.5 g of ginger was significantly better than placebo. For maximum nausea severity, both 0.5 g and 1 g were significantly better than placebo. Effects for delayed N&V were not significant. This trial did not control for emetogenicity of the chemotherapy regimens. Adverse events were infrequent and were not severe.[23]

Conversely, data on ginger used to prevent N&V have not been as promising. A randomized, double-blind, placebo-controlled study evaluated the use of ginger 160 mg per day in patients receiving high-dose cisplatin ($>50 \text{ mg/m}^2$). Patients (N = 251) were assigned to receive either ginger or placebo. The incidence of delayed nausea, intercycle nausea, and anticipatory nausea did not differ between the two treatment arms.[140]

Multiday Chemotherapy

Regimens that include chemotherapy doses on multiple sequential days (multiday chemotherapy) present a unique challenge to preventing CINV because after the first dose of chemotherapy, nausea may be both acute and delayed. Although there is no standard antiemetic regimen for multiday chemotherapy, a corticosteroid and a 5-HT₃ receptor antagonist should be given with each day of highly and moderately emetogenic chemotherapy.[7,141] Evidence demonstrates benefit for the addition of an NK-1 antagonist to highly and moderately emetogenic multiday chemotherapy.[2,7,13,141] The choice of antiemetic drugs and their schedule should be matched to the emetogenicity of the individual chemotherapy agents and their sequence. In addition, the length of delayed nausea varies and will depend on the emetogenicity of the last day's chemotherapy.

Dexamethasone is scheduled on each day of a multiday chemotherapy regimen and for 2 to 3 days after if there is a risk of delayed nausea. Additional dexamethasone is not necessary if the chemotherapy regimen contains a corticosteroid. It is not known whether dexamethasone 20 mg given each day of a 5-day cisplatin regimen provides additional antiemetic benefit, and it may add toxicity.[13,142] Therefore, an alternative dexamethasone schedule (20 mg on days 1 and 2, followed by 8 mg twice daily on days 6 and 7, and 4 mg twice daily on day 8), based on the timing of CINV and to reduce the total steroid dose, has been studied in patients receiving 5-day cisplatin regimens.[12,13]

Standard antiemetic prophylaxis includes a 5-HT₃ receptor antagonist given before the first chemotherapy dose each day of a multiday chemotherapy regimen.[2,7,13,141] No 5-HT₃ receptor antagonist is favored over other agents in the class for multiday chemotherapy.

Palonosetron is a 5-HT₃ receptor antagonist with a longer half-life and higher receptor-binding affinity than other members in its class, allowing it to be given less frequently.[71] A prospective, uncontrolled trial demonstrated that palonosetron, as a single IV dose with dexamethasone 20 mg before two 3-day chemotherapy regimens, resulted in an 80% CR. [143] Palonosetron was also studied with dexamethasone as prophylaxis for a 5-day cisplatin-based regimen for germ cell tumors.[12] When palonosetron plus dexamethasone was given on days 1, 3, and 5, 51% of patients experienced no emesis on days 1 to 5, and 83% experienced no emesis on days 6 to 9.

Alternative methods of 5-HT₃ receptor antagonist delivery have been studied. Granisetron as a 7-day continuous transdermal patch was compared with daily oral granisetron in patients receiving multiday chemotherapy in a double-blind, phase III, noninferiority study.[63] The patch

demonstrated complete control in 60% of patients, while the oral formulation did so in 65% of patients, achieving noninferiority.

The NK-1 antagonist aprepitant and its IV formulation, fosaprepitant, have been studied with multiday chemotherapy in dosing schedules that differ from their FDA-approved schedules. A nonrandomized trial evaluated the use of aprepitant, granisetron, and dexamethasone for CINV prophylaxis with 3- and 5-day highly and moderately emetogenic chemotherapy.[144] Aprepitant was given at 125 mg orally before the first dose of chemotherapy, then 80 mg orally on each day of chemotherapy and for 2 following days (total, 5–7 days). CR was seen in 57.9% and 72.5% of patients receiving highly and moderately emetogenic chemotherapy, respectively. Similarly promising results were found in a subsequent single-arm trial looking at a 7-day oral aprepitant regimen with dexamethasone and a 5-HT₃

receptor antagonist for 5-day cisplatin-based chemotherapy.[145]

A randomized, double-blind, placebo-controlled crossover trial of aprepitant, a 5-HT₃ receptor antagonist, and dexamethasone was conducted in patients receiving 5-day cisplatin-based chemotherapy for germ cell tumors.[86] Oral aprepitant 125 mg was given on day 3, followed by oral aprepitant 80 mg daily on days 4 to 7. More patients achieved CR with aprepitant than with placebo, 42% versus 13% ($P < .001$). IV fosaprepitant 150 mg given on days 3 and 5 was studied in a small phase II trial evaluating its use with a 5-HT₃ receptor antagonist and dexamethasone in 5-day cisplatin-based chemotherapy.[146] Preliminary results showed a CR rate of 28.1%, lower than results of the oral aprepitant trial conducted by the same institution.

High-Dose Chemotherapy With Stem Cell Transplant

Prevention of emesis during high doses of chemotherapy, with or without total-body irradiation, continues to be a challenging area of patient care.[147] Current [guidelines](#) primarily address single-day therapies. In addition, while emesis prevention for the multiple days of chemotherapy or radiation therapy used in this setting is based on single-day experiences, additional research is needed to improve symptom control for these patients.[147] This has led to the addition of NK-1 antagonists to the daily dosing of a serotonin antagonist plus dexamethasone.[147-149] Additional evidence is needed to determine optimal combinations, as CR rates range as low as 30%.[149] Also, experience has primarily been with aprepitant. The newer NK-1 antagonists may offer additional benefit.

Overall, these antiemetic combinations are well tolerated, with most side effects involving the dexamethasone component. In addition, while drug interactions were originally a concern, they

do not appear to be clinically significant.[150] Also, emesis is controlled to a much greater extent than is nausea, which continues to be challenging for many patients.[147,151] Finally, a randomized phase III trial studied the use of aprepitant, granisetron, and dexamethasone for the prevention of CINV in multiple myeloma patients receiving high-dose melphalan with autologous stem cell transplant. A statistically positive benefit, without an increase in side effects, was seen in patients who received the three-drug regimen. [148]

Current Clinical Trials

Use our [advanced clinical trial search](#) to find NCI-supported cancer clinical trials that are now enrolling patients. The search can be narrowed by location of the trial, type of treatment, name of the drug, and other criteria. [General information](#) about clinical trials is also available.

References

1. García Gómez J, Pérez López ME, García Mata J, et al.: SEOM clinical guidelines for the treatment of antiemetic prophylaxis in cancer patients receiving chemotherapy. Clin Transl Oncol 12 (11): 770-4, 2010. [\[PUBMED Abstract\]](#)
2. Hesketh PJ, Kris MG, Basch E, et al.: Antiemetics: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol 35 (28): 3240-3261, 2017. [\[PUBMED Abstract\]](#)
3. Dupuis LL, Boodhan S, Holdsworth M, et al.: Guideline for the prevention of acute nausea and vomiting due to antineoplastic medication in pediatric cancer patients. Pediatr Blood Cancer 60 (7): 1073-82, 2013. [\[PUBMED Abstract\]](#)
4. Dupuis LL, Robinson PD, Boodhan S, et al.: Guideline for the prevention and treatment of anticipatory nausea and vomiting due to chemotherapy in pediatric cancer patients.

Pediatr Blood Cancer 61 (8): 1506-12,
2014. [\[PUBMED Abstract\]](#)

5. Hesketh PJ: Understanding the pathobiology of chemotherapy-induced nausea and vomiting. Providing a basis for therapeutic progress. Oncology (Williston Park) 18 (10 Suppl 6): 9-14, 2004. [\[PUBMED Abstract\]](#)
6. Rudd JA, Andrews PLR: Mechanisms of acute, delayed, and anticipatory emesis induced by anticancer therapies. In: Hesketh PJ, ed.: Management of Nausea and Vomiting in Cancer and Cancer Treatment. Jones and Bartlett Publishers, Inc., 2005, pp 15-66.
7. National Comprehensive Cancer Network: NCCN Clinical Practice Guidelines in Oncology: Antiemesis. Version 2.2024. Plymouth Meeting, Pa: National Comprehensive Cancer Network, 2024. [Available online with free registration](#). Last accessed March 4, 2025.

8. Navari RM: Pathogenesis-based treatment of chemotherapy-induced nausea and vomiting--two new agents. J Support Oncol 1 (2): 89-103, 2003 Jul-Aug. [\[PUBMED Abstract\]](#)
9. Grunberg SM, Deuson RR, Mavros P, et al.: Incidence of chemotherapy-induced nausea and emesis after modern antiemetics. Cancer 100 (10): 2261-8, 2004. [\[PUBMED Abstract\]](#)
10. Fabi A, Barduagni M, Lauro S, et al.: Is delayed chemotherapy-induced emesis well managed in oncological clinical practice? An observational study. Support Care Cancer 11 (3): 156-61, 2003. [\[PUBMED Abstract\]](#)
11. Colagiuri B, Roscoe JA, Morrow GR, et al.: How do patient expectancies, quality of life, and postchemotherapy nausea interrelate? Cancer 113 (3): 654-61, 2008. [\[PUBMED Abstract\]](#)
12. Einhorn LH, Brames MJ, Dreicer R, et al.: Palonosetron plus dexamethasone for prevention of chemotherapy-induced nausea

and vomiting in patients receiving multiple-day cisplatin chemotherapy for germ cell cancer. Support Care Cancer 15 (11): 1293-300, 2007. [\[PUBMED Abstract\]](#)

13. Roila F, Molassiotis A, Herrstedt J, et al.: 2016 MASCC and ESMO guideline update for the prevention of chemotherapy- and radiotherapy-induced nausea and vomiting and of nausea and vomiting in advanced cancer patients. Ann Oncol 27 (suppl 5): v119-v133, 2016. [\[PUBMED Abstract\]](#)
14. Lexicomp Online. Hudson, Ohio: Lexi-Comp, Inc., 2025. [Available online with subscription.](#) Last accessed Jan. 15, 2025.
15. Gez E, Brufman G, Kaufman B, et al.: Methylprednisolone and chlorpromazine in patients receiving cancer chemotherapy: a prospective non-randomized study. J Chemother 1 (2): 140-3, 1989. [\[PUBMED Abstract\]](#)

16. ASHP Therapeutic Guidelines on the Pharmacologic Management of Nausea and Vomiting in Adult and Pediatric Patients Receiving Chemotherapy or Radiation Therapy or Undergoing Surgery. Am J Health Syst Pharm 56 (8): 729-64, 1999. [\[PUBMED Abstract\]](#)
17. Hurley JD, Eshelman FN: Trimethobenzamide HCl in the treatment of nausea and vomiting associated with antineoplastic chemotherapy. J Clin Pharmacol 20 (5-6 Pt 1): 352-6, 1980 May-Jun. [\[PUBMED Abstract\]](#)
18. Schwartzberg L, Roeland E, Andric Z, et al.: Phase III safety study of intravenous NEPA: a novel fixed antiemetic combination of fosnetupitant and palonosetron in patients receiving highly emetogenic chemotherapy. Ann Oncol 29 (7): 1535-1540, 2018. [\[PUBMED Abstract\]](#)
19. Mori K, Saito Y, Tominaga K: Antiemetic efficacy of alprazolam in the combination of

metoclopramide plus methylprednisolone.
Double-blind randomized crossover study in
patients with cisplatin-induced emesis. Am J
Clin Oncol 16 (4): 338-41, 1993. [\[PUBMED
Abstract\]](#)

20. Navari RM, Qin R, Ruddy KJ, et al.: Olanzapine for the Prevention of Chemotherapy-Induced Nausea and Vomiting. N Engl J Med 375 (2): 134-42, 2016. [\[PUBMED Abstract\]](#)
21. Navari RM, Nagy CK, Gray SE: The use of olanzapine versus metoclopramide for the treatment of breakthrough chemotherapy-induced nausea and vomiting in patients receiving highly emetogenic chemotherapy. Support Care Cancer 21 (6): 1655-63, 2013. [\[PUBMED Abstract\]](#)
22. Todaro B: Cannabinoids in the treatment of chemotherapy-induced nausea and vomiting. J Natl Compr Canc Netw 10 (4): 487-92, 2012. [\[PUBMED Abstract\]](#)

23. Ryan JL, Heckler CE, Roscoe JA, et al.: Ginger (*Zingiber officinale*) reduces acute chemotherapy-induced nausea: a URCC CCOP study of 576 patients. *Support Care Cancer* 20 (7): 1479-89, 2012. [\[PUBMED Abstract\]](#)
24. Marx WM, Teleni L, McCarthy AL, et al.: Ginger (*Zingiber officinale*) and chemotherapy-induced nausea and vomiting: a systematic literature review. *Nutr Rev* 71 (4): 245-54, 2013. [\[PUBMED Abstract\]](#)
25. Olver IN, Webster LK, Bishop JF, et al.: A dose finding study of prochlorperazine as an antiemetic for cancer chemotherapy. *Eur J Cancer Clin Oncol* 25 (10): 1457-61, 1989. [\[PUBMED Abstract\]](#)
26. Carr BI, Bertrand M, Browning S, et al.: A comparison of the antiemetic efficacy of prochlorperazine and metoclopramide for the treatment of cisplatin-induced emesis: a prospective, randomized, double-blind study.

J Clin Oncol 3 (8): 1127-32, 1985. [\[PUBMED Abstract\]](#)

27. Olver IN, Wolf M, Laidlaw C, et al.: A randomised double-blind study of high-dose intravenous prochlorperazine versus high-dose metoclopramide as antiemetics for cancer chemotherapy. Eur J Cancer 28A (11): 1798-802, 1992. [\[PUBMED Abstract\]](#)
28. Levinson DF, Simpson GM: Neuroleptic-induced extrapyramidal symptoms with fever. Heterogeneity of the 'neuroleptic malignant syndrome'. Arch Gen Psychiatry 43 (9): 839-48, 1986. [\[PUBMED Abstract\]](#)
29. Wampler G: The pharmacology and clinical effectiveness of phenothiazines and related drugs for managing chemotherapy-induced emesis. Drugs 25 (Supl 1): 35-51, 1983. [\[PUBMED Abstract\]](#)
30. American Geriatrics Society 2012 Beers Criteria Update Expert Panel: American Geriatrics Society updated Beers Criteria for

potentially inappropriate medication use in older adults. J Am Geriatr Soc 60 (4): 616-31, 2012. [\[PUBMED Abstract\]](#)

31. Hardy JR, O'Shea A, White C, et al.: The efficacy of haloperidol in the management of nausea and vomiting in patients with cancer. J Pain Symptom Manage 40 (1): 111-6, 2010. [\[PUBMED Abstract\]](#)
32. Digges M, Hussein A, Wilcock A, et al.: Pharmacovigilance in Hospice/Palliative Care: Net Effect of Haloperidol for Nausea or Vomiting. J Palliat Med 21 (1): 37-43, 2018. [\[PUBMED Abstract\]](#)
33. Navari RM: Comparison of intermittent versus continuous infusion metoclopramide in control of acute nausea induced by cisplatin chemotherapy. J Clin Oncol 7 (7): 943-6, 1989. [\[PUBMED Abstract\]](#)
34. Agostinucci WA, Gannon RH, Golub GR, et al.: Continuous i.v. infusion versus multiple bolus doses of metoclopramide for prevention of

- cisplatin-induced emesis. Clin Pharm 7 (6): 454-7, 1988. [\[PUBMED Abstract\]](#)
35. Kris MG, Gralla RJ, Tyson LB, et al.: Improved control of cisplatin-induced emesis with high-dose metoclopramide and with combinations of metoclopramide, dexamethasone, and diphenhydramine. Results of consecutive trials in 255 patients. Cancer 55 (3): 527-34, 1985. [\[PUBMED Abstract\]](#)
36. Kris MG, Tyson LB, Gralla RJ, et al.: Extrapyramidal reactions with high-dose metoclopramide. N Engl J Med 309 (7): 433-4, 1983. [\[PUBMED Abstract\]](#)
37. Hsu ES: A review of granisetron, 5-hydroxytryptamine₃ receptor antagonists, and other antiemetics. Am J Ther 17 (5): 476-86, 2010 Sep-Oct. [\[PUBMED Abstract\]](#)
38. Trammel M, Roederer M, Patel J, et al.: Does pharmacogenomics account for variability in control of acute chemotherapy-induced nausea and vomiting with 5-

hydroxytryptamine type 3 receptor antagonists? Curr Oncol Rep 15 (3): 276-85, 2013. [\[PUBMED Abstract\]](#)

39. Hatoum HT, Lin SJ, Buchner D, et al.: Comparative clinical effectiveness of various 5-HT₃ RA antiemetic regimens on chemotherapy-induced nausea and vomiting associated with hospital and emergency department visits in real world practice. Support Care Cancer 20 (5): 941-9, 2012. [\[PUBMED Abstract\]](#)
40. Aogi K, Sakai H, Yoshizawa H, et al.: A phase III open-label study to assess safety and efficacy of palonosetron for preventing chemotherapy-induced nausea and vomiting (CINV) in repeated cycles of emetogenic chemotherapy. Support Care Cancer 20 (7): 1507-14, 2012. [\[PUBMED Abstract\]](#)
41. Hesketh PJ: Comparative review of 5-HT₃ receptor antagonists in the treatment of acute chemotherapy-induced nausea and

vomiting. Cancer Invest 18 (2): 163-73, 2000. [\[PUBMED Abstract\]](#)

42. Navari R, Gandara D, Hesketh P, et al.: Comparative clinical trial of granisetron and ondansetron in the prophylaxis of cisplatin-induced emesis. The Granisetron Study Group. J Clin Oncol 13 (5): 1242-8, 1995. [\[PUBMED Abstract\]](#)
43. Eisenberg P, Figueroa-Vadillo J, Zamora R, et al.: Improved prevention of moderately emetogenic chemotherapy-induced nausea and vomiting with palonosetron, a pharmacologically novel 5-HT₃ receptor antagonist: results of a phase III, single-dose trial versus dolasetron. Cancer 98 (11): 2473-82, 2003. [\[PUBMED Abstract\]](#)
44. Gralla R, Lichinitser M, Van Der Veegt S, et al.: Palonosetron improves prevention of chemotherapy-induced nausea and vomiting following moderately emetogenic chemotherapy: results of a double-blind

randomized phase III trial comparing single doses of palonosetron with ondansetron. Ann Oncol 14 (10): 1570-7, 2003. [\[PUBMED Abstract\]](#)

45. Boccia R, Grunberg S, Franco-Gonzales E, et al.: Efficacy of oral palonosetron compared to intravenous palonosetron for the prevention of chemotherapy-induced nausea and vomiting associated with moderately emetogenic chemotherapy: a phase 3 trial. Support Care Cancer 21 (5): 1453-60, 2013. [\[PUBMED Abstract\]](#)
46. Hickok JT, Roscoe JA, Morrow GR, et al.: 5-Hydroxytryptamine-receptor antagonists versus prochlorperazine for control of delayed nausea caused by doxorubicin: a URCC CCOP randomised controlled trial. Lancet Oncol 6 (10): 765-72, 2005. [\[PUBMED Abstract\]](#)
47. Kris MG: Why do we need another antiemetic? Just ask. J Clin Oncol 21 (22):

4077-80, 2003. [\[PUBMED Abstract\]](#)

48. Tyers MB: Pharmacology and preclinical antiemetic properties of ondansetron. Semin Oncol 19 (4 Suppl 10): 1-8, 1992. [\[PUBMED Abstract\]](#)
49. Kaasa S, Kvaløy S, Dicato MA, et al.: A comparison of ondansetron with metoclopramide in the prophylaxis of chemotherapy-induced nausea and vomiting: a randomized, double-blind study. International Emesis Study Group. Eur J Cancer 26 (3): 311-4, 1990. [\[PUBMED Abstract\]](#)
50. Hainsworth J, Harvey W, Pendergrass K, et al.: A single-blind comparison of intravenous ondansetron, a selective serotonin antagonist, with intravenous metoclopramide in the prevention of nausea and vomiting associated with high-dose cisplatin chemotherapy. J Clin Oncol 9 (5): 721-8, 1991. [\[PUBMED Abstract\]](#)

51. De Mulder PH, Seynaeve C, Vermorken JB, et al.: Ondansetron compared with high-dose metoclopramide in prophylaxis of acute and delayed cisplatin-induced nausea and vomiting. A multicenter, randomized, double-blind, crossover study. *Ann Intern Med* 113 (11): 834-40, 1990. [\[PUBMED Abstract\]](#)
52. Marty M, Pouillart P, Scholl S, et al.: Comparison of the 5-hydroxytryptamine₃ (serotonin) antagonist ondansetron (GR 38032F) with high-dose metoclopramide in the control of cisplatin-induced emesis. *N Engl J Med* 322 (12): 816-21, 1990. [\[PUBMED Abstract\]](#)
53. Hesketh PJ: Comparative trials of ondansetron versus metoclopramide in the prevention of acute cisplatin-induced emesis. *Semin Oncol* 19 (4 Suppl 10): 33-40, 1992. [\[PUBMED Abstract\]](#)
54. Pectasides D, Mylonakis A, Varthalitis J, et al.: Comparison of two different doses of

ondansetron plus dexamethasone in the prophylaxis of cisplatin-induced emesis.

Oncology 54 (1): 1-6, 1997 Jan-Feb. [[PUBMED Abstract](#)]

55. Hasler SB, Hirt A, Ridolfi Luethy A, et al.: Safety of ondansetron loading doses in children with cancer. Support Care Cancer 16 (5): 469-75, 2008. [[PUBMED Abstract](#)]
56. U.S. Food and Drug Administration: FDA Drug Safety Communication: New information regarding QT prolongation with ondansetron (Zofran). Silver Spring, Md: U.S. Food and Drug Administration, 2012. [Available online](#). Last accessed March 4, 2025.
57. Beck TM, Hesketh PJ, Madajewicz S, et al.: Stratified, randomized, double-blind comparison of intravenous ondansetron administered as a multiple-dose regimen versus two single-dose regimens in the prevention of cisplatin-induced nausea and

vomiting. J Clin Oncol 10 (12): 1969-75, 1992. [\[PUBMED Abstract\]](#)

58. Finn AL: Toxicity and side effects of ondansetron. Semin Oncol 19 (4 Suppl 10): 53-60, 1992. [\[PUBMED Abstract\]](#)

59. Coates AS, Childs A, Cox K, et al.: Severe vascular adverse effects with thrombocytopenia and renal failure following emetogenic chemotherapy and ondansetron. Ann Oncol 3 (9): 719-22, 1992. [\[PUBMED Abstract\]](#)

60. Ruff P, Paska W, Goedhals L, et al.: Ondansetron compared with granisetron in the prophylaxis of cisplatin-induced acute emesis: a multicentre double-blind, randomised, parallel-group study. The Ondansetron and Granisetron Emesis Study Group. Oncology 51 (1): 113-8, 1994 Jan-Feb. [\[PUBMED Abstract\]](#)

61. Jantunen IT, Muhonen TT, Kataja VV, et al.: 5-HT3 receptor antagonists in the prophylaxis

of acute vomiting induced by moderately emetogenic chemotherapy--a randomised study. Eur J Cancer 29A (12): 1669-72, 1993. [\[PUBMED Abstract\]](#)

62. Gebbia V, Cannata G, Testa A, et al.: Ondansetron versus granisetron in the prevention of chemotherapy-induced nausea and vomiting. Results of a prospective randomized trial. Cancer 74 (7): 1945-52, 1994. [\[PUBMED Abstract\]](#)
63. Boccia RV, Gordan LN, Clark G, et al.: Efficacy and tolerability of transdermal granisetron for the control of chemotherapy-induced nausea and vomiting associated with moderately and highly emetogenic multi-day chemotherapy: a randomized, double-blind, phase III study. Support Care Cancer 19 (10): 1609-17, 2011. [\[PUBMED Abstract\]](#)
64. Raftopoulos H, Cooper W, O'Boyle E, et al.: Comparison of an extended-release formulation of granisetron (APF530) versus

palonosetron for the prevention of chemotherapy-induced nausea and vomiting associated with moderately or highly emetogenic chemotherapy: results of a prospective, randomized, double-blind, noninferiority phase 3 trial. Support Care Cancer 23 (3): 723-32, 2015. [\[PUBMED Abstract\]](#)

65. U.S. Food and Drug Administration: FDA Drug Safety Communication: Abnormal heart rhythms associated with use of Anzemet (dolasetron mesylate). Silver Spring, Md: U.S. Food and Drug Administration, 2010. [Available online](#). Last accessed March 4, 2025.
66. Fauser AA, Duclos B, Chemaissani A, et al.: Therapeutic equivalence of single oral doses of dolasetron mesilate and multiple doses of ondansetron for the prevention of emesis after moderately emetogenic chemotherapy. European Dolasetron Comparative Study Group. Eur J Cancer 32A (9): 1523-9, 1996. [\[PUBMED Abstract\]](#)

67. Eisenberg P, MacKintosh FR, Ritch P, et al.: Efficacy, safety and pharmacokinetics of palonosetron in patients receiving highly emetogenic cisplatin-based chemotherapy: a dose-ranging clinical study. *Ann Oncol* 15 (2): 330-7, 2004. [\[PUBMED Abstract\]](#)
68. Tricco AC, Soobiah C, Antony J, et al.: Safety of serotonin (5-HT₃) receptor antagonists in patients undergoing surgery and chemotherapy: protocol for a systematic review and network meta-analysis. *Syst Rev* 2: 46, 2013. [\[PUBMED Abstract\]](#)
69. Faria C, Li X, Nagl N, et al.: Outcomes Associated with 5-HT₃-RA Therapy Selection in Patients with Chemotherapy-Induced Nausea and Vomiting: A Retrospective Claims Analysis. *Am Health Drug Benefits* 7 (1): 50-8, 2014. [\[PUBMED Abstract\]](#)
70. Schwartzberg L, Barbour SY, Morrow GR, et al.: Pooled analysis of phase III clinical studies of palonosetron versus ondansetron,

dolasetron, and granisetron in the prevention of chemotherapy-induced nausea and vomiting (CINV). Support Care Cancer 22 (2): 469-77, 2014. [\[PUBMED Abstract\]](#)

71. Affronti ML, Bubalo J: Palonosetron in the management of chemotherapy-induced nausea and vomiting in patients receiving multiple-day chemotherapy. Cancer Manag Res 6: 329-37, 2014. [\[PUBMED Abstract\]](#)

72. Navari RM: Palonosetron for the prevention of chemotherapy-induced nausea and vomiting: approval and efficacy. Cancer Manag Res 1: 167-76, 2009. [\[PUBMED Abstract\]](#)

73. Aapro MS, Grunberg SM, Manikhas GM, et al.: A phase III, double-blind, randomized trial of palonosetron compared with ondansetron in preventing chemotherapy-induced nausea and vomiting following highly emetogenic chemotherapy. Ann Oncol 17 (9): 1441-9, 2006. [\[PUBMED Abstract\]](#)

74. Navari RM: Palonosetron: a second-generation 5-hydroxytryptamine receptor antagonist. *Future Oncol* 2 (5): 591-602, 2006. [\[PUBMED Abstract\]](#)
75. Humphreys S, Pellissier J, Jones A: Cost-effectiveness of an aprepitant regimen for prevention of chemotherapy-induced nausea and vomiting in patients with breast cancer in the UK. *Cancer Manag Res* 5: 215-24, 2013. [\[PUBMED Abstract\]](#)
76. Warr DG, Street JC, Carides AD: Evaluation of risk factors predictive of nausea and vomiting with current standard-of-care antiemetic treatment: analysis of phase 3 trial of aprepitant in patients receiving adriamycin-cyclophosphamide-based chemotherapy. *Support Care Cancer* 19 (6): 807-13, 2011. [\[PUBMED Abstract\]](#)
77. Hesketh PJ, Aapro M, Street JC, et al.: Evaluation of risk factors predictive of nausea and vomiting with current standard-of-care

antiemetic treatment: analysis of two phase III trials of aprepitant in patients receiving cisplatin-based chemotherapy. Support Care Cancer 18 (9): 1171-7, 2010. [\[PUBMED Abstract\]](#)

78. Hu W, Fang J, Nie J, et al.: Addition of aprepitant improves protection against cisplatin-induced emesis when a conventional anti-emetic regimen fails. Cancer Chemother Pharmacol 73 (6): 1129-36, 2014. [\[PUBMED Abstract\]](#)
79. Poli-Bigelli S, Rodrigues-Pereira J, Carides AD, et al.: Addition of the neurokinin 1 receptor antagonist aprepitant to standard antiemetic therapy improves control of chemotherapy-induced nausea and vomiting. Results from a randomized, double-blind, placebo-controlled trial in Latin America. Cancer 97 (12): 3090-8, 2003. [\[PUBMED Abstract\]](#)
80. Hesketh PJ, Grunberg SM, Gralla RJ, et al.: The oral neurokinin-1 antagonist aprepitant for

the prevention of chemotherapy-induced nausea and vomiting: a multinational, randomized, double-blind, placebo-controlled trial in patients receiving high-dose cisplatin--the Aprepitant Protocol 052 Study Group. J Clin Oncol 21 (22): 4112-9, 2003. [\[PUBMED Abstract\]](#)

81. Schmoll HJ, Aapro MS, Poli-Bigelli S, et al.: Comparison of an aprepitant regimen with a multiple-day ondansetron regimen, both with dexamethasone, for antiemetic efficacy in high-dose cisplatin treatment. Ann Oncol 17 (6): 1000-6, 2006. [\[PUBMED Abstract\]](#)
82. Warr DG, Hesketh PJ, Gralla RJ, et al.: Efficacy and tolerability of aprepitant for the prevention of chemotherapy-induced nausea and vomiting in patients with breast cancer after moderately emetogenic chemotherapy. J Clin Oncol 23 (12): 2822-30, 2005. [\[PUBMED Abstract\]](#)

83. Bubalo JS, Herrington JD, Takemoto M, et al.: Phase II open label pilot trial of aprepitant and palonosetron for the prevention of chemotherapy-induced nausea and vomiting (CINV) in patients receiving moderately emetogenic FOLFOX chemotherapy for the treatment of colorectal cancer. Support Care Cancer 26 (4): 1273-1279, 2018. [\[PUBMED Abstract\]](#)
84. Rapoport BL, Jordan K, Boice JA, et al.: Aprepitant for the prevention of chemotherapy-induced nausea and vomiting associated with a broad range of moderately emetogenic chemotherapies and tumor types: a randomized, double-blind study. Support Care Cancer 18 (4): 423-31, 2010. [\[PUBMED Abstract\]](#)
85. Aapro MS, Schmoll HJ, Jahn F, et al.: Review of the efficacy of aprepitant for the prevention of chemotherapy-induced nausea and vomiting in a range of tumor types. Cancer

Treat Rev 39 (1): 113-7, 2013. [\[PUBMED Abstract\]](#)

86. Albany C, Brames MJ, Fausel C, et al.: Randomized, double-blind, placebo-controlled, phase III cross-over study evaluating the oral neurokinin-1 antagonist aprepitant in combination with a 5HT3 receptor antagonist and dexamethasone in patients with germ cell tumors receiving 5-day cisplatin combination chemotherapy regimens: a hoosier oncology group study. J Clin Oncol 30 (32): 3998-4003, 2012. [\[PUBMED Abstract\]](#)
87. Grunberg S, Chua D, Maru A, et al.: Single-dose fosaprepitant for the prevention of chemotherapy-induced nausea and vomiting associated with cisplatin therapy: randomized, double-blind study protocol--EASE. J Clin Oncol 29 (11): 1495-501, 2011. [\[PUBMED Abstract\]](#)

88. EMEND (Fosaprepitant Dimeglumine) for Injection, for Intravenous Use. Whitehouse Station, NJ: Merck & Co., Inc., 2022. [Available online](#). Last accessed March 4, 2025.
89. Shelley WB, Talanin N, Shelley ED: Polysorbate 80 hypersensitivity. Lancet 345 (8960): 1312-3, 1995. [\[PUBMED Abstract\]](#)
90. CINVANTI (aprepitant): highlights of prescribing information. San Diego, Calif: Heron Therapeutics, 2024. [Available online](#). Last accessed March 4, 2025.
91. Schwartzberg L, Navari R, Clark-Snow R, et al.: Phase IIIb Safety and Efficacy of Intravenous NEPA for Prevention of Chemotherapy-Induced Nausea and Vomiting (CINV) in Patients with Breast Cancer Receiving Initial and Repeat Cycles of Anthracycline and Cyclophosphamide (AC) Chemotherapy. Oncologist 25 (3): e589-e597, 2020. [\[PUBMED Abstract\]](#)

92. Hesketh PJ, Rossi G, Rizzi G, et al.: Efficacy and safety of NEPA, an oral combination of netupitant and palonosetron, for prevention of chemotherapy-induced nausea and vomiting following highly emetogenic chemotherapy: a randomized dose-ranging pivotal study. *Ann Oncol* 25 (7): 1340-6, 2014. [\[PUBMED Abstract\]](#)
93. Aapro M, Rugo H, Rossi G, et al.: A randomized phase III study evaluating the efficacy and safety of NEPA, a fixed-dose combination of netupitant and palonosetron, for prevention of chemotherapy-induced nausea and vomiting following moderately emetogenic chemotherapy. *Ann Oncol* 25 (7): 1328-33, 2014. [\[PUBMED Abstract\]](#)
94. Aapro M, Karthaus M, Schwartzberg L, et al.: NEPA, a fixed oral combination of netupitant and palonosetron, improves control of chemotherapy-induced nausea and vomiting (CINV) over multiple cycles of chemotherapy: results of a randomized, double-blind, phase

3 trial versus oral palonosetron. Support Care Cancer 25 (4): 1127-1135, 2017. [[PUBMED Abstract](#)]

95. Hata A, Okamoto I, Inui N, et al.: Randomized, Double-Blind, Phase III Study of Fosnetupitant Versus Fosaprepitant for Prevention of Highly Emetogenic Chemotherapy-Induced Nausea and Vomiting: CONSOLE. J Clin Oncol 40 (2): 180-188, 2022. [[PUBMED Abstract](#)]
96. Zhang L, Lu S, Feng J, et al.: A randomized phase III study evaluating the efficacy of single-dose NEPA, a fixed antiemetic combination of netupitant and palonosetron, versus an aprepitant regimen for prevention of chemotherapy-induced nausea and vomiting (CINV) in patients receiving highly emetogenic chemotherapy (HEC). Ann Oncol 29 (2): 452-458, 2018. [[PUBMED Abstract](#)]
97. Schwartzberg LS, Modiano MR, Rapoport BL, et al.: Safety and efficacy of rolapitant for prevention of chemotherapy-induced nausea

and vomiting after administration of moderately emetogenic chemotherapy or anthracycline and cyclophosphamide regimens in patients with cancer: a randomised, active-controlled, double-blind, phase 3 trial. *Lancet Oncol* 16 (9): 1071-8, 2015. [\[PUBMED Abstract\]](#)

98. Rapoport BL, Chasen MR, Gridelli C, et al.: Safety and efficacy of rolapitant for prevention of chemotherapy-induced nausea and vomiting after administration of cisplatin-based highly emetogenic chemotherapy in patients with cancer: two randomised, active-controlled, double-blind, phase 3 trials. *Lancet Oncol* 16 (9): 1079-89, 2015. [\[PUBMED Abstract\]](#)
99. VARUBI (rolapitant): highlights of prescribing information. Lake Forest, Ill: TerSera Therapeutics LLC, 2020. [Available online](#). Last accessed March 4, 2025.

100. U.S. Food and Drug Administration:
Postmarket Drug Safety Information for
Patients and Providers: Varubi (rolapitant)
injectable emulsion. Silver Spring, Md: U.S.
Food and Drug Administration, 2018.
[Available online](#). Last accessed March 4, 2025.
101. Bishop JF, Matthews JP, Wolf MM, et al.: A
randomised trial of dexamethasone,
lorazepam and prochlorperazine for emesis
in patients receiving chemotherapy. Eur J
Cancer 28 (1): 47-50, 1992. [\[PUBMED
Abstract\]](#)
102. Chiara S, Campora E, Lionetto R, et al.:
Methylprednisolone for the control of CMF-
induced emesis. Am J Clin Oncol 10 (3): 264-7,
1987. [\[PUBMED Abstract\]](#)
103. Parry H, Martin K: Single-dose i.v.
dexamethasone--an effective anti-emetic in
cancer chemotherapy. Cancer Chemother
Pharmacol 28 (3): 231-2, 1991. [\[PUBMED
Abstract\]](#)

104. Cassileth PA, Lusk EJ, Torri S, et al.: Antiemetic efficacy of high-dose dexamethasone in induction therapy in acute nonlymphocytic leukemia. *Ann Intern Med* 100 (5): 701-2, 1984. [\[PUBMED Abstract\]](#)
105. Aapro MS, Plezia PM, Alberts DS, et al.: Double-blind crossover study of the antiemetic efficacy of high-dose dexamethasone versus high-dose metoclopramide. *J Clin Oncol* 2 (5): 466-71, 1984. [\[PUBMED Abstract\]](#)
106. Schallier D, Van Belle S, De Greve J, et al.: Methylprednisolone as an antiemetic drug. A randomised double blind study. *Cancer Chemother Pharmacol* 14 (3): 235-7, 1985. [\[PUBMED Abstract\]](#)
107. Kris MG, Gralla RJ, Clark RA, et al.: Consecutive dose-finding trials adding lorazepam to the combination of metoclopramide plus dexamethasone: improved subjective effectiveness over the

combination of diphenhydramine plus metoclopramide plus dexamethasone. Cancer Treat Rep 69 (11): 1257-62, 1985. [\[PUBMED Abstract\]](#)

108. Greenberg DB, Surman OS, Clarke J, et al.: Alprazolam for phobic nausea and vomiting related to cancer chemotherapy. Cancer Treat Rep 71 (5): 549-50, 1987. [\[PUBMED Abstract\]](#)

109. Hockenberry-Eaton M, Benner A: Patterns of nausea and vomiting in children: nursing assessment and intervention. Oncol Nurs Forum 17 (4): 575-84, 1990 Jul-Aug. [\[PUBMED Abstract\]](#)

110. Münstedt K, Borces D, Bohlmann MK, et al.: Glucocorticoid administration in antiemetic therapy: is it safe? Cancer 101 (7): 1696-702, 2004. [\[PUBMED Abstract\]](#)

111. Roila F, Tonato M, Cognetti F, et al.: Prevention of cisplatin-induced emesis: a double-blind multicenter randomized

crossover study comparing ondansetron and ondansetron plus dexamethasone. J Clin Oncol 9 (4): 675-8, 1991. [\[PUBMED Abstract\]](#)

112. Zaglama NE, Rosenblum SL, Sartiano GP, et al.: Single, high-dose intravenous dexamethasone as an antiemetic in cancer chemotherapy. Oncology 43 (1): 27-32, 1986. [\[PUBMED Abstract\]](#)
113. Klygis LM: Dexamethasone-induced perineal irritation in head injury. Am J Emerg Med 10 (3): 268, 1992. [\[PUBMED Abstract\]](#)
114. More on dexamethasone-induced perineal irritation. N Engl J Med 314 (25): 1643-4, 1986. [\[PUBMED Abstract\]](#)
115. Baharav E, Harpaz D, Mittelman M, et al.: Dexamethasone-induced perineal irritation. N Engl J Med 314 (8): 515-6, 1986. [\[PUBMED Abstract\]](#)
116. Triozzi PL, Goldstein D, Laszlo J: Contributions of benzodiazepines to cancer therapy. Cancer Invest 6 (1): 103-11, 1988. [\[PUBMED Abstract\]](#)

117. Laszlo J, Clark RA, Hanson DC, et al.:
Lorazepam in cancer patients treated with cisplatin: a drug having antiemetic, amnesic, and anxiolytic effects. J Clin Oncol 3 (6): 864-9, 1985. [\[PUBMED Abstract\]](#)
118. Bishop JF, Olver IN, Wolf MM, et al.:
Lorazepam: a randomized, double-blind, crossover study of a new antiemetic in patients receiving cytotoxic chemotherapy and prochlorperazine. J Clin Oncol 2 (6): 691-5, 1984. [\[PUBMED Abstract\]](#)
119. Henry DW, Burwinkle JW, Klutman NE:
Determination of sedative and amnestic doses of lorazepam in children. Clin Pharm 10 (8): 625-9, 1991. [\[PUBMED Abstract\]](#)
120. van Hoff J, Olszewski D: Lorazepam for the control of chemotherapy-related nausea and vomiting in children. J Pediatr 113 (1 Pt 1): 146-9, 1988. [\[PUBMED Abstract\]](#)
121. Bymaster FP, Falcone JF, Bauzon D, et al.:
Potent antagonism of 5-HT(3) and 5-HT(6)

receptors by olanzapine. Eur J Pharmacol 430 (2-3): 341-9, 2001. [\[PUBMED Abstract\]](#)

122. ZYPREXA (olanzapine): highlights of prescribing information. Indianapolis, Ind: Eli Lilly and Company, 2021. [Available online](#). Last accessed March 4, 2025.
123. Hocking CM, Kichenadasse G: Olanzapine for chemotherapy-induced nausea and vomiting: a systematic review. Support Care Cancer 22 (4): 1143-51, 2014. [\[PUBMED Abstract\]](#)
124. Passik SD, Lundberg J, Kirsh KL, et al.: A pilot exploration of the antiemetic activity of olanzapine for the relief of nausea in patients with advanced cancer and pain. J Pain Symptom Manage 23 (6): 526-32, 2002. [\[PUBMED Abstract\]](#)
125. Navari RM, Einhorn LH, Passik SD, et al.: A phase II trial of olanzapine for the prevention of chemotherapy-induced nausea and vomiting: a Hoosier Oncology Group study.

Support Care Cancer 13 (7): 529-34,
2005. [\[PUBMED Abstract\]](#)

126. Navari RM, Einhorn LH, Loehrer PJ, et al.: A phase II trial of olanzapine, dexamethasone, and palonosetron for the prevention of chemotherapy-induced nausea and vomiting: a Hoosier oncology group study. Support Care Cancer 15 (11): 1285-91, 2007. [\[PUBMED Abstract\]](#)
127. Tan L, Liu J, Liu X, et al.: Clinical research of Olanzapine for prevention of chemotherapy-induced nausea and vomiting. J Exp Clin Cancer Res 28: 131, 2009. [\[PUBMED Abstract\]](#)
128. Navari RM, Gray SE, Kerr AC: Olanzapine versus aprepitant for the prevention of chemotherapy-induced nausea and vomiting: a randomized phase III trial. J Support Oncol 9 (5): 188-95, 2011 Sep-Oct. [\[PUBMED Abstract\]](#)
129. Mizukami N, Yamauchi M, Koike K, et al.: Olanzapine for the prevention of chemotherapy-induced nausea and vomiting

in patients receiving highly or moderately emetogenic chemotherapy: a randomized, double-blind, placebo-controlled study. J Pain Symptom Manage 47 (3): 542-50, 2014. [\[PUBMED Abstract\]](#)

130. Ben Amar M: Cannabinoids in medicine: A review of their therapeutic potential. J Ethnopharmacol 105 (1-2): 1-25, 2006. [\[PUBMED Abstract\]](#)

131. Davis MP: Oral nabilone capsules in the treatment of chemotherapy-induced nausea and vomiting and pain. Expert Opin Investig Drugs 17 (1): 85-95, 2008. [\[PUBMED Abstract\]](#)

132. Ahmedzai S, Carlyle DL, Calder IT, et al.: Anti-emetic efficacy and toxicity of nabilone, a synthetic cannabinoid, in lung cancer chemotherapy. Br J Cancer 48 (5): 657-63, 1983. [\[PUBMED Abstract\]](#)

133. Chan HS, Correia JA, MacLeod SM: Nabilone versus prochlorperazine for control of cancer chemotherapy-induced emesis in children: a

double-blind, crossover trial. Pediatrics 79 (6): 946-52, 1987. [\[PUBMED Abstract\]](#)

134. Johansson R, Kilkku P, Groenroos M: A double-blind, controlled trial of nabilone vs. prochlorperazine for refractory emesis induced by cancer chemotherapy. Cancer Treat Rev 9 (Suppl B): 25-33, 1982. [\[PUBMED Abstract\]](#)
135. Niiranen A, Mattson K: A cross-over comparison of nabilone and prochlorperazine for emesis induced by cancer chemotherapy. Am J Clin Oncol 8 (4): 336-40, 1985. [\[PUBMED Abstract\]](#)
136. Tramèr MR, Carroll D, Campbell FA, et al.: Cannabinoids for control of chemotherapy induced nausea and vomiting: quantitative systematic review. BMJ 323 (7303): 16-21, 2001. [\[PUBMED Abstract\]](#)
137. Meiri E, Jhangiani H, Vredenburg JJ, et al.: Efficacy of dronabinol alone and in combination with ondansetron versus

ondansetron alone for delayed chemotherapy-induced nausea and vomiting. Curr Med Res Opin 23 (3): 533-43, 2007. [\[PUBMED Abstract\]](#)

138. Strasser F, Luftner D, Possinger K, et al.: Comparison of orally administered cannabis extract and delta-9-tetrahydrocannabinol in treating patients with cancer-related anorexia-cachexia syndrome: a multicenter, phase III, randomized, double-blind, placebo-controlled clinical trial from the Cannabis-In-Cachexia-Study-Group. J Clin Oncol 24 (21): 3394-400, 2006. [\[PUBMED Abstract\]](#)
139. Coffman KL: The debate about marijuana usage in transplant candidates: recent medical evidence on marijuana health effects. Curr Opin Organ Transplant 13 (2): 189-95, 2008. [\[PUBMED Abstract\]](#)
140. Bossi P, Cortinovis D, Fatigoni S, et al.: A randomized, double-blind, placebo-controlled, multicenter study of a ginger

extract in the management of chemotherapy-induced nausea and vomiting (CINV) in patients receiving high-dose cisplatin. Ann Oncol 28 (10): 2547-2551, 2017. [\[PUBMED Abstract\]](#)

141. Einhorn LH, Rapoport B, Koeller J, et al.: Antiemetic therapy for multiple-day chemotherapy and high-dose chemotherapy with stem cell transplant: review and consensus statement. Support Care Cancer 13 (2): 112-6, 2005. [\[PUBMED Abstract\]](#)
142. Baltzer L, Pisters KM, Kris MG, et al.: High dose ondansetron (OND) plus dexamethasone (DEX) for the prevention of nausea and vomiting with multiple day cisplatin chemotherapy. [Abstract] Proceedings of the American Society of Clinical Oncology 12: A-1607, 462, 1993.
143. Lorusso V, Giampaglia M, Petrucelli L, et al.: Antiemetic efficacy of single-dose palonosetron and dexamethasone in patients

receiving multiple cycles of multiple day-based chemotherapy. Support Care Cancer 20 (12): 3241-6, 2012. [[PUBMED Abstract](#)]

144. Jordan K, Kinitz I, Voigt W, et al.: Safety and efficacy of a triple antiemetic combination with the NK-1 antagonist aprepitant in highly and moderately emetogenic multiple-day chemotherapy. Eur J Cancer 45 (7): 1184-7, 2009. [[PUBMED Abstract](#)]
145. Olver IN, Grimison P, Chatfield M, et al.: Results of a 7-day aprepitant schedule for the prevention of nausea and vomiting in 5-day cisplatin-based germ cell tumor chemotherapy. Support Care Cancer 21 (6): 1561-8, 2013. [[PUBMED Abstract](#)]
146. Adra N, Albany C, Brames MJ, et al.: Phase II study of fosaprepitant + 5HT3 receptor antagonist + dexamethasone in patients with germ cell tumors undergoing 5-day cisplatin-based chemotherapy: a Hoosier Cancer Research Network study. Support Care

Cancer 24 (7): 2837-42, 2016. [\[PUBMED Abstract\]](#)

147. Stiff PJ, Fox-Geiman MP, Kiley K, et al.: Prevention of nausea and vomiting associated with stem cell transplant: results of a prospective, randomized trial of aprepitant used with highly emetogenic preparative regimens. Biol Blood Marrow Transplant 19 (1): 49-55.e1, 2013. [\[PUBMED Abstract\]](#)
148. Schmitt T, Goldschmidt H, Neben K, et al.: Aprepitant, granisetron, and dexamethasone for prevention of chemotherapy-induced nausea and vomiting after high-dose melphalan in autologous transplantation for multiple myeloma: results of a randomized, placebo-controlled phase III trial. J Clin Oncol 32 (30): 3413-20, 2014. [\[PUBMED Abstract\]](#)
149. Sakurai M, Mori T, Kato J, et al.: Efficacy of aprepitant in preventing nausea and vomiting due to high-dose melphalan-based

conditioning for allogeneic hematopoietic stem cell transplantation. Int J Hematol 99 (4): 457-62, 2014. [\[PUBMED Abstract\]](#)

150. Bubalo JS, Cherala G, McCune JS, et al.: Aprepitant pharmacokinetics and assessing the impact of aprepitant on cyclophosphamide metabolism in cancer patients undergoing hematopoietic stem cell transplantation. J Clin Pharmacol 52 (4): 586-94, 2012. [\[PUBMED Abstract\]](#)
151. Yeh SP, Lo WC, Hsieh CY, et al.: Palonosetron and dexamethasone for the prevention of nausea and vomiting in patients receiving allogeneic hematopoietic stem cell transplantation. Support Care Cancer 22 (5): 1199-206, 2014. [\[PUBMED Abstract\]](#)

Nonpharmacological Management of Nausea and

Vomiting

Nonpharmacological strategies are also used to manage nausea and vomiting (N&V). These strategies include the following:

- Dietary alterations. For more information, see the [Behavioral strategies for symptom management](#) section in Nutrition in Cancer Care.
- Hypnosis.
- Acupuncture. For more information, see [Acupuncture](#).
- Acupressure.
- Relaxation techniques.
- Behavioral therapy.
- Guided imagery.

Guided imagery, hypnosis, and systematic desensitization as means to progressive muscle relaxation have been the most frequently studied

treatments for anticipatory N&V (ANV). They are the recommended treatments for this classically conditioned response. For more information, see the [Treatment of ANV](#) section.

Radiation-Induced Nausea and Vomiting

Radiation therapy (RT) is an important cause of nausea and vomiting (N&V) in patients with cancer. Observational studies suggest that some degree of N&V occurs in 80% of patients undergoing RT.[1] Risk factors for developing N&V are known. Radiation-induced N&V (RINV) worsen quality of life, leading to treatment delays and cancelled appointments and compromising cancer control. [2,3]

Epidemiology

Two large prospective observational studies provide information on the frequency of RINV and antiemetic measures. The Italian Group for Antiemetic Research in Radiotherapy analyzed the incidence of RINV in 1,020 patients receiving various kinds of RT.[4] Overall, 28% of patients reported nausea, vomiting, or both. The median time to the first episode of vomiting was 3 days. Antiemetic drugs were administered to 17% of the patients, including 12% treated prophylactically and 5% given rescue therapy. In a second cohort of 368 patients receiving RT, the overall incidence rate for nausea was 39% and for vomiting, 7%.[5] Nausea was more frequent in patients receiving RT to the lower abdomen or pelvis (66%), compared with patients receiving RT to the head-and-neck area (48%). Antiemetics during RT are underprescribed.[6]

Pathophysiology of RINV

The pathophysiology of RINV is incompletely understood. Serotonin, substance P, and

dopamine are neurotransmitters involved in radiation-induced emesis.[7] RINV bears a close similarity to chemotherapy-induced N&V (CINV). The effectiveness of serotonin antagonists in RINV supports a role for serotonin in radiation-induced emesis.[7] Substance P antagonists have not been used in RINV as extensively as in CINV. Preclinical work suggests a role for substance P in RINV.[8] Substance P antagonists are only beginning to be studied for RINV. Substance P may play a role in prolonged N&V after the administration of RT.

Risk Stratification

The incidence and severity of RINV are determined by:

- Radiation site.
- Volume.
- Fractionation schedule.
- Single and total dose.

The most important factor appears to be the radiation field. The risk of N&V for a patient being treated with RT depends on multiple other factors in addition to the emetogenicity of the specific RT regimen. Patient-specific factors include the following:[3]

- Simultaneous administration of chemotherapy.
- Age.
- Gender.
- Alcohol consumption.
- Anxiety.
- Previous experience of RINV or CINV.

Prevention and Treatment of RINV

The body of literature describing treatments for RINV is much smaller than for CINV.[9] Most of the studies were for patients with moderate- to high-risk features for RINV.

Antiemetic therapy: Prevention and treatment of N&V

Several studies show the superiority of serotonin antagonists for the prophylaxis of RINV.[10-15] For example, ondansetron and dolasetron have shown superiority over placebo or metoclopramide.

Dosing of the serotonin antagonists has been single-dose pretreatment or for consecutive days (up to 5–7 days total). Most studies have been conducted in patients at moderate to high risk of RINV.

Recommended dosing is ondansetron 8 mg, regardless of schedule given.[3] Granisetron dosing is 2 mg orally per day.[3] A recent meta-analysis covering nine clinical trials showed differing rates of control when emesis versus nausea is considered. Compared with placebo, fewer patients had residual emesis (40% vs. 57%; relative risk [RR], 0.7), and fewer patients required rescue medication (6.5% vs. 36%; RR, 0.18).[16] The control of nausea seems to be more difficult. Most

patients developed RT-induced nausea despite treatment (70% vs. 83% with placebo; RR, 0.84).[17] In summary, these trials show that patients receiving upper-abdomen irradiation are more likely to control RINV with 5-hydroxytryptamine-3 (5-HT₃) receptor antagonists than metoclopramide, phenothiazines, or placebo.[10-15]

The adverse effects of 5-HT₃ receptor antagonists are generally mild, consisting mainly of headache, constipation, and asthenia.[18] Randomized trials in RINV have examined the use of different 5-HT₃ receptor antagonists, but there are no data comparing them and no consensus on optimal dosing for RINV.[19] A systematic review of 25 randomized and nonrandomized trials revealed that 5-HT₃ receptor antagonists were most commonly administered for the entire duration of a course of RT. Optimal duration and timing of 5-HT₃ use before, during, and after RT administration needs to be determined.[20] With regard to palonosetron, the appropriate dosing

and frequency in the RINV setting are still unclear, with once-weekly dosing possible when the drug is combined with other agents.[21]

Corticosteroids

Corticosteroids are an attractive therapeutic antiemetic option because of their widespread availability and low cost. For short-term use, the side effects are few and do not outweigh the benefit of these agents. One randomized trial showed that dexamethasone was significantly more effective than placebo in patients receiving RT to the upper abdomen.[22] Combining corticosteroids with a 5-HT₃ receptor antagonist was assessed in a well-designed randomized trial, in which a 5-day course of dexamethasone plus ondansetron was compared with ondansetron plus placebo in 211 patients who received RT to the upper abdomen.[23] During the first 5 days, there was a statistically nonsignificant trend toward complete control of nausea (50% vs. 38% with placebo) and vomiting (78% vs. 71%), which

was the primary objective of the trial. The effects of dexamethasone extended beyond the initial 5-day period, and significantly more patients had complete control of emesis over the entire course of RT (23% vs. 12% with placebo), a secondary objective of the trial. The addition of dexamethasone has a modest effect on RINV and is potentially a useful addition to a 5-HT₃ receptor antagonist in this setting.[23]

Neurokinin-1 (NK-1) receptor antagonists

NK-1 receptor antagonists have an established role in the management of CINV; however, no studies have evaluated their impact solely on the risk of RINV. Although preclinical data indicate that RINV is mediated in part by substance P,[8] recommendation of these agents is premature, and NK-1 receptor antagonists are not included in the antiemetic guidelines for RINV.[3] A phase III, randomized, placebo-controlled trial compared an NK-1 receptor antagonist, fosaprepitant, combined with palonosetron and dexamethasone, with

palonosetron and dexamethasone alone in the prevention of N&V in patients who received concomitant RT and cisplatin.[21][[Level of evidence: I](#)] Patients received fractionated radiation therapy with weekly cisplatin, 40 mg/m². All patients received dexamethasone on the same schedule: 16 mg on day 1, 8 mg twice a day on day 2, 4 mg twice a day on day 3, and 4 mg once on day 4. More patients who received the three-drug regimen reached a complete response (65.7% for the fosaprepitant group vs. 48.7% for the placebo group).

Fosaprepitant has also been compared with olanzapine for the prevention of N&V in patients with head and neck or esophageal cancer who received RT concurrently with highly emetogenic chemotherapy.[24] For those who received olanzapine, palonosetron, and dexamethasone (OPD), dosing was as follows: dexamethasone 20 mg and palonosetron 0.25 mg intravenously (IV) on day 1 of chemotherapy, and olanzapine 10 mg

on days 1 to 4 of chemotherapy. For those who received fosaprepitant, palonosetron, and dexamethasone (FPD), dosing was as follows: dexamethasone 12 mg, palonosetron 0.25 mg IV, and fosaprepitant 150 mg IV on day 1 of chemotherapy, followed by dexamethasone 4 mg bid on days 2 to 3 of chemotherapy. Complete response was similar between the two groups, with a rate of 76% overall in the OPD arm and 74% overall in the FPD arm. This suggests that NK-1 receptor antagonists may play a role in patients receiving highly emetogenic chemotherapy.[24]

Other agents

Older, less-specific antiemetic drugs, such as prochlorperazine, metoclopramide, and cannabinoids, have shown limited efficacy in the prevention or treatment of RINV, although they may have a role in treating patients with milder symptoms and as rescue agents.[25]

Duration of Prophylaxis

The appropriate duration of antiemetic prophylaxis for patients receiving fractionated RT is not clear. There have been no randomized trials using 5-HT₃ receptor antagonists that compared a 5-day course of treatment with a more protracted course.[\[7\]](#) A systematic review that included 25 randomized and nonrandomized trials revealed that 5-HT₃ receptor antagonists were most commonly administered for the entire duration of a course of RT.[\[20\]](#)

Rescue Therapy

Studies suggest the benefit of 5-HT₃ receptor antagonists once nausea or vomiting occurs, but there are no trials specifically in this setting.[\[26\]](#) The emerging role of olanzapine in breakthrough emesis in patients with CINV has not been studied in RINV.[\[27\]](#)

Guidelines and Patient Management

For patients at high risk of developing RINV, prophylaxis with a 5-HT₃ receptor antagonist is

recommended in the clinical practice guidelines from the Multinational Association of Supportive Care in Cancer (MASCC) and American Society of Clinical Oncology (ASCO). Based on results from patients receiving highly emetogenic chemotherapy, the addition of dexamethasone to the 5-HT₃ receptor antagonist is suggested. The antiemetic clinical practice guidelines from both MASCC and ASCO also recommend that patients receiving moderately emetogenic RT be given with a 5-HT₃ receptor antagonist, with or without a short course of dexamethasone.[7] There are no fully published comparative clinical trials on the use of NK-1 receptor antagonists in preventing RINV; therefore, its use cannot be recommended.

Antiemetic dosing suggestions for the prevention of RINV are summarized in Table 6.

Table 6. Antiemetic Dosing for Radiation Therapy^a

Drug Category	Antiemetic	Dose
Serotonin (5-HT ₃) receptor antagonists	Granisetron	2 mg PO daily
	Ondansetron	8 mg PO or 0.15 mg/kg IV daily

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; IV prn = as needed; RT = radiation therapy; TBI = total-body irradiation.

^aAdapted from Roila et al.[3] and Hesketh et al.[28]

Drug Category	Antiemetic	Dose
	Palonosetron	0.25 mg IV or 0.5 mg PO
	Dolasetron	100 mg PO only
Corticosteroids	Dexamethasone	4 mg PO or IV

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; IV prn = as needed; RT = radiation therapy; TBI = total-body irradiation.

^aAdapted from Roila et al.[3] and Hesketh et al.[28]

Drug Category	Antiemetic	Dose
Dopamine receptor antagonists	Metoclopramide	20 mg PO
	Prochlorperazine	10 mg PO or IV

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; IV prn = as needed; RT = radiation therapy; TBI = total-body irradiation.

^aAdapted from Roila et al.[3] and Hesketh et al.[28]

References

1. Dennis K, Maranzano E, De Angelis C, et al.: Radiotherapy-induced nausea and vomiting.

Expert Rev Pharmacoecon Outcomes Res 11 (6): 685-92, 2011. [\[PUBMED Abstract\]](#)

2. Maranzano E: Radiation-induced emesis: a problem with many open questions. Tumori 87 (4): 213-8, 2001 Jul-Aug. [\[PUBMED Abstract\]](#)
3. Roila F, Molassiotis A, Herrstedt J, et al.: 2016 MASCC and ESMO guideline update for the prevention of chemotherapy- and radiotherapy-induced nausea and vomiting and of nausea and vomiting in advanced cancer patients. Ann Oncol 27 (suppl 5): v119-v133, 2016. [\[PUBMED Abstract\]](#)
4. Maranzano E, De Angelis V, Pergolizzi S, et al.: A prospective observational trial on emesis in radiotherapy: analysis of 1020 patients recruited in 45 Italian radiation oncology centres. Radiother Oncol 94 (1): 36-41, 2010. [\[PUBMED Abstract\]](#)
5. Enblom A, Bergius Axelsson B, Steineck G, et al.: One third of patients with radiotherapy-

- induced nausea consider their antiemetic treatment insufficient. Support Care Cancer 17 (1): 23-32, 2009. [\[PUBMED Abstract\]](#)
6. Horiot JC: Prophylaxis versus treatment: is there a better way to manage radiotherapy-induced nausea and vomiting? Int J Radiat Oncol Biol Phys 60 (4): 1018-25, 2004. [\[PUBMED Abstract\]](#)
 7. Feyer P, Jahn F, Jordan K: Radiation induced nausea and vomiting. Eur J Pharmacol 722: 165-71, 2014. [\[PUBMED Abstract\]](#)
 8. Yamamoto K, Nohara K, Furuya T, et al.: Ondansetron, dexamethasone and an NK1 antagonist block radiation sickness in mice. Pharmacol Biochem Behav 82 (1): 24-9, 2005. [\[PUBMED Abstract\]](#)
 9. Chow E, Meyer RM, Ding K, et al.: Dexamethasone in the prophylaxis of radiation-induced pain flare after palliative radiotherapy for bone metastases: a double-blind, randomised placebo-controlled, phase

- 3 trial. Lancet Oncol 16 (15): 1463-72, 2015. [\[PUBMED Abstract\]](#)
10. Aass N, Håtun DE, Thoresen M, et al.: Prophylactic use of tropisetron or metoclopramide during adjuvant abdominal radiotherapy of seminoma stage I: a randomised, open trial in 23 patients. Radiother Oncol 45 (2): 125-8, 1997. [\[PUBMED Abstract\]](#)
11. Bey P, Wilkinson PM, Resbeut M, et al.: A double-blind, placebo-controlled trial of i.v. dolasetron mesilate in the prevention of radiotherapy-induced nausea and vomiting in cancer patients. Support Care Cancer 4 (5): 378-83, 1996. [\[PUBMED Abstract\]](#)
12. Volk A, Kersting S, Konopke R, et al.: Surgical therapy of intrapancreatic metastasis from renal cell carcinoma. Pancreatology 9 (4): 392-7, 2009. [\[PUBMED Abstract\]](#)
13. Franzén L, Nyman J, Hagberg H, et al.: A randomised placebo controlled study with

- ondansetron in patients undergoing fractionated radiotherapy. Ann Oncol 7 (6): 587-92, 1996. [[PUBMED Abstract](#)]
14. Lanciano R, Sherman DM, Michalski J, et al.: The efficacy and safety of once-daily Kytril (granisetron hydrochloride) tablets in the prophylaxis of nausea and emesis following fractionated upper abdominal radiotherapy. Cancer Invest 19 (8): 763-72, 2001. [[PUBMED Abstract](#)]
 15. Priestman TJ, Dunn J, Brada M, et al.: Final results of the Royal College of Radiologists' trial comparing two different radiotherapy schedules in the treatment of cerebral metastases. Clin Oncol (R Coll Radiol) 8 (5): 308-15, 1996. [[PUBMED Abstract](#)]
 16. Priestman TJ, Roberts JT, Upadhyaya BK: A prospective randomized double-blind trial comparing ondansetron versus prochlorperazine for the prevention of nausea and vomiting in patients undergoing

fractionated radiotherapy. Clin Oncol (R Coll Radiol) 5 (6): 358-63, 1993. [[PUBMED Abstract](#)]

17. Chow E, Zeng L, Salvo N, et al.: Update on the systematic review of palliative radiotherapy trials for bone metastases. Clin Oncol (R Coll Radiol) 24 (2): 112-24, 2012. [[PUBMED Abstract](#)]

18. Goodin S, Cunningham R: 5-HT(3)-receptor antagonists for the treatment of nausea and vomiting: a reappraisal of their side-effect profile. Oncologist 7 (5): 424-36, 2002. [[PUBMED Abstract](#)]

19. Salvo N, Doble B, Khan L, et al.: Prophylaxis of radiation-induced nausea and vomiting using 5-hydroxytryptamine-3 serotonin receptor antagonists: a systematic review of randomized trials. Int J Radiat Oncol Biol Phys 82 (1): 408-17, 2012. [[PUBMED Abstract](#)]

20. Dennis K, Nguyen J, Presutti R, et al.: Prophylaxis of radiotherapy-induced nausea

and vomiting in the palliative treatment of bone metastases. Support Care Cancer 20 (8): 1673-8, 2012. [\[PUBMED Abstract\]](#)

21. Ruhlmann CH, Christensen TB, Dohn LH, et al.: Efficacy and safety of fosaprepitant for the prevention of nausea and emesis during 5 weeks of chemoradiotherapy for cervical cancer (the GAND-emesis study): a multinational, randomised, placebo-controlled, double-blind, phase 3 trial. Lancet Oncol 17 (4): 509-18, 2016. [\[PUBMED Abstract\]](#)
22. Kirkbride P, Bezjak A, Pater J, et al.: Dexamethasone for the prophylaxis of radiation-induced emesis: a National Cancer Institute of Canada Clinical Trials Group phase III study. J Clin Oncol 18 (9): 1960-6, 2000. [\[PUBMED Abstract\]](#)
23. Wong RK, Paul N, Ding K, et al.: 5-hydroxytryptamine-3 receptor antagonist with or without short-course dexamethasone

- in the prophylaxis of radiation induced emesis: a placebo-controlled randomized trial of the National Cancer Institute of Canada Clinical Trials Group (SC19). *J Clin Oncol* 24 (21): 3458-64, 2006. [\[PUBMED Abstract\]](#)
24. Navari RM, Nagy CK, Le-Rademacher J, et al.: Olanzapine versus fosaprepitant for the prevention of concurrent chemotherapy radiotherapy-induced nausea and vomiting. *J Community Support Oncol* 14 (4): 141-7, 2016. [\[PUBMED Abstract\]](#)
25. Roila F, Herrstedt J, Gralla RJ, et al.: Prevention of chemotherapy- and radiotherapy-induced nausea and vomiting: guideline update and results of the Perugia consensus conference. *Support Care Cancer* 19 (Suppl 1): S63-5, 2011. [\[PUBMED Abstract\]](#)
26. Mystakidou K, Katsouda E, Linou A, et al.: Prophylactic tropisetron versus rescue tropisetron in fractionated radiotherapy to moderate or high emetogenic areas: a

- prospective randomized open label study in cancer patients. Med Oncol 23 (2): 251-62, 2006. [\[PUBMED Abstract\]](#)
27. Navari RM, Nagy CK, Gray SE: The use of olanzapine versus metoclopramide for the treatment of breakthrough chemotherapy-induced nausea and vomiting in patients receiving highly emetogenic chemotherapy. Support Care Cancer 21 (6): 1655-63, 2013. [\[PUBMED Abstract\]](#)
28. Hesketh PJ, Kris MG, Basch E, et al.: Antiemetics: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol 35 (28): 3240-3261, 2017. [\[PUBMED Abstract\]](#)

Pediatric Chemotherapy-Induced Acute Nausea and Vomiting

Pediatric Guidelines for Acute Nausea and Vomiting (N&V)

Chemotherapy-induced N&V (CINV) is an important problem in the pediatric population. As in adults, nausea in children is more of a problem than vomiting. Parents of children who received active antineoplastic therapy in Ontario, Canada, identified nausea as the fourth most prevalent and bothersome treatment-related symptom.[\[1\]](#)

Current approaches to prevent CINV are based on an accurate description of the potential of antineoplastic therapies to cause N&V. Current recommendations, based on published guidelines, [\[2\]](#) include patients aged 1 month to 18 years who are about to receive their first-ever course of antineoplastic therapy. These recommendations focus on the prevention of acute CINV (i.e., within 24 hours of administration of an antineoplastic agent).

Guidelines define optimal control of acute CINV as no vomiting, no retching, no nausea, no use of

antiemetic agents other than those given for CINV prevention, and no nausea-related change in the child's usual appetite and diet. This level of CINV control is to be achieved on each day that antineoplastic therapy is administered and for 24 hours after administration of the last agent in the antineoplastic therapy cycle.

Emetic Risk

In children receiving antineoplastic agents who were not given antiemetic prophylaxis or who were given ineffective prophylaxis, expected rates of complete CINV control were as follows:[[2](#)]

- High emetic risk, less than 10%.
- Moderate emetic risk, 10% to less than 30%.
- Low emetic risk, 30% to less than 90%.
- Minimal emetic risk, more than 90%.

The expected rate of complete CINV control in children receiving antiemetic prophylaxis (5-

hydroxytryptamine-3 [5-HT₃] receptor antagonist with or without dexamethasone) is more than 70% to 80%.[2] Each chemotherapy agent carries an inherent risk of emesis, which is the first issue to consider when planning chemotherapy treatment. For more information about preventing acute or delayed CINV, see [Table 5](#).

Acute Chemotherapy-Induced Nausea and Vomiting—Antiemetic Prophylaxis

Highly emetogenic chemotherapy

Guidelines [2,3] recommend that children aged 6 months and older who are receiving antineoplastic agents of high emetic risk that are not known or suspected to interact with aprepitant receive aprepitant, a 5-HT₃ receptor antagonist, and dexamethasone. Children older than 6 months who cannot receive dexamethasone should receive a 5-HT₃ receptor antagonist plus aprepitant. Children who cannot receive

aprepitant should receive a 5-HT₃ receptor antagonist plus dexamethasone.[4][[Level of evidence: IV](#)]

Moderately emetogenic chemotherapy

Children receiving antineoplastic agents of moderate emetogenicity should receive ondansetron, granisetron, or palonosetron plus dexamethasone. Children aged 6 months and older and whose antineoplastic agents do not interact with aprepitant and who cannot receive dexamethasone should receive a 5-HT₃ receptor antagonist plus aprepitant.[3,4][[Level of evidence: IV](#)]

Low emetogenic chemotherapy

Children receiving antineoplastic agents of low emetogenicity should receive a 5-HT₃ receptor antagonist.[3]

Minimal emetogenic potential

Children receiving antineoplastic agents of minimal emetogenicity should receive no routine prophylaxis.[3]

Other Antiemetic Modalities

Current consensus is that the following modalities may be effective in children receiving antineoplastic agents:[2]

- Acupuncture.
- Acupressure.
- Guided imagery.
- Music therapy.
- Progressive muscle relaxation.
- Psychoeducational support and information.

In addition, virtual reality may convey some benefit. Other recommendations (low level of evidence) include the following:

- Eating smaller, more-frequent meals.

- Reducing food aromas and other stimuli with strong odors.
- Avoiding foods that are spicy, fatty, or highly salty.
- Taking antiemetics before meals so that the effect is present during and after meals.
- Using measures and foods (e.g., “comfort foods”) that previously helped minimize nausea.

Despite a lack of strong evidence, most experts think that these recommendations are unlikely to result in undesirable effects or to adversely affect quality of life, and they may convey benefit.[\[2\]](#)

Antiemetics

Prophylaxis with a 5-HT₃ receptor antagonist alone leads to poor CINV control in pediatric patients receiving antineoplastic agents of moderate and high emetic risk. A synthesis of three studies that evaluated alternative antiemetic agents

(chlorpromazine and metoclopramide) in children receiving highly emetogenic chemotherapy observed a complete CINV control rate of 9% (95% confidence interval: 0, 20).[2] When corticosteroids are contraindicated, it is recommended that nabilone or chlorpromazine be administered together with ondansetron or granisetron to children receiving highly emetogenic chemotherapy. Metoclopramide is a third option for children receiving moderately emetogenic chemotherapy. Corticosteroids combined with a serotonin antagonist are recommended for patients receiving highly and moderately emetogenic chemotherapy.[5]

Antiemetic dosing suggestions for pediatric patients are summarized in Table 7.

Table 7. Pediatric Antiemetic Dosing

Drug Category	Medication	
Phenothiazines	Chlorpromazine	0.5 m q6h; r to 1 n q6h; r dose:
	Prochlorperazine	9–13 l PO qc maxir 7.5 m
		13–18 PO bi

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B = intramuscular; IV = intravenous; NK-1 = neurokinin-1; l = sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		maxir 10 mg
		18–39 tid or maxir 15 mg
	Promethazine	Age > mg/kg 6h; m dose:

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B₁ = intramuscular; IV = intravenous; NK-1 = neurokinin-1; l = sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
Substituted benzamides	Metoclopramide	Mode emet chem mg/kg once prech then (mg/kg q6h
Serotonin (5-HT ₃) receptor antagonists	Granisetron	40 µg 40 µg maxir mg/d
	Ondansetron	Age 0 mg/kg

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B = intramuscular; IV = intravenous; NK-1 = neurokinin-1; l = sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		mg/m prech then c highly or q1: mode emet chem
		Low e chem 0.3 m (10 m once prech
		Maxir dose:

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B: intramuscular; IV = intravenous; NK-1 = neurokinin-1; l sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		maxim 16 mg
	Palonosetron	Age 1 µg/kg dose:
Substance P antagonists (NK- 1 receptor antagonists)	Aprepitant	Capsu y: 125 prech day 1, qd x2

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B: intramuscular; IV = intravenous; NK-1 = neurokinin-1; l sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		Suspe mo-1 kg): 3 prech day 1, mg/kg
	Fosaprepitant	Age 1 mg
Corticosteroids	Dexamethasone	Highly chem mg/m

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B: intramuscular; IV = intravenous; NK-1 = neurokinin-1; l sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		Mode emet chem BSA ≤ q12h
		BSA > q12h

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B₁ = intramuscular; IV = intravenous; NK-1 = neurokinin-1; l = sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		Maxim mg/d
	Methylprednisolone	4–10

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B: intramuscular; IV = intravenous; NK-1 = neurokinin-1; l sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
Benzodiazepines	Lorazepam	Antici 0.05 r (maxi mg/d bedti befor chem and o prech
		Break 0.02–1 mg/kg (maxi q6h p

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B: intramuscular; IV = intravenous; NK-1 = neurokinin-1; l sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
Atypical antipsychotics	Olanzapine	0.1–0.4 mg/kg maxir
Other pharmacological agents	Dronabinol	Age 6 mg/m ² prech
	Nabilone	Age >

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B₁ = intramuscular; IV = intravenous; NK-1 = neurokinin-1; l = sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Drug Category	Medication	
		<18 kg q12h
		18–30 kg q12h
		>30 kg 12h
		Maxir 0.06 r

5-HT₃ = 5-hydroxytryptamine-3; bid = twice a day; B = buccal; IM = intramuscular; IV = intravenous; NK-1 = neurokinin-1; l = sublingual; tid = 3 times a day.

^aPalonosetron prescribing information lists the pedia

Multiagent, single-day chemotherapy regimens

Experience in pediatrics and guidelines recommend basing the emetogenicity of combination antineoplastic regimens on that of the agent of highest emetic risk.[\[20\]](#) The emetogenicity of the antineoplastic combinations in the following list appears to be higher than would be appreciated by assessment of the emetic risk of the individual agents.[\[21\]](#)

High Level of Emetic Risk (>90% Frequency of Emesis in Absence of Prophylaxis)

- Cyclophosphamide + anthracycline.
- Cyclophosphamide + etoposide.
- Cytarabine ($150\text{--}200\text{ mg/m}^2$) + daunorubicin.
- Cytarabine (300 mg/m^2) + etoposide.
- Cytarabine (300 mg/m^2) + teniposide.
- Doxorubicin + ifosfamide.

- Doxorubicin + methotrexate (5 g/m^2).
- Etoposide + ifosfamide.

References

1. Dupuis LL, Milne-Wren C, Cassidy M, et al.: Symptom assessment in children receiving cancer therapy: the parents' perspective. Support Care Cancer 18 (3): 281-99, 2010. [\[PUBMED Abstract\]](#)
2. Dupuis LL, Boodhan S, Holdsworth M, et al.: Guideline for the prevention of acute nausea and vomiting due to antineoplastic medication in pediatric cancer patients. Pediatr Blood Cancer 60 (7): 1073-82, 2013. [\[PUBMED Abstract\]](#)
3. Dupuis LL, Sung L, Molassiotis A, et al.: 2016 updated MASCC/ESMO consensus recommendations: Prevention of acute chemotherapy-induced nausea and vomiting in children. Support Care Cancer 25 (1): 323-331, 2017. [\[PUBMED Abstract\]](#)

4. Patel P, Robinson PD, Cohen M, et al.: Prevention of acute and delayed chemotherapy-induced nausea and vomiting in pediatric cancer patients: A clinical practice guideline. *Pediatr Blood Cancer* 69 (12): e30001, 2022. [\[PUBMED Abstract\]](#)
5. White L, Daly SA, McKenna CJ, et al.: A comparison of oral ondansetron syrup or intravenous ondansetron loading dose regimens given in combination with dexamethasone for the prevention of nausea and emesis in pediatric and adolescent patients receiving moderately/highly emetogenic chemotherapy. *Pediatr Hematol Oncol* 17 (6): 445-55, 2000. [\[PUBMED Abstract\]](#)
6. Lexicomp Online. Hudson, Ohio: Lexi-Comp, Inc., 2025. [Available online with subscription.](#) Last accessed Jan. 15, 2025.
7. Relling MV, Mulhern RK, Fairclough D, et al.: Chlorpromazine with and without lorazepam

as antiemetic therapy in children receiving uniform chemotherapy. J Pediatr 123 (5): 811-6, 1993. [\[PUBMED Abstract\]](#)

8. Chan HS, Correia JA, MacLeod SM: Nabilone versus prochlorperazine for control of cancer chemotherapy-induced emesis in children: a double-blind, crossover trial. Pediatrics 79 (6): 946-52, 1987. [\[PUBMED Abstract\]](#)
9. Köseoglu V, Kürekçi AE, Sarici U, et al.: Comparison of the efficacy and side-effects of ondansetron and metoclopramide-diphenhydramine administered to control nausea and vomiting in children treated with antineoplastic chemotherapy: a prospective randomized study. Eur J Pediatr 157 (10): 806-10, 1998. [\[PUBMED Abstract\]](#)
10. Berrak SG, Ozdemir N, Bakirci N, et al.: A double-blind, crossover, randomized dose-comparison trial of granisetron for the prevention of acute and delayed nausea and emesis in children receiving moderately

emetogenic carboplatin-based chemotherapy. Support Care Cancer 15 (10): 1163-8, 2007. [\[PUBMED Abstract\]](#)

11. Kadota R, Shen V, Messinger Y: Safety, pharmacokinetics, and efficacy of palonosetron in pediatric patients: a multicenter, stratified, double-blind, phase 3, randomized study. [Abstract] J Clin Oncol 25 (18 suppl): A-9570, 2007.
12. Kang HJ, Loftus S, Taylor A, et al.: Aprepitant for the prevention of chemotherapy-induced nausea and vomiting in children: a randomised, double-blind, phase 3 trial. Lancet Oncol 16 (4): 385-94, 2015. [\[PUBMED Abstract\]](#)
13. Siddiqui MA, Ghaznawi HI: Some observations on intestinal parasites in Hajjis visiting Saudi Arabia, during 1983 G (1403 H.) Pilgrimage. J Egypt Soc Parasitol 15 (2): 705-12, 1985. [\[PUBMED Abstract\]](#)

14. Small BE, Holdsworth MT, Raisch DW, et al.: Survey ranking of emetogenic control in children receiving chemotherapy. J Pediatr Hematol Oncol 22 (2): 125-32, 2000 Mar-Apr. [\[PUBMED Abstract\]](#)
15. Hirota T, Honjo T, Kuroda R, et al.: [Antiemetic efficacy of granisetron in pediatric cancer treatment--(2). Comparison of granisetron and granisetron plus methylprednisolone as antiemetic prophylaxis]. Gan To Kagaku Ryoho 20 (15): 2369-73, 1993. [\[PUBMED Abstract\]](#)
16. Dupuis LL, Nathan PC: Options for the prevention and management of acute chemotherapy-induced nausea and vomiting in children. Paediatr Drugs 5 (9): 597-613, 2003. [\[PUBMED Abstract\]](#)
17. Flank J, Thackray J, Nielson D, et al.: Olanzapine for treatment and prevention of acute chemotherapy-induced vomiting in children: a retrospective, multi-center review.

- Pediatr Blood Cancer 62 (3): 496-501,
2015. [\[PUBMED Abstract\]](#)
18. Elder JJ, Knoderer HM: Characterization of Dronabinol Usage in a Pediatric Oncology Population. J Pediatr Pharmacol Ther 20 (6): 462-7, 2015 Nov-Dec. [\[PUBMED Abstract\]](#)
 19. Dalzell AM, Bartlett H, Lilleyman JS: Nabilone: an alternative antiemetic for cancer chemotherapy. Arch Dis Child 61 (5): 502-5, 1986. [\[PUBMED Abstract\]](#)
 20. Dupuis LL, Boodhan S, Sung L, et al.: Guideline for the classification of the acute emetogenic potential of antineoplastic medication in pediatric cancer patients. Pediatr Blood Cancer 57 (2): 191-8, 2011. [\[PUBMED Abstract\]](#)
 21. Holdsworth MT, Raisch DW, Frost J: Acute and delayed nausea and emesis control in pediatric oncology patients. Cancer 106 (4): 931-40, 2006. [\[PUBMED Abstract\]](#)

Pediatric Delayed Nausea and Vomiting

The nature and prevalence of delayed nausea and vomiting (N&V) in children after administration of antineoplastic agents have not been well described.[1] Additionally, most pediatric chemotherapy regimens are given over multiple days, making the onset and duration of risk of delayed versus acute N&V unclear.

Research on chemotherapy-induced N&V (CINV) in children has been limited in part by the lack of assessment tools and the subjective nature of nausea. In the pediatric population, vomiting is more easily recognizable and measurable than nausea.[1] Difficulties in assessing nausea in young children may contribute to the common perception that young children experience CINV less frequently than older children. In addition, caregivers may have a higher tolerance for vomiting in young children and may miss detecting

nausea.[1] In view of these limitations, studies often use dietary intake to assess the extent of nausea.

Several investigators have attempted to determine the prevalence of delayed N&V in the pediatric population. One early study suggested a low incidence.[2] A large study assessed the nature and prevalence of delayed CINV in children.[1] Nausea was self-assessed daily using a numeric scale reflecting the effect of nausea on activities and a faces scale for children aged 3 to 6 years. Diet was also assessed daily. Results showed a 33% rate of delayed vomiting in patients who received cyclophosphamide, cisplatin, or carboplatin and an 11% rate in those who received other antineoplastic agents. No antiemetics were given on 412 (79%) of 522 study days. Nevertheless, on 381 (93%) of those 412 study days, patients were completely free from vomiting. Antiemetics were most often given as single agents (ondansetron, on 54 study days; dimenhydrinate,

on 17 study days; dexamethasone, on 6 study days). Diet was not affected. The authors concluded that antineoplastic-induced delayed N&V may be less prevalent in children than in adults.[1] The high percentage of children who did not experience delayed vomiting may reflect a lack of significant emetogenic potential among many of the regimens in the study. In 100 of 174 chemotherapy cycles, no antiemetics were administered. In addition, there was no characterization of antiemetic response in moderate and severe chemotherapy regimens.

Another study evaluated the incidence of delayed N&V in pediatric patients receiving moderately and highly emetogenic chemotherapy as well as premedications (ondansetron alone or with dexamethasone, depending on a treatment's emetogenic potential).[3] Investigators measured nausea severity and duration, vomiting severity, the number of vomiting episodes, interference with daily activities, and assessment of appetite.

The authors found that delayed N&V occurred with both moderately and highly emetogenic chemotherapy regimens, but that the severity of N&V varied. In addition, toddlers had better antiemetic control than older children, which may be the result of anxiety differences between the age groups. The reasons for toddler patients' greater complete control are unclear but are consistent with a previous study of N&V control rates in children.[4] Anxiety and patient perception may be important contributors to N&V in older children; a relationship between control of acute N&V and the occurrence of delayed N&V was found.

Another study suggested a higher incidence of delayed N&V than was previously found in a pediatric population.[5] In 40 pediatric cancer patients receiving chemotherapy, N&V was measured from the child's perspective using the Adapted Rhodes Index of Nausea and Vomiting for Pediatrics; from the primary caregiver's

perspective using the Adapted Rhodes Index of Nausea and Vomiting for Parents; and from the nurses' perspective using the National Cancer Institute Nausea and Vomiting Grading Criteria. The highest frequency of nausea occurred in the delayed period, with 60% of patients (n = 24) reporting delayed nausea. The authors concluded that CINV occurred throughout the chemotherapy course, with delayed N&V occurring most frequently and with greater severity and distress. Delayed N&V in the pediatric population requires further study.

Because well-designed studies on the prevention of delayed N&V in children are not available, the best available evidence comes from adult data and a pediatric clinical practice guideline.[6][[Level of evidence: IV](#)]; [7]

Delayed Chemotherapy-Induced Nausea and Vomiting—Antiemetic Prophylaxis

Highly emetogenic chemotherapy

Palonosetron is the preferred 5-hydroxytryptamine-3 (5-HT₃) receptor antagonist in the acute phase in patients at high risk of delayed CINV. Guidelines recommend that children who can receive aprepitant and dexamethasone continue to do so during the delayed phase. If dexamethasone cannot be used, aprepitant should be continued. If aprepitant cannot be used, dexamethasone should be continued. If olanzapine was started during the acute phase, it should be continued during the delayed phase.[6][[Level of evidence: IV](#)]

In a phase III, double-blind, randomized controlled trial, 128 patients aged 3 to 18 years receiving highly emetogenic chemotherapy were randomly assigned to receive intravenous ondansetron and dexamethasone plus olanzapine or placebo on days 1 and 2. More patients in the olanzapine group had complete control of vomiting in the delayed phase (73% vs. 48%; $P = .005$), although

there was no difference in control in the acute phase or overall. More patients in the placebo group required rescue medications for vomiting than in the olanzapine group (29% vs. 14%; $P = .025$). Grade 1/2 sedation was greater in the olanzapine group than in the placebo group (46% vs. 14%).[8]

Moderately emetogenic chemotherapy

In the delayed phase, children receiving antineoplastic agents of moderate emetogenicity who received a 5-HT₃ receptor inhibitor and dexamethasone in the acute phase should consider dexamethasone during the delayed phase.

Children receiving a 1-day regimen of antineoplastic agents of moderate emetogenicity who received a 5-HT₃ receptor inhibitor and fosaprepitant or aprepitant in the acute phase should continue oral aprepitant in the delayed phase. Children receiving a multiday regimen of

antineoplastic agents of moderate emetogenicity who received a 5-HT₃ receptor inhibitor and fosaprepitant or aprepitant in the acute phase should consider not using oral aprepitant in the delayed phase.

Children receiving a 5-HT₃ receptor inhibitor with olanzapine during the acute phase should consider continuing olanzapine during the delayed phase.[6][[Level of evidence: IV](#)]

Low emetogenic chemotherapy

Children receiving antineoplastic agents of low emetogenicity should not receive routine prophylaxis during the delayed phase.[6][[Level of evidence: IV](#)]

Minimal emetogenic potential

Children receiving antineoplastic agents of minimal emetogenicity should not receive routine prophylaxis during the delayed phase.[6][[Level of evidence: IV](#)]

References

1. Dupuis LL, Lau R, Greenberg ML: Delayed nausea and vomiting in children receiving antineoplastics. *Med Pediatr Oncol* 37 (2): 115-21, 2001. [\[PUBMED Abstract\]](#)
2. Foot AB, Hayes C: Audit of guidelines for effective control of chemotherapy and radiotherapy induced emesis. *Arch Dis Child* 71 (5): 475-80, 1994. [\[PUBMED Abstract\]](#)
3. Holdsworth MT, Raisch DW, Frost J: Acute and delayed nausea and emesis control in pediatric oncology patients. *Cancer* 106 (4): 931-40, 2006. [\[PUBMED Abstract\]](#)
4. Small BE, Holdsworth MT, Raisch DW, et al.: Survey ranking of emetogenic control in children receiving chemotherapy. *J Pediatr Hematol Oncol* 22 (2): 125-32, 2000 Mar-Apr. [\[PUBMED Abstract\]](#)
5. Rodgers C, Kollar D, Taylor O, et al.: Nausea and vomiting perspectives among children receiving moderate to highly emetogenic

chemotherapy treatment. Cancer Nurs 35 (3): 203-10, 2012 May-Jun. [\[PUBMED Abstract\]](#)

6. Patel P, Robinson PD, Cohen M, et al.: Prevention of acute and delayed chemotherapy-induced nausea and vomiting in pediatric cancer patients: A clinical practice guideline. Pediatr Blood Cancer 69 (12): e30001, 2022. [\[PUBMED Abstract\]](#)
7. Dupuis LL, Sung L, Molassiotis A, et al.: 2016 updated MASCC/ESMO consensus recommendations: Prevention of acute chemotherapy-induced nausea and vomiting in children. Support Care Cancer 25 (1): 323-331, 2017. [\[PUBMED Abstract\]](#)
8. Moothedath AW, Meena JP, Gupta AK, et al.: Efficacy and Safety of Olanzapine in Children Receiving Highly Emetogenic Chemotherapy: A Randomized, Double-blind Placebo-controlled Phase 3 Trial. J Pediatr Hematol Oncol 44 (8): 446-453, 2022. [\[PUBMED Abstract\]](#)

Pediatric Anticipatory Nausea and Vomiting

Patients with cancer who have received chemotherapy may experience nausea and vomiting (N&V) when anticipating further chemotherapy. Study differences in methodology, timing, and assessment instruments; small samples; and a focus on nausea or vomiting but not both has led to difficulties in capturing the prevalence of anticipatory N&V (ANV) in children. Accurate prevalence is also stymied by using parent or caregiver proxy reports of nausea and nonvalidated nausea assessment tools.

In patients receiving 5-hydroxytryptamine-3 (5-HT₃) receptor antagonists and corticosteroids as antiemetic agents, approximately one-third of adults experienced ANV, while 6% to 11% reported anticipatory vomiting.^[1] A study of children in the pre-5-HT₃ receptor antagonist era reported anticipatory nausea in 23 (29%) of 80 children and

anticipatory vomiting in 16 (20%) of 80 children who had received 11 cycles of antineoplastic therapy, on average, before evaluation.[2] In the post-5-HT₃ receptor antagonist era, the reported prevalence of anticipatory nausea in children has ranged from 0% to 59%.[3] Similar to observations in adult patients, the reported prevalence of anticipatory nausea is always higher than that of anticipatory vomiting in children, although one study reported an equivalent prevalence (5 [26%] of 19 patients) for these conditions.[4]

This section focuses on the management of ANV in children aged 1 month to 18 years who are receiving antineoplastic medication. Optimal control of ANV is defined as no vomiting, no retching, no nausea, no use of antiemetic agents other than those given for the prevention or treatment of chemotherapy-induced N&V (CINV), and no nausea-related change in the child's usual appetite and diet. This level of ANV control is to be achieved during the 24 hours before

administration of the first antineoplastic agent of the upcoming planned antineoplastic cycle.

Approaches to Prevent ANV in Children

ANV appears to be a conditioned response to CINV experienced in the acute phase (24 hours after administration of chemotherapy) and delayed phase (more than 24 hours after and within 7 days of administration of chemotherapy).[3] The anxiety and distress attendant to CINV reinforce the conditioned response.[3] It follows, then, that a higher rate of complete control of acute and delayed CINV would result in lower rates of ANV. Adherence to evidence-based recommendations for CINV prevention has been shown to substantially improve complete control of acute CINV.[5]

Optimized control of acute and delayed CINV may help minimize exposure to the negative stimuli required for conditioning to occur. Consensus

recommendations call for antiemetic interventions to be based on published guidelines for the prevention of acute CINV in children receiving antineoplastic agents,[6,7] including antineoplastic agent-naïve patients. Once antineoplastic therapy has been initiated, the selection of antiemetic interventions should be informed by evidence-based guidelines and tailored to the patient's CINV control and any adverse effects associated with antiemetic agents.

Interventions to Control ANV in Children

Hypnosis

Hypnosis has been defined as an intervention that “provides suggestions for changes in sensation, perception, cognition, affect, mood, or behavior.”[8] Two trials evaluated the role of hypnosis in controlling ANV in children. One study recruited 54 children aged 5 to 17 years who had reported experiencing anticipatory nausea,

anticipatory vomiting, or both in a previous study and who were about to receive at least two identical antineoplastic treatment courses.[9] On average, children were 15.8 months (range, 0.5–118 months) from their cancer diagnosis at the time of the study. The control group had received antineoplastic therapy for much longer than the other two groups (29.5 months vs. 8 or 11.5 months).

Although it is not possible to precisely ascertain the emetogenicity of the antineoplastic therapy these children received, it appears that most received highly emetogenic treatment. The antiemetic agents taken for prophylaxis were not reported, but children's antiemetic regimens were unchanged during the trial. The severity of N&V was assessed through semistructured interviews. Children were randomly assigned to receive one of three possible interventions: hypnosis training (imagination-focused therapy), active cognitive distraction (relaxation), or contact with a therapist

(control). The authors reported a significant improvement in complete control of anticipatory vomiting in the group who received hypnosis training (12 [57%] of 21 patients at baseline vs. 18 [86%] of 21 patients after hypnosis training; $P < .05$). Complete control of anticipatory nausea increased from 5 (24%) of 21 patients at baseline to 8 (38%) of 21 patients after hypnosis training.[9]

Another study evaluated hypnosis as a means of preventing ANV in 20 children aged 6 to 18 years who were naïve to chemotherapy.[10] Controls were matched for age (± 3 years) and the emetogenicity of their antineoplastic treatment. Insufficient information is available to determine the emetogenicity of the antineoplastic regimens. Children randomly assigned to receive hypnosis did not receive antiemetic prophylaxis but did receive antiemetic agents as needed. Children in the control group received standard antiemetic prophylaxis for 4 to 6 hours after antineoplastic therapy. Ondansetron was given to more children

in the control group (7 of 10 patients) than in the hypnosis group (3 of 10 patients).

Children randomly assigned to receive hypnosis were taught self-hypnosis during the initial antineoplastic treatment, while children in the control group spent equivalent time in conversation with a therapist. Researchers used a daily structured interview with the children to assess ANV at 1 to 2 months, and at 4 to 6 months after diagnosis. At the time of first assessment, children who had been taught self-hypnosis reported significantly less anticipatory nausea than did the control group, although the incidence was not reported. The rate of anticipatory vomiting was identical in each group (1 of 10 patients). By the time of the second assessment, there was no difference between the groups in the rate of anticipatory nausea. The rate of anticipatory vomiting between the groups was also similar (hypnosis, 0 of 10 patients vs. control, 2 of 10 patients).[\[10\]](#)

Pharmacological interventions

Studies of pharmacological interventions for ANV have been conducted only in adults and are limited to benzodiazepines. Because patients who experience ANV have been observed to be more anxious than patients who do not experience ANV, [11] anxiolytics have been studied. Studies in adults have evaluated benzodiazepines as a treatment for ANV.[12,13] In one randomized trial, adult patients with cancer received a placebo or lorazepam 2 mg by mouth the night before antineoplastic treatment, the morning of treatment, and at bedtime for the next 5 days during 180 antineoplastic treatment courses containing cisplatin.[12] Patients also received metoclopramide 2 mg/kg per dose, clemastine, and dexamethasone for antiemetic prophylaxis. At the time of randomization, approximately two-thirds of patients were naïve to antineoplastic agents. ANV was defined as nausea, vomiting, or both that occurred within 12 hours before antineoplastic therapy or 1 hour after the start of

antineoplastic therapy. A significantly higher proportion of treatments with lorazepam were associated with complete ANV control, compared with the control group (52% vs. 32%; $P < .05$). Few adverse effects occurred; 76% of the patients who received lorazepam and 32% of the controls had mild sedation.

Women with breast cancer who were naïve to antineoplastic treatment were enrolled in a double-blind placebo-controlled trial comparing the incidence of ANV after relaxation training and either alprazolam ($n = 29$) or placebo ($n = 28$). Alprazolam 0.25 mg or placebo was given twice daily by mouth for 6 to 12 months. Triazolam was also given as needed to patients in both study arms to manage insomnia. The proportion of patients who experienced complete control of anticipatory nausea and anticipatory vomiting before the fourth antineoplastic treatment was similar in both study arms (26% vs. 25% and 4% vs. 0%, respectively). Diazepam 5 mg twice daily was

given to 29 adult cancer patients with ANV for 3 days before each of four consecutive antineoplastic treatment courses.[13] Thirteen patients (45%) experienced complete ANV control at some time over the four antineoplastic treatment courses.

Conclusions

While the improvement in complete control of ANV provided by psychological interventions such as hypnosis or systematic desensitization may not be dramatic, these interventions may benefit individual patients with minimal risk. For this reason, one guideline development panel recommends that such interventions be offered to age-appropriate patients who experience ANV where the expertise and resources exist to deliver them.[6]

Despite the lack of evidence supporting the use of benzodiazepines to treat ANV in children, guidelines based on clinical experience

recommend using lorazepam for ANV in children. [14] The recommended initial dose was based on current pediatric dosing recommendations, with the usual adult dose as the maximum dose.[15] This dose should be titrated to the needs of each child, with dose lowering recommended for excessive sedation.

References

1. Morrow GR, Roscoe JA, Hynes HE, et al.: Progress in reducing anticipatory nausea and vomiting: a study of community practice. Support Care Cancer 6 (1): 46-50, 1998. [\[PUBMED Abstract\]](#)
2. Dolgin MJ, Katz ER, McGinty K, et al.: Anticipatory nausea and vomiting in pediatric cancer patients. Pediatrics 75 (3): 547-52, 1985. [\[PUBMED Abstract\]](#)
3. Tyc VL, Mulhern RK, Bieberich AA: Anticipatory nausea and vomiting in pediatric cancer patients: an analysis of conditioning

and coping variables. J Dev Behav Pediatr 18 (1): 27-33, 1997. [\[PUBMED Abstract\]](#)

4. Stockhorst U, Spennes-Saleh S, Körholz D, et al.: Anticipatory symptoms and anticipatory immune responses in pediatric cancer patients receiving chemotherapy: features of a classically conditioned response? Brain Behav Immun 14 (3): 198-218, 2000. [\[PUBMED Abstract\]](#)
5. Aapro M, Molassiotis A, Dicato M, et al.: The effect of guideline-consistent antiemetic therapy on chemotherapy-induced nausea and vomiting (CINV): the Pan European Emesis Registry (PEER). Ann Oncol 23 (8): 1986-92, 2012. [\[PUBMED Abstract\]](#)
6. Dupuis LL, Boodhan S, Holdsworth M, et al.: Guideline for the prevention of acute nausea and vomiting due to antineoplastic medication in pediatric cancer patients. Pediatr Blood Cancer 60 (7): 1073-82, 2013. [\[PUBMED Abstract\]](#)

7. Patel P, Robinson PD, Devine KA, et al.: Prevention and treatment of anticipatory chemotherapy-induced nausea and vomiting in pediatric cancer patients and hematopoietic stem cell recipients: Clinical practice guideline update. *Pediatr Blood Cancer* 68 (5): e28947, 2021. [\[PUBMED Abstract\]](#)
8. Montgomery GH, Schnur JB, Kravits K: Hypnosis for cancer care: over 200 years young. *CA Cancer J Clin* 63 (1): 31-44, 2013. [\[PUBMED Abstract\]](#)
9. Zeltzer LK, Dolgin MJ, LeBaron S, et al.: A randomized, controlled study of behavioral intervention for chemotherapy distress in children with cancer. *Pediatrics* 88 (1): 34-42, 1991. [\[PUBMED Abstract\]](#)
10. Jacknow DS, Tschann JM, Link MP, et al.: Hypnosis in the prevention of chemotherapy-related nausea and vomiting in children: a

prospective study. J Dev Behav Pediatr 15 (4): 258-64, 1994. [\[PUBMED Abstract\]](#)

11. Andrykowski MA: The role of anxiety in the development of anticipatory nausea in cancer chemotherapy: a review and synthesis. Psychosom Med 52 (4): 458-75, 1990 Jul-Aug. [\[PUBMED Abstract\]](#)
12. Malik IA, Khan WA, Qazilbash M, et al.: Clinical efficacy of lorazepam in prophylaxis of anticipatory, acute, and delayed nausea and vomiting induced by high doses of cisplatin. A prospective randomized trial. Am J Clin Oncol 18 (2): 170-5, 1995. [\[PUBMED Abstract\]](#)
13. Razavi D, Delvaux N, Farvacques C, et al.: Prevention of adjustment disorders and anticipatory nausea secondary to adjuvant chemotherapy: a double-blind, placebo-controlled study assessing the usefulness of alprazolam. J Clin Oncol 11 (7): 1384-90, 1993. [\[PUBMED Abstract\]](#)

14. van Hoff J, Olszewski D: Lorazepam for the control of chemotherapy-related nausea and vomiting in children. J Pediatr 113 (1 Pt 1): 146-9, 1988. [\[PUBMED Abstract\]](#)
15. National Comprehensive Cancer Network: NCCN Clinical Practice Guidelines in Oncology: Antiemesis. Version 2.2024. Plymouth Meeting, Pa: National Comprehensive Cancer Network, 2024. [Available online with free registration](#). Last accessed March 4, 2025.

Latest Updates to This Summary (03/10/2025)

The PDQ cancer information summaries are reviewed regularly and updated as new information becomes available. This section describes the latest changes made to this summary as of the date above.

Editorial changes were made to this summary.

This summary is written and maintained by the [PDQ Supportive and Palliative Care Editorial Board](#), which is editorially independent of NCI. The summary reflects an independent review of the literature and does not represent a policy statement of NCI or NIH. More information about summary policies and the role of the PDQ Editorial Boards in maintaining the PDQ summaries can be found on the [About This PDQ Summary](#) and [PDQ® Cancer Information for Health Professionals](#) pages.

About This PDQ Summary

Purpose of This Summary

This PDQ cancer information summary for health professionals provides comprehensive, peer-reviewed, evidence-based information about the prevention and control of treatment-related

nausea and vomiting in cancer patients. It is intended as a resource to inform and assist clinicians in the care of their patients. It does not provide formal guidelines or recommendations for making health care decisions.

Reviewers and Updates

This summary is reviewed regularly and updated as necessary by the [PDQ Supportive and Palliative Care Editorial Board](#), which is editorially independent of the National Cancer Institute (NCI). The summary reflects an independent review of the literature and does not represent a policy statement of NCI or the National Institutes of Health (NIH).

Board members review recently published articles each month to determine whether an article should:

- be discussed at a meeting,
- be cited with text, or

- replace or update an existing article that is already cited.

Changes to the summaries are made through a consensus process in which Board members evaluate the strength of the evidence in the published articles and determine how the article should be included in the summary.

The lead reviewers for Nausea and Vomiting Related to Cancer Treatment are:

- Jared R. Lowe, MD, HMDC (University of North Carolina School of Medicine)
- Maria Petzel, RD, CSO, LD, CNSC, FAND (University of TX MD Anderson Cancer Center)

Any comments or questions about the summary content should be submitted to Cancer.gov through the NCI website's [Email Us](#). Do not contact the individual Board Members with questions or comments about the summaries. Board members will not respond to individual inquiries.

Levels of Evidence

Some of the reference citations in this summary are accompanied by a level-of-evidence designation. These designations are intended to help readers assess the strength of the evidence supporting the use of specific interventions or approaches. The PDQ Supportive and Palliative Care Editorial Board uses a [formal evidence ranking system](#) in developing its level-of-evidence designations.

Permission to Use This Summary

PDQ is a registered trademark. Although the content of PDQ documents can be used freely as text, it cannot be identified as an NCI PDQ cancer information summary unless it is presented in its entirety and is regularly updated. However, an author would be permitted to write a sentence such as “NCI’s PDQ cancer information summary about breast cancer prevention states the risks succinctly: [include excerpt from the summary].”

The preferred citation for this PDQ summary is:

PDQ® Supportive and Palliative Care Editorial Board. PDQ Nausea and Vomiting Related to Cancer Treatment. Bethesda, MD: National Cancer Institute. Updated <MM/DD/YYYY>. Available at: <https://www.cancer.gov/about-cancer/treatment/side-effects/nausea/nausea-hp-pdq>. Accessed <MM/DD/YYYY>. [PMID: 26389491]

Images in this summary are used with permission of the author(s), artist, and/or publisher for use within the PDQ summaries only. Permission to use images outside the context of PDQ information must be obtained from the owner(s) and cannot be granted by the National Cancer Institute. Information about using the illustrations in this summary, along with many other cancer-related images, is available in [Visuals Online](#), a collection of over 2,000 scientific images.

Disclaimer

The information in these summaries should not be used as a basis for insurance reimbursement determinations. More information on insurance coverage is available on Cancer.gov on the [Managing Cancer Care](#) page.

Contact Us

More information about contacting us or receiving help with the Cancer.gov website can be found on our [Contact Us for Help](#) page. Questions can also be submitted to Cancer.gov through the website's [Email Us](#).

Updated: March 10, 2025

If you would like to reproduce some or all of this content, see [Reuse of NCI Information](#) for guidance about copyright and permissions. In the case of permitted digital reproduction, please credit the National Cancer Institute as the source and link to the original NCI product using the original product's title;

e.g., "Nausea and Vomiting Related to Cancer Treatment (PDQ®)–Health Professional Version was originally published by the National Cancer Institute."