

Optimal Block Time for AMM Liquidity Providers under Jump-Diffusion Prices

Nils Bundi^[0000-0003-3576-3289]

Zurich University of Applied Sciences, School of Engineering,
Winterthur, Switzerland
`bund@zhaw.ch`

Abstract. Loss-versus-Rebalancing (LVR) is the dominant adverse-selection cost borne by liquidity providers on automated market makers. Under geometric Brownian motion, arbitrage profit scales with the probability of a profitable block, which vanishes as the block time $\Delta t \rightarrow 0$ — the standing argument for ever-shorter blocks. Modeling the reference price instead as a jump-diffusion, I show that the constant-product LVR rate splits into a diffusion channel carrying the known multiplier $F(\gamma/(\sigma\sqrt{\Delta t}))$ and a jump channel $\lambda V \cdot G(\gamma; m, \delta^2)$ carrying no Δt , the two interacting only through an explicitly bounded remainder. The block schedule therefore governs only one channel. For *symmetric* jump laws the jump channel is moreover an exact lower bound, $\ell(\Delta t) \geq \lambda V G > 0$, so the rate does not vanish as $\Delta t \rightarrow 0$, and descends *slowly* as $\sqrt{\Delta t}$. At Ethereum’s calibrated 12-second slot the rate is 471 bp/yr against a floor of 125, so only three quarters of LP loss is schedule-addressable. At Solana’s 400 ms slot the jump channel already dominates. Netting the rate against per-block consensus cost, the LP-side optimal block time is invariant in pool size and in every jump parameter (λ, m, δ) : jumps shift the level of LP loss but not the planner’s marginal tradeoff. Volatility, the fee tier, and consensus cost set the optimum, near 8 s. However, LVR is only one input to block-time welfare, so this bounds the LP-side contribution rather than settling the design question.

Keywords: Automated market makers · Loss-versus-Rebalancing · Jump-diffusion · Block time · Decentralized finance · Blockchain protocol design

1 Introduction

Automated market makers (AMMs) intermediate the bulk of on-chain spot trading, yet the passive liquidity providers who fund them systematically cede value to better-informed arbitrageurs whenever the external price moves. How fast a blockchain produces blocks—the cadence at which arbitrageurs can act—directly shapes this loss, and the prevailing intuition holds that faster blocks are better. This paper asks whether price *jumps* overturn that intuition, and finds that they impose a floor no block speed can breach.

Milionis et al. [22] formalize the adverse-selection cost of passive liquidity provision on a constant-function market maker (CFMM) as *loss-versus-rebalancing*

(LVR): the gap between a passive liquidity provider’s (LP) mark-to-market return and the return of a continuously rebalanced portfolio holding the LP’s instantaneous exposure. For a constant-product market maker (CPMM), a special case of the CFMM, with pool value $V(P)$ and external reference price S_t following a GBM with volatility σ , the LVR rate equals $\sigma^2 V/8$. Hence, it is independent of the block-time, the rate at which arbitrageurs are able to re-align the pool. Milionis et al. [21] extend this to swap fees γ and Poisson block arrivals of mean block-time Δt . In their low-fee fast-block regime, the LVR rate is multiplied by a strictly increasing function of $\sigma\sqrt{\Delta t}/\gamma$, so shorter blocks increase welfare.

This GBM-based prescription—“shorten blocks to reduce LVR”—is the implicit case for sub-second block times deployed on Solana, modern Ethereum L2s, and a host of app-chains. Empirically, however, crypto reference prices are not GBM but exhibit pronounced jumps [25,10,24]. A jump produces an instantaneous, Δt -independent price gap which is cleared in the very next block and the LP eats the concavity loss.

This paper asks: *what does the LVR/block-time relationship look like once prices jump?* The answer is that it splits in two: one channel the block schedule governs, and one it cannot touch — jumps drive the mispricing across the no-arbitrage band at a rate the protocol does not choose, so faster blocks clear each gap sooner without making it smaller. Calibrated to ETH/USDT, the split is stark: at Ethereum’s 12-second slot the rate is 471 bp/yr against a jump floor of 125; at Solana’s 400 ms the floor already dominates; and weighing LVR against Ethereum consensus cost puts the LP-side optimum near 8 s, above today’s sub-second targets of other chains.

Contributions. For a CPMM under jump-diffusion with any finite-activity jump law — Merton $(\mu, \sigma, \lambda, m, \delta)$ the running instance — Theorem 1 gives an additive zero-fee decomposition of the LVR rate into the classical diffusion term $\sigma^2 V/8$ and a nonnegative jump term $\frac{\lambda}{2} V \mathbb{E}[(e^{J/2} - 1)^2]$ that carries no Δt . Reintroducing a swap fee γ and discrete block time Δt , Theorem 2 shows that

$$\ell(\Delta t) = \frac{\sigma^2}{8} V \cdot F\left(\frac{\gamma}{\sigma\sqrt{\Delta t}}\right) + \lambda V \cdot G(\gamma; m, \delta^2) + \mathcal{E}(\Delta t), \quad (1)$$

where $F : \mathbb{R}_+ \rightarrow (0, 1]$ is the trade-probability multiplier P_{trade} of Milionis et al. [21] (Lemma 1, Remark 5), $G = \frac{1}{8} \mathbb{E}[(|J| - \gamma)^2 \mathbf{1}\{|J| > \gamma\}]$, and the remainder $\mathcal{E}$ — the only place the two channels meet — is bounded on both sides in Proposition 1 under a stated mixing condition. For symmetric ν satisfying a mild aggregation hypothesis (Merton does), Theorem 3 establishes exactly, with no appeal to (1), that $\ell(\Delta t) \geq \lambda V G > 0$ at every Δt : an irreducible jump floor that no block-time reduction can eliminate, approached only as $\sqrt{\Delta t}$ (Corollary 1). Finally, netting $\ell_0(\Delta t)$, the first two terms of (1), against pool-attributable block-production cost, Theorem 4 characterizes the LP-side planner’s optimum: invariant in pool size, and in the jump parameters (λ, m, δ) to the order at which (1) separates the channels — jumps shift the welfare level but not the planner’s marginal tradeoff.

2 Related Work

CFMMs are the dominant trading mechanism in decentralized finance, with the CPMM invariant first deployed at scale by Uniswap [1]. Angeris et al. [2] characterize arbitrage equilibria, reserve dynamics and the convergence of pool prices to external references. Milionis et al. [22] introduce *loss-versus-rebalancing* (LVR), the LP’s underperformance against a continuously-rebalanced benchmark, and derive under GBM a measure-independent rate—independent of the LP’s risk-neutral probability and of hedging considerations—with the full-range Uniswap-V2 specialization $\sigma^2 V/8$. Milionis et al. [21] extend this to proportional fees with Poisson-arrival blocks. Their key structural result, and the one this paper builds on, is that the log mispricing between pool and reference is ergodic with stationary law uniform on the no-arbitrage band and exponential tails outside it, so that arbitrage profits scale down from the frictionless LVR by the stationary probability of a profitable block—the first formal link between protocol-level block-time and LP loss. Nezlobin and Tassy [23] extend this to general block-time distributions and Fritsch and Canidio [18] validate the scaling empirically across major Uniswap pools. Cartea, Drissi and Monga [9] frame the counterpart notion of *predictable loss* in continuous-time wealth dynamics and solve for optimal concentrated-liquidity provision against fee income and concentration risk, while Capponi and Jia [7] micro-found the loss channel through a deposit game exhibiting a “liquidity freeze” regime.

On price dynamics, jump-diffusion has been standard in derivatives pricing since Merton [20], with Kou’s asymmetric double-exponential law [19] and the Carr–Geman–Madan–Yor pure-jump Lévy model [8] as flexible alternatives, and Cont and Tankov [12] the reference for the Itô–Lévy machinery used below. Empirically, crypto jumps are pervasive: Scaillet et al. [25] document frequent, clustered and order-flow-predictable jumps in Bitcoin at high frequency; Saef et al. [24] find in tick data from seven exchanges over April 2019–September 2021 that Bitcoin jumps on $\sim 58\%$ and Ethereum on $\sim 32\%$ of trading days, with systematic temporal clustering; and Chaim and Laurini [10] document nonlinear dependence and tail co-movement across major cryptocurrencies. Cont [11] surveys the broader stylized facts of asset returns.

Block-time choice balances several welfare components beyond LP loss. On *consensus cost*, Decker and Wattenhofer [15] identify block propagation delay as the primary cause of forks and Croman et al. [13] per-block bandwidth overhead as the binding scaling constraint. On *finality*, Buterin [6] surveys paths to Ethereum single-slot finality, where BLS aggregation over committee attestations is the per-slot bottleneck. On *decentralization and censorship resistance*, Benhaim et al. [4] show committee-based consensus attains equivalent security with committees one to two orders of magnitude smaller—legitimizing the architecture behind sub-second blocks, but exposing how small committees increase censorship vulnerability. On *MEV redistribution*, Daian et al. [14] document the maximal-extractable-value economy that block-time mediates: longer blocks expand per-block extractable value, shorter blocks fragment it and erode the searcher timing advantage. *Oracle freshness*, the gap between on-chain feeds

and live markets, scales directly with Δt . LP-side adverse selection, the subject of this paper, is one input among these.

Two gaps emerge, and this paper closes both. The LVR construct is well-developed under GBM but has not been extended to jump-diffusion reference prices despite the empirical prevalence of price discontinuities in crypto markets, and the LP-side literature lacks an explicit planner's optimum that exposes how the jump parameters enter, or fail to enter, that optimum.

3 Model

The model is that of Milionis et al. [22,21] with a single primitive changed: the reference price follows a Merton jump-diffusion rather than pure GBM, capturing the discontinuous price moves documented empirically in cryptocurrency markets.

3.1 Price Process

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbb{P})$ be a filtered probability space supporting a standard Brownian motion W and an independent Poisson random measure with intensity $\lambda dt \nu(dy)$, where ν is the distribution of the log jump size J , assumed to have finite activity and to satisfy $\int (e^{j/2} - 1)^2 \nu(dj) < \infty$; the quadratic results of Section 5 need only a finite second moment. Merton's $J \sim \mathcal{N}(m, \delta^2)$ is the running instance and the one calibrated in Section 7. Section 4 needs no more than the moment condition, while from Section 5 on I assume in addition that ν is *symmetric* — for Merton, $m = 0$. Remark 4 isolates exactly where symmetry is used. The reference price S_t follows

$$\frac{dS_t}{S_{t-}} = \mu dt + \sigma dW_t + (e^{J_t} - 1) dN_t, \quad (2)$$

with N a Poisson process of intensity λ . Time is measured in years throughout, so σ , λ and Δt are all in annual units; block times are quoted in seconds in Section 7 for readability only. The notation $S_{t-} := \lim_{s \uparrow t} S_s$ denotes the left limit, i.e., the pre-jump price at time t ; using S_{t-} rather than S_t in the denominator makes the relative increment well-defined at jump times and ensures the integrand is predictable.

3.2 CPMM

A CPMM with reserves (x_t, y_t) maintains the invariant $x_t y_t = L$. The marginal pool price is $P_t = y_t/x_t$, and the pool's holdings as functions of P are $x(P) = \sqrt{L/P}$ and $y(P) = \sqrt{LP}$. The pool value (in numeraire units) at marginal price P is $V(P) := y(P) + P x(P) = 2\sqrt{LP}$. By the envelope identity, $V'(P) = x(P)$ and $V''(P) = -\frac{1}{2}\sqrt{L}P^{-3/2} = -x(P)/(2P) < 0$. Note that the pool value V is concave and this concavity is the source of LVR. I work with the full-range

Uniswap specialization throughout for concreteness, but per Milionis et al. [22, Remark 1] the LVR construct depends only on a locally-smooth demand curve $x^*(P)$ and applies with no modification to general CFMMs and concentrated-liquidity AMMs (within active tick ranges).

3.3 Arbitrageur

A risk-neutral, capital-unconstrained arbitrageur observes the external price S_t continuously and the pool price P_t . Trading on the pool incurs a proportional fee $\gamma \in [0, 1)$ in log-price units (i.e., a swap effectively moves the marginal price by an extra factor $e^{\pm\gamma}$), so the no-arbitrage band is $\{|\log S - \log P| \leq \gamma\}$. Blocks arrive as a Poisson process of rate $1/\Delta t$, so Δt is the mean interblock time, and the arbitrageur can trade only when a block arrives. The reference price itself moves continuously between blocks: S is an off-chain venue price, so the arbitrageur's information is continuous even though on-chain reserves settle only at block inclusion. Mempool visibility of pending transactions extends the same continuity to the on-chain order flow, so the pool price is not the only price a participant can observe between blocks. Being competitive, the arbitrageur trades myopically whenever it is profitable [21, §3].

3.4 LVR

Following Milionis et al. [22], define the LVR over $[0, T]$ as

$$\text{LVR}(T) := \int_0^T V'(P_{s-}) dP_s - [V(P_T) - V(P_0)]. \quad (3)$$

The first term is the return of a rebalancing strategy holding $V'(P_{s-}) = x(P_{s-})$ units of the risky asset at each instant; the second is the LP's mark-to-market return. The *LVR rate* is $\ell(t) = \mathbb{E}_t[d\text{LVR}_t]/dt$, evaluated at the current pool state.

In the continuous-arbitrage limit ($\gamma = 0$, $\Delta t \rightarrow 0$), the pool price tracks the reference price, $P_t \equiv S_t$, and (3) is the Itô–Lévy remainder of $V(S_t)$.

4 LVR under Jump-Diffusion: Zero-Fee Decomposition

Theorem 1 (LVR decomposition). *Under the zero-fee continuous-arbitrage limit with S_t following (2) and the CPMM pool value $V(P) = 2\sqrt{LP}$, the instantaneous LVR rate is*

$$\ell = \underbrace{\frac{\sigma^2}{8} V(S_t)}_{\text{diffusion-LVR}} + \underbrace{\frac{\lambda}{2} V(S_t) \mathbb{E}[(e^{J/2} - 1)^2]}_{\text{jump-LVR}}, \quad (4)$$

and for $J \sim \mathcal{N}(m, \delta^2)$, $\mathbb{E}[(e^{J/2} - 1)^2] = e^{m+\delta^2/2} - 2e^{m/2+\delta^2/8} + 1 \geq 0$. This expression is 0 if and only if $J \equiv 0$.

Proof. See Appendix A.1 of the extended version [5].

Remark 1 (Measure). The drift enters (3) only through $\int V'(S_{s-}) dS_s$, which appears in both terms and cancels, so (4) involves σ and the jump measure alone and no normalization of μ is required. Under GBM the rate is in this sense measure-independent [22]; with jumps it is not, since an equivalent change of measure rescales λ and reshapes ν , so Theorem 1 is a statement under the physical measure and calibration of the model is versus realized-price observations (Section 7).

Remark 2. The proof depends on the jump distribution only through $\mathbb{E}[(e^{J/2} - 1)^2]$. For any compound-Poisson jump measure ν with $\int (e^{j/2} - 1)^2 \nu(dj) < \infty$ and $\nu \neq \delta_0$, the jump-LVR rate is $\frac{\lambda V}{2} \int (e^{j/2} - 1)^2 \nu(dj) > 0$. Merton is illustrative; Kou's double-exponential law admits the same structure with different moments, as does any finite-activity ν . The results of Section 5 likewise use the shape of ν only through symmetry and the aggregation hypothesis of Theorem 3, not through Gaussianity.

Remark 3. Theorem 1 does not involve Δt . At $\gamma = 0$, the arbitrageur eliminates any pricing gap within the next block, so block-time refinement only redistributes *when* the loss is realized, not its rate. The block-time becomes economically relevant only once fees create a no-arbitrage band.

5 Fees and Block Time

I now allow $\gamma > 0$ and finite $\Delta t > 0$. Write $z_t := \log(S_t/P_t)$ for the log mispricing between reference and pool. At a block arrival τ_i the arbitrageur trades iff $|z| > \gamma$ and moves the pool to the near edge of the no-arbitrage band, so $z_{\tau_i} = \text{bound}(z_{\tau_i-}, -\gamma, \gamma)$; between blocks P is frozen and z inherits the dynamics of $\log S$. The mispricing is therefore a *persistent state*: a trade leaves it at $\pm\gamma$, not at 0, and absent a trade it carries forward. Writing $\mathcal{J}_i := \text{bound}(z_{\tau_i-}, -\gamma, \gamma) - z_{\tau_i-}$ for the arbitrage resets and T_k for the jump times of N , the Itô-Lévy formula applied to $\log S$ under (2) gives

$$z_t = \left(\mu - \frac{1}{2}\sigma^2\right)t + \sigma W_t + \underbrace{\sum_{k: T_k \leq t} J_k}_{\text{exogenous, rate } \lambda} + \underbrace{\sum_{i: \tau_i \leq t} \mathcal{J}_i}_{\text{endogenous, rate } 1/\Delta t}. \quad (5)$$

The mispricing is driven by two point processes: the exogenous price jumps, which push z *away* from the band at a rate the protocol cannot choose, and the block-triggered resets, which push it back. Every result below follows from that asymmetry (illustrated in Appendix B of the extended version [5]). As in [21, Assumption 2] I set $\mu = \frac{1}{2}\sigma^2 - \lambda m$.

At a block with pre-trade mispricing z the pool's marginal log-price moves by $\xi = \text{sign}(z)(|z| - \gamma)_+$, and by the computation in the proof of Theorem 1 the

LP's loss is $h(\xi) := \frac{V}{2}(e^{\xi/2} - 1)^2$. Blocks being Poisson and independent of z , the pre-block law of z is its stationary law π , and

$$\ell(\Delta t) = \frac{1}{\Delta t} \mathbb{E}_\pi[h(\xi)], \quad h(\xi) = \frac{V}{8}\xi^2 + \frac{V}{2}\rho(\xi), \quad \rho(\xi) := (e^{\xi/2} - 1)^2 - \frac{\xi^2}{4}. \quad (6)$$

Two facts about the split in (6) carry the results below. The quadratic surrogate $\frac{V}{8}\xi^2$ is a *convex* function of z , which h itself is not — a downward move of arbitrary size costs at most $V/2$ — and convexity is what Theorem 3 needs. And the curvature residual ρ is signed under symmetry: pairing $\pm\xi$,

$$\frac{1}{2}[\rho(\xi) + \rho(-\xi)] = \cosh \xi - 2 \cosh \frac{\xi}{2} + 1 - \frac{\xi^2}{4} = \sum_{n \geq 2} \frac{1 - 2^{1-2n}}{(2n)!} \xi^{2n} \geq 0, \quad (7)$$

so for any symmetric law of ξ the exact loss is at least the quadratic surrogate, with leading term $\frac{7}{192}\xi^4$. Both statements are exact, for every $\xi \in \mathbb{R}$.

Remark 4 (Driftlessness and symmetry). Setting $\mu = \frac{1}{2}\sigma^2 - \lambda m$ makes $\log S_t$ driftless. Additionally setting $m = 0$ makes z symmetric. The two properties are used differently below: Driftlessness is a modelling restriction rather than a normalization, since the rate at $\gamma > 0$ depends on the full law of z (its cost is small, as a non-zero drift Δ tilts the stationary law by only $2\Delta/\sigma^2$ [21, Thm. 8], worth under 0.01% of ℓ for any $|\Delta| \leq 1.6/\text{yr}$, which covers the empirical ETH drift by a wide margin); Symmetry of ν is used only to sign the remainder and to prove the floor, and it is symmetry that is needed, not Gaussianity: Theorems 1, 2 and 4 and Lemmas 1–2 hold for arbitrary ν , while Proposition 1 and Theorem 3 require it. Everything the block schedule governs — the separated rate ℓ_0 , the planner's cubic and its invariances — is therefore general; symmetry enters only in transferring those properties to the exact rate ℓ .

5.1 Diffusion Contribution with Fees

Everything therefore turns on π . For $\lambda = 0$ it is given in closed form by [21, Thm. 1]: with $\theta := \sqrt{2/\Delta t}/\sigma$ and $\eta := \theta\gamma = \sqrt{2}\gamma/(\sigma\sqrt{\Delta t})$,

$$p(x) = c \quad (|x| \leq \gamma), \quad p(x) = c e^{-\theta(|x|-\gamma)} \quad (|x| > \gamma), \quad c = \frac{\theta}{2(1+\eta)}, \quad (8)$$

uniform on the band with mass $\eta/(1+\eta)$ and exponential tails of mass $\frac{1}{2}(1+\eta)^{-1}$ each, so $\mathbb{P}_\pi(|z| > \gamma) = (1+\eta)^{-1}$. Two features of (8) carry the argument: the band holds most of the mass when blocks are fast, which is why the multiplier below vanishes; and the tails decay on the scale $1/\theta = \sigma\sqrt{\Delta t}/2$, so a jump with $|J| \gg \gamma$ lands far outside both band and tail and is cleared in full at the next block however small Δt is. With $\lambda > 0$ the stationary law solves an integro-differential rather than an ordinary differential equation, the compound-Poisson generator adding a non-local term of relative order $\lambda\Delta t$; that perturbation is carried through Proposition 1 and illustrated in Appendix B of the extended version [5].

Lemma 1 (Diffusion LVR with fees). *Let $\lambda = 0$. For the quadratic surrogate in (6) the LVR rate is exactly*

$$\ell_{\text{diff}}(\Delta t) = \frac{\sigma^2}{8} V \cdot F(\kappa), \quad F(\kappa) = \frac{1}{1 + \sqrt{2}\kappa} = \mathbb{P}_\pi(|z| > \gamma), \quad \kappa := \frac{\gamma}{\sigma\sqrt{\Delta t}}, \quad (9)$$

the stationary probability that an arriving block offers a profitable trade. F maps $\mathbb{R}_+$ onto $(0, 1]$, satisfies $F(0) = 1$ and $F(\kappa) \sim 1/(\sqrt{2}\kappa)$, and is strictly decreasing, with $F'(\kappa) = -\sqrt{2}(1 + \sqrt{2}\kappa)^{-2} < 0$.

Proof. See Appendix A.2 of the extended version [5].

Fritsch and Canidio [18] test this scaling empirically.

Remark 5 (Relation to Milionis et al.). Under a GBM reference price the mispricing is a regulated Brownian motion driven by the single point process of block arrivals, whose marks always push z towards the band [21, Eq. (9)]. Equation (5) adds a second, exogenous driver whose marks push z away from it; the floor of Corollary 1 is the consequence. The per-trade loss $\frac{V}{8}\xi^2$ entering (6) is their Lemma 2 profit up to $e^\gamma = 1 + O(\gamma)$, and (9) is exact for the quadratic loss.

5.2 Jump Contribution with Fees

When the reference price jumps by J , the arbitrageur captures the concavity loss only if the jump exceeds the fee threshold (otherwise the no-arbitrage band absorbs it without realization).

Lemma 2 (Jump LVR with fees). *Under (2), and for the same quadratic surrogate underlying (9), the separated contribution of jumps to the LVR rate is $\lambda V \cdot G_\nu(\gamma)$, where*

$$G_\nu(\gamma) := \frac{1}{8} \mathbb{E}_\nu \left[(|J| - \gamma)^2 \cdot \mathbf{1}\{|J| > \gamma\} \right], \quad (10)$$

a functional of ν alone; I write $G(\gamma; m, \delta^2)$ for its Merton value. For $J \sim \mathcal{N}(m, \delta^2)$, G is strictly positive whenever $\mathbb{P}(|J| > \gamma) > 0$, satisfies $G(0; m, \delta^2) = (m^2 + \delta^2)/8$, and is strictly decreasing in γ .

Proof. See Appendix A.3 of the extended version [5].

A closed-form evaluation for symmetric Merton jumps ($m = 0$) follows from the truncated-normal second-moment identity $\Psi(x) := \mathbb{E}[(|Z| - x)_+^2] = 2[(1 + x^2)\Phi(-x) - x\varphi(x)]$ for $Z \sim \mathcal{N}(0, 1)$. Letting $\zeta := \gamma/\delta$, this yields

$$G(\gamma; 0, \delta^2) = \frac{\delta^2}{4} [(1 + \zeta^2)\Phi(-\zeta) - \zeta\varphi(\zeta)] = \frac{\delta^2}{8} \Psi(\zeta). \quad (11)$$

The jump term thus carries a fee discount of its own, Ψ at the jump-size scale δ ; crucially it is Ψ , not F , and it does not involve Δt , because the size of a jump is set by δ rather than by how long the block lasted. At $\gamma = 0$, $G(0; 0, \delta^2) = \delta^2/8$. For asymmetric Merton ($m \neq 0$) and other jump distributions, G admits analogous truncated-moment representations.

Remark 6 (The curvature residual in closed form). Lemmas 1 and 2 evaluate the quadratic surrogate of (6). The residual $\frac{V}{2}\mathbb{E}_\pi[\rho(\xi)]$ they omit is available in closed form on both channels. On the diffusion channel the stationary tails are exponential with rate θ , so the exact- h rate is $\frac{V}{2\Delta t}\Lambda(\theta)/(1+\eta)$ with $\Lambda(\theta) = \theta^2/(\theta^2-1) - 2\theta^2/(\theta^2-\frac{1}{4}) + 1$, and $2\theta^2\Lambda(\theta) = 1 + \frac{7}{4}\theta^{-2} + O(\theta^{-4}) = 1 + \frac{7}{8}\sigma^2\Delta t + O((\sigma^2\Delta t)^2)$, a relative correction of 2.2×10^{-7} at $\Delta t = 12$ s that vanishes as $\Delta t \rightarrow 0$. On the jump channel the same substitution turns G into a truncated Gaussian moment-generating expression G^h in Φ , with $G^h/G - 1 = \frac{7}{16}\delta^2 + O(\delta^4) = 1.6 \times 10^{-4}$ at $\delta = 0.0192$. Both corrections are nonnegative by (7) whenever π is symmetric, and both are carried as exactly-evaluated terms of $\mathcal{E}$ in Proposition 1 — so nothing below requires δ or $\sigma\sqrt{\Delta t}$ to be small. The unbounded support of J is likewise immaterial: $\mathbb{E}[\rho(\xi)]$ is finite.

Remark 7. Equation (11) reveals a clean equivalence: to leading order in δ , a Poisson stream of jumps with standard deviation δ contributes the same LVR rate as a diffusion of effective volatility $\sigma_J := \delta\sqrt{\lambda}$, both giving $\sigma_J^2 V/8$ at $\gamma = 0$. Fees break it asymmetrically: the diffusion suffers the $F(\gamma/(\sigma\sqrt{\Delta t}))$ discount, which vanishes as $\Delta t \rightarrow 0$, while jumps suffer only the Δt -invariant $\Psi(\gamma/\delta)$ discount. The jump floor's persistence is the formal expression of this asymmetry.

5.3 Main Formula

Combining the diffusion contribution (9) with the jump contribution of Lemma 2 yields the LVR rate as a sum of a Δt -dependent diffusion term and a Δt -invariant jump floor.

Theorem 2 (LVR rate under jump-diffusion with fees). *Under (2) with swap fee $\gamma \in [0, 1)$ and mean block time $\Delta t > 0$, the LVR rate of a CPMM satisfies*

$$\ell(\Delta t) = \ell_0(\Delta t) + \mathcal{E}(\Delta t), \quad \ell_0(\Delta t) := \frac{\sigma^2}{8} V \cdot F\left(\frac{\gamma}{\sigma\sqrt{\Delta t}}\right) + \lambda V \cdot G(\gamma; m, \delta^2), \quad (12)$$

with F as in (9) and G as in (10). Equation (12) is a definition of $\mathcal{E}$ and carries no content until $\mathcal{E}$ is controlled; Proposition 1 bounds it on both sides. I call ℓ_0 the separated rate: it is the object in which the two channels do not interact, and the properties established for it below are properties of ℓ_0 unless stated otherwise. The one property that transfers to ℓ itself with no remainder is the floor, Theorem 3.

Proposition 1 (Two-sided bound on $\mathcal{E}$). *Let ν be symmetric, and let n_* denote the least n with $\sup_{m, m'} \|K_0^n(m, \cdot) - K_0^n(m', \cdot)\|_{\text{TV}} \leq \frac{1}{2}$, where K_0 is the post-block kernel at $\lambda = 0$. Assume the mixing condition*

$$n_* \leq C(1 + \eta^2), \quad C = 2. \quad (13)$$

Only η enters (13); it is checked numerically over every block time, volatility and Uniswap fee tier considered here. It is used solely to bound the finite- λ remainder

in the separated expression. With $g(d) := (|d| - \gamma)_+^2$, the separated rate ℓ_0 differs from the exact stationary rate (6) in four ways: it charges $\mathbb{E}[g(J)]$ in place of $\mathbb{E}[g(M + X + J)]$ on the jump term, it charges that term at the full jump rate λ rather than the thinned single-jump block rate $\lambda(1 + \lambda\Delta t)^{-2}$, it evaluates the diffusion term against the $\lambda = 0$ law (8) and block clock, and it replaces h by its quadratic surrogate. Each is signed, so the remainder is bounded asymmetrically (with $\delta^2 := \mathbb{E}_\nu[J^2]$ for non-Merton ν):

$$\begin{aligned} & -\underbrace{\frac{5}{2}\lambda\Delta t\ell_{\text{diff}}(\Delta t)}_{\text{block clock}} - \underbrace{\frac{\lambda V}{4}(2\gamma^2 + \sigma^2\Delta t)}_{\text{stationary law}} - \underbrace{2\lambda^2\Delta tVG_\nu(\gamma)}_{\text{thinning}} \leq \mathcal{E}(\Delta t) \\ & \leq \underbrace{\frac{\lambda V}{8}(2\sigma^2\Delta t + \gamma^2)}_{\text{interaction}} + \underbrace{\frac{\lambda V}{4}(2\gamma^2 + \sigma^2\Delta t)}_{\text{stationary law}} + \underbrace{\mathcal{E}_{\text{curv}}}_{\text{curvature}} + \underbrace{\frac{V\lambda^2\Delta t}{8}(\gamma^2 + 3\sigma^2\Delta t + 2\delta^2)}_{\text{multi-jump}}, \end{aligned}$$

where the interaction term uses only $|M| \leq \gamma$, which holds pathwise since a block leaves the pool inside the band, and $\mathcal{E}_{\text{curv}} = \frac{V}{2}\mathbb{E}_\pi[\rho(\xi)] \geq 0$ is the curvature residual, evaluated exactly in closed form in Remark 6. The stationary-law term covers the one place ℓ_{diff} is evaluated against (8) rather than the true law of the carried state, which jump feedback shifts. Its leading piece $\lambda V\gamma^2/2$ does not vanish with Δt , because the band relaxes on the timescale γ^2/σ^2 whatever the block schedule, so an $O(\lambda\Delta t)$ perturbation per block accumulates over $O(\Delta t^{-1})$ blocks. The thinning term is the deficit of charging every jump at rate λ when a fraction $1 - (1 + \lambda\Delta t)^{-2}$ of them land in intervals with other jumps, whose aggregate losses appear, nonnegative, in the last term of the upper side. At the calibration of Section 7 this gives $\mathcal{E}(\Delta t) \in [-0.653, +0.849]$ bp/yr at $\Delta t = 12$ s and $[-0.355, +0.464]$ at 50 ms, never exceeding 0.29% of the rate over the deployed range 50 ms–12 s. Evaluating (6) by quadrature puts the realised remainder at +0.165 and +0.052 bp/yr respectively, inside the bound and of the predicted sign. As $\Delta t \rightarrow 0$ the block-clock term vanishes while the interaction tends to $\lambda V\gamma^2/8$, so $\ell(\Delta t) - \lambda VG_\nu(\gamma)$ is squeezed into $[0, 0.463]$ bp/yr, at most 0.37% of the floor.

Proof. See Appendix A.4 of the extended version [5].

The floor claim holds exactly, it is not subject to $\mathcal{E}$ and Proposition 1, and the argument never touches the stationary law — which is what makes it available, since with $\lambda > 0$ that law has no closed form. Two limits should be kept apart here: $\Delta t \rightarrow 0$ shortens the interval between blocks, but the jump law is exogenous to the block schedule, so it does not shrink J . Fast blocks do not imply small jumps, and the floor exists precisely because of that.

Theorem 3 (Exact jump floor). *Let ν be symmetric and satisfy the aggregation hypothesis $\mathbb{E}[g(\sum_1^k J)] \geq k\mathbb{E}[g(J)]$ for every $k \geq 1$, as Merton with $m = 0$ does. Then for every $\Delta t > 0$,*

$$\ell(\Delta t) \geq \lambda V \cdot G_\nu(\gamma) = \inf_{\Delta t' > 0} \ell_0(\Delta t') > 0,$$

with no approximation and no remainder, the middle equality being immediate from $F > 0$ and $F(\kappa) \rightarrow 0$. The floor that the separated rate (12) locates is therefore a floor for the exact rate, not an artifact of the separation.

Proof. See Appendix A.5 of the extended version [5].

Corollary 1 (Approach to the floor). *The separated rate $\Delta t \mapsto \ell_0(\Delta t)$ is strictly increasing on $(0, \infty)$, with*

$$\lim_{\Delta t \rightarrow 0} \ell_0(\Delta t) = \lambda V \cdot G(\gamma; m, \delta^2), \quad \lim_{\Delta t \rightarrow \infty} \ell_0(\Delta t) = \frac{\sigma^2}{8} V + \lambda V \cdot G(\gamma; m, \delta^2),$$

and the floor is approached at the algebraic rate $\ell_0(\Delta t) - \lambda V G \sim \sigma^3 V \sqrt{\Delta t} / (8\sqrt{2}\gamma)$. For the exact rate, $\lambda V G \leq \liminf_{\Delta t \rightarrow 0} \ell(\Delta t) \leq \limsup_{\Delta t \rightarrow 0} \ell(\Delta t) \leq \lambda V G + \lambda V \gamma^2 / 8 + \lambda V \gamma^2 / 2 + \mathcal{E}_{\text{curv}}$, the lower bound by Theorem 3 and the upper by Proposition 1; the two differ by 0.37% of the floor at the calibration of Section 7. Monotonicity likewise transfers to ℓ , under (13), wherever the separated increment clears the width of the remainder envelope: $\ell(\Delta t_2) > \ell(\Delta t_1)$ for any $\Delta t_1 < \Delta t_2$ with $\ell_0(\Delta t_2) - \ell_0(\Delta t_1) > 1.50$ bp/yr, which holds for every pair reported in Section 7, and quadrature of (6) confirms monotonicity of ℓ itself across that range.

Proof. See Appendix A.6 of the extended version [5].

Remark 8 (The floor as un-recoverable LVR). In the accounting of [21, Thms. 3–4], arbitrage profit and fee income sum to the frictionless LVR of Theorem 1: fees split LVR rather than reduce it, and $\ell(\Delta t)$ is the arbitrage half, LP loss net of the fees LPs receive back. On the diffusion channel that split is $F(\kappa) : 1 - F(\kappa)$ with $F(\kappa) \rightarrow 0$, so shorter blocks return diffusion-LVR to LPs as fee income. On the jump channel (11) makes it $\Psi(\gamma/\delta) : 1 - \Psi(\gamma/\delta)$, a function of γ/δ alone with no Δt in it, because a jump exceeding the band is cleared at the next block whatever the block rate. Fees recover only $1 - \Psi(\gamma/\delta)$ of jump-LVR—4.1% at the calibration below—and the rest is the floor. The lever against it is γ/δ , i.e. fee-tier design, not block time.

6 The Planner’s Block-Time Optimum

Shorter blocks impose real costs that must be funded somehow. Each block requires fixed per-block work—block gossip and propagation [15], BLS signature aggregation over committee attestations [6], and view-change/commit overhead in committee-based consensus [4]—so the true *validator operating cost* is a resource cost of hardware, bandwidth, and energy that scales with the number of blocks produced. Since this resource cost is not directly observable, I rely on the chain’s native-token issuance as a proxy for it. Because issuance is a chosen compensation policy rather than a measured cost, it overstates the resource cost where it acts as a growth subsidy and understates it where fees and MEV fund validators [6,3]. This bias only shifts the level of the cost, an effect discussed in Section 8. I thus adopt the simplest specification with fixed per-block cost rate $R(\Delta t) = r/\Delta t$, calibrating the per-block chain-level cost r from observed issuance.

A representative pool with TVL V inherits a share of the chain’s marginal resource cost in proportion to value secured: $c(\Delta t) = R(\Delta t) \cdot V/V_{\text{chain}} = c/\Delta t$ where $c := r \cdot V/V_{\text{chain}}$ and V_{chain} is the chain’s native-token market capitalization. The LP-side planner chooses Δt to minimize the total per-unit-time cost rate combining the LVR rate from Theorem 2 with the per-pool consensus cost, $W(\Delta t) = \ell_0(\Delta t) + c/\Delta t$, written against the separated rate ℓ_0 of (12). Theorem 4 is therefore a statement about ℓ_0 ; Remark 10 quantifies what the remainder does to it.

Theorem 4 (Optimal block time). *Let $c > 0$. Any interior critical point Δt^{opt} of $W(\Delta t)$ satisfies $\ell'_0(\Delta t^{\text{opt}}) = c/(\Delta t^{\text{opt}})^2$. With $\kappa = \gamma/(\sigma\sqrt{\Delta t})$, $d\kappa/d\Delta t = -\kappa/(2\Delta t)$, and the derivative in Lemma 1,*

$$\ell'_0(\Delta t) = \frac{\sigma^2 V}{8} F'(\kappa) \cdot \left(-\frac{\kappa}{2\Delta t}\right) = \frac{\sigma^2 V}{8} \cdot \frac{\sqrt{2}\kappa}{2\Delta t(1 + \sqrt{2}\kappa)^2} > 0. \quad (14)$$

The first-order condition, after substituting $\Delta t = \gamma^2/(\sigma^2\kappa^2)$ and writing $\eta := \sqrt{2}\kappa$, reduces to a cubic:

$$\boxed{\eta^{\text{opt}}(1 + \eta^{\text{opt}})^2 = \frac{V\gamma^2}{8c}, \quad \Delta t^{\text{opt}} = \frac{2\gamma^2}{\sigma^2(\eta^{\text{opt}})^2}.} \quad (15)$$

Since $\eta \mapsto \eta(1 + \eta)^2$ is a strictly increasing bijection of $(0, \infty)$ onto itself, (15) admits a unique positive solution for every $c > 0$, and W attains its minimum there. The root is available in closed form by Cardano’s formula: with $A := V\gamma^2/(8c)$, $B := A/2 + 1/27$ and $u := B + \sqrt{B^2 - 1/729}$, $\eta^{\text{opt}} = u^{1/3} + \frac{1}{9}u^{-1/3} - \frac{2}{3}$.

Proof. See Appendix A.7 of the extended version [5].

Remark 9. Differentiating (15) yields four sensitivities. First, $\partial\Delta t^{\text{opt}}/\partial\lambda = \partial\Delta t^{\text{opt}}/\partial m = \partial\Delta t^{\text{opt}}/\partial\delta = 0$: the jump parameters do not enter the first-order condition, shifting the level of W by λVG but leaving the marginal tradeoff untouched. This is the paper’s main qualitative message. It is exact for ℓ_0 and holds up to an explicitly bounded shift for ℓ (Remark 10). Second, $\partial\Delta t^{\text{opt}}/\partial c > 0$: more expensive consensus lengthens the optimum, the planner amortizing the per-block cost over longer intervals and accepting more LVR per block. Third, $\partial\Delta t^{\text{opt}}/\partial\gamma$ is U-shaped, since at very low γ the multiplier F stays close to 1 and block-shortening cannot suppress diffusion-LVR, while at very high γ it is already crushed and shortening offers no further benefit; the interior minimum sits where shortening is an effective lever. Fourth, $\partial\Delta t^{\text{opt}}/\partial V = 0$: since $c = r V/V_{\text{chain}}$ is proportional to V , the right-hand side of (15) equals $\gamma^2 V_{\text{chain}}/(8r)$ and V cancels, so the optimum is a property of the chain and the fee tier, not of the pool. This one is exact: V cancels from (15) identically, with no appeal to the separation.

Remark 10 (What the remainder does to the optimum). The invariance in (λ, m, δ) is a property of ℓ_0 , whose jump term is an additive constant in Δt . In the exact

model $\mathcal{E}$ carries both λ and Δt , so the exact first-order condition is not jump-free and the invariance is not a theorem about ℓ . Its failure is small enough to bound directly. Under (13), any minimizer of $\ell + c/\Delta t$ lies in $\{\Delta t : W(\Delta t) + \mathcal{E}^-(\Delta t) \leq \min_s [W(s) + \mathcal{E}^+(s)]\}$ with $\mathcal{E}^\pm$ the envelope of Proposition 1; at the calibration of Section 7 this set is $[7.3, 9.9]$ s around the separated optimum $\Delta t^{\text{opt}} = 8.4$ s. So the remainder can move the optimum by about a second — which is also the range over which W sits within 1 bp/yr of its minimum anyway, so the shift is not resolvable against the flatness of the objective. The honest statement is therefore quantitative rather than absolute: jump parameters move Δt^{opt} by less than the flatness of the objective can resolve.

7 Numerical Illustration

I illustrate the qualitative structure of Theorems 2 and 4 for a hypothetical full-range LP on an ETH/USDT Uniswap pool. Model parameters are summarized in Table 1.

Table 1. Baseline model parameters for the ETH/USDT full-range LP illustration

Parameter	Value
σ (diffusion volatility)	0.8156
λ (jump intensity)	283.3/yr
m (mean log jump)	0
δ (std of log jump)	0.0192
γ (swap fee)	0.0005
V (pool TVL)	1,000,000 USD
c (consensus cost)	260×10^{-5} USD/block

The price parameters $(\sigma, \lambda, \delta)$ are estimated from Binance ETH/USDT 5-minute closes over January 2020–June 2026, 2,373 days. I take σ from the continuous part of realised variance, estimated by bipower variation, and obtain λ and δ from the jump-detection test of Lee and Mykland at the 1% level, correcting the detected jump sizes for their diffusive component. The per-pool consensus cost c follows from Ethereum’s gross issuance ($\sim \$1.7\text{B}/\text{yr}$ at $V_{\text{ETH}} \approx \$208\text{B}$), scaled by $25\%/30\% \approx 0.83$ to reflect that current staking ($\sim 30\%$ of supply) exceeds Drake’s security-optimal $1/4$ [16,17], then attributed pro-rata by $V/V_{\text{chain}} = 4.8 \times 10^{-6}$, yielding $c \approx \$260 \times 10^{-5}/\text{block}$. The remaining parameters are modeling choices: $m = 0$ (symmetric jumps — the sample gives $\hat{m} = -0.0012$ with centred Lee–Mykland local-volatility window; the 99% interval from the asymptotic standard error of the mean detected jump, $[-0.0026, +0.0003]$, contains zero, consistent with the choice), $\gamma = 5$ bps (Uniswap 0.05% tier), $V = \$1\text{M}$ (pool TVL). Sensitivities below cover $\sigma \in \{0.30, 0.816, 1.50\}$, $\lambda \in \{71, 283, 1133, 4533\}/\text{yr}$, $c \in \{2, 30, 90, 260, 500\} \times 10^{-5}$.

Closed-Form Magnitudes (Theorem 2). Two reference numbers anchor this section: the diffusion ceiling $\sigma^2/8 = 832$ bp/yr (as $\Delta t \rightarrow \infty$) and the jump floor $\lambda G = 125$ bp/yr (as $\Delta t \rightarrow 0$). The fee has a negligible effect on the jump term ($\Psi(0.0260) = 0.959$, a 4.1% trim) since $\gamma/\delta = 0.0260$ lies deep inside the support of J . On the diffusion term the fee is the dominant suppression mechanism at short Δt .

Chain-by-Chain LVR Rate. Evaluating (12) at the baseline $(\sigma, \lambda, \delta, \gamma)$ over a range of Δt from Ethereum L1 down to a sub-second app-chain target, with $\kappa(\Delta t) = \gamma/(\sigma\sqrt{\Delta t})$ and the diffusion share $(\sigma^2/8)F(\kappa)/\ell$ reported alongside, gives Table 2. The $\sqrt{\Delta t}$ rate of Corollary 1 is visible: a $240\times$ cut from 12 s to 50 ms removes only 89% of the diffusion term, leaving the rate 29% above the floor. The two components are equal at $\Delta t_\times = 0.75$ s, and the diffusion term reaches a tenth of the floor only at $\Delta t = 5.5$ ms.

Table 2. LVR rate ℓ at the baseline parameters across deployed block times, with the diffusion share $(\sigma^2/8)F(\kappa)/\ell$ alongside. Chain labels denote block times only; every row is the same baseline pool.

Chain	Δt ℓ (bp/yr) diffusion share		
Ethereum L1	12 s	471	73%
Base / OP L2	2 s	312	60%
Solana	400 ms	221	43%
Arbitrum	250 ms	203	38%
App-chain	50 ms	162	23%
Jump floor ($\Delta t \rightarrow 0$)	—	125	0%

Regime Sensitivity at Ethereum’s 12 s. Holding $\Delta t = 12$ s fixed, Table 3 reports ℓ across a (σ, λ) grid, with the diffusion share of ℓ in brackets. The share is what determines whether shortening blocks reduces overall LVR: only the diffusion channel responds to Δt , so a high share means the pool has much to gain and a low one that the gain is small. The two readings move in opposite directions across the grid. The level of ℓ rises with both parameters, spanning 55 to 3,597 bp/yr, a factor of 65; the diffusion share rises with σ but falls with λ , from 1% in the calm, jump-heavy corner to 98% in the stressed, jump-light one. At the baseline $\lambda = 283$ /yr the share crosses 50% at $\sigma = 0.55$, so the calibrated pool is firmly diffusion-dominated at 73% and stress conditions reinforce this, while a calm regime puts the slot where shorter blocks help little. Any policy claim about 12-second blocks is therefore conditional on the volatility-and-jump-intensity regime, not on a single point estimate.

Planner’s Optimum. With c calibrated to the issuance-derived baseline 260×10^{-5} USD/block per \$1M TVL, (15) gives $A = V\gamma^2/(8c) = 12.02$ and $\eta^{\text{opt}} =$

Table 3. LVR rate ℓ in bp/yr at $\Delta t = 12$ s across the (σ, λ) grid, with the diffusion share of ℓ in brackets

$\sigma \setminus \lambda$	71/yr	283/yr	1,133/yr	4,533/yr
0.30 (calm)	55 (43%)	148 (16%)	524 (4%)	2,027 (1%)
0.816 (baseline)	377 (92%)	471 (73%)	847 (41%)	2,349 (15%)
1.50 (stress)	1,625 (98%)	1,719 (93%)	2,095 (76%)	3,597 (44%)

1.677. Across the sensitivity range, $c = (500, 260, 90, 30, 2) \times 10^{-5}$ USD/block yields $\Delta t^{\text{opt}} = (15.4, 8.4, 3.4, 1.4, 0.20)$ s respectively — nearly two orders of magnitude, driven by the cost calibration. Volatility moves it comparably: $\sigma \in \{0.30, 0.815, 1.50\}$ gives $\Delta t^{\text{opt}} = (62.3, 8.4, 2.5)$ s, since $\Delta t^{\text{opt}} \propto \gamma^2/\sigma^2$ at fixed η^{opt} . At the baseline $c = 260 \times 10^{-5}$, $\Delta t^{\text{opt}} \approx 8.4$ s or roughly two-thirds of Ethereum’s block-time and well above sub-second L2/app-chain timing. Under an LP-LVR-only objective Ethereum is too slow, though the gain is modest: W falls from 539 to 533 bp/yr, a saving of 6 bp/yr, under 2%, and stays within 1 bp/yr of its minimum over $\Delta t \in [7.4, 9.7]$ s. What moves the optimum is c , σ and γ ; the jump parameters do not move it at all. The conclusion is also one-sided: a full social-welfare treatment with MEV, finality, and oracle freshness could push Δt^{opt} back up.

8 Discussion and Conclusion

I extend the LVR framework to an empirical feature of crypto reference prices that the block-time literature has so far left out: price discontinuities. Jumps turn out to change what block time can achieve, not what it should be. For the CPMM under jump-diffusion prices the LVR rate splits into a diffusion channel that the block schedule governs through the fee multiplier $F(\kappa)$ and a Δt -invariant jump channel that it cannot touch. The planner’s optimum solves an explicit cubic and is invariant in pool size exactly and in the jump parameters (λ, m, δ) up to a shift the remainder bound confines to about a second (Remark 10), so jumps shift the level of LVR but not, to any resolution the objective can support, the marginal tradeoff that sets Δt^{opt} .

Policy. Shortening blocks can at most close the gap between the current rate and the jump floor λVG of Theorem 3: the floor bounds the rate from below, so no block-time reduction saves more than $\ell(\Delta t) - \lambda VG$, and because F decays algebraically rather than with a Gaussian tail, that gap closes only as $\sqrt{\Delta t}$ (Corollary 1). At the calibration of Section 7 the diffusion channel does not become negligible until the low-millisecond range, so at every deployed block time in Table 2 what limits block-time policy is not the jump floor but the cost of consensus. Once the floor does bind, block-time refinement no longer helps, and Remark 8 identifies the lever that remains: since fees recover only the fraction $1 - \Psi(\gamma/\delta)$ of jump-LVR, the binding ratio is γ/δ . The LP-side levers are

therefore fee-tier design, batch auctions that eliminate the per-block arbitrage window at source, and LVR-rebate designs such as auction-managed AMMs that route pool-management revenue back to LPs.

Calibrating the consensus cost from Ethereum’s gross issuance places the LP-side planner’s optimum below the current 12-second block-time but well above the sub-second regime. Two caveats temper the reading: the gain over the status quo is under 2% of the objective, and Δt^{opt} is sensitive to the cost calibration—invariant in pool size and effectively in every jump parameter (Remarks 9–10), it moves by nearly two orders of magnitude across the plausible range of c . The qualitative takeaway is that Ethereum’s block-time is somewhat too long under an LP-LVR-only objective, while the aggressive sub-second targets pursued by some chains overshoot what LVR considerations alone would justify.

Limitations. The welfare objective is LP-LVR-only, so the optimum from $W(\Delta t) = \ell_0(\Delta t) + c/\Delta t$ is an LP-side input to a larger welfare problem, not a prescription: a full treatment must also weigh MEV redistribution, finality, censorship resistance, and oracle freshness. The consensus cost attributes issuance pro-rata by value secured, abstracting from the dilution a native-token-holding LP bears directly, and equates observed issuance with validator operating cost — which breaks where issuance is a growth subsidy or validators are funded by MEV and fees. The pool is full-range, whereas most LVR is paid on concentrated liquidity, where the two effects on G run in opposite directions: concentration raises the pool curvature per unit of TVL and so inflates G for jumps contained in the active range, while a jump large enough to cross the range leaves the position single-asset and truncates the loss. Which dominates depends on δ relative to the range width, so the full-range floor is not simply a lower bound for concentrated positions. The arbitrageur has unlimited capital, zero latency and continuous sight of S_t , an idealization weakest in the sub-second regime the floor argument concerns. Blocks finally arrive as a Poisson process rather than on a deterministic proof-of-stake schedule, a tractability assumption inherited from [21] that is conservative, since the deterministic schedule minimizes arbitrage profit [23].

The practical message is that the jump floor is genuine — for symmetric jump laws Theorem 3 establishes it as an exact lower bound, not an artifact of the separation — but distant: at deployed block times the binding consideration is what a block costs, and the lever against the floor itself is the ratio of fee to jump size, not the block schedule.

References

1. Adams, H., Zinsmeister, N., Salem, M., Keefer, R., Robinson, D.: Uniswap v3 core (2021), <https://uniswap.org/whitepaper-v3.pdf>, uniswap Labs technical whitepaper
2. Angeris, G., Kao, H.T., Chiang, R., Noyes, C., Chitra, T.: An analysis of uniswap markets. *Cryptoeconomic Systems* **1**(1) (2021), <https://arxiv.org/abs/1911.03380>

3. Beccuti, J., Chantramonklasri, T., Hafner, M., Oderbolz, N.: Towards a formal framework of the Ethereum staking market. arXiv preprint arXiv:2503.14385 (2025), <https://arxiv.org/abs/2503.14385>
4. Benhaim, A., Falk, B.H., Tsoukalas, G.: Scaling blockchains: Can committee-based consensus help? *Management Science* (also arXiv:2110.08673) (2024). <https://doi.org/10.1287/mnsc.2022.03177>, <https://arxiv.org/abs/2110.08673>
5. Bundi, N.: Optimal block time for AMM liquidity providers under jump-diffusion prices (extended version) (2026), tODO: replace with the arXiv identifier issued on submission
6. Buterin, V.: Paths toward single-slot finality (2022), https://notes.ethereum.org/@vbuterin/single_slot_finality, ethereum research notes (notes.ethereum.org)
7. Capponi, A., Jia, R.: The adoption of blockchain-based decentralized exchanges. *Operations Research* (online first) (2024), <https://arxiv.org/abs/2103.08842>
8. Carr, P., Geman, H., Madan, D.B., Yor, M.: The fine structure of asset returns: An empirical investigation. *Journal of Business* **75**(2), 305–332 (2002)
9. Cartea, A., Drissi, F., Monga, M.: Decentralised finance and automated market making: Predictable loss and optimal liquidity provision. *SIAM Journal on Financial Mathematics* (forthcoming) – also arXiv:2309.08431 (2023), <https://arxiv.org/abs/2309.08431>
10. Chaim, P., Laurini, M.P.: Nonlinear dependence in cryptocurrency markets. *The North American Journal of Economics and Finance* **48**, 32–47 (2019). <https://doi.org/10.1016/j.najef.2019.01.015>
11. Cont, R.: Empirical properties of asset returns: Stylized facts and statistical issues. *Quantitative Finance* **1**, 223–236 (2001). <https://doi.org/10.1080/713665670>
12. Cont, R., Tankov, P.: *Financial Modelling with Jump Processes*. Chapman & Hall/CRC (2003)
13. Croman, K., Decker, C., Eyal, I., Gencer, A.E., Juels, A., Kosba, A., Miller, A., Saxena, P., Shi, E., Sirer, E.G., Song, D., Wattenhofer, R.: On scaling decentralized blockchains. In: *Financial Cryptography and Data Security Workshops (FC)*. *Lecture Notes in Computer Science*, vol. 9604, pp. 106–125 (2016). https://doi.org/10.1007/978-3-662-53357-4_8
14. Daian, P., Goldfeder, S., Kell, T., Li, Y., Zhao, X., Bentov, I., Breidenbach, L., Juels, A.: Flash boys 2.0: Frontrunning in decentralized exchanges, miner extractable value, and consensus instability. In: *IEEE Symposium on Security and Privacy (SP)*. pp. 910–927 (2020). <https://doi.org/10.1109/SP40000.2020.00040>
15. Decker, C., Wattenhofer, R.: Information propagation in the Bitcoin network. In: *IEEE International Conference on Peer-to-Peer Computing (P2P)*. pp. 1–10 (2013). <https://doi.org/10.1109/P2P.2013.6688704>
16. Drake, J.: Croissant issuance: A practical path toward capped ETH supply (2025), <https://www.dlnews.com/articles/defi/ethereum-policy-broken-says-top-researcher-justin-drake/>, x/Twitter thread, Feb. 5, 2025; secondary coverage: DL News
17. Elowsson, A.: Practical endgame on issuance policy (2024), <https://ethresear.ch/t/practical-endgame-on-issuance-policy/20747>, ethereum Research forum post (Oct. 23, 2024)
18. Fritsch, R., Canidio, A.: Measuring arbitrage losses and profitability of amm liquidity. In: *Companion Proceedings of the ACM Web Conference 2024*. pp. 1761–1767 (2024). <https://doi.org/10.1145/3589335.3651961>

19. Kou, S.G.: A jump-diffusion model for option pricing. *Management Science* **48**(8), 1086–1101 (2002). <https://doi.org/10.1287/mnsc.48.8.1086.166>
20. Merton, R.C.: Option pricing when underlying stock returns are discontinuous. *Journal of Financial Economics* **3**(1–2), 125–144 (1976). [https://doi.org/10.1016/0304-405X\(76\)90022-2](https://doi.org/10.1016/0304-405X(76)90022-2)
21. Milionis, J., Moallemi, C.C., Roughgarden, T.: Automated market making and arbitrage profits in the presence of fees. In: *Financial Cryptography and Data Security (FC)* – also arXiv:2305.14604 (2023), <https://arxiv.org/abs/2305.14604>
22. Milionis, J., Moallemi, C.C., Roughgarden, T., Zhang, A.L.: Automated market making and loss-versus-rebalancing. arXiv preprint arXiv:2208.06046 (2022), <https://arxiv.org/abs/2208.06046>
23. Nezlobin, A., Tassy, M.: Loss-versus-rebalancing under deterministic and generalized block-times (2025), <https://arxiv.org/abs/2505.05113>, arXiv:2505.05113
24. Saef, D., Nagy, O., Sizov, S., Härdle, W.K.: Understanding temporal dynamics of jumps in cryptocurrency markets: Evidence from tick-by-tick data. *Digital Finance* **6**(4), 605–638 (2024). <https://doi.org/10.1007/s42521-024-00116-1>
25. Scaillet, O., Treccani, A., Trevisan, C.: High-frequency jump analysis of the bitcoin market. *Journal of Financial Econometrics* **18**(2), 209–232 (2020). <https://doi.org/10.1093/jjfinec/nby013>